

# Explosion dynamics from strainmeter and microbarometer observations, Soufrière hills Volcano, Montserrat : 2008–2009

Chardot, L.; Foroozan, R.; Sacks, S.; Linde, A.; Stewart, R.; Fournier, N.; Komorowski, J. C.;  
Sparks, R. S. J.; Voight, Barry.; Clarke, A. B.; Hidayat, Dannie; Elsworth, Derek; Mattioli, Glen;  
Widiwijayanti, Christina

2010

Chardot, L., Voight, B., Foroozan, R., Sacks, S., Linde, A., Stewart, R., et al. (2010). Explosion  
dynamics from strainmeter and microbarometer observations, Soufrière Hills Volcano,  
Montserrat: 2008–2009. *Geophysical Research Letters*, 37.

<https://hdl.handle.net/10356/95520>

<https://doi.org/10.1029/2010GL044661>

---

© 2010 American Geophysical Union. This paper was published in *Geophysical Research Letters* and is made available as an electronic reprint (preprint) with permission of American Geophysical Union. The paper can be found at the following official URL: [<http://dx.doi.org/10.1029/2010GL044661>]. One print or electronic copy may be made for personal use only. Systematic or multiple reproduction, distribution to multiple locations via electronic or other means, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper is prohibited and is subject to penalties under law.

# Explosion dynamics from strainmeter and microbarometer observations, Soufrière Hills Volcano, Montserrat: 2008–2009

L. Chardot,<sup>1,2</sup> B. Voight,<sup>3</sup> R. Foroozan,<sup>3</sup> S. Sacks,<sup>4</sup> A. Linde,<sup>4</sup> R. Stewart,<sup>1</sup> D. Hidayat,<sup>3,5</sup> A. Clarke,<sup>6</sup> D. Elsworth,<sup>3</sup> N. Fournier,<sup>1,7</sup> J.-C. Komorowski,<sup>1,8</sup> G. Mattioli,<sup>9</sup> R. S. J. Sparks,<sup>10</sup> and C. Widiwijayanti<sup>5</sup>

Received 13 July 2010; accepted 27 July 2010; published 24 September 2010.

[1] Vulcanian explosions with plumes to 12 km occurred at Soufrière Hills volcano (SHV) between July 2008 and January 2009. We report strainmeter and barometric data, featuring quasi-linear strain changes that correlate with explosive evacuation of the conduit at rates of  $\sim 0.9\text{--}2 \times 10^7 \text{ kg s}^{-1}$ . July and January explosion-generated strains were similar,  $\sim 20$  nanostrain at  $\sim 5$  km, and interpreted as contractions of a quasi-cylindrical conduit, with release of magmatic pressure, and exsolution-generated overpressure of order 10 MPa. The 3 December 2008 event was distinctive with larger signals ( $\sim 140\text{--}200$  nanostrain at 5–6 km) indicating that a rapid pressurization preceded and triggered the explosion. Modeling suggests a dike with ENE trend, implying that feeder dikes at SHV had diverse attitudes at different times during the eruption. All explosions were associated with acoustic pulses and remarkable atmospheric gravity waves. **Citation:** Chardot, L., et al. (2010), Explosion dynamics from strainmeter and microbarometer observations, Soufrière Hills Volcano, Montserrat: 2008–2009, *Geophys. Res. Lett.*, 37, L00E24, doi:10.1029/2010GL044661.

## 1. Introduction

[2] The fourth episode of eruptive activity at Soufrière Hills volcano (SHV) between 29 July 2008 and 3 Jan 2009 began and terminated with pumice-bearing explosions. The Vulcanian explosions in this episode were captured by the CALIPSO strainmeter/microbarometer array installed in 2002–03 to investigate the dynamics of the SHV magmatic system [Mattioli et al., 2004]. We use volumetric strain, barometric, and broadband seismic data to constrain explosion source mechanisms and dynamic parameters. Such high-precision strain measurements of explosions are valu-

able but scarce. The innovation of using strain data acquired in 2008–09 to explore explosion dynamics is shared with data procured in 2003 [Voight et al., 2010], but significant new features are also recognized here, including explosion precursors, pre-explosion pressurization for one case that is used to deduce feeder-dike geometry, and acoustic waves that define explosion onsets. We present unique pressure-corrected-strain and microbarometric gravity wave records for all events and all operating stations.

## 2. Background on Explosions of July 2008 to January 2009

[3] Prior to the explosion on 29 July 2008, a 38-h swarm of intense seismicity occurred with hypocenters at 0–3.4 km depth [Stewart et al., 2008]. Then, following a quiet interval of a few hours, intense seismicity restarted at 03:32 29 July 2008 (all times are UTC), probably indicating a minor partial collapse of the dome carapace. The explosion followed at 03:38, generated a plume to 12 km a.s.l. (Washington VAAC), and produced pyroclastic flows in several drainages [Stewart et al., 2008; Komorowski et al., 2010]. The pyroclastic flows contained vesicular pumice and were generated by column collapse. Disregarding deposits in the Tar River valley due to a partial dome collapse [Stewart et al., 2008], pumice deposits on the west flank covered  $1.9 \text{ km}^2$ , and assuming a thickness of 0.5–1 m, comprised a volume of  $0.5\text{--}1.0 \text{ Mm}^3$  DRE, plus co-pyroclastic flow (co-PF) ash [Komorowski et al., 2010; G. Wadge, written communication, 2010].

[4] The second extrusive phase a few months later was more complex [Stewart et al., 2009]. Following volcano-tectonic seismicity starting 28 Nov, an explosion occurred at 01:37 on 3 Dec 2008. It was preceded by 2 min of seismicity with rockfall signatures. Dome-rock ballistics were propelled 2 km from the vent, a plume rose 12 km high, and pyroclastic currents split around St. Georges Hill west of the volcano [Komorowski et al., 2010]. Of the three events considered here, the Dec deposits are least well understood, because of burial during further events in Dec and Jan, and rainfall erosion. A surge possibly caused by a small directed blast from the dome base occurred in the Gages area. The deposit limits were carefully mapped [Komorowski et al., 2010], yielding a volume of  $1.34 \text{ Mm}^3$  DRE including co-PF ash, or  $1.17 \text{ Mm}^3$  excluding co-PF ash. However, the proportions of lithics and pumice are not well defined. Komorowski et al. [2010] presumed most ejecta were from hot pre-Dec 2008 dome rock, in essence not fresh juvenile products; but Wadge et al. [2010] included the DRE volume as fresh lava.

<sup>1</sup>Montserrat Volcano Observatory, Flemmings, Montserrat, West Indies.

<sup>2</sup>Ecole et Observatoire des Sciences de la Terre, Strasbourg, France.

<sup>3</sup>Earth and Mineral Sciences, Pennsylvania State University, University Park, Pennsylvania, USA.

<sup>4</sup>Department of Terrestrial Magnetism, Carnegie Institute of Washington, Washington, D. C., USA.

<sup>5</sup>Earth Observatory of Singapore, Singapore.

<sup>6</sup>School of Earth and Space Exploration, Arizona State University, Tempe, Arizona, USA.

<sup>7</sup>Now at GNS Science, Wairakei Research Center, Taupo, New Zealand.

<sup>8</sup>Institut de Physique du Globe de Paris, Paris, France.

<sup>9</sup>Department of Geosciences, University of Arkansas, Fayetteville, Arizona, USA.

<sup>10</sup>Department of Earth Sciences, Bristol University, Bristol, UK.

[5] Three subsequent smaller explosions (4 Dec, 0:43; 4 Dec, 20:22; 5 Dec, 9:41) caused plumes of limited height, but only the last generated a pyroclastic flow. The dome grew rapidly after 8 Dec, and after 21 Dec cyclic tremor was observed [Stewart *et al.*, 2009], similar to tilt/seismic cycles in 1997 and 1999–2000 [Voight *et al.*, 1999; Young *et al.*, 2003]. The cyclic tremor amplitude increased after 31 Dec, suggesting increased pressure [Voight *et al.*, 1999; Young *et al.*, 2003]. On 2 Jan 2009 a small pyroclastic flow developed, and on 3 Jan explosions occurred from the dome top at 6:55, 8:13, 8:47 and 11:07 [Komorowski *et al.*, 2010]. The last two began with jets to ~500 m, and generated pumice flows by column collapse. The last event produced a plume to 11 km. The pumice deposit volume is not so well defined as for the July explosion. For all 3 Jan events, Komorowski *et al.* [2010] reckoned a total volume of 0.95 Mm<sup>3</sup> DRE, but constraints are poor and partitioning between events is uncertain. For the largest event at 11:07, we take 0.5 Mm<sup>3</sup> DRE as a rough estimate of pumice deposits.


[6] Complete computations of eruption volumes including plume-height based estimates are given in the auxiliary material.<sup>1</sup> The plume is modeled as a buoyant thermal in a humid tropical environment [Woods and Kienle, 1994; Sparks *et al.*, 1997; Druitt *et al.*, 2002; Tupper *et al.*, 2009]. The above volumes are approximate, but they provide useful quantitative constraints for analyses based on strain observations.

### 3. Instrumentation and Observations

[7] Data were obtained from the CALIPSO array with instruments in ~200-m-deep boreholes and surface sites, at distances of about 5.1 to 9.6 km from the vent [Mattioli *et al.*, 2004] (Figure 1). We furnish observations of strain and air pressure linked to the explosions, and compare these data with broadband seismic records from surface station MBGH. Three sites (AIRS, TRNT, GERD) were operating for the 3 Jan 2009 explosion, and two (AIRS, TRNT) for 29 July and 3 Dec 2008. Each borehole contains a Sacks-Evertson high-dynamic-range volumetric strainmeter recording data at 50 samples per second (sps) with a precision of ~0.1 nanostrain (nS) in periods from 0–50 Hz [Linde and Sacks, 1995]. In this paper we round-off values to the nearest nanostrain.

[8] The strainmeters are affected by changes in atmospheric pressure, requiring corrections based on microbarometric data [e.g., Nakao *et al.*, 1989] (see auxiliary material). The explosions generated complicated airwaves recorded at 1 sps on barographs, with a shock wave spike at explosion onset, followed by a strong pressure decrease that evolved into long-period oscillations (Figures 2 and S1–S5). Similar distinctive ~800 s oscillations were observed with SHV explosions in 2003 [Voight *et al.*, 2010]. They propagated at about 30 m s<sup>-1</sup> beyond 5 km as gravity waves, i.e., atmospheric oscillations generated by the rising plume, and were recorded at various azimuths and distances. Such observations of gravity waves in the near-field are unprecedented for explosions at this scale [cf. Ripepe *et al.*, 2010; Kanamori *et al.*, 1994].


<sup>1</sup>Auxiliary materials are available in the HTML. doi:10.1029/2010GL044661.



**Figure 1.** Montserrat map with location of strainmeter sites (red triangles), the broadband seismic station MBGH (red dot) and the approximate location of the active vent of Soufrière Hills Volcano (red star). Colored pattern is surface volumetric strain for modeled dike with N77°E trend, 600 m width, -2 to -3 km depth, 10 MPa pressure, 5 GPa rock modulus (see text for details). Blue is field of volumetric contraction, orange is field of expansion. Scale is nanostrains.

[9] The data from AIRS and GERD show high signal/noise ratios (Figure 2). TRNT, located near the shoreline, yielded noisy data for July 2008 and Jan 2009 explosions (amplitudes ~10 nS), but was crucial for identifying strain polarity (Figures S1–S3). For the July and Jan events the strainmeters at all operating sites expanded. Because AIRS and the other sites are far beyond the nodal distance for a shallow axisymmetrical source [Linde & Sacks, 1995], the source had to contract, and this indicates a source pressure reduction. The data at AIRS, TRNT and GERD have identical polarity and can be satisfied by a shallow cylindrical conduit source. Thus the July and Jan explosion patterns (Figures 2, S1, and S3–S5) confirm the same four strain stages recognized in 2003 explosion data [Voight *et al.*, 2010]: 1) a brief transition between disturbance onset and a profound change in strain; 2) a quasi-linear ramp producing most of the strain change; 3) a more gentle decrease of strain to a minimum value; and 4) a recovery lasting hours to approach a background magnitude. In several 2008 cases we recognize also a precursor stage (*p*). No precursor stage is recognized for 3 Jan, but for the 29 July explosion, seismic activity initiated 6 min prior to the explosion, interpreted as a minor rockfall.

[10] A different pattern developed on 3 Dec 2008, quite unlike any of the explosions of 2003 [Voight *et al.*, 2010]. In this case a strong and rapid (~100 s) strain pulse preceded the explosion, accompanied by bursts of seismicity (Stage *p* (Figure 2)). AIRS recorded peak values of ~143 nS, and the



**Figure 2.** (a) Raw strain data and raw barometer data for three explosive events (from top to bottom: 29 July 2008, 3 Dec 2008, 3 Jan 2009) recorded at AIRS. Traces have been aligned with the approximate onset of the strain signal (29 July 2008 at 3:38:30; 3 Dec 2008 at 1:35:50; 3 Jan 2009 at 11:07:20; all times UTC), with the onsets marked by a vertical line on the two panels. In the Dec event, the explosion followed a rapid large strain buildup. (b) Strain signals corrected for atmospheric pressure for the same three explosions recorded at AIRS (black), TRNT (red) and GERD (blue). Strain signals are superposed on the respective seismic signals recorded at MGBH (green), with normalized amplitudes. Stages as explained in text are separated by red lines, with [p] [1] [2] [3] [4] the successive stages.

more distant TRNT  $\sim 202$  nS (Figure S2), both with high signal/noise ratios and the same polarity (contraction). During the following  $\sim 10$  minutes both sites showed reduction in strain (expansion), accompanied by explosive

surface manifestations (acoustic and gravity waves, and plume generation). Seismic activity virtually coincided with the initial strain pulse (Figure 2), but before onset of the

rapid pulse we detected no strain or seismic changes that could be related to either the strain pulse or explosion.

[11] The Stage *p* changes for the Dec event are inconsistent with solely a spherical pressure source at any depth under SHV, or a shallow cylindrical source, or a single-sensed combination of the two [Lisowski, 2007]. The observed strain polarities can be produced by such sources, but not strain magnitudes at TRNT that exceed those at the nearer station AIRS; also for a cylindrical source the implied pressures exceed conduit strength (see auxiliary material). Thus we sought to see if a dike source could help to satisfy the data. As developed in Section 4, our analyses strongly support inclusion of a dike source, with Stage *p* comprising mainly an intense, rapid dike pressurization. This pattern is comparable with that observed during the 3 March 2004 event, also modeled using a dominant dike source [Linde et al., 2010].

[12] We judge Stage 1 of the Dec explosion as having started shortly before the AIRS strain peak was reached (the TRNT peak occurred slightly earlier), no doubt triggered by the pressure increase and indicated by a reduction in strain rate (slight break-in-slope) and generation of an acoustic wave that originated at the vent  $\sim 15$  s before its arrival at AIRS ( $5.1 \text{ km}/340 \text{ m s}^{-1}$ ). The peak of the AIRS strain curve ( $\sim 143 \text{ nS}$ ) marks the point where incremental strain released by the explosion exceeds the dike source strain build-up. This defines the Stage 2 onset, and thereafter the stages appear similar to those of the other explosions. The strain recoveries at AIRS and TRNT are  $-88$  and  $-135 \text{ nS}$ , respectively, approximately  $0.6$ – $0.7$  of the peak values.

[13] For all explosions, Stage 1 is a brief period with small strain and small seismicity that occurs when the pressurized gas under a strong plug reaches the strength of the cap [Voight et al., 2010; Alidibirov and Dingwell, 1996]. The acoustic pulse is generated within Stage 1 and indicates that fragmentation and material expulsion is initiated in this stage. Stage 2 is a swift contraction of the pressure source that correlates with energetic expulsion of the pyroclast/gas mixture in the conduit, with durations of 105 and 230 s for the July and Jan events, respectively. Voight et al. [2010] propose that most of the ejecta are expelled during the quasi-linear strain-ramp, and that the further gradual reduction of strain to a minimum value in Stage 3 reflects combined effects of discharge of ash and gas from the top of the magma column, pressure adjustments within the deeper conduit system, and rise of magma. Stage 4 then represents a gradual strain recovery over several hours due to magma ascent and exsolution/gas pressure effects.

#### 4. Analyses: Procedures, Results, and Discussion

[14] For the July and Jan events the strain-ramp is interpreted as a rapid reduction in internal pressure about a quasi-cylindrical conduit wall, associated with pyroclast/gas evacuation. We assume a conduit with diameter of 30 m [Voight et al., 1999; Melnik and Sparks, 2002], positioned over a planar feeder dike that rises from the magma chamber at  $\sim 5 \text{ km}$  depth [cf. Mattioli et al., 1998; Costa et al., 2007; Hautmann et al., 2009; Linde et al., 2010]. We assume the cylinder/dike transition occurs roughly at 2 km beneath the vent at  $\sim 1000 \text{ m a.s.l.}$  Our data suggest strong involvement of a dike in the Dec event.

[15] Following Voight et al. [2010] we estimate conduit drawdown from erupted DRE volume adjusted for porosity, divided by conduit area. We assume an average density of  $\sim 1500 \text{ kg m}^{-3}$ , based on clast data from 1997 and 2003 explosions [Clarke et al., 2002; Edmonds et al., 2006]. For a 30-m conduit the total volume estimates of  $0.92$ – $1.42 \text{ Mm}^3$  DRE for July, and  $0.76 \text{ Mm}^3$  DRE for Jan, suggest drawdowns of  $2.3$ – $3.5 \text{ km}$  (July), and  $1.9 \text{ km}$  (Jan). The  $2.3 \text{ km}$  estimate for July is roughly compatible with our proposed  $2 \text{ km}$  nominal depth to the top of the dike, but the larger volume implies a  $3.5 \text{ km}$  drawdown. Because strain data for July suggest no dike signature, we prefer the lower volume and drawdown estimates. For the Dec explosion, discussed below, the issue is more complex.

[16] We then use our strain-ramp amplitudes to estimate average conduit pressures released in the explosions [Voight et al., 2010]. Using an axisymmetric elastic finite element model (see auxiliary material), we apply a magmatic load using an average density of  $1500 \text{ kg m}^{-3}$  and a superposed uniform overpressure (pressure exceeding magmatic) over the drawdown lengths of conduit (see auxiliary material). We developed sub-models for homogeneous and inhomogeneous cases, in the latter instance applying a velocity/modulus increase with depth as based on the SEA-CALIPSO experiment [Shalev et al., 2010]. For the homogeneous case we used Young's modulus  $E = 5 \text{ GPa}$ , the average local modulus at the strainmeter sites, and a Poisson ratio of  $0.25$ . Comparable moduli have been used in modeling other volcanoes [Lisowski, 2007; Houlié and Montagner, 2007] (and others cited in auxiliary material) and suggest a compliant, fractured and/or hydrothermally altered rock mass. For inhomogeneous media and AIRS data (best signal/noise ratio), the Stage 2 strain-ramps for July and Jan explosions ( $24$  and  $18 \text{ nS}$ ) suggest preliminary average overpressures of  $\sim 30$  and  $\sim 25 \text{ MPa}$ ; homogeneous-case values are slightly larger. These results are dependent on magma density and modulus. Increasing magma density can reduce overpressures  $<20\%$ , whereas assuming  $E = 3 \text{ GPa}$  (arguably typical for volcanic areas [see Lisowski, 2007]) yields overpressures of order  $10 \text{ MPa}$ . Although Robertson et al. [1998] proposed an overpressure of  $27.5 \text{ MPa}$  for a 1996 SHV explosion, we nevertheless regard our larger estimates with discomfort. Typically, overpressures  $<15$ – $20 \text{ MPa}$  are given by conduit models for magma attributes appropriate for SHV, consistent with our reduced-modulus results [de' Michieli Vitturi et al., 2010; Melnik and Sparks, 2002; Voight et al., 2010]. For Jan 2009 we can also calculate the ratio of measured strain changes at AIRS and GERD sites (Figure 2) as  $18/4.5 = 4$ . The corresponding modeled values are  $>5$ ; this discrepancy, also recognized for 2003 explosion data [Voight et al., 2010], might reflect modulus heterogeneity [cf. Shalev et al., 2010]. Associated mass evacuation rates are given by erupted mass divided by Stage 2 strain-ramp durations, yielding  $2.3 \times 10^7$ , and  $0.86 \times 10^7 \text{ kg s}^{-1}$  for the July and Jan explosions, respectively. These results are similar to those reported for explosions at SHV in 1997 [Druitt et al., 2002] and 2003 [Voight et al., 2010].

[17] For the Dec explosion we propose that a dominant dike source (with lesser contributions from conduit and chamber sources) can explain the observations. Stage *p* and Stage 1 strains at AIRS, TRNT ( $c.143,202 \text{ nS}$ ) correspond mainly to pressurization of a dike; the contribution from the



upper cylindrical conduit is only of the order 20 or 10 nS at AIRS, TRNT (e.g., July and Jan data), with chamber contributions uncertain but likely relatively small. During Stage 2, with ejection of the gas/pyroclast mixture, the cylindrical conduit contracts and the dike partially closes. To test this hypothesis we calculated strain changes using several finite-element models (Figure 1 and auxiliary material). The top of the vertical dike was held to the depth range 1 to 2 km. The modeled strains at AIRS and TRNT vary rapidly with dike azimuth and their polarity requires an ENE dike orientation (best fit,  $\sim N77^\circ E$ ). For the March 2004 event, Linde *et al.* [2010] found evidence to constrain a dike of WNW orientation. These and other results [e.g., Hautmann *et al.*, 2009] suggest that feeder dikes at SHV had diverse attitudes at different times during the eruption.

[18] We use a model dike width of 600 m but our conclusions are independent of this choice, as model strain values depend on the product of dike area and opening (or pressure). With  $E = 5$  GPa, and pressure 10 MPa, Figure 1 shows model contractive strains at AIRS and TRNT as 125 and 185 nS, respectively. With addition of the upper conduit contributions at these sites as  $\sim 20$  and 10 nS (contractive during Stage *p* pressurization), a nearly exact match is produced of the observed strains. This result demonstrates the plausibility of a dike source although the solution is not unique, and no contribution from a magma chamber is included in the calculation. For the March 2004 event, Linde *et al.* [2010] modeled a dike and a deep spherical pressure source, and neglected a shallow source. The pressure loss assumed for the deep source contributed contractions of 50 and 20 nS at AIRS and TRNT, respectively. However, we think it is not certain that a deep source involved a pressure loss. An increase in chamber pressure applied instantaneously to the bottom of a (cylindrical) conduit has been modeled by de' Michieli Vitturi *et al.* [2010]. For magma properties based on SHV, a pressure increase is transmitted through the lower 3/5 of the conduit in only about 1 min;  $\sim 10$  min are required to pressurize the upper 4/5, and  $\sim 30$  min to affect also the uppermost conduit. These results are germane to the Dec 2008 and March 2004 events, as they show that deep pressure increases can in principle be transmitted in very short times (few minutes) to mid-level portions of the transport system, and thereby be detected by our strainmeter array.

[19] **Acknowledgments.** This research was supported by NSF Continental Dynamics and I & F Programs, and NERC (UK). We thank L. Johnson and R. Kelz at NSF. Additional support was provided by the CALIPSO Consortium, and by Montserrat Volcano Observatory. BV is grateful to G. Swanson of NOAA. RSJS is supported by an ERC grant. We appreciated the reviews by A. Neri and an anonymous person, which helped us to improve the paper.

## References

- Alidibirov, M. A., and D. B. Dingwell (1996), Magma fragmentation by rapid decompression, *Nature*, **380**, 146–148, doi:10.1038/380146a0.
- Clarke, A. B., et al. (2002), Computational modeling of the transient dynamics of the August 1997 Vulcanian explosions at Soufrière Hills Volcano, Montserrat, in *The Eruption of Soufrière Hills Volcano, From 1995 to 1999*, edited by T. H. Druitt and B. P. Kokelaar, *Mem. Geol. Soc. London*, **21**, 319–348.
- Costa, A., et al. (2007), Control of magma flow in dykes on cyclic lava dome extrusion, *Geophys. Res. Lett.*, **34**, L02303, doi:10.1029/2006GL027466.
- de' Michieli Vitturi, M., A. B. Clarke, A. Neri, and B. Voight (2010), Transient effects of magma ascent dynamics along a geometrically variable dome-feeding conduit, *Earth Planet. Sci. Lett.*, **295**, 541–553, doi:10.1016/j.epsl.2010.04.029.
- Druitt, T. H., et al. (2002), Episodes of cyclic Vulcanian explosive activity with fountain collapse at Soufrière Hills Volcano, Montserrat, in *The Eruption of Soufrière Hills Volcano, From 1995 to 1999*, edited by T. H. Druitt and B. P. Kokelaar, *Mem. Geol. Soc. London*, **21**, 281–306.
- Edmonds, M., R. A. Herd, and M. H. Strutt (2006), Tephra deposits associated with a large lava dome collapse, Soufrière Hills Volcano, Montserrat, 12–15 July 2003, *J. Volcanol. Geotherm. Res.*, **153**, 313–330, doi:10.1016/j.jvolgeores.2005.12.008.
- Hautmann, S. J., et al. (2009), Modelling ground deformation caused by an oscillating overpressure in a dyke conduit at Soufrière Hills Volcano, Montserrat, *Tectonophysics*, **471**, 87–95, doi:10.1016/j.tecto.2008.10.021.
- Houlié, N., and J.-P. Montagner (2007), Hidden dykes detected on ultra long period seismic signals at Piton de Fournaise Volcano?, *Earth Planet. Sci. Lett.*, **261**, 1–8, doi:10.1016/j.epsl.2007.04.018.
- Kanamori, H., et al. (1994), Excitation of atmospheric oscillations by volcanic eruptions, *J. Geophys. Res.*, **99**(B11), 21,947–21,961, doi:10.1029/94JB01475.
- Komorowski, J.-C., et al. (2010), Insights into processes and deposits of hazardous vulcanian explosions at Soufrière Hills Volcano during 2008 and 2009 (Montserrat, West Indies), *Geophys. Res. Lett.*, **37**, L00E19, doi:10.1029/2010GL042558.
- Linde, A. T., and L. S. Sacks (1995), Continuous monitoring of volcanoes with borehole strainmeters, in *Mauna Loa Revealed: Structure, Composition, History, and Hazards*, *Geophys. Monogr. Ser.*, vol. 92, edited by J. M. Rhodes and J. P. Lockwood, pp. 171–185, AGU, Washington, D. C.
- Linde, A. T., et al. (2010), The Vulcanian explosion at Soufrière Hills Volcano, Montserrat on March 2004 as revealed by strain data, *Geophys. Res. Lett.*, **37**, L00E07, doi:10.1029/2009GL041988.
- Lisowski, M. (2007), Analytical volcano deformation source models, in *Volcano Deformation*, edited by D. Dzurisin, pp. 279–304, Springer, Berlin.
- Mattioli, G., et al. (1998), GPS measurement of surface deformation around Soufrière Hills Volcano, Montserrat, from October 1995 to July 1996, *Geophys. Res. Lett.*, **25**(18), 3417–3420, doi:10.1029/98GL00931.
- Mattioli, G., et al. (2004), Prototype PBO instrumentation of CALIPSO Project captures world-record lava dome collapse on Montserrat, *Eos Trans. AGU*, **85**(34), doi:10.1029/2004EO340001.
- Melnik, O. E., and R. S. J. Sparks (2002), Modelling of conduit flow dynamics during explosive activity at Soufrière Hills Volcano, Montserrat, in *The Eruption of Soufrière Hills Volcano, From 1995 to 1999*, edited by T. H. Druitt and B. P. Kokelaar, *Mem. Geol. Soc. London*, **21**, 307–317.
- Nakao, S., A. Linde, S. Miura, M. Mishina, and A. Takagi (1989), Non-linear barometric response of borehole strainmeters and its interpretation, *J. Phys. Earth*, **37**, 357–383.
- Ripepe, M., S. De Angelis, G. Lacanna, and B. Voight (2010), Observation of infrasonic and gravity waves at Soufrière Hills Volcano, Montserrat, *Geophys. Res. Lett.*, **37**, L00E14, doi:10.1029/2010GL042557.
- Robertson, R. E. A., et al. (1998), The explosive eruption of Soufrière Hills Volcano, Montserrat, September 17, 1996, *Geophys. Res. Lett.*, **25**, 3429–3432, doi:10.1029/98GL01442.
- Shalev, E., et al. (2010), Three-dimensional seismic velocity tomography of Montserrat from the SEA-CALIPSO Offshore/Onshore Experiment, *Geophys. Res. Lett.*, **37**, L00E17, doi:10.1029/2010GL042498.
- Sparks, R. S. J., M. L. Bursik, S. N. Carey, J. S. Gilbert, L. S. Glaze, H. Sigurdsson, and A. W. Woods (1997), *Volcanic Plumes*, 574 pp., John Wiley, Chichester, U. K.
- Stewart, R., et al. (2008), Report for the Scientific Advisory Committee on Montserrat Volcanic Activity, Montserrat Volcano Obs., Flemmings, Montserrat, 13 Oct.
- Stewart, R., et al. (2009), Report for the Scientific Advisory Committee on Montserrat Volcanic Activity, Montserrat Volcano Obs., Flemmings, Montserrat, 4 March.
- Tupper, A., et al. (2009), Tall clouds from small eruptions: The sensitivity of eruption height and fine ash content to tropospheric instability, *Nat. Hazards*, **51**, 375–401, doi:10.1007/s11069-009-9433-9.
- Voight, B., et al. (1999), Magma flow instability and cyclic activity at Soufrière Hills Volcano, Montserrat, British West Indies, *Science*, **283**, 1138–1142, doi:10.1126/science.283.5405.1138.
- Voight, B., et al. (2010), Unique strainmeter observations of Vulcanian explosions, Soufrière Hills Volcano, Montserrat, July 2003, *Geophys. Res. Lett.*, **37**, L00E18, doi:10.1029/2010GL042551.
- Wadge, G., et al. (2010), Lava production at Soufrière Hills Volcano, Montserrat: 1995–2009, *Geophys. Res. Lett.*, **37**, L00E03, doi:10.1029/2009GL041466.
- Woods, A. W., and J. Kienle (1994), The dynamics and thermodynamics of volcanic clouds: Theory and observations from the April 15 and April 21 1990 eruptions of Redoubt Volcano, Alaska, *J. Volcanol. Geotherm. Res.*, **62**, 273–299, doi:10.1016/0377-0273(94)90037-X.
- Young, S. R., B. Voight, and H. J. Duffell (2003), Magma extrusion dynamics revealed by high-frequency gas monitoring at Soufrière Hills

Volcano, Montserrat, *Spec. Publ. Geol. Soc. London*, 213, 219–230, doi:10.1144/GSL.SP.2003.213.01.13.

---

L. Chardot, Ecole et Observatoire des Sciences de la Terre, 5, rue Rene Descartes, F-67084 Strasbourg CEDEX, France.

A. Clarke, School of Earth and Space Exploration, Arizona State University, Box 871404, Tempe, AZ 85287, USA.

D. Elsworth, R. Foroozan, and B. Voight, Earth and Mineral Sciences, Pennsylvania State University, 503 Deike Bldg., University Park, PA 16802, USA. (voight@ems.psu.edu)

N. Fournier, GNS Science, Wairakei Research Center, 114 Karetoto Rd., Taupo 3377, New Zealand.

D. Hidayat and C. Widiwijayanti, Earth Observatory of Singapore, 50 Nanyang Ave., Block N2-01a-15, Singapore 639798.

J.-C. Komorowski, Institut de Physique du Globe de Paris, 4, place Jussieu, F-75252 Paris CEDEX 05, France.

A. Linde and S. Sacks, Department of Terrestrial Magnetism, Carnegie Institute of Washington, 5241 Broad Branch Rd., NW, Washington, DC 20015, USA.

G. Mattioli, Department of Geosciences, University of Arkansas, 113 Ozark Hall, Fayetteville, AR 72701, USA.

R. S. J. Sparks, Department of Earth Sciences, Bristol University, Queen's Road, Bristol BS8 1RJ, UK.

R. Stewart, Montserrat Volcano Observatory, Flemmings, Montserrat, West Indies.