Charge transfer and charge separation dynamics in efficient non-fullerene solar cells

Ma, Chao; Chan, Christopher C. S.; Chow, Philip C. Y.; Yan, He; Wong, Kam Sing

2020

Ma, C., Chan, C. C. S., Chow, P. C. Y., Yan, H., & Wong, K. S. (2020). Charge transfer and charge separation dynamics in efficient non-fullerene solar cells. Proc. Of the 7th Asian Spectroscopy Conference (ASC 2020). doi:10.32655/ASC_8-10_Dec2020.15

<https://hdl.handle.net/10356/144220>

https://doi.org/10.32655/ASC_8-10_Dec2020.15

© 2020 Nanyang Technological University.

Downloaded on 26 Apr 2023 21:40:32 SGT

Charge Transfer and Charge Separation Dynamics in Efficient Non-fullerene Solar Cells

Chao Ma¹, Christopher C. S. Chan¹, Philip C. Y. Chow², He Yan³ and Kam Sing Wong¹

¹ Department of Physics, Hong Kong University of Science and Technology, Hong Kong, P. R. China.

² Department of Mechanical Engineering, University of Hong Kong, Hong Kong, P. R. China

³ Department of Chemistry, Hong Kong University of Science and Technology, Hong Kong, P. R. China

E-mail: phkswong@ust.hk

Abstract

Recently, Non-fullerene organic solar cells (OSCs) has been making rapid progress and reaching record breaking efficiency [1]. In the conventional idea for the fullerene based OSCs, one need a large donor-acceptor (D/A) offsets to cause rapid (i.e., <1ps) and effective charge separation from exciton at the D/A heterojunction [2]. However, recent work showed that non-fullerene OSC with small D/A offset can also achieved very high power conversion efficiency (PCE) [1,3-4]. The explanation for this is that charge separation occurs over hundreds of ps from thermalized charge transfer exciton (CTEs) at the D/A heterojunction for the non-fullerene system [5], thus charge separation in efficient non-fullerene OSC with small D/A offset is an endothermic process.

Here we will present our recent transient absorption (TA) spectroscopy studies to further elucidate the temperature dependence of charge transfer and charge separation dynamics in a number of non-fullerene OSC model systems with negligible D/A LUMO offset but variable HOMO offsets. We selectively create acceptor excitons and showed that acceptor excitons dissociate into CTEs at a similar rate (~100ps) as donor excitons despite the significantly large driving force provided by the HOMO offset. In contrast, by tracing the electro-absorption signal, it was demonstrated the generation of free charges after hole transfer required additional thermal excitation. Our result show the charge separation is indeed an endothermic behavior in non-fullerene OSCs, but charge transfer is not. Our result give a deeper understanding of the working mechanism of non-fullerene OSCs, which will have important fundamental implications on the development of OSCs.

Keywords: Ultrafast spectroscopy, organic solar cells, charge generation and separation.

References:

- [1] J. Yuan et al., *Joule*, **2000**, 3, 1.
- [2] S. Gelinas, et al., *Science*, **2014**, 343, 512.
- [3] S. Chen, et al., *J. Am. Chem.Soc.*, **2017**, 139, 6298.
- [4] G. Zhang et al., *Chem. Rev.* **2018**, 17, 703.
- [5] P. C. Y. Chow et al, *Nat. Commun.* (accepted)