The gap in surgical footprints : the role of connexin43 during post-surgical peritoneal adhesion formation

Chua, Jia Wang

2019

Chua, J. W. (2019). The gap in surgical footprints : the role of connexin43 during post-surgical peritoneal adhesion formation. Doctoral thesis, Nanyang Technological University, Singapore.

<https://hdl.handle.net/10356/89476>

<https://doi.org/10.32657/10220/48042>

**THE GAP IN SURGICAL FOOTPRINTS:
THE ROLE OF CONNEXIN43
DURING POST-SURGICAL
PERITONEAL ADHESION FORMATION**

CHUA JIA WANG

**Interdisciplinary Graduate School
NTU Institute for Health Technologies**

A thesis submitted to the Nanyang Technological University
in partial fulfillment of the requirement for the degree of
Doctor of Philosophy

2018

Statement of Originality

I hereby certify that the work embodied in this thesis is the result of original research and has not been submitted for a higher degree to any other University or Institution.

[12 Feburary 2019]

.....
..

Date

.....

[Chua Jia Wang]

Supervisor Declaration Statement

I have reviewed the content and presentation style of this thesis and declare it is free of plagiarism and of sufficient grammatical clarity to be examined. To the best of my knowledge, the research and writing are those of the candidate except as acknowledged in the Author Attribution Statement. I confirm that the investigations were conducted in accord with the ethics policies and integrity standards of Nanyang Technological University and that the research data are presented honestly and without prejudice.

[12 February 2019]

.....
Date

.....
Prof. David L. Becker

Authorship Attribution Statement

This thesis **does not** contain any materials from papers published in peer-reviewed journals or from papers accepted at conferences in which I am listed as an author.

[12 Feb 2019]

.....

..

Date

.....

[Chua Jia Wang]

Acknowledgements

I would like to take this opportunity to thank the Ministry of Education for funding the Nanyang Technological University Research Scholarship. In addition, I would like to express my gratitude to Interdisciplinary Graduate School and Nanyang Institute of Health Technologies for conferring this scholarship award to me so that I may pursue my research interests.

I wish to express my appreciation to Prof. David Becker for this wonderful opportunity to work in his laboratory and his trust in my capabilities of taking on various projects. His scientific counsel and encouragement has been key in propelling my interest in research. I would also like to thank my co-supervisor and my mentor, Assoc. Prof Kimberly Kline and Dr. Anthony Philips, for their thoughtful feedback and guidance throughout my project. For the committee's unwavering support throughout my candidature, I am sincerely grateful. I would also like to thank Dr. Rao Nandini Chilagondanahalli Lakshmi Narayana from Tan Tock Seng Hospital for all her precious advice on histopathology.

I would also wish to thank some members of the laboratory: Dr. Rebecca Hu and Dr. Allyson Soon for showing me the ropes at the start of journey; Mr. Leigh Madden, for managing the laboratory smoothly – thank you for your assistance with our furry friends; Mr. Milton Kwek, for being there even before the very start - thank you for for being someone I can always count on, always helping out with surgeries, and being someone I can easily bounce ideas off; Ms. Erica Teo and Ms. Ong Zi Xin, for being my younger sisters that I never had, and for making the laboratory my second home; and also to all students that I had the privilege of mentoring. Also, I would like to thank Dr. Ng Lai Guan and Dr. Sapna Devi, for the sparking my interest to pursue this entire endeavour.

Above all, I would like to thank and dedicate this thesis to my family for their unequivocal love and support throughout my journey, and my wife, Abigail Loh, for being my lifeline, my pillar of strength, and for always believing in me.

Contents

Scope of Work and Structure of Thesis	14
Hypothesis	17
Chapter 1: Literature Review.....	18
1.1 Post-surgical peritoneal adhesions.....	18
1.1.1 Definition.....	18
1.1.2 Epidemiology, complications, and economic burden.....	19
1.1.3 Early pathogenesis of post-surgical peritoneal adhesion formation.....	21
1.1.3.1 Early responders to the scene.....	21
1.1.3.2 Release of cytokine cocktails.....	22
1.1.3.3 The coagulation cascade, fibrinolytic balance, and matrix deposition	24
1.1.4 Experimental models	27
1.1.4.1 Cell culture systems.....	27
1.1.4.2 Ex vivo tissue culture system.....	28
1.1.4.3 In vivo models	29
1.1.5 Adhesion-prevention strategies	33
1.1.5.1 Technical measures.....	33
1.1.5.2 Barrier measures	34
1.1.5.3 Pharmacological agents.....	35
1.2 Connexins	38
1.2.1 Structure and nomenclature	38
1.2.2 Connexin43 in post-surgical peritoneal adhesions	41
Chapter 2: Materials and Methods	43
2.1 Ex vivo Work	43
2.1.1 <i>Ex vivo</i> media preparation	43
2.1.2 Tissue collection.....	43
2.1.3 Induction of <i>ex vivo</i> adhesion formation and experimental design	44
2.2 In vivo Work	46
2.2.1 Experimental animals	46
2.2.2 Induction of adhesion formation	47
2.2.2.1 Surgery preparation and entry.....	47
2.2.2.2 Ischaemic button model of post-surgical peritoneal adhesions	47
2.2.2.3 Parietal peritoneum excision model of post-surgical peritoneal adhesions.....	48
2.2.2.4 Small bowel loop model of post-surgical peritoneal adhesions	48
2.2.2.5 Caecal abrasion model of post-surgical peritoneal adhesions	48
2.2.2.6 Surgery closure and post-recovery regime	49
2.2.3 Experimental Designs.....	50
2.2.3.1 Assessing various in vivo models of post-surgical peritoneal adhesions	50
2.2.3.2 Studying the role of Cx43 during peritoneal adhesion formation	50
2.2.3.3 Studying the role of endothelial-Cx43 during peritoneal adhesion formation	51
2.2.4 Peritoneal lavage	51
2.2.5 Peritoneal fluid tPA and PAI-1 total antigen ELISA	51
2.2.6 Macroscopic assessment of peritoneal adhesions.....	52
2.3 Histology	53
2.3.1 Tissue processing and microtome sectioning.....	53
2.3.2 Haematoxylin and eosin (H&E) staining	54
2.3.3 Picro-Sirius red (PSR) staining	54
2.3.4 Immunofluorescence staining.....	54
2.3.4.1 Heat-induced antigen retrieval (HIAR)	55
2.3.4.2 Proteolytic-induced antigen retrieval (PIAR)	56
2.4 Brightfield and confocal microscopy	57
2.5 Microscopic Assessment.....	58
2.5.1 Inclusion and exclusion criteria.....	58
2.5.2 Extent of inflammation.....	59
2.5.3 Fibrosis extent	60
2.5.4 Fibrosis severity.....	62

2.5.5 Protein levels of α SMA	63
2.5.6 Protein levels of Cx43 (ischaemic buttons)	64
2.5.7 Protein levels of TGF- β 1 (ischaemic buttons)	64
2.5.8 Protein levels of Cx43 (blood vessels)	64
2.7 Statistical Analysis.....	65
2.8 List of materials and reagent used.....	66
Chapter 3: Exploration of an ex vivo model of peritoneal adhesion formation.....	67
3.1 Introduction	67
3.1.1 Cell culture systems.....	67
3.1.2 <i>Ex vivo</i> tissue culture systems	68
3.2 Aims and objectives.....	70
3.3 Materials and Methods	71
3.3.1 Ex vivo work	71
3.3.1.1 Ex vivo media preparation.....	71
3.3.1.2 Tissue collection	71
3.3.1.3 Induction of ex vivo adhesion formation and experimental design.....	72
3.3.2 Histology	73
3.3.2.1 Tissue processing and microtome sectioning	73
3.3.2.2 Haematoxylin and eosin (H&E) staining.....	74
3.3.2.3 Picro-Sirius red (PSR) staining.....	74
3.3.2.4 Immunofluorescence staining.....	74
3.3.3 Brightfield and confocal microscopy	76
3.4 Results	77
3.4.1 Development of <i>ex vivo</i> adhesions required the presence of abrasion, clotted blood, and contact.....	77
3.4.2 Comparison of peritoneal strips cultured under different conditions.....	81
3.4.3 Development of <i>ex vivo</i> adhesions over time	83
3.5 Discussion	85
3.5.1. Condition for creating adhesions required the presence of abrasion, clotted blood, and contact and its relevance to the human system.....	85
3.5.2. Refinements for the mouse model.....	87
3.5.3. Microscopic assessments.....	88
3.5.4. Model suitability for studying the role of Cx43 during adhesion formation.....	90
3.6 Conclusion.....	93
Chapter 4: In search of an <i>in vivo</i> model of post-surgical peritoneal adhesion formation	95
4.1 Introduction	95
4.1.1 Selection of species	97
4.1.2 Selection of anatomic site.....	100
4.1.3 Selection of Adhesiogenic Stimulus.....	102
4.1.3.1 Abrasion injury	103
4.1.3.2 Excision injury.....	104
4.1.3.3 Presence of foreign bodies/materials	105
4.1.3.4 Ischaemic injury.....	106
4.1.4 Selection of timing and macroscopic assessment.....	107
4.1.5 Other considerations	111
4.1.6 Selection of model for studying the role of Cx43 during adhesion formation.....	112
4.2 Aims and Objectives.....	113
4.3 Materials and Method.....	114
4.3.1 Experimental animals	114
4.3.2 Induction of adhesion formation	114
4.3.2.1 Surgery preparation and entry.....	114
4.3.2.2 Ischaemic button model of post-surgical peritoneal adhesions	115
4.3.2.3 Parietal peritoneum excision model of post-surgical peritoneal adhesions.....	115
4.3.2.4 Small bowel loop model of post-surgical peritoneal adhesions	115
4.3.2.5 Caecal abrasion model of post-surgical peritoneal adhesions	116
4.3.2.6 Surgery closure and post-recovery regime	116
4.3.3 Macroscopic assessment of peritoneal adhesions.....	117
4.4 Results	118

4.4.1 Mice subjected to SBL did not survive until intended date of inspection.	118
4.4.2 Adhesions were observed in CA, IB, and PPE adhesion models.	120
4.5 Discussion	124
4.6 Conclusions	128
Chapter 5: The role of connexin43 during post-surgical peritoneal adhesion formation	129
5.1 Introduction	129
5.1.1 Skin.....	130
5.1.2 Liver	131
5.1.3 Heart	132
5.1.4 Kidney	133
5.1.5 Connexin43 in post-surgical peritoneal adhesions	134
5.1.6 Connexin43 antisense oligodeoxynucleotides.....	135
5.2 Aims and Objectives.....	137
5.3 Materials and Methods	139
5.3.1 In vivo Work.....	139
5.3.1.1 Experimental animals	139
5.3.1.2 Ischaemic button model of post-surgical peritoneal adhesions	139
5.3.1.3 Studying the role of Cx43 during peritoneal adhesion formation	140
5.3.1.4 Peritoneal lavage.....	141
5.3.1.5 Peritoneal fluid tPA and PAI-1 total antigen ELISA.....	141
5.3.1.6 Macroscopic assessment of peritoneal adhesions.....	142
5.3.2 Histology	143
5.3.2.1 Tissue processing and microtome sectioning	143
5.3.2.2 Haematoxylin and eosin (H&E) staining.....	144
5.3.2.3 Picro-Sirius red (PSR) staining.....	144
5.3.2.4 Immunofluorescence staining.....	144
5.3.2.5 Heat-induced antigen retrieval (HIAR)	145
5.3.3 Brightfield and confocal microscopy	146
5.3.4 Microscopic Assessment	147
5.3.4.1 Inclusion and exclusion criteria	147
5.3.4.2 Extent of inflammation	148
5.3.4.3 Fibrosis extent.....	149
5.3.4.4 Fibrosis severity	151
5.3.4.5 Protein levels of α SMA	152
5.3.4.6 Protein levels of Cx43 (ischaemic buttons)	153
5.3.4.7 Protein levels of TGF- β 1 (ischaemic buttons).....	153
5.3.5 Statistical Analysis	154
5.4 Results	155
5.4.1 Adhesions were observed as early as 6 hours post-surgery during early phase adhesion formation	155
5.4.2 Significant upregulation of Cx43 at the IB during early adhesion formation.	158
5.4.3 Significant increase in extent of inflammation at the IB during early adhesion formation.	160
5.4.4 Treatment with Cx43asODN significantly reduced pathological increases of Cx43 in the mesothelial region within the IB.	162
5.4.5 Treatment with Cx43asODN significantly reduced adhesion strength and severity at 7 days post-surgery.....	165
5.4.6 Treatment with Cx43asODN significantly reduced the fibrotic extent and severity of adhesions.	167
5.4.7 Effect of Cx43asODN treatment on the plasminogen-activating activity at 7 days post-surgery.	170
5.4.8 Treatment with Cx43asODN significantly reduced the degree of fibroblast activation at 7 days post-surgery.....	172
5.4.9 Treatment with Cx43asODN reduced the extent of adhesion formation during early adhesion formation.	174
5.4.10 Effect of Cx43asODN treatment on the extent of inflammation during early adhesion formation.	175
5.5 Discussion	178

5.5.1 Brief comparison of the <i>ex vivo</i> mouse peritoneal strip model and <i>in vivo</i> IB model	178
5.5.2 Adhesions formed by the IB model and their distribution	179
5.5.3 Increases of Cx43 protein levels during early adhesion formation	181
5.5.4 Role of Cx43 during adhesion formation	185
5.5.5.1 Role of Cx43 in inflammation and cell migration	185
5.5.5.2 Role of Cx43 in plasminogen-activator activity	189
5.6 Conclusions	190
Chapter 6: Exploring the role of endothelial-connexin43 in peritoneal adhesion formation	192
6.1 Introduction	192
6.1.1 Endothelial connexins and inflammation	193
6.1.2 Endothelial connexins and coagulation	194
6.1.3 Endothelial connexins and leukocyte adhesion	195
6.1.4 Endothelial-Cx43 transgenic mice	197
6.2 Aims and objectives.....	199
6.3 Materials and Methods	200
6.3.1 In vivo Work.....	200
6.3.1.1 Experimental animals	200
6.3.1.2 Ischaemic button model of post-surgical peritoneal adhesions	201
6.3.1.3 Peritoneal lavage.....	202
6.3.1.4 Peritoneal fluid tPA and PAI-1 total antigen ELISA.....	202
6.3.1.5 Macroscopic assessment of peritoneal adhesions.....	203
6.3.2 Histology	204
6.3.2.1 Tissue processing and microtome sectioning	204
6.3.2.2 Haematoxylin and eosin (H&E) staining.....	205
6.3.2.3 Picro-Sirius red (PSR) staining.....	205
6.3.2.4 Immunofluorescence staining	205
6.3.2.5 Proteolytic-induced antigen retrieval (PIAR).....	206
6.3.3 Brightfield and confocal microscopy	207
6.3.4 Microscopic Assessment	208
6.3.4.1 Inclusion and exclusion criteria	208
6.3.4.2 Fibrosis extent.....	209
6.3.4.3 Fibrosis severity	211
6.3.4.4 Protein levels of α SMA	212
6.3.5 Statistical Analysis	213
6.4 Results	213
6.4.1 Endothelial-Cx43 protein levels in tamoxifen-inducible endothelial cell-specific Cx43 transgenic mice were significantly reduced upon tamoxifen treatment.....	213
6.4.2 Knockdown of endothelial-Cx43 protein levels did not reduce adhesion formation.	216
6.4.3 The role of endothelial-Cx43 on adhesion formation.....	218
6.4 Discussion	223
6.4.1 Potential effects of Tamoxifen on the model	223
6.4.2 Endothelial Cx43 does not appear to be important for adhesion formation.....	225
6.6 Conclusions	229
Chapter 7: Summary and Conclusion	230
Publications	234
Appendix.....	235
APPENDIX I: SUPPLEMENTARY MATERIALS AND METHODS	235
References.....	237

Abstract

Little is known about the role of connexin43 in post-surgical peritoneal adhesion formation. Hence, this study sets out to investigate this through the use of *ex vivo* and *in vivo* models of adhesion formation. A novel mouse peritoneal strip *ex vivo* model was established and demonstrated to reproducibly generate adhesions. Through this model, it was shown that regions of adhesion had higher Cx43 protein levels. This phenomenon was further investigated, *in vivo*, in an ischaemic button model of adhesion formation after determining it to be the best model for the specific inquiry. Adhesions generated by this model were achieved by creating an ischaemic button through a ligation of an out-pouching of the abdominal sidewall with a suture. Assessment during early adhesion formation found pathological increase of Cx43 levels in the mesothelial regions within the ischaemic button. The significance of this pathological increase was further pursued through the use of Cx43 antisense oligodeoxynucleotides. Treatment with Cx43 antisense oligodeoxynucleotides saw a reduction in subsequent macroscopic assessment of adhesion strength and severity. Later microscopic assessment validated this and was further verified with the measurement of the overall plasminogen-activating activity in the peritoneal cavity. Treatment also saw a notable reduction in leukocyte recruitment and TGF- β 1 levels during early adhesion formation. In an attempt to further tease out the source of cellular contribution of Cx43 to adhesion formation, tamoxifen-induced endothelial-Cx43 specific transgenic mice were generated and subjected to the ischaemic button model. No difference in subsequent adhesion formation was observed between transgenic and their wild type controls. Overall, this study suggests Cx43 plays a significant role in the events of adhesion pathogenesis by regulating early leukocyte recruitment and TGF- β 1 levels. Findings also indicate this regulatory role is likely to be mediated by Cx43 derived from cells other than endothelial cells. Further investigation into the contribution of Cx43 from other major cell types involved in this series of events will open up new strategies for fine-tuning connexin-based therapeutics for post-surgical peritoneal adhesions.

Abbreviations

AMP	Adenosine monophosphate
ANOVA	Analysis of variance
ASBO	Adhesive small bowel obstruction
αSMA	Alpha smooth muscle actin
asODN	Antisense oligodeoxynucleotides
ATP	Adenosine triphosphate
CA	Caecal abrasion
CCL1	C-C motif ligand 1
CCR8	C-C motif receptor 8
CKD	Chronic kidney disease
Cx	Connexin
DAPI	4',6-Diamindino-2-phenylindole, dihyrdochloride
DNA	Deoxyribonucleic acid
EC-Cx43^{-/-}	Tamoxifen-treated Tie2-CreER ^{T2+/-} -Cx43 ^{fl/fl} mice
EC-Cx43^{fl/fl}	Tamoxifen-treated Cx43 ^{fl/fl} mice
ECM	Extracellular matrix
EDTA	Ethylenediamineettraacetic acid
EF	Entire fibrotic
FDA	US Food and drug administration
GJ	Gap junction
H&E	Haematoxylin and eosin
HIER	Heat induced antigen retrieval
IB	Ischaemic buttons
IBS	Ischaemic button and suture
IF	Immediate fibrotic
IHC	Immunohistochemistry staining
IL	Interleukin
mRNA	Messenger ribonucleic acid
NAD⁺	Nicotinamide adenine dinucleotide
PAA	Plasminogen-activating activity
PAI	Plasminogen activating inhibitors
PBS	Phosphate-buffered saline
PIER	Proteolytic-induced antigen retrieval
PMSF	Phenylmethylsulfonyl fluoride
PPE	Parietal peritoneum excision
PSR	Picro-Sirius red
ROI	Region of interest
RNA	Ribonucleic acid
RT	Room temperature
SBO	Small bowel obstruction
SBL	Small bowel loop
TGF-β1	Transforming growth factor beta 1
TNF-α	Tumor necrosis factor alpha
tPA	Tissue plasminogen activators
VEGF	Vascular endothelial growth factor

List of Figures and Tables

TABLE 1.1 POST-SURGICAL ADHESION CLASSIFICATION	19
FIGURE 1.1 KEY EVENTS DURING EARLY PATHOGENESIS OF POST-SURGICAL PERITONEAL ADHESIONS.	27
FIGURE 1.2 POPULAR MODELS OF POST-SURGICAL ADHESION FORMATION.....	32
FIGURE 1.3 MODEL OF A CONNEXIN MONOMER, HEMICHANNEL AND GAP JUNCTION CHANNEL.....	40
TABLE 1.2 CONNEXIN FAMILY MEMBERS IN HUMAN AND MOUSE	41
FIGURE 2.1 MAJOR STEPS FOR INDUCING <i>EX VIVO</i> ADHESIONS FORMATION.....	45
TABLE 2.1 EXPLORING CONDITIONS FOR INDUCING <i>EX VIVO</i> ADHESIONS	45
FIGURE 2.2 MAJOR STEPS DURING VARIOUS ADHESION-INDUCING OPERATIONS IN MICE.	49
TABLE 2.2 SCORING MATRIX FOR ASSESSING PERITONEAL ADHESIONS	53
TABLE 2.3 LIST OF PRIMARY ANTIBODIES USED IN THIS STUDY	56
TABLE 2.4 LIST OF SECONDARY ANTIBODIES USED IN THIS STUDY	57
FIGURE 2.3 INCLUSION AND EXCLUSION CRITERIA.....	58
FIGURE 2.4 DESIGNATING ROIs FOR SCORING THE EXTENT OF INFLAMMATION.....	59
FIGURE 2.5 SCORING MATRIX FOR GRADING EXTENT OF INFLAMMATION	60
FIGURE 2.6. DEFINING REGIONS AROUND THE ISCHAEMIC BUTTON	62
FIGURE 2.7. EXAMPLE OF IMAGE ANALYSIS WORKFLOW FOR PSR STAINED IMAGES.....	63
FIGURE 3.1 MAJOR STEPS FOR INDUCING <i>EX VIVO</i> ADHESIONS FORMATION.....	73
TABLE 3.1 EXPLORING CONDITIONS FOR INDUCING <i>EX VIVO</i> ADHESIONS	73
TABLE 3.2 LIST OF PRIMARY ANTIBODIES USED IN THIS STUDY	76
TABLE 3.3 LIST OF SECONDARY ANTIBODIES USED IN THIS STUDY	76
FIGURE 3.2 EXAMPLES OF <i>EX VIVO</i> ADHESIONS FORMED IN PERITONEAL STRIPS CULTURED UNDER CONDITION 1	79
FIGURE 3.3 MICROSCOPIC ASSESSMENT OF PERITONEAL STRIPS THAT WERE CULTURED WITH A PLACEMENT OF BLOOD CLOT BETWEEN ABRADED SURFACES.....	80
FIGURE 3.4 H&E ASSESSMENT OF CULTURED PERITONEAL STRIPS CULTURED UNDER DIFFERENT CONDITIONS	82
FIGURE 3.5 MICROSCOPIC ASSESSMENT OF PERITONEAL STRIPS CULTURED UNDER CONDITION 1 OVER 72 HOURS	84
FIGURE 4.1 PRISMA FLOWCHAT OF LITERATURE SEARCH STRATEGY.....	97
TABLE 4.1 OVERVIEW OF RATS AND MICE PERITONEAL ADHESIONS MODELS	99
TABLE 4.2 TYPES OF STIMULI EMPLOYED TO INDUCE ADHESIONS.....	103
TABLE 4.3 SCORING MATRIX FOR ASSESSING STRENGTH OF PERITONEAL ADHESIONS	111
FIGURE 4.2 MAJOR STEPS DURING VARIOUS ADHESION-INDUCING OPERATIONS IN MICE.	116
TABLE 4.4 SCORING MATRIX FOR ASSESSING PERITONEAL ADHESIONS	118
FIGURE 4.3 SMALL BOWEL OBSTRUCTION OBSERVED IN SMALL BOWEL LOOP MODEL.....	120
FIGURE 4.4 OVERVIEW OF THE ADHESIONS OBSERVED IN VARIOUS ADHESION MODELS UPON NECROPSY.	122
FIGURE 4.5 COMPARING THE PERFORMANCE OF POPULAR ADHESION MODELS.....	123
TABLE 4.5 OVERVIEW OF THE PERFORMANCE OF ADHESION MODELS EXPLORED	128
FIGURE 5.1 MAJOR STEPS DURING VARIOUS ADHESION-INDUCING OPERATIONS IN MICE.	140
TABLE 5.1 SCORING MATRIX FOR ASSESSING PERITONEAL ADHESIONS	143
TABLE 5.2 LIST OF PRIMARY ANTIBODIES USED IN THIS STUDY	146
TABLE 5.3 LIST OF SECONDARY ANTIBODIES USED IN THIS STUDY	146
FIGURE 5.2 INCLUSION AND EXCLUSION CRITERIA.....	147
FIGURE 5.3 DESIGNATING ROIs FOR SCORING THE EXTENT OF INFLAMMATION.....	148
FIGURE 5.4 SCORING MATRIX FOR GRADING EXTENT OF INFLAMMATION	149
FIGURE 5.5 DEFINING REGIONS AROUND THE ISCHAEMIC BUTTON	151
FIGURE 5.6 EXAMPLE OF IMAGE ANALYSIS WORKFLOW FOR PSR STAINED IMAGES.....	152
FIGURE 5.7 TYPES OF ADHESIONS OBSERVED AT EARLY AND LATE PHASES OF ADHESION DEVELOPMENT	157
FIGURE 5.8 Cx43 WAS SIGNIFICANTLY UPREGULATED AT THE IB REGIONS DURING EARLY ADHESION FORMATION	159
FIGURE 5.9 EXTENT OF INFLAMMATION WAS SIGNIFICANTLY INCREASED DURING EARLY ADHESION FORMATION	161

FIGURE 5.10 Cx43asODN SIGNIFICANTLY REDUCED Cx43 PROTEIN LEVELS IN THE REGIONS WITHIN THE IB AT 24 HOURS POST-SURGERY.....	164
FIGURE 5.11 TREATMENT WITH Cx43asODN SIGNIFICANTLY REDUCED SEVERITY AND STRENGTH OF ADHESIONS DEVELOPED AT 7 DAYS POST-SURGERY.....	166
FIGURE 5.12 TREATMENT WITH Cx43asODN SIGNIFICANTLY REDUCED FIBROSIS EXTENT OF THE ADHESIONS DEVELOPED AT 7 DAYS POST-SURGERY.....	169
FIGURE 5.13 TREATMENT WITH Cx43asODN SIGNIFICANTLY REDUCED FIBROSIS SEVERITY OF THE ADHESIONS DEVELOPED AT 7 DAYS POST-SURGERY.....	170
FIGURE 5.14 EFFECT OF Cx43asODN TREATMENT ON TISSUE PLASMINOGEN ACTIVATOR (tPA) AND PLASMINOGEN ACTIVATOR INHIBITOR-1 (PAI) PROTEINS IN THE PERITONEAL FLUID AT DAY 7 POST-SURGERY.....	171
FIGURE 5.15 TREATMENT WITH Cx43asODN SIGNIFICANTLY REDUCED ASMA PROTEIN LEVELS WITHIN THE FIBROTIC REGION AT 7 DAYS POST-SURGERY.....	173
FIGURE 5.16 Cx43asODN TREATMENT APPEARED TO REDUCE THE EXTENT OF ADHESIONS DURING EARLY ADHESION FORMATION	174
FIGURE 5.17 Cx43 asODN TREATMENT SIGNIFICANTLY REDUCED THE EXTENT OF INFLAMMATION WITHIN THE IB AT 6 AND 24 HOURS	176
FIGURE 5.18 TREATMENT WITH Cx43asODN APPEARED TO REDUCE TGF- β 1 PROTEIN LEVELS WITHIN THE IB REGION DURING EARLY PHASE OF ADHESION FORMATION.....	177
FIGURE 5.19 EPIDIDYMAL FAT PAD DISTRIBUTION IN THE CAUDAL ABDOMINAL REGION	180
FIGURE 5.20. SCHEMATIC DIAGRAM OF Cx43 PROTEIN LEVELS DURING EARLY ADHESION FORMATION.....	183
FIGURE 5.21 NEUTROPHILS IN THE MESOTHELIAL REGIONS.....	184
FIGURE 6.1 Cx43 SUPPORTS PRO-INFLAMMATORY SIGNALING IN STIMULATED ENDOTHELIAL CELLS BY FACILITATING THE INITIAL ADHESIONS EVENTS OF LEUKOCYTE RECRUITMENT.....	197
FIGURE 6.2 MAJOR STEPS DURING VARIOUS ADHESION-INDUCING OPERATIONS IN MICE.	202
TABLE 6.1 SCORING MATRIX FOR ASSESSING PERITONEAL ADHESIONS	204
TABLE 6.2 LIST OF PRIMARY ANTIBODIES USED IN THIS STUDY	207
TABLE 6.3 LIST OF SECONDARY ANTIBODIES USED IN THIS STUDY	207
FIGURE 6.3 INCLUSION AND EXCLUSION CRITERIA.....	209
FIGURE 6.4 DEFINING REGIONS AROUND THE ISCHAEMIC BUTTON	211
FIGURE 6.5 EXAMPLE OF IMAGE ANALYSIS WORKFLOW FOR PSR STAINED IMAGES.....	212
FIGURE 6.6 ENDOTHELIAL-Cx43 PROTEIN LEVELS IN THE TAMOXIFEN-TREATED ENDOTHELIAL-SPECIFIC Cx43 TRANSGENIC MICE EC-Cx43 ^{-/-} WAS REDUCED UPON TAMOXIFEN TREATMENT.....	214
FIGURE 6.7 ENDOTHELIAL-Cx43 PROTEIN LEVELS IN THE TAMOXIFEN-TREATED EC-Cx43 ^{-/-} TRANSGENIC MICE WAS SIGNIFICANTLY REDUCED UPON TAMOXIFEN TREATMENT.....	215
FIGURE 6.8 KNOCKDOWN OF ENDOTHELIAL-Cx43 PROTEIN LEVELS DID NOT APPEAR TO AFFECT ADHESION DEVELOPED AT 7 DAYS POST-SURGERY.....	218
FIGURE 6.9 KNOCKDOWN OF ENDOTHELIAL-Cx43 PROTEIN LEVELS DID NOT APPEAR TO AFFECT FIBROSIS EXTENT OF THE ADHESIONS DEVELOPED AT 7 DAYS POST-SURGERY.....	220
FIGURE 6.10 KNOCKDOWN OF ENDOTHELIAL-Cx43 PROTEIN LEVELS DID NOT APPEAR TO AFFECT FIBROSIS SEVERITY OF THE ADHESIONS DEVELOPED AT 7 DAYS POST-SURGERY.....	221
FIGURE 6.11 KNOCKDOWN OF ENDOTHELIAL-Cx43 PROTEIN LEVELS DID NOT APPEAR TO AFFECT ASMA PROTEIN LEVELS WITHIN THE FIBROTIC REGION AT 7 DAYS POST-SURGERY.....	222
FIGURE 6.12 EFFECT OF ENDOTHELIAL-Cx43 KNOCKDOWN ON TISSUE PLASMINOGEN ACTIVATOR (tPA) AND PLASMINOGEN ACTIVATOR INHIBITOR-1 (PAI) PROTEINS IN THE PERITONEAL FLUID AT DAY 7 POST-SURGERY.	223
SUPPLEMENTARY TABLE 1. <i>Ex vivo</i> MEDIA AND TISSUE COLLECTION REAGENT COMPONENTS.....	235
SUPPLEMENTARY TABLE 2. PRIMERS USED FOR GENOTYPING	236
SUPPLEMENTARY TABLE 3. SEQUENCE OF MOUSE Cx43 ANTISENSE OLIGODEOXYNUCLEOTIDES ...	236
SUPPLEMENTARY TABLE 4. SET PROGRAMME FOR H&E STAINING.....	236

Scope of Work and Structure of Thesis

Post-surgical peritoneal adhesions are pathological connections found between tissue surfaces at non-anatomic locations of the peritoneal cavity, usually as a result of injury sustained during surgery (Braun & Diamond, 2014; diZerega & Campeau, 2001). Adhesions have been reported to develop in at least 93% of patients undergoing intra-abdominal operations (Ellis, 1997) and their presence may give rise to a series of clinical issues, such as adhesive small bowel obstruction (SBO), secondary infertility, chronic abdominal pain, adhesiolysis and related complications (ten Broek, Bakkum, Laarhoven, & van Goor, 2016).

Connexins (Cx) are proteins that facilitate the direct cytoplasmic exchange of ions and small molecules between adjacent cells through the formation of intercellular channels, or gap junctions (Nielsen et al., 2012). Connexins have an important role in supporting the sharing of metabolic demands across cells and tissues (Evans, De Vuyst, & Leybaert, 2006). Despite the potential involvement of connexins in various facets of adhesion pathogenesis (i.e. inflammation, coagulation, mesenchymal transition), their contribution in it has yet to be explored (Cogliati et al., 2016; Okamoto & Suzuki, 2017; Soon, Chua, & Becker, 2016). In particular, little is known about the contribution of Cx43 to adhesion pathogenesis, which is also the most comprehensively studied Cx and by far the most abundant and ubiquitous isoform in mammalian cells (Nielsen et al., 2012).

This project aims to fill these research gaps by approaching it systematically by first reviewing existing literature (detailed in **Chapter 1**) before investigating the role of

Cx43 in adhesion formation using *ex vivo* and *in vivo* adhesion models. Overall descriptions of the materials and methodology used in this endeavour are detailed in **Chapter 2**.

Recently, a rat peritoneal strip *ex vivo* adhesion model established by Saed and colleagues has been described to form adhesive bands within 72 hours (Saed, Fletcher, & Diamond, 2016). One advantage of this model is that it allows tracking and sampling following the identification of desired events (i.e. vascularisation or adhesion creation). This allows molecular events of adhesion formation to be characterised. Furthermore, multiple strips may be obtained from each rat, allowing multiple arms of a study to be tested using one animal. Adaptation of the rat *ex vivo* model to mouse is explored and detailed in **Chapter 3**.

Despite the advantages of an *ex vivo* model, studying the role of Cx43 in an *in vivo* model is still more physiologically relevant. To achieve this, it is essential to first identify the post-surgical peritoneal adhesion model best suited for this purpose. Work in **Chapter 4** addresses this question through two phases. First, a systematic literature search was first performed to explore the types of models available. The findings were then discussed with respect to the selection of species, anatomical sites, adhesiogenic stimuli, and macroscopic assessment. Next, concluding from these findings, relevant models are selected and explored to identify the model most ideal for investigating this specific inquiry. Models were assessed based on established macroscopic assessment scoring for the quality and quantity of adhesions, and also evaluated based other considerations (e.g. degree of model standardisation, identification and definition of site trauma).

After identifying the ideal model, the role of Cx43 in post-surgical peritoneal adhesion formation is explored and detailed in **Chapter 5**. To investigate whether or not Cx43 is implicated during adhesion formation, the potential pathological changes to Cx43 levels during early phase adhesion formation is investigated via immunohistochemistry. Next, to determine the significance of the contribution of Cx43 levels to adhesion formation, an experimental group of mice were treated with Cx43 antisense oligodeoxynucleotides to significantly reduce its protein levels. Treatment effects were then evaluated at both early and late phase of adhesion formation. Effects on late phase adhesion formation were evaluated using macroscopic assessment based on established scoring matrices that were later verified with microscopic assessment and validated with measurements of the overall plasminogen-activating activity in the peritoneal cavity. Effects on early phase revealed potential mechanisms of which Cx43 may have affected adhesion formation.

Attempts to further dissect the contribution of Cx43 are detailed in **Chapter 6**. The role of endothelial-Cx43 in post-surgical peritoneal adhesion formation is explored using our tamoxifen-inducible endothelial-Cx43 specific transgenic mice. Effects of tamoxifen-induced endothelial-Cx43 knockdown on late phase adhesion formation were evaluated in a similar manner as in the previous chapter.

Finally, **Chapter 7** summarises the previous chapters and concludes this work.

Hypothesis

This thesis hypothesises that Cx43 has an important role during early adhesion pathogenesis, and that a single treatment with Cx43asODN during this period will transiently reduce Cx43 protein levels, resulting in better adhesion formation outcomes due to fewer recruited leukocytes to the injury site, reduced activated fibroblasts, and promotion of plasminogen-activating activity.

Chapter 1: Literature Review

1.1 Post-surgical peritoneal adhesions

1.1.1 Definition

Post-surgical peritoneal adhesions are pathological connections found between tissue surfaces at non-anatomic locations of the peritoneal cavity, usually as a result of injury sustained during surgery (Braun & Diamond, 2014; diZerega & Campeau, 2001). These pathological connections are heterogenous; the time taken to develop symptoms varies and the clinical presentation ranges from “strands, bands, or even sheets of tissues of different thickness or tenacity” (Beyene, Kavalukas, & Barbul, 2015; Moris et al., 2017; Okabayashi et al., 2014). In general, formation of post-surgical adhesions may be categorised based on the classification provided by Diamond and Nezhat (**Table 1.1**) (Beyene et al., 2015; Diamond & Nezhat, 1993). Under this classification, adhesions may be distinguished as “Type 1” or “Type 2”. The first category “Type 1”, encompasses adhesions that have developed at sites previously devoid of adhesions. The second category “Type 2”, encompasses adhesions that have reformed after adhesiolysis. These different types of adhesions can be further categorised as “A”, if the found adhesions are located at sites where no surgical procedures were previously performed; or “B”, if the found adhesions are located at sites where surgical procedures were previously performed (Diamond & Nezhat, 1993).

Table 1.1 Post-surgical adhesion classification

Classification	Description
Type 1	De novo adhesions
A	No surgical procedures were previously performed at site
B	Surgical procedures were previously performed at site
Type 2	Reformed adhesions after adhesiolysis
A	No surgical procedures were previously performed at site
B	Surgical procedures were previously performed at site

Adapted from (Diamond & Nezhat, 1993)

1.1.2 Epidemiology, complications, and economic burden

Almost like a surgical footprint, 93% of patients undergoing intra-abdominal operations have been reported to develop adhesions (Ellis, 1997). Epidemiological data provided by the Surgical and Clinical Adhesions Research group reported that lower gastrointestinal procedures involving the small intestine, colon, appendix, or the uterus, were more likely to result in adhesions (Ellis et al., 1999; Lower et al., 2004; Parker et al., 2001; Parker et al., 2005). It is estimated that 33% of patients who received abdominal or pelvic surgery, were subsequently readmitted for adhesion-related complications (Ellis et al., 1999). Lower gastrointestinal procedures such as total proctocolectomy (15.4%) and ileostomy (10.6%) were reported to have the highest risk of adhesion-related hospital readmission (Parker et al., 2005). On the contrary, procedures performed in the upper abdominal region (e.g. stomach, pancreas, gallbladder) were reported to have lower rates of adhesion formation (ten Broek et al., 2013a).

The presence of adhesions may spawn a series of clinical issues, such as adhesive small bowel obstruction (SBO), secondary infertility, chronic abdominal pain,

adhesiolysis and related complications (ten Broek et al., 2016). Adhesions account for up to 96% of patients with chronic abdominal and/or pelvic pain (Swank & Jeekel, 2004), 60% to 70% of patients with SBO (Ellis, 1998; Parker et al., 2001; ten Broek et al., 2013a) and 20% to 40% of patients with secondary infertility during laparoscopic evaluation (Kaminski et al., 2006; Milingos et al., 2000). These numbers are also likely to be underestimated because diagnosis of adhesions can only be confirmed at a second laparotomy or laparoscopy. Considering the incidence of post-surgical adhesions, it is unethical to perform a repeated surgery on patients undergoing an abdominal procedure just to evaluate post-surgical adhesion formation (Okabayashi et al., 2014).

The presence of post-surgical adhesions also causes other clinical issues that impact both doctors and patients. Adhesion presence increases the difficulty of repeated operations, prolonging the surgery time by almost an average of 25 minutes (Beck et al., 2000; Coleman, McLain, & Moran, 2000). Adhesiolysis further increases the risk of bowel injury (Van Der Krabben et al., 2000). Furthermore, it may increase the difficulty of future laparoscopic surgeries or make it not even possible (Coleman et al., 2000; Van Der Krabben et al., 2000). In addition, the length of hospital stay after peritoneal adhesion-related surgery is usually longer due to an extended recovery period (Beck et al., 2000; Coleman et al., 2000).

Indeed, post-surgical adhesions have a significant economic impact. In the USA, the total cost of hospital and surgical expenditures for the management of adhesion-related complications has increased from \$1.3 billion in 1994 to \$5 billion in 2001 (Parker et al., 2001). In 2005, \$3.45 billion was estimated to be the direct hospital

costs of SBO in USA alone (Wiseman, 2008). Despite of an on-going and compelling need, there is still a paucity in the existence of an ideal method or agent for preventing intra-abdominal adhesion (Becker & Stucchi, 2004; ten Broek et al., 2016).

1.1.3 Early pathogenesis of post-surgical peritoneal adhesion formation

Post-surgical peritoneal adhesions are considered to be a consequence of the pathological process of peritoneal healing following injury, usually arising from abdominal surgery (Arung, Meurisse, & Detry, 2011; diZerega & Campeau, 2001; Mutsaers et al., 2015). Almost immediately, the peritoneal injury incites an inflammatory response that is characterised by a simultaneous release of a cocktail of cytokines and chemokines, recruitment of inflammatory cells, and the activation of the coagulation cascade (Beyene et al., 2015; Schnuriger et al., 2011). In the following sections, these processes are described in further detail.

1.1.3.1 Early responders to the scene

Following peritoneal injury, the release of cocktail of cytokines and chemokines sets the stage for the biochemical and cellular cascades that ensue. While released cytokines kickstarts the repair process, chemokines promote the recruitment of leukocytes to the site of injury. The early responders are predominantly neutrophils, with a gradual shift to macrophages after 24 hours (Maciver, McCall, & James Shapiro, 2011).

These recruited leukocytes are suggested to play a crucial role in the outcome of adhesion formation (Beyene et al., 2015; Maciver et al., 2011). The reduction of

neutrophils through the use of cyclophosphamide was shown to reduce adhesion quantity and severity (Vural et al., 1999). In another study, abrogating the interaction between chemokine (CCL1) and its receptor (CCR8) interaction in peritoneal macrophages also hampered the migration of peritoneal macrophages, resulting in a significant reduction of peritoneal adhesions (Hoshino et al., 2007).

Also, activated stromal mast cells increase vascular permeability through the release of histamine and kinins (Sammour, Kahokehr, Soop, & Hill, 2010). The increase of vascular permeability results in the accumulation of exudates made up of plasma protein and cells (Beyene et al., 2015). Furthermore, vascular endothelial growth factor (VEGF) release from mast cells contributes to angiogenesis in areas of peritoneal injury. Its contribution to adhesion formation is underscored by the reduction in adhesion formation upon anti-VEGF interventions (Ignjatovic et al., 2010; Saltzman, Olson, Mohanraj, Carson, & Ramakrishnan, 1996). Finally, these early responders are thought to contribute to the outcome of adhesion formation through the initiation of another robust cytokine cascade upon recruitment (Beyene et al., 2015; Maciver et al., 2011).

1.1.3.2 Release of cytokine cocktails

Amid the milieu of cytokines released in the inflamed peritoneal cavity, the earliest cytokines present are interleukin (IL)-1 and tumor necrosis factor (TNF)- α (Beyene et al., 2015). The early role for IL-1 is supported by the subsequent increase in adhesion scores when IL-1a is immediately administered following caecal abrasion in rats (Hershlag, Otterness, Bliven, Diamond, & Polan, 1991). Conversely, reduced adhesion formation is subsequently observed in mice receiving a prophylactic

treatment of IL-1 antibodies before caecal abrasion (Hershlag et al., 1991). On the same note, early increases of TNF- α (measured at 30, 90 and 180 minutes) were correlated with the subsequent development of more severe adhesions (Kaidi et al., 1995). The stimulation of human mesothelial cells with these early cytokines results in pathological changes in cell morphology accompanied with the release of IL-6 and 8 (Sylvia et al., 2006; Topley et al., 1993). Although these cytokines are thought to play a role in recruitment of leukocytes, their role in adhesion formation is still not well defined (Beyene et al., 2015).

One of the best characterised cytokines is the transforming growth factor- β 1 (TGF- β 1), due to its strong association to adhesion formation (Beyene et al., 2015). Proposed to be the main pro-fibrotic mediator of adhesion formation, it is found to be elevated in peritoneal adhesion tissue and fluids sampled from patients (Chegini, 2008; Delaney, Kasprzycka, Ciemerych, & Zimowska, 2017; Saed, Zhang, & Diamond, 2001). Patient-isolated adhesion fibroblasts, which are myofibroblasts isolated from adhesion tissues, are found to have elevated levels of TGF- β 1 (Saed et al., 2001). Furthermore, a significant increase in TGF- β 1 was observed in these isolated adhesion fibroblasts in response to treatment of hypoxia, a strong adhesiogenic stimulus (Saed et al., 2001). Conversely, lower adhesion scores were recorded in heterozygous TGF- β 1 transgenic mice at 7 days post-challenge with talc, another potent adhesiogenic stimulus (Krause, Katz, Wheeler, Ebner, & McKinnon, 1999).

Stimulation of normal fibroblasts with TGF- β 1 also induces their differentiation into the adhesion phenotype, which is characterised by the acquisition of a *de novo*

expression of α smooth muscle actin (α SMA) (Asazuma-Nakamura et al., 2009; Desmouliere, Geinoz, Gabbiani, & Gabbiani, 1993). Apart from regulating fibroblast differentiation, TGF- β 1 is also recently demonstrated to be involved in mesothelial-to-mesenchymal transition (Sandoval et al., 2016). This transition is described to be one of the possible origins for peritoneal adhesion fibroblasts during adhesion formation (Aoki et al., 2013; López-Cabrera, 2014; Sandoval et al., 2016; Yang, Chen, & Lin, 2003). Mesothelial cells stimulated with TGF- β 1 are able to induce acquisition of a *de novo* expression of α SMA over time, which is characteristic of myofibroblastic phenoconversion (Asazuma-Nakamura et al., 2009). Moreover, blockage of TGF- β 1 in an ischaemic button adhesion mouse model hampered mesenchymal transition of mesothelial cells, and reduced overall adhesion severity (Sandoval et al., 2016).

1.1.3.3 The coagulation cascade, fibrinolytic balance, and matrix deposition

Following almost simultaneously with the inflammatory phase, the coagulation cascade generally results from vascular injury, which exposes tissue factor and von Willebrand factor (Gaertner & Massberg, 2016). The elicited coagulation cascade initiates the mobilisation of coagulation factors that ultimately results in the generation of thrombin, which is responsible for the conversion of fibrinogen to fibrin (Engelmann & Massberg, 2013). Secondary vasodilation and capillary leak induced by cytokine release from recruited leukocytes in the peritoneal cavity eventually results the intraperitoneal accumulation of fibrin-rich exudates (Beyene et al., 2015). Forming as early as 3 hours, these fibrin deposits help form the first provisional bridge between two unrelated surfaces (Maciver et al., 2011; Rout, Saed, & Diamond, 2002). On this note, many investigators consider adhesion formation to

be a result of the imbalance between fibrin deposition and degradation (Arung et al., 2011; Beyene et al., 2015; Braun & Diamond, 2014; Maciver et al., 2011; Moris et al., 2017).

In the event fibrin deposits are degraded completely, peritoneal healing resumes without adhesion formation (Arung et al., 2011). However, if these fibrin deposits persist, its presence facilitates the invasion and proliferation of fibroblasts. Consequently, this leads to the deposition of extracellular matrix (ECM), which includes collagen and fibronectin (Braun & Diamond, 2014). Eventually, the matrix develops into a more permanent fibrous connection, leading to the development of adhesions (Arung et al., 2011; Braun & Diamond, 2014; Maciver et al., 2011). This highlights the importance of fibrinolysis for peritoneal healing without adhesion formation.

Plasmin, an active serine protease that is essential for breaking down fibrin, is converted from its inactive substrate form, plasminogen, by the action of plasminogen activators. In the peritoneal cavity, 95% of the plasminogen-activating activity (PAA) is contributed by tissue plasminogen activators (tPA) (Holmdahl, Eriksson, al-Jabreen, & Risberg, 1996). The expression of tPA can be found on mesothelial cells, endothelial cells, fibroblasts, and macrophages (Arung et al., 2011; Braun & Diamond, 2014). It is important to point out that tPA is a strong plasminogen activator only in the presence of fibrin. This results in its activation only at sites where it is required, thus preventing systemic activation (Ichinose, Takio, & Fujikawa, 1986; Norrman, Wallen, & Ranby, 1985). However, the ability

of tPA in mounting an effective fibrinolytic response also depends on the presence of plasminogen activating inhibitors (PAIs).

One of the most potent inhibitors of tPA is PAI-1 (Moris et al., 2017). It inhibits tPA by antagonising and forming inactive complexes, resulting in reduced PAA (Beyene et al., 2015). Of note, significantly higher levels of PAI-1 are detected in peritoneal tissues of patients with adhesions, which have suggested its use as a biomarker to identify patients with a higher likelihood of forming adhesions (Ivarsson, Bergstrom, Eriksson, Risberg, & Holmdahl, 1998). Along these lines, other proteins, such as α 2-anti-plasmin, α 2-macroglobulin, and α 1-anti-trypsin, may also directly inhibit the fibrolytic action of plasmin, although their contribution in peritoneal fibrinolysis is still unclear (Holmdahl, 2000).

Overall, the outcome of peritoneal healing (normal healing or adhesion formation) appears to hinge on the balance between the plasminogen activators and their inhibitors (**Fig. 1.1**) (Moris et al., 2017). Pathological conditions such as surgery trauma, in which PAA is markedly decreased due to increasing concentrations of PAI-1 and reduced oxygenation of the tissues, leads to the persistence of the fibrin matrix (diZerega, 1997). This matrix gradually becomes more organised, providing the necessary scaffolding that leads to the development of adhesions (Arung et al., 2011; Braun & Diamond, 2014).

Figure 1.1 Key events during early pathogenesis of post-surgical peritoneal adhesions.

Peritoneal injury incites an inflammatory response that is characterised by a release of a cocktail of cytokines, leukocyte recruitment, and the activation of the coagulation cascade. This leads to a build up of fibrin deposits, which forms the first provisional bridge between two surfaces. Following on, adhesion formation hinges on the level of plasminogen activating activity, which is determined by the levels of plasminogen activators (**tPA**) and plasminogen activator inhibitors (**PAI-1**). (**A**) If fibrin deposits are degraded completely, peritoneal healing resumes without adhesion formation. (**B**) However, if these fibrin deposits persist, its presence facilitates the invasion and proliferation of fibroblasts leading to the deposition of extracellular matrix. Eventually, the persistence of the fibrin deposits establishes the matrix into a more permanent fibrous connection, leading to the ensuing development of adhesions

1.1.4 Experimental models

1.1.4.1 Cell culture systems

Cell culture systems shed light on how cells function and respond to stimuli. For instance, cultures of fibroblasts taken from either normal or adhesion-prone peritoneum may be useful substrates to determine the potential of various test agents in limiting, arresting, or rescuing the adhesion phenotype (Diamond, El-Hammady,

Munkarah, Bieber, & Saed, 2005; Saed, Al-Hendy, Salama, & Diamond, 2008; Saed & Diamond, 2004b).

However, these single cell systems have their limitations. Due to their nature, these systems are isolated cultures and so do not take into account the effects of the surrounding cells or the matrix in their physiological environment that would be found *in vivo*. This environment may sometimes have a strong influence over their response to stimuli (Yin, Meng, Zhang, & Sun, 2009) or may impact the cell's differentiation fate and function (Ravi, Paramesh, Kaviya, Anuradha, & Solomon, 2015). On this note, changes in cell morphology can be quickly observed in primary cultures of mesothelial cells, in which they become vacuolated and flattened after 2-3 passages (Beyene et al., 2015). This short window may make it challenging to investigate how they function and respond to stimuli.

1.1.4.2 Ex vivo tissue culture system

Only up to recent, one *ex vivo* model system has been described (Saed et al., 2016). Briefly, this model involves culturing rat peritoneal strips that receive a combination of abrasion injury and clotted blood. Under these conditions, a single adhesive band found to be dense, opaque, and vascularized, is produced as early as 24 hours (Saed et al., 2016). The authors also describe adhesive bands being created by 48 hours when abraded strips were cultured in the absence of clotted blood (Saed et al., 2016). This *ex vivo* model presents a cost-effective solution for assessing or screening the efficacy of anti-adhesion agents. In contrast, *in vivo* studies are usually a costly venture, requiring around large numbers of animals per test group (Wiseman, 2000). These costs can easily snowball when the calibration of the employed model,

exploration of multiple variants, and comparing dose-responses are involved. Fortunately, in this *ex vivo* model, multiple strips may be obtained from each rat, allowing multiple arms of a study to be tested using one animal, drastically reducing the study costs.

In addition, an *ex vivo* model offers a less isolated system compared to conventional cell culture based systems. The lack of cues from their surrounding cells or matrix in their physiological environment in the latter systems may influence their response to drug stimuli differently (Yin et al., 2009). Furthermore, questions about the long-term effects of drugs that are challenging to assess using cell culture based systems due to culture confluence limitations, may be answered easily with the *ex vivo* model. Hence, this *ex vivo* model offers cost-effective solution to bridge the gap between cell culture systems and animal drug testing models for the assessment of potential anti-adhesions agents.

1.1.4.3 In vivo models

Despite the advantages of the *ex vivo* model, *in vivo* models still generate adhesions closest to its native physiological environment. This is necessary for the better understanding of the pathogenesis of adhesion formation, which is critical for identifying new therapeutic targets. Since the first induction of adhesions in an animal model in 1889 (von Dembowski, 1889), many post-surgical adhesion animal models have been developed and refined to achieve this. In general, most of these models share a similar design. A chosen adhesiogenic stimulus is exacted on a designated organ of interest in the selected animal species. Next, depending on experimental design, a test article (i.e. liquids, gels, or barriers) may be explored.

Animals are then allowed to recover for about one week post-operation before the developed adhesions are assessed, usually by gross macroscopic and/or histological methods (Wiseman, 2000).

One of the most frequently employed methods of inducing adhesion is by exacting abrasion injury to the caecum (**Fig. 1.2a**). This is commonly employed to model intestinal surgery. Abrasion injury is usually achieved through gauze (Du, Liu, Xin, & Liu, 2015; Karaca et al., 2016; Koca, Tarhan, Kaya, & Gokce Ceylan, 2016; Lee et al., 2016), but may also be achieved using scalpel (Andrews, Ford, & Martin, 2012; Aysan, Ayar, Aren, & Cifter, 2002), and sometimes using brush (209^h), or abrasive paper (Whang et al., 2011). The end point of abrasion is usually described by the number of times the site is abraded, although not practiced widely. Instead, petechial haemorrhage appears to be the general consensus for the end point of abrasion (Aysan et al., 2002; Irkorucu, Ferahkose, Memis, Ekinici, & Akin, 2009; Karaca et al., 2016; Koca et al., 2016; Lai et al., 2005; Xu et al., 2002a). The fibrin from the bleed facilitates the initial adhesion between the two surfaces during early adhesiogenesis (Wiseman, 2000). Considering this, if the amount of bleed is not standardised, it may present a confounding variable to the development of adhesion (diZerega & Campeau, 2001; Kraemer et al., 2014). Desiccation for up to five minutes following haemorrhage is also sometimes included and is suggested to be a potent combination for the creation of adhesions (Ozel, Avsar, Topaloglu, & Sahin, 2005). Variations following abrasion also include the addition of chemical irritants such as iodine (e.g. iodine and (Osada et al., 1999) alcohol (Liebman, Langer, Marshall, & Collins, 1993)). Abrasive injuries to the caecum is also sometimes combined with a parietal peritoneal abrasion (Ignjatovic et al., 2010; Townsend et

al., 2011; Wei et al., 2016a; Wei et al., 2015; Wei et al., 2016b) or a peritoneal excisional defect (Bayhan et al., 2016; Poehnert, Abbas, Kreipe, Klemptner, & Winny, 2015, 2015; Poehnert et al., 2015; Wu et al., 2015), sometimes with/without approximation of the two sites with sutures.

Another frequently employed method of inducing adhesion is the creation of a parietal peritoneal excisional defect (**Fig. 1.2b**). This is usually achieved by removing the parietal peritoneum and the underlying muscle with scalpel or surgical scissors. This models surgical trauma during abdominal or gynaecologic surgery. Standardisation of the area of excision may be achieved through the use of similar-sized biopsy punches. Although repeated findings from earlier work have demonstrated that different-sized defects heal in the same time frame without the formation of adhesions (Ellis, 1962; Hubbard, Khan, Carag, Albites, & Hricko, 1967; Robbins, Brunschwig, & Foote, 1949), later work have suggested that larger defects usually results in greater adhesion formation (reviewed by (Ozel et al., 2005). Nevertheless, a general consensus is that adhesiogenesis is aggravated when suturing is performed to repair the defects (Falk et al., 1998; Hellebrekers et al., 2000). This is likely to be contributed by secondary ischaemia and foreign body reaction caused by tight suturing (Ellis, 1971; Kraemer et al., 2014). On this note, the creation of ischaemic buttons is another popular method that also encompasses a combination of injuries from ischaemia and foreign body reaction for modeling surgical trauma during abdominal or gynaecologic surgery (**Fig. 1.2c**). Ligating an out-pouching of the abdominal sidewall with a suture creates these buttons. One major advantage of this model is that the button area is well defined and easily

identified due to placement of the suture. This facilitates the study of both early and late molecular events of adhesiogenesis by allowing tracking of the trauma area.

While the above-described methods have been the main staple of adhesion research for the past few decades, a persisting struggle in the field is the lack of universally accepted models (Beyene et al., 2015). An array of different species of animals and adhesion-inducing operations exists. In addition, the lack of a standardised timing for evaluating adhesion formation and scoring systems further complicate this, making it challenging to perform direct comparison of results from different reports. Although post-surgical adhesion is a shared interest across the fields of gynecology, surgery, and peritoneal dialysis, limited communication between publication venues used for research in this area hamper forward progress (Beyene et al., 2015). These issues will be discussed in greater detail in **Chapter 4**.

Figure 1.2 Popular models of post-surgical adhesion formation.

(A) Caecal abrasion model involves abrading the anterior portion of the caecum; usually until petechial subserosal haemorrhages (blue arrowheads) develop. (B) Parietal peritoneum excision model involves making an excision through the musculo-peritoneal layer (yellow arrowhead). (C) Ischaemic button model involves ligating an out-pouching of the abdominal sidewall, usually with a suture, to create an ischaemic button (green arrowheads).

1.1.5 Adhesion-prevention strategies

Preventing the formation of post-surgical peritoneal adhesions is best strategy to reduce morbidity and its associated economic costs. Over the years, many preventions strategies have been proposed and can be grouped into these main categories: technical measures, barriers, and pharmacological agents.

1.1.5.1 Technical measures

The overarching goal in this prevention strategy is to limit surgery trauma to the peritoneal cavity as far as possible. To achieve this, it is fundamental that basic surgical principles are to be observed at all times; limiting bowel exposure and desiccation, careful hemostasis, gentle handling of tissues, and dissecting should be carried out delicately when possible (Arung et al., 2011; Moris et al., 2017). Also, it is critical to avoid the introduction of foreign bodies such as: lint from drapes or gowns, fibres from paper disposables, powders from gloves, and suture materials. When possible, suture materials that stimulate lesser tissue reactivity (i.e. polyglactin) should be chosen over fibrogenic suture materials (i.e. silk, chromic catgut) (Elkins, Stovall, Warren, Ling, & Meyer, 1987; O'Leary & Coakley, 1992).

Ten Broek and colleagues performed a meta-analysis of 17 studies in an attempt to determine surgical techniques to reduce post-surgical adhesion formation (ten Broek, Kok-Krant, Bakkum, Bleichrodt, & van Goor, 2013b). The group revealed that reduced incidence of adhesions was observed when abdominal closure with the peritoneum left open, and when laparoscopy was performed (ten Broek et al., 2013b). The former practice is thought to be due to the reduction in secondary ischaemia and necrosis due to suturing or grafting of peritoneal defects (Gomel,

Urman, & Gurgan, 1996). If suturing or grafting of peritoneal defects are performed, secondary ischaemia and necrosis predisposes the defect site to reduced fibrinolytic activity, increasing the likelihood of adhesion formation (Gomel et al., 1996). The latter practice of employing a laparoscopic approach is well validated by several studies (Burns et al., 2013; Smolarek et al., 2016; Takase-Sanchez, Brooks, Hale, & Heit, 2015). On this note, it is important to point out that heated humidified CO₂ is reported to result in lower post-operative discomfort, faster recovery, and reduced adhesion formation, when compared to cold dry CO₂ that is used for insufflation in usual laparoscopic practices (DiZerega, 2000; Ordoñez, Domínguez, Evrard, & Koninckx, 1997; Rout et al., 2002).

1.1.5.2 Barrier measures

Over the years, clinical applications of several physical barriers have been approved (Moris et al., 2017). In general, the application of physical barriers (solid and liquid) is performed to cover the injured tissues and to maintain their separation until the reepithelialisation is completed (DiZerega, 2000; Risberg, 1997). Although, liquid type barriers appear to be superior, considering their ability to cover surfaces better, the majority of these barriers failed to demonstrate significant clinical improvement, usually due to their rapid absorption (Beyene et al., 2015; Brochhausen et al., 2011). Conversely, efficacy of physical barriers is naturally limited due to their inability to cover all of the visceral and peritoneal surfaces. While studies suggest that the formation of adhesions is minimised, there is no significant clinical improvement due to complications arising from adhesions (e.g. rate of reoperations, secondary infertility, chronic abdominal pain (ten Broek et al., 2014).

Seprafilm, a chemically modified sodium hyaluronate/carbomethylcellulose absorbable barrier, is shown to reduce adhesion incidence and ASBO that requires reoperation (Fazio et al., 2006). However, overall SBO incidence remained unchanged (Robb & Mariette, 2014). If not used properly, the application of Seprafilm may increase the risk of fistula formation and anastomotic leak (Beck et al., 2003; Robb & Mariette, 2014). Other notable barriers, Gynecare Interceed and Gore Preclude, are both shown to reduce the adhesion incidence and extent (Ahmad, O'Flynn, Hindocha, & Watson, 2015; Wiseman, Trout, Franklin, & Diamond, 1999). The former is an oxidised regenerated cellulose while the latter is a polytetrafluorethylene membrane; both are similar in terms of safety and efficacy (Ahmad et al., 2015; Al-Jabri & Tulandi, 2011; Wiseman et al., 1999). However, there are certain drawbacks. In the presence of blood Interceed becomes ineffective, which may complicate or limit its application in certain clinical situations (Ahmad et al., 2015). Also, for Preclude, which is used to prevent pelvic adhesions from developing, a strip of Gore-Tex is permanently left in the pelvis potentially increasing the risk of infertility (Al-Jabri & Tulandi, 2011).

1.1.5.3 Pharmacological agents

Given the wide array of agents that have been explored to determine their efficacy in preventing adhesion formation, it is challenging to cover all of these agents (Beyene et al., 2015). In general, these agents are used in attempt to preclude the adhesion pathway early on or potentiate physiological fibrinolysis and adhesion resorption efforts (Maciver et al., 2011). An inexhaustible list includes, fibrinolytic agents, anti-coagulants, antibiotics, corticosteroids, and non-steroidal anti-

inflammatory drugs (NSAIDs) (Arung et al., 2011; Beyene et al., 2015). Below, some of these agents are reviewed, but this does not include everything.

The importance of the fibrinolytic system in the pathogenesis of adhesion formation has led much interest in the exploration of fibrinolytic drugs and anti-coagulants. While fibrinolytic agents, such as recombinant tPA, have demonstrated to reduce adhesions in animal models (Nappi et al., 2007), they are also observed to impede wound healing (Evans, McAree, Guyton, Hawkins, & Stakleff, 1993). Further limiting their acceptance for use in preventing adhesion formation, is their likelihood to induce haemorrhagic complications (Arung et al., 2011; Nappi et al., 2007). Anti-coagulants are also a favourite adhesion-reducing agent explored among investigators (Beyene et al., 2015). Heparin, one of the popular anti-coagulants explored for reducing adhesions, is shown to reduce adhesions in a rabbit model (Fukasawa, Girgis, & diZerega, 1991). However, this success is not observed in clinical trials, whether administered independently, or together with Interceed barrier (Turkcapar et al., 1995).

Considering that increased peritoneal adhesion may be borne out of increased peritoneal infection, the use of antibiotics has also been explored to reduce adhesion formation (Arung et al., 2011). On the contrary, intra-abdominal application of antibiotics and its use in irrigation solutions were shown to increase peritoneal adhesion formation (Nappi et al., 2007; Sortini et al., 2006).

Thought to reduce adhesion formation by reducing cytokines and vascular permeability, corticosteroid therapy has also yielded disappointing results.

Methylprednisolone and progesterone were both found to be ineffective in reducing the incidence of post-surgical adhesions (Confino, Friberg, Vermesh, Thomas, & Gleicher, 1988; Kirdak, Uysal, & Korun, 2008). Furthermore, although corticosteroid therapy is immunosuppressive, it may also delay wound healing (Arung et al., 2011; Risberg, 1997). Treatment of NSAIDs has been widely explored and appears to be more promising (Beyene et al., 2015). Peritoneal adhesions in some animal models are reduced through the inhibition of prostaglandin and thromboxane synthesis (Menzies & Ellis, 1990). Pre-operative or post-operative administration of nimesulide, a selective cyclooxygenase-2 inhibitor, is shown to reduce post-surgical adhesion in a rat model (Nappi et al., 2007). Although some of these anti-inflammatory drugs are shown to be promising in preventing/reducing adhesions, side effects are yet to be ascertained and current clinical evidence is not sufficient to recommend their use in humans (Arung et al., 2011; Beyene et al., 2015; Nappi et al., 2007).

In summary, there is still a lack of an ideal method or agent for preventing post-surgical peritoneal adhesion (Becker & Stucchi, 2004; ten Broek et al., 2016). Considering that post-surgical peritoneal adhesions still represent a challenging clinical problem of significant magnitude and economic burden, much more research is warranted. Particularly, research into previously unexplored areas in this complex clinical manifestation may breathe new life into existing strategies for preventing post-surgical peritoneal adhesion. Of note, little is understood about the role connexins in post-surgical peritoneal adhesions formation. Therefore, understanding how they contribute to adhesion pathogenesis may help to identify

novel avenues or bolster existing strategies for preventing post-surgical peritoneal adhesions formation.

1.2 Connexins

Connexins (Cx) are proteins that facilitate the direct cytoplasmic exchange of ions and small molecules between adjacent cells through the formation of intercellular channels, or gap junctions (Nielsen et al., 2012). Connexins have an important role in supporting the sharing of metabolic demands across cells and tissues (Evans et al., 2006). They are also multidimensional proteins that associate with cytoskeletal elements, adhesion molecules, tight junctions, adherens junctions and cellular machinery, which facilitate their function, transport, assembly, internalization and turnover (Hervé, Derangeon, Sarrouilhe, Giepmans, & Bourmeyster, 2012; Nielsen et al., 2012; Wright, Richards, & Becker, 2012).

1.2.1 Structure and nomenclature

One Cx comprises four transmembrane domain proteins with intracellular N- and C-terminus that are connected by two extracellular loops and one cytoplasmic loop (**Fig. 1.3a**) (Nielsen et al., 2012). An annular hemichannel or “connexon” is formed when six Cx units oligomerise in the endoplasmic reticulum (**Fig. 1.3b**) (Wright et al., 2012). A functional gap junction forms with the apposition of the extracellular portion of a hemichannel from adjacent cells (Wright et al., 2012). Aside from the formation of gap junctions, nonjunctional channels (hemichannels) also provide a pathway for the release of paracrine messengers (i.e. adenosine triphosphate (ATP), glutamate, nicotinamide adenine dinucleotide (NAD⁺)) from cells (Evans et al., 2006). On this note, a hemichannel can be classified either homomeric or

heteromeric depending on whether they contain one Cx type or different Cxs. Similarly, gap junctions are termed either homotypic or heterotypic depending on whether identical hemichannels form cell-cell channels (**Fig. 1.3c**). This confers further diversity in both their composition and resulting function (Nielsen et al., 2012).

Although 21 different Cxs have been identified in human and 20 in mouse (**Table 1.2**), Cxs generally share significant structure homogeneity (50-80%), particularly the TM and extracellular domains (Goodenough, Goliger, & Paul, 1996; Nielsen et al., 2012). Instead, the main structural differences are mostly found in the cytoplasmic portion, in which the length of C-terminal tail varies and is a major determinant of the molecular mass of Cx (Evans et al., 2006). In general, Cx are named according to their predicted molecular weight (e.g., Cx43 is approximately 43kDa in size). This may sometimes be confusing when homologous connexins of different species (i.e. human and mouse) have different names (i.e. molecular weight) (Nielsen et al., 2012). It is useful to point out that no homologous connexins mentioned in this thesis have differing names.

Connexins can be further classified under subgroups α , β , γ , δ , or ϵ , depending on the degree of sequence identity and cytoplasmic loop length (Nielsen et al., 2012). Following on, connexins are abbreviated with “GJ” for gap junction, and are indexed in order of their discovery (i.e. Cx43 is the first connexin of the α -group to be discovered and hence named GJ α 1) (**Table 1.2**) (Nielsen et al., 2012).

Figure 1.3 Model of a connexin monomer, hemichannel and gap junction channel.

(A) One connexin monomer contains four transmembrane domain proteins with intracellular N- and C-terminus that are connected by two extracellular loops and one cytoplasmic loop. (B) Six connexin subunits form a hemichannel. The yellow circle indicates the location of the channel pore. (C) A hemichannel can be classified either homomeric or heteromeric depending on whether they contain one Cx type or different Cxs. Similarly, gap junctions are termed either homotypic or heterotypic depending on whether identical hemichannels form cell-cell channels. Diagram adapted from Nielsen et al., 2012.

Table 1.2 Connexin family members in human and mouse

GJ	Human connexins	Mouse connexins
GJϵ1	Cx23	Cx23
GJβ7	Cx25	
GJβ2	Cx26	Cx26
GJγ3	Cx30.2	Cx29
GJβ6	Cx30	Cx30
GJβ4	Cx30.3	Cx30.3
GJβ3	Cx31	Cx31
GJβ5	Cx31.1	Cx31.1
GJδ3	Cx31.9	Cx30.2
GJβ1	Cx32	Cx32
-	-	Cx33
GJδ2	Cx36	Cx36
GJα4	Cx37	Cx37
GJα5	Cx40	Cx40
GJδ4	Cx40.1	Cx39
GJα1	Cx43	Cx43
GJγ1	Cx45	Cx45
GJα3	Cx46	Cx46
GJγ2	Cx47	Cx47
GJα8	Cx50	Cx50
GJα9	Cx59	-
GJα10	Cx62	Cx62

Adapted from Nielsen, 2012.

1.2.2 Connexin43 in post-surgical peritoneal adhesions

Connexins have an important role in supporting homeostasis and the efficient coordination of many cellular processes in the body. It does so by providing a conduit for intercellular communication, sharing the metabolic demands across cells and tissues (Evans et al., 2006). Also, Cx may also participate in extracellular communications through hemichannels via the release of paracrine messengers (Evans et al., 2006). Despite the potential involvement of connexins in various facets

of adhesion pathogenesis (i.e. inflammation, coagulation, mesenchymal transition), their contribution in it has yet to be explored (Cogliati et al., 2016; Okamoto & Suzuki, 2017; Soon et al., 2016). In particular, little is known about the contribution of Cx43 in the pathogenesis, which is also the most comprehensively studied Cx and by far the most abundant and ubiquitous isoform in mammalian cells (Nielsen et al., 2012).

Findings from different fibrosis-related diseases appear to suggest that Cx43 is essential for fibroblast proliferation, apoptosis, migration, and for regulating the production of ECM (Mendoza-Naranjo et al., 2012; Mori, Power, Wang, Martin, & Becker, 2006; Qiu et al., 2003). Furthermore, it has been shown to be able to regulate leukocyte recruitment by reducing the expression of adhesion molecules (Abed et al., 2014; Soon et al., 2016). In addition, Cx43 has been demonstrated to regulate in mesenchymal transition under inflammatory conditions (i.e. diabetes) (Abed et al., 2014; Chilton, Giles, & Smith, 2007; Uyama et al., 2003), or in the presence of TGF- β 1, a potent profibrotic mediator (Asazuma-Nakamura et al., 2009). These findings will be discussed in greater detail in **Chapter 5**. Seeing how Cx43 has a potential role in many facets of adhesion pathogenesis, the role of Cx43 in post-surgical peritoneal adhesion formation is explored in this study.

Chapter 2: Materials and Methods

2.1 Ex vivo Work

2.1.1 *Ex vivo* media preparation

To prepare 1 L of culture media, the components and supplements were combined as listed in **Appendix II, Supplementary Table 1**. This initial mixture was then adjusted to a pH of 7.1 to 7.3 before the final volume was brought up to 1 L with deionised water. The prepared media was then filter-sterilised (0.2 µm) and aliquots were stored at 4°C.

2.1.2 Tissue collection

All animal procedures were performed in accordance with the animal protocols approved by the Institutional Animal Care and Use Committee of the Animal Research Facility of Nanyang Technological University, protocol number A0372. Before euthanasia with CO₂, fur around the peritoneum was removed using an electric trimmer and depilatory cream. After euthanasia, blood was collected immediately into a 1.5 mL microfuge tube after cardiac puncture. To standardise the amount of clotted blood, 200 µL of the freshly harvested blood was straightaway pipetted into a fresh 600 µL microfuge tube and left to clot on ice. The peritoneum site was sterilised with three rounds of 7.5% (v/v) providone-iodine solution and 70% (v/v) alcohol. Whole tissue from the abdomen was harvested and full thickness peritoneal tissue strips (1.0 x 2.0 cm) comprising of the skin, underlying muscle, and peritoneum, were collected from the same animal (refer to **Fig. 2.1**) from which

blood was collected, and stored in Solution A (refer to **Appendix II, Supplementary Table 1**).

2.1.3 Induction of *ex vivo* adhesion formation and experimental design

In **Chapter 3**, a method to induce *ex vivo* adhesion formation was modified from originally described by (Saed et al., 2016) and is summarised in **Figure 2.1**. Harvested peritoneal tissue strips in Solution A were rinsed in 70% (v/v) alcohol for 3 seconds and transferred to washing Solutions B and C (refer to **Appendix II, Supplementary Table 1**), for 3 minutes each. Experimental conditions involving the presence of abrasion trauma and/or clotted blood were explored and are summarised in **Table 2.1**. Under conditions involving abrasion trauma, peritoneum on the tissue strips was abraded 20 times evenly end-to-end by passing a sterile size 10 scalpel in one direction with the sharp side down relative to the peritoneum. Care was taken to only hold one end of the strip while performing the abrasion so that additional trauma was not introduced. The edge portion of the tissue, which had been grasped by forceps, was trimmed away before culture. In those experimental conditions where the presences of clotted blood were explored, 200 µL of clotted blood was added to the abraded site. Peritoneal strips without abrasion and/or clotted blood provided controls.

Strips were folded in half with peritoneal surfaces facing each other and padded with sterile cotton gauze before being placed in a sterile tissue histology cassette. The cotton gauze sandwiched between cassette of the lid and the folded peritoneal strip provides contact pressure. The contact pressure ensured abraded surfaces remained in contact with one another without the use of sutures, which is a major change from

the published method by (Saed et al., 2016). The cassettes were then placed in a 100-mm cell culture dish, full submerged with media and incubated in a cell culture incubator at 37°C and 5% CO₂ for up to 72 hours unless otherwise mentioned. Strips were unfolded slowly and inspected thoroughly for presence of adhesion bands at 72 hours.

Figure 2.1 Major steps for inducing *ex vivo* adhesions formation

(A) Peritoneal strips were harvested and trimmed. Depending on the condition explored (refer to **Table 2**), each strip was abraded and/or had a clotted blood placed on the peritoneal surface. (B) Example of condition 1, where a blood clot is placed on an abraded portion of the strip. (C) Strips were then folded in half with peritoneal surfaces facing each other and padded with cotton gauze before being placed in a tissue histology cassette. (D) The cassettes were then placed in a 100-mm cell culture dish with media and incubated in a cell culture incubator at 37°C and 5% CO₂ for up to 72 hours.

Table 2.1 Exploring conditions for inducing *ex vivo* adhesions

Conditions	Abrasion	Blood
1	+	+
2	+	-
3	-	+
4	-	-

Conditions 1 to 4 (n=5 each, 2 experimental repeats)

2.2 In vivo Work

2.2.1 Experimental animals

All animal procedures were performed in accordance with the animal protocols approved by the Institutional Animal Care and Use Committee of the Animal Research Facility of Nanyang Technological University, protocol number A0372. Tie2-CreER^{T2} transgenic mice and Cx43^{fl/fl} transgenic mice were purchased from The European Mouse Mutant Archive and The Jackson Laboratory respectively. Mice were bred under specific pathogen-free conditions under a 12-hour light-dark cycle and were supplied standard rodent pellet diet and water *ad libitum*. For the generation of experimental tamoxifen-inducible endothelial cell-specific Cx43 transgenic mice (Tie2-CreER^{T2+/-}Cx43^{fl/fl}), Tie2-CreER^{T2} and Cx43^{fl/fl} transgenic mice were mated in-house and maintained on the C57BL/6 background. Briefly, the breeding strategy used to achieve this starts with the mating homozygous parents of both strains to obtain heterozygous genotype Tie2-CreER^{T2+/-}Cx43^{fl/-}, which was identified through genotyping. Next, mating of this genotype with the Cx43^{fl/fl} genotype allowed the generation of the Tie2-CreER^{T2+/-}Cx43^{fl/fl} genotype, which were identified through genotyping and used as experimental mice. Conversely, Tie2-CreER^{T2-/-}Cx43^{fl/fl} mice were used as their controls. The primers used for genotyping the mice are listed in **Appendix II, Supplementary Table 2**. Only age-matched (7 to 12 weeks) males were used for experiments.

2.2.2 Induction of adhesion formation

The major steps for the induction of adhesions and the different models of post-surgical peritoneal adhesions are described in the following sections and summarised in **Figure 2.2**.

2.2.2.1 Surgery preparation and entry

Prior to the procedure, fur around the incision site was removed using an electric trimmer and depilatory cream before returning the mice to their cages and allowing them to rest for at least 24 hours. Briefly, mice were anaesthetised with isoflurane and placed on a warm heating mat throughout the surgery. All surgeons used non-powdered surgical gloves during the surgery and all instruments were sterilised. Next, the midline incision site was sterilised with 7.5% (v/v) providone-iodine solution (**Fig. 2.2a**). A midline laparotomy of 4 cm was performed to facilitate the exposure of the parietal peritoneum (**Fig. 2.2b**). Depending on experiment design, the following four adhesion-inducing procedures may be employed on the mice.

2.2.2.2 Ischaemic button model of post-surgical peritoneal adhesions

Ischaemic buttons (IB) were created to induce adhesion formation as previously described by Sandoval et al., (2016) with minor modifications. Each ischaemic button was created by placing a 4/0 non-absorbable silk suture at the base of ~3 mm of the sidewall parietal peritoneum, which was grasped with a needle holder. Two ischaemic buttons, spaced at least 1 cm apart, were created on the peritoneum wall of each side of the midline (**Fig. 2.2c**).

2.2.2.3 Parietal peritoneum excision model of post-surgical peritoneal adhesions

The parietal peritoneum excision (PPE) procedure was adapted with modifications (Falk et al., 1998). After the midline laparotomy, the skin was carefully separated from the musculo-peritoneal layer to facilitate excision made on the latter layer. A defect was then created using a 6 mm biopsy punch on the left side of the midline incision. The parietal peritoneum, together with the underlying layer of muscle, was removed when the defect was created with the punch (**Fig. 2.2d**).

2.2.2.4 Small bowel loop model of post-surgical peritoneal adhesions

The small bowel loop (SBL) procedure was adapted with modifications (Andrews et al., 2012). A loop of small bowel was selected in the mid jejunal region and folded back into a juxtaposed position. A size 10 scalpel abraded the serosal surfaces of the selected bowel, which were to be placed in contact, for 20 times in the same direction, over a length of 10 mm. Two 7/0 polyglyconate sutures were later placed at each end of the abraded portion, holding the bowel in apposition (**Fig. 2.2e**).

2.2.2.5 Caecal abrasion model of post-surgical peritoneal adhesions

The caecal abrasion (CA) procedure is adapted as described by Hemadeh and colleagues (Hemadeh, Chilukuri, Bonet, Hussein, & Chaudry, 1993). Following midline laparotomy, the caecum was located and exposed. The anterior portion was abraded with a size 10 scalpel until petechial subserosal hemorrhages developed (**Fig. 2.2f**).

2.2.2.6 Surgery closure and post-recovery regime

The laparotomy site was then closed in two layers, the inner layer with simple interrupted polyglycolic acid absorbable 6.0 sutures (**Fig. 2.2g**) and the outer layer using metallic wound clips (**Fig. 2.2h**). After closure, mice were regularly monitored until euthanasia at various time points of interest (e.g. 6, 24 hours, and 7 days) post-surgery by euthanasia with CO₂. For post-recovery, the mice received buprenorphine (0.1 mL per 10 g of animal weight) into the loose skin over the interscapular region.

Figure 2.2 Major steps during various adhesion-inducing operations in mice.

(A) Disinfection of the incision site after shaving. This was followed by a (B) midline incision of the skin. (C) Ischaemic button (IB) model. Two IBs, spaced approximately 1 cm apart, were created on the peritoneum on each side of the midline incision. Each button (yellow arrowheads) was created by placing a silk ligature at the base of 3 mm of parietal tissue that is grasped with a needle holder. (D) Small bowel loop (SBL) model. A loop of small bowel was selected at middle small bowel level to be folded back into a juxtaposed position. The serosal surfaces of the selected bowel were scalpel abraded and two absorbable sutures (yellow arrowheads) were later placed at each end of the abraded portion, holding the bowel in apposition. (E) Parietal peritoneum excision (PPE) model. An excision is made through the musculo-peritoneal layer using a 6 mm biopsy punch on each side of the midline incision. (F) Caecal abrasion (CA) model. Caecum is located and exposed, before abrading the anterior portion with a scalpel blade until petechial subserosal haemorrhages (yellow arrowheads) develop. (G) Wound closure by applying simple interrupted sutures to the peritoneal musculature and (H) metallic clips to the skin.

2.2.3 Experimental Designs

2.2.3.1 Assessing various in vivo models of post-surgical peritoneal adhesions

In Chapter 4, four post-surgical peritoneal adhesions models were explored to determine which is the most ideal model for studying the role of connexin 43 in adhesion formation: the IB model, the PPE model, the SBL model, and the CA model (n=3, each). Mice were regularly monitored until euthanasia on day 7 post-surgery.

2.2.3.2 Studying the role of Cx43 during peritoneal adhesion formation

In **Chapter 5**, mouse Cx43 antisense oligodeoxynucleotides (asODN) (refer to **Appendix II, Supplementary Table 3** for sequence) was used in an IB model of peritoneal adhesion to investigate the role of Cx43 in adhesion formation. Surgery was performed as described earlier. Prior to surgery closure, each mouse received a two-formulation treatment only after all the buttons were created for that mouse. In the first formulation, 25 μ L of 300 μ M of Cx43asODN in 30% Pluronic[®] F-127 in DI H₂O was delivered to each of the 4 buttons. To ensure further coverage of the peritoneal cavity, an additional 400 μ L of 300 μ M of Cx43asODN in 20% Pluronic[®] F-127 in DI H₂O was delivered to the left and right flank of the peritoneal cavity. For control, a vehicle group received the same treatment regime with DI water instead of Cx43asODN delivered in the same respective concentrations of Pluronic[®] F-127. After the respective treatments were administered, surgery closure and post-recovery regime was performed as described earlier. Mice were regularly monitored until euthanasia at various time points of interest (e.g. 6 (n=5, each group), 24 hours

(n=5, each group), and 7 days (n=9 and 10, vehicle and treatment groups respectively)).

2.2.3.3 Studying the role of endothelial-Cx43 during peritoneal adhesion formation

In **Chapter 6**, transgenic mice were used in an IB model of peritoneal adhesion to investigate the role of endothelial Cx43 in adhesion formation. For induction of Cre-mediated DNA recombination (EC-Cx43^{-/-}), 6 weeks old Cx43^{fl/fl} mice carrying the Tie2-CreER^{T2} transgene received daily oral gavage of 4 mg tamoxifen (Sigma-Aldrich, USA) for 5 days (Anastassiadis, Glaser, Kranz, Berhardt, & Stewart, 2010; Forde, Constien, Grone, Hammerling, & Arnold, 2002) (n=6). Age matched tamoxifen-treated mice Cx43^{fl/fl} (EC-Cx43^{fl/fl}) served as a control group (n=9). All tamoxifen-treated mice were rested at least a week prior to the surgery. Mice were regularly monitored until euthanasia on day 7 post-surgery.

2.2.4 Peritoneal lavage

Prior to necropsy, 3 mL of cold phosphate-buffered saline (PBS) was injected into the peritoneal cavity and the abdomen was gently agitated for 1 min. Approximately 2 mL of the fluid was recovered. Samples with blood contamination were omitted. Cellular debris was removed by microcentrifugation at 9000 g for 15 min at 4°C, before freezing the supernatant in 250 µL aliquots at -80°C.

2.2.5 Peritoneal fluid tPA and PAI-1 total antigen ELISA

Thawed aliquots of the peritoneal lavage were analysed for the total antigen concentration of tPA and PAI-1 present, according to manufacturer's instructions

(Molecular Innovations, USA). Samples that had coefficient of variation values more than 15% were excluded. Detected concentrations were normalised using total protein content determined from a bicinchoninic acid assay performed on the same aliquot (ThermoFisher Scientific, USA).

2.2.6 Macroscopic assessment of peritoneal adhesions

An inversed U-shaped incision was made across the peritoneal cavity for necropsy and the adhesions in each animal were evaluated according to established scoring matrices. These matrices assessed peritoneal adhesions based on qualities such as, extent, involvement, strength, and severity (refer to **Table 2.2**) (Adhesion Scoring Group, 1994; Cassidy, Sherburne, Heydrick, & Stucchi, 2015; Hoffmann et al., 2009; Nair, Bhat, & Aurora, 1974). Three observers blinded to the experimental groups scored each mouse and their scores were later averaged to give a single score per mouse. Inter-observer scores did not differ more than 2. Scores are represented as a percentage of maximum score possible for each parameter (e.g. if 2 out of 4 buttons were found to have adhesions, a score of 50% will be awarded to the mouse for the extent of adhesion). After which, the mean parameter scores for each of the groups were calculated.

Table 2.2 Scoring matrix for assessing peritoneal adhesions

Variable	Score	Description
Extent (Cassidy et al., 2015)	0	Complete absence of adhesions
	1	Only 1 button has an adhesion band
	2	2 buttons have adhesion bands
	3	3 buttons have adhesion bands
	4	4 buttons have adhesion bands
Involvement (Quantity) (Nair et al., 1974)	0	Complete absence of adhesions
	1	1 adhesion band between the organs or between one organ and abdominal wall
	2	2 adhesion bands between the organs or between one organ and abdominal wall
	3	>2 adhesion bands between the organs or between one organ and abdominal wall or adhesions of intestinal loops without any adhesion to the abdominal wall. Significant adhesion.
	4	Adhesion of all viscera to the abdominal wall
Severity (Quality) (Adhesion Scoring Group, 1994)	0	Complete absence of adhesions
	1	Filmy, avascular
	2	Dense, opaque and/or vascular
	3	Very thick cohesive attachment
Strength (Hoffmann et al., 2009)	0	Complete absence of adhesions
	1	Gentle traction requiring little effort to break adhesion
	2	Blunt dissection requiring a blunt instrument to break adhesion
	3	Sharp dissection requiring a sharp instrument to break adhesion
	4	Adhesion of all viscera to the abdominal wall

2.3 Histology

2.3.1 Tissue processing and microtome sectioning

Tissue samples are collected and stored in 4% (w/v) paraformaldehyde at 4°C overnight. On the following day, the tissues were transferred into 70% (v/v) ethanol and stored at 4°C overnight. After, tissues were processed using the HistoCore PEARL (Leica, Germany) tissue processor and embedded in paraffin with the HistoCore Arcadia C (Leica, Germany). Following, tissue samples were sectioned at 5 µm thickness using a Leica RM2245 microtome (Leica, Germany) and mounted

onto Polysine[®] microscope slides and dried at 40°C for at least an hour before staining.

2.3.2 Haematoxylin and eosin (H&E) staining

Slides were loaded into Leica Autostainer XL (Leica, Germany) and a set programme was run (refer to **Appendix II, Supplementary Table 4**). At the end of the programme, glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at room temperature (RT).

2.3.3 Picro-Sirius red (PSR) staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and DI water. Slides were then immersed in Picro-Sirius red (Abcam, United Kingdom) for 1 hour. After, slides were transferred into two changes of acetic acid followed by two changes of 100% (v/v) ethanol. Slides were cleared again with two changes of Clearene (Leica, Germany) before glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at RT.

2.3.4 Immunofluorescence staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and PBS. Depending on the antibody, either heat-induced or proteolytic-induced antigen retrieval methods may be performed (refer to below sections). If no antigen retrieval was required, slides were then immersed in permeabilisation/blocking buffer (0.3% (v/v) Triton-X100, 1M Lysine in PBS) for 1 hour at RT before incubation in antibody

buffer (1% (w/v) BSA in PBS) for 45 minutes. After incubation, the boundaries of the tissue samples were dried and a hydrophobic perimeter was drawn around the sample with a hydrophobic pen (Agilent Technologies, United States). Tissue samples were then immunolabeled with primary antibody diluted in blocking solution (refer to **Table 2.3** for dilutions and incubation conditions regarding primary antibodies) and incubated overnight at 4°C. Controls without primary antibodies were performed similarly with antibody buffer in the absence of any primary antibody. After primary antibody incubation, tissues were washed twice, 3 minutes each, using washing buffer (0.05% (v/v) Tween-20 in PBS) before 1 hour incubation at RT with the appropriate secondary antibody (refer to **Table 2.4** for dilutions regarding secondary antibodies). After incubation, tissues were washed twice, 3 minutes each, with washing buffer before staining with DAPI (Life Technologies, USA) diluted at 1 in 10000 with antibody buffer. Finally, slides were washed twice, 3 minutes each, in PBS prior to applying Citiflour™ AF1 (Electron Microscopy Sciences, London) anti-fade mounting medium before mounting coverslips and sealing with nail varnish.

2.3.4.1 Heat-induced antigen retrieval (HIAR)

After rehydration, slides were submerged in Tris-EDTA antigen retrieval solution (10 mM Tris Base, 1 mM EDTA, 0.05% (v/v) Tween 20, pH 9.0) and incubated at 60°C for 30 minutes. Slides were rested in the same solution for 5 minutes at RT. After, slides were immersed in antibody buffer and the remainder of standard staining protocol was performed as described earlier.

2.3.4.2 Proteolytic-induced antigen retrieval (PIAR)

A 20X proteinase K stock solution (400 ug/ml) was prepared by dissolving 8 mg of proteinase K (Bioline, Singapore) in 50% (v/v) TE buffer (50 mM Tris Base, 1 mM EDTA, 0.5% (v/v) Triton X-100, pH 8.0), and 50% (v/v) glycerol. This stock solution was then diluted to a 1X working solution with TE Buffer. After rehydration, slides were submerged in the working solution and incubated for 5 minutes at 37°C. Following, slides were rested in the same solution for 5 minutes at RT before being washed in twice, 5 minutes each, with a proteinase K inactivating wash buffer (1 mM PMSF in PBS). After, slides were immersed in antibody buffer and the standard staining protocol was performed. Slides were washed with proteinase K inactivating wash buffer instead of PBS prior to applying the anti-fade mounting medium. The remainder of standard staining protocol was performed as described earlier.

Table 2.3 List of primary antibodies used in this study

Name	Protocol	Reactivity	Dilution	Catalogue No.
Rabbit polyclonal anti-Cx43	IHC	Cx43	1:1000	C6219 (Sigma, USA)
Rabbit polyclonal anti- α SMA	IHC	α SMA	1:500	AB5694 (Abcam, UK)
Rabbit monoclonal anti-TGF- β 1	IHC + HIAR	TGF- β 1	1:400	AB215715 (Abcam, UK)
Rat polyclonal anti-CD31	IHC + PIAR	CD31	1:25	553370 (BD Pharmingen, USA)

IHC – Immunohistochemistry staining, **HIAR** – Heat induced antigen retrieval, **PIAR** – Proteolytic-induced antigen retrieval.

Table 2.4 List of secondary antibodies used in this study

Name	Reactivity	Dilution	Catalogue No.
Goat anti-rat AF568	Rat	1:500	A11077 (ThermoFischer Scientific, USA)
Goat anti-rabbit AF488	Rabbit	1:500	A11008 (ThermoFischer Scientific, USA)
Goat anti-rabbit AF555	Rabbit	1:500	A21422 (ThermoFischer Scientific, USA)

2.4 Brightfield and confocal microscopy

Histological stained (H&E and PSR) tissue samples were automatically scanned and digitised into 24-bit images with an AxioScan.Z1 slide scanner (Zeiss, Germany) equipped with a 20X 0.8NA Plan-Apochromat air objective and a Hitachi HV-F202SCL. Scaling per pixel for these images is 0.22 μm by 0.22 μm . For the visualisation of immunohistochemical stained tissue samples, Z-stacks of 10 optical sections were acquired using an upright confocal microscope TCS SP8 (Leica, Germany) with a 40X 1.2NA Plan-Apochromat or a 63X 1.32NA oil objective. Z-stacks were taken by first finding the centre position before proceeding to take slices above and below this centre point. The thickness of each Z is 0.699 μm . Fluorophores were excited sequentially using a 405 nm, 488 nm and 532 nm wavelength laser. The images were acquired at 8-bit greyscale at 1024 x 1024 pixels with a Z-series frame average of 2 or 3. Maximum projections were calculated using Leica Application Suite X (Leica, Germany) for each of the Z-stacks recorded. All images were captured with identical parameters and camera settings to allow for direct comparison of staining intensity.

2.5 Microscopic Assessment

2.5.1 Inclusion and exclusion criteria

Tissues samples were first subjected to a simple inclusion/exclusion criterion to identify samples with features that allowed fair comparison with one another. The inclusion criterion included only tissues sections with a visible button (refer to **Fig. 2.3a**) or with a visible button accompanied with sutures only at the base of the button (refer to **Fig. 2.3b**). The exclusion criterion rejected tissues sections with no visible button (refer to **Fig. 2.3c**) or with a visible button accompanied with sutures found at other locations other than at the base of the button (refer to **Fig. 2.3d**).

Figure 2.3 Inclusion and exclusion criteria

The inclusion criterion included only tissues sections with a visible button (**A**) or with a visible button accompanied with sutures only at the base of the button (**B**). The exclusion criterion rejected tissues sections with no visible button (**C**) or with a visible button accompanied with sutures found at other locations other than at the base of the button (**D**). **IB** – Ischaemic Button region, **S** – Suture region.

2.5.2 Extent of inflammation

For scoring the extent of inflammation, each IB sample was first divided into two regions, referred to as the “Within IB” and “Outside IB” regions (refer to **Fig. 2.4**). The “Within IB” region was defined as the area spanning 500 μm away from the periphery of the button. The “Outside IB” region was defined as the area encompassing 1500 μm away from both left and right flanks of the perimeter of the “Within IB” region. In each region, seven region-of-interests (ROIs), 62,500 μm^2 each, were evenly and objectively distributed along the mesothelial edge of the tissue. Each ROI was checked to ensure that more than half of the sampled area was covered by tissue containing both the mesothelium and the underlying muscle layer. Next, three observers blinded to the experimental groups scored the extent of inflammation in each ROI on a 5-point scale based on a scoring matrix described in **Figure 2.5**. Scores from the three observers were then averaged. Scores between observers did not differ more than 1.

Figure 2.4 Designating ROIs for scoring the extent of inflammation

Seven ROIs (black squares) were taken each from “Within IB” and “Outside IB” regions. The “Within IB” region was defined as the area spanning 500 μm away from the periphery of the button (green box). The “Outside IB” region was defined as the area encompassing 1500 μm away from both left and right flanks of the perimeter of the “Within IB” region (blue box). **IB** – Ischaemic Button region, **S** – Suture region.

Figure 2.5 Scoring matrix for grading extent of inflammation

Regions-of-interests were graded for the extent of inflammation based the above scoring matrix. Three representative images accompany the description of each grade, detailed in the following.

Grade 0, little/no recruitment of leukocytes to the edge or muscle area; **Grade 1**, some recruitment of leukocytes to the edge but little/none to the muscle area; **Grade 2**, moderate recruitment of leukocytes to the edge and/or to the muscle area (no focal accumulation of leukocyte was observed); **Grade 3**, large recruitment of leukocyte to the edge and/or to muscle area (some focal accumulation of leukocytes was observed); **Grade 4**, extensive recruitment of leukocyte, which is limited to a particular region.

2.5.3 Fibrosis extent

During adhesion formation, fibrin deposits that persist allow the invasion and proliferation of fibroblasts, leading to the deposition of extracellular matrix proteins (e.g. collagen). As the adhesion develops and matures, the matrix becomes increasingly filled with more extracellular matrix proteins over time, eventually transforming the matrix into a more permanent fibrous connection. At an earlier stage (e.g. 7 days as opposed to 21 days) during adhesion development, the entire matrix may not necessarily be “filled” up with collagen, or other matrix proteins. The assessment of the extent of the fibrotic area (or matrix) is a good measure of area of the matured adhesion. In order to facilitate this analysis, several regions of

the IB sample were further identified. The entire fibrotic (EF) region was identified as the overall region that included the immediate fibrotic (IF) region and the regions of adhesion-inducing determinants, such as ischaemic button and suture (IBS) (refer to **Fig. 2.6**). The EF region was defined by region occupied by the abnormal encroaching of extracellular matrix that is observed across the entire ischaemic button tissue region and is bordered at the base of the ischaemic button by thickened mesothelium. The IBS region includes the ischaemic button region and the suture material region. The ischaemic button can be defined by the “ball of muscle” (due to the out-pouching) and can be distinguished from the matrix due to its pinkish characteristic. The suture material regions are determined by the presence of the suture material (greyish brown). The IF region is of the main interest and can be calculated by subtracting the area of the IBS region from the calculated area of the EF region. Note that these terms are also used in the following sections when determining fibrosis severity and protein levels of α SMA. Two ROIs were drawn on digital stitched images of tissue sections stained with H&E. One was drawn around the EF region and the other, around the IBS region. Areas of these regions were measured using image analysis software Zen Lite 2.3 (Zeiss, Germany) and used to calculate the area of IF region by subtracting the IBS region from EF region. Since the size of IBS region may influence the size of the IF region, the calculated area of the IF region is normalised to the area of IBS region to ensure fair comparison between the samples.

Figure 2.6. Defining regions around the ischaemic button

Representative digital stitched image of a section of the IB sample stained with H&E. The entire fibrotic (EF) region was identified as the total fibrotic region (area encompassed by red dashed lines). This is made up of the immediate fibrotic (IF) region and the regions of adhesion-inducing determinants such as the ischemic button and suture (IBS) (black dashed lines). The immediate fibrotic region (IF) may be determined by drawing ROIs to exclude the FR region from the IBS region or by the subtracting the FR region from IBS region. **B** – Ischaemic Button region, **S** – Suture region.

2.5.4 Fibrosis severity

ROIs were first drawn around the IF regions of the digital stitched images of tissue sections stained with PSR using ImageJ analysis software (version 1.51w; NIH) (**Fig. 2.7b**). These ROIs were saved for later use and documentation. Next, the amounts of collagen deposition were estimated using the greyscale threshold method described by Schipke et al., (2017) using ImageJ analysis software. Briefly, conversion of the acquired images to the RGB stacks generated greyscale images for red, green, and blue channels. Manual threshold was performed on the greyscale image corresponding to the green channel to detect collagen. Specific threshold values (80 to 255) were determined using at least 10 sections to ensure values were representative and then kept constant throughout image analysis. Next, an output mask is generated through manual thresholding at determined values (**Fig. 2.7c**).

The corresponding ROIs were loaded onto the respective output masks and the detected area was measured using the measure tool (**Fig. 2.7d**) and later normalised to the IBS area.

Figure 2.7. Example of image analysis workflow for PSR stained images.

(A) Example of PSR stained image to be analysed. (B) ROIs were first drawn around the IF regions of the digital stitched images of tissue sections stained with PSR using ImageJ. Next, the stained image to be analysed were converted to the RGB stacks, which generates greyscale images for red, green, and blue channels. Manual threshold was performed on the greyscale image corresponding to the green channel to detect collagen, which would produce an output mask. (C) Example of output mask after manual threshold (80 to 255). Specific threshold values were kept constant throughout analysis. (D) Example of the loading of corresponding ROIs for the respective sections so that the measure tool can be used to detect the area of interest in ImageJ.

2.5.5 Protein levels of α SMA

ROIs were drawn around the IF region in the digital stitched images of tissue sections immunostained with α SMA. Areas of blood vessels in which strong α SMA staining can be found, were excluded from the ROIs. Protein levels of α SMA in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the α SMA levels were determined by quantifying the area in the binary mask that is generated after performing manual threshold. Specific threshold values (80 to 255) were kept constant throughout image analysis. The detected area was measured using the measure tool and later normalised to the IBS area.

2.5.6 Protein levels of Cx43 (ischaemic buttons)

Tissue sections were immunostained with Cx43 and five fields-of-view were each acquired from regions within and away from the ischaemic button. From the acquired images, ROIs were drawn around the mesothelial and muscle regions. Protein levels of Cx43 in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the Cx43 levels were determined by quantifying the area in the binary mask generated after performing manual threshold. Specific threshold values were kept constant throughout image analysis (100 to 255). The measured Cx43 pixel area was then normalised to the pixel area of the ROI measured.

2.5.7 Protein levels of TGF- β 1 (ischaemic buttons)

Tissue sections were immunostained with TGF- β 1 and five fields-of-view were each acquired from regions within the ischaemic button. From the acquired images, ROIs were drawn around the muscle regions. Protein levels of TGF- β 1 in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the TGF- β 1 levels were determined by quantifying the area in the binary mask generated after performing manual threshold. Specific threshold values were kept constant throughout image analysis (90 to 255). The measured TGF- β 1 pixel area was then normalised to the pixel area of the ROI measured.

2.5.8 Protein levels of Cx43 (blood vessels)

Skin harvested using 6 mm sized biopsy punch were processed, sectioned and immunostained with Cx43. Five fields-of-view were each acquired from regions in

between the hypodermis and the panniculus carnosus. From the acquired images, ROIs were drawn around the blood vessels. Protein levels of Cx43 in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the Cx43 levels were determined by quantifying the area in the binary mask generated after performing manual threshold. Specific threshold values were kept constant throughout image analysis (100 to 255). The measured Cx43 pixel area was then normalised to the pixel area of the ROI measured.

2.7 Statistical Analysis

All data is presented as mean value $\pm$ standard deviation. In general, all statistical comparisons made between vehicle and Cx43asODN treatment groups or between EC-Cx43^{fl/fl} and EC-Cx43^{-/-} groups were made using a Mann-Whitney *U* test, unless mentioned otherwise. Statistical comparisons made between EC-Cx43^{fl/fl} and EC-Cx43^{-/-} for the analysis of endothelial-Cx43 knockdown in the blood vessels was performed using a Student's *t*-test. A two-way analysis of variance (ANOVA) followed by a post-hoc Tukey's multiple comparisons test was used for analysing the protein levels of Cx43 or the extent of inflammation in regions within and outside the IB regions at 6 and 24 hours post-surgery between vehicle and Cx43asODN treatment groups. All statistical comparisons were made using GraphPad Prism version 6.0f for Mac, (GraphPadTM Software, San Diego, USA).

2.8 List of materials and reagent used

Materials/Reagents	Catalogue Number, Brand
4',6-Diamindino-2-Phenylindole, Dihydrochloride (DAPI)	D1306 (ThermoFisher Scientific, USA)
4/0 non-absorbable silk sutures	C0762083 (Aesculap, USA)
7/0 absorbable polyglyconate sutures	C0022003 (Aesculap, USA)
6 mm biopsy punch	33-36 (Integra LifeSciences, USA)
Bicinchoninic acid assay, Pierce TM	23225 (ThermoFisher Scientific, USA)
CitiFluor TM AF-1 mounting medium	17970-100 (Electron Microscopy Sciences, London)
Corn Oil	C8267 (Sigma, USA)
Clearene TM	3803600 (Leica, Germany)
Eosin	3801615 (Leica, Germany)
Ethylenediamine tetraacetic acid (EDTA)	03690 (Sigma, USA)
Glycerol	G6279 (Sigma, USA)
Hematoxylin	3801570 (Leica, Germany)
Lysine	L5626 (Sigma, USA)
Non-powdered surgical gloves	330061070 (Ansell, Australia)
Organo/Limonene mount TM	08015 (Sigma, USA)
Paraformaldehyde	P6148 (Sigma, USA)
Phenylmethylsulfonyl fluoride	36978 (ThermoFisher Scientific, USA)
Phosphate-buffered saline (PBS)	BUF-2040-10X (1 st Base, Singapore)
Picro-Sirius Red	Ab150681 (Abcam, UK)
Plasminogen activator inhibitor-1 ELISA	MPAIK-TOT, L/N 916 (Molecular Innovations, USA)
Pluronic [®] F-127	P2443 (Sigma, USA)
Polysine TM microscope slides	P4981 (ThermoFisher Scientific, USA)
Proteinase K	BIO-37037 (Bioline, Singapore)
Tamoxifen	T5648 (Sigma, USA)
Tissue histology cassettes	39LC-500-1 (Leica, Germany)
Tissue plasminogen activator ELISA	MTPAKT-TOT, L/N 917 (Molecular Innovations, USA)
Trion-X100	T8787 (Sigma, USA)
Tris Base	T1503 (Sigma, USA)
Tween-20	P9416 (Sigma, USA)
Scalpel blades	BB510 #10 (Aesculap, USA)

Chapter 3: Exploration of an ex vivo model of peritoneal adhesion formation

3.1 Introduction

Despite many advances across the surgical sciences, post-surgical peritoneal adhesions still plague the outcomes of even the most experienced surgeons. In an attempt to assess the efficacy of anti-adhesion agents, various experimental models of peritoneal adhesions have been employed. However, the disease models are inherently variable and so this usually becomes a costly venture requiring large numbers of animals per test group (Wiseman, 2000). Added to this are the associated requirement for calibration of the employed model, testing of multiple treatment variants and the comparison of dose-responses, which all further increase the number of animals used and the related costs.

3.1.1 Cell culture systems

Cell culture systems may represent a cost-effective option for the initial screen for potential anti-adhesion agent discovery. These systems have the capacity for high throughput testing of multiple conditions. In addition, such systems can also shed light on how cells function and respond to stimuli. For instance, cultures of fibroblasts taken from either normal or adhesion-prone peritoneum may be useful substrates to determine the potential of various test agents in limiting, arresting, or rescuing the adhesion phenotype (Diamond et al., 2005; Saed et al., 2008; Saed & Diamond, 2004b). However, these single cell systems have their limitations. Due to their nature, these systems are isolated cultures and so do not take into account the

effects of the surrounding cells or the matrix in their physiological environment that would be found *in vivo*. This environment may sometimes have a strong influence over their response to stimuli (Yin et al., 2009) or may impact the cell's differentiation fate and function (Ravi et al., 2015). Furthermore, long-term effects of drugs are challenging to assess using cell culture based systems as cultured cell confluence often limits the duration of method. On the contrary, 3D cultures have a generally lower proliferation rates and may be better suited for this purpose (Ravi et al., 2015). In the light of these limitations, there remains a major demand for new model systems to bridge the gap between single cell culture systems and *in vivo* animal drug testing models for the assessment of potential anti-adhesions agents.

3.1.2 *Ex vivo* tissue culture systems

One solution is the use of explanted tissue cultures. Recent work by Saed and colleagues has described the first *ex vivo* rat model system of peritoneal adhesion formation (Saed et al., 2016). In this model, an abrasion injury is first performed on strips of tissue comprising peritoneum and its overlying abdominal muscle that have been harvested from rats. Following this, clotted blood from the same animal is placed on the abraded portion before the strips are folded in half, with the ends sutured to prevent unfolding of the strip. The folded strips are then cultured for up to 72 hours. Each strip usually produces a single adhesive band and their studies show it was detected as early as 24 hours (Saed et al., 2016). The adhesions generated under these experimental conditions were found to be consistent with the clinical observations of “Type 1B adhesions” (Saed et al., 2016). This subtype is classified as: “*de novo* adhesions that are produced at sites where a surgical procedure was performed and previously clear of adhesions” (Diamond, 1997). The

adhesive bands generated by the model were described to be dense, opaque, and vascularised.

This model appears encouraging, although, several variables are not standardised (e.g. the quantity of blood and the abrasion injury). There are other considerations in the model's methods that may introduce variability. For example, the suturing of the folded tissue strips to keep the ends from unfolding may introduce additional grasping and/or ischaemic injury. Furthermore, the presence of the suture may cause tissue reaction. These factors may introduce variables that may hamper the reproducibility of the model.

The size of the rat peritoneal cavity allows multiple strips to be obtained from each rat, significantly reducing the number of animals required. This allows multiple arms of a study to be tested on strips obtained from one animal. This is advantageous as it is not possible to test multiple liquids, gels, or barriers in a single animal using *in vivo* methods, due to the obvious potential crossover effects (Wiseman, 2000). Furthermore, paired dose-response comparisons and intra-animal paired comparisons for different test articles can be accomplished with the availability of multiple strips (Saed et al., 2016). Also, another advantage of the model is that it can be monitored and sampled following identification of desired events such as vascularisation or adhesion creation. This allows molecular events of adhesion formation to be characterised and the effect of test articles on these processes to be assessed. This makes the model a cost-effective option for assessing the efficacy of a wide range of possible test articles.

Adapting the existing model to be performed using mice further open up a valuable option for future researchers to use genetically modified transgenic strains. Therefore, an adapted mouse model was explored to determine its suitability for use for studying the role of Cx43 during adhesion formation.

3.2 Aims and objectives

Work in this chapter aims to explore a mouse peritoneal strip *ex vivo* adhesions model for studying the role of Cx43 during adhesion formation. This would be achieved by first determining the culture conditions required for the reproducible creation of *ex vivo* adhesions. Next, adhesion formation would be then confirmed macroscopically, and verified using microscopic assessment of histological (H&E and Picro-Sirius Red) or immunohistochemical stains (α SMA). Following on, immunohistochemical evaluation of Cx43 protein levels would determine if Cx43 is implicated in adhesion formation. Finally, the feasibility of the model for studying the *ex vivo* adhesion over time would then be assessed to determine its suitability for studying Cx43 in adhesion formation.

3.3 Materials and Methods

3.3.1 Ex vivo work

3.3.1.1 Ex vivo media preparation

To prepare 1 L of culture media, the components and supplements were combined as listed in **Appendix II, Supplementary Table 1**. This initial mixture was then adjusted to a pH of 7.1 to 7.3 before the final volume was brought up to 1 L with deionised water. The prepared media was then filter-sterilised (0.2 µm) and aliquots were stored at 4°C.

3.3.1.2 Tissue collection

Before euthanasia with CO₂, fur around the mouse peritoneum was removed using an electric trimmer and depilatory cream. After euthanasia, blood was collected immediately into a 1.5 mL microfuge tube after cardiac puncture. To standardise the amount of clotted blood, 200 µL of the freshly harvested blood was straightaway pipetted into a fresh 600 µL microfuge tube and left to clot on ice. The peritoneum site was sterilised with three rounds of 7.5% (v/v) providone-iodine solution and 70% (v/v) alcohol. Whole tissue from the abdomen was harvested and full thickness peritoneal tissue strips (1.0 x 2.0 cm) comprising of the skin, underlying muscle, and peritoneum, were collected from the same mouse (refer to **Fig. 3.1**) from which blood was collected, and stored in Solution A (refer to **Appendix II, Supplementary Table 1**).

3.3.1.3 Induction of ex vivo adhesion formation and experimental design

A method to induce *ex vivo* adhesion formation was modified from originally described by (Saed et al., 2016). This method was adapted using mice and is summarised in **Figure 3.1**. Harvested mouse peritoneal tissue strips in Solution A were rinsed in 70% (v/v) alcohol for 3 seconds and transferred to washing Solutions B and C (refer to **Appendix II, Supplementary Table 1**), for 3 minutes each. Experimental conditions involving the presence of abrasion trauma and/or clotted blood were explored and are summarised in **Table 3.1**. Under conditions involving abrasion trauma, peritoneum on the tissue strips was abraded 20 times evenly end-to-end by passing a sterile size 10 scalpel in one direction with the sharp side down relative to the peritoneum. Care was taken to only hold one end of the strip while performing the abrasion so that additional trauma was not introduced. The edge portion of the tissue, which had been grasped by forceps, was trimmed away before culture. In those experimental conditions where the presences of clotted blood were explored, 200 µL of clotted blood was added to the abraded site. Peritoneal strips without abrasion and/or clotted blood provided controls.

Strips were folded in half with peritoneal surfaces facing each other and padded with sterile cotton gauze before being placed in a sterile tissue histology cassette. The cotton gauze sandwiched between cassette of the lid and the folded peritoneal strip provides contact pressure. The contact pressure ensured abraded surfaces remained in contact with one another without the use of sutures, which is a major change from the published method by (Saed et al., 2016). The cassettes were then placed in a 100-mm cell culture dish, full submerged with media and incubated in a cell culture incubator at 37°C and 5% CO₂ for up to 72 hours unless otherwise mentioned. Strips

were unfolded slowly and inspected thoroughly for presence of adhesion bands at 72 hours.

Figure 3.1 Major steps for inducing *ex vivo* adhesions formation

(A) Peritoneal strips were harvested and trimmed. Depending on the condition explored (refer to **Table 2**), each strip was abraded and/or had a clotted blood placed on the peritoneal surface. (B) Example of condition 1, where a blood clot is placed on an abraded portion of the strip. (C) Strips were then folded in half with peritoneal surfaces facing each other and padded with cotton gauze before being placed in a tissue histology cassette. (D) The cassettes were then placed in a 100-mm cell culture dish with media and incubated in a cell culture incubator at 37°C and 5% CO₂ for up to 72 hours.

Table 3.1 Exploring conditions for inducing *ex vivo* adhesions

Conditions	Abrasion	Blood
1	+	+
2	+	-
3	-	+
4	-	-

Conditions 1 to 4 (n=5 each, 2 experimental repeats)

3.3.2 Histology

3.3.2.1 Tissue processing and microtome sectioning

Tissue samples are collected and stored in 4% (w/v) paraformaldehyde at 4°C overnight. On the following day, the tissues were transferred into 70% (v/v) ethanol

and stored at 4°C overnight. After, tissues were processed using the HistoCore PEARL (Leica, Germany) tissue processor and embedded in paraffin with the HistoCore Arcadia C (Leica, Germany). Following, tissue samples were sectioned at 5 µm thickness using a Leica RM2245 microtome (Leica, Germany) and mounted onto Polysine® microscope slides and dried at 40°C for at least an hour before staining.

3.3.2.2 Haematoxylin and eosin (H&E) staining

Slides were loaded into Leica Autostainer XL (Leica, Germany) and a set programme was run (refer to **Appendix II, Supplementary Table 4**). At the end of the programme, glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at room temperature (RT).

3.3.2.3 Picro-Sirius red (PSR) staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and DI water. Slides were then immersed in Picro-Sirius red (Abcam, United Kingdom) for 1 hour. After, slides were transferred into two changes of acetic acid followed by two changes of 100% (v/v) ethanol. Slides were cleared again with two changes of Clearene (Leica, Germany) before glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at RT.

3.3.2.4 Immunofluorescence staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and PBS. Depending

on the antibody, either heat-induced or proteolytic-induced antigen retrieval methods may be performed (refer to below sections). If no antigen retrieval was required, slides were then immersed in permeabilisation/blocking buffer (0.3% (v/v) Triton-X100, 1M Lysine in PBS) for 1 hour at RT before incubation in antibody buffer (1% (w/v) BSA in PBS) for 45 minutes. After incubation, the boundaries of the tissue samples were dried and a hydrophobic perimeter was drawn around the sample with a hydrophobic pen (Agilent Technologies, United States). Tissue samples were then immunolabeled with primary antibody diluted in blocking solution (refer to **Table 3.2** for dilutions and incubation conditions regarding primary antibodies) and incubated overnight at 4°C. Controls without primary antibodies were performed similarly with antibody buffer in the absence of any primary antibody. After primary antibody incubation, tissues were washed twice, 3 minutes each, using washing buffer (0.05% (v/v) Tween-20 in PBS) before 1 hour incubation at RT with the appropriate secondary antibody (refer to **Table 3.3** for dilutions regarding secondary antibodies). After incubation, tissues were washed twice, 3 minutes each, with washing buffer before staining with DAPI (Life Technologies, USA) diluted at 1 in 10000 with antibody buffer. Finally, slides were washed twice, 3 minutes each, in PBS prior to applying Citiflour™ AF1 (Electron Microscopy Sciences, London) anti-fade mounting medium before mounting coverslips and sealing with nail varnish.

Table 3.2 List of primary antibodies used in this study

Name	Protocol	Reactivity	Dilution	Catalogue No.
Rabbit polyclonal anti-Cx43	IHC	Cx43	1:1000	C6219 (Sigma, USA)
Rabbit polyclonal anti- α SMA	IHC	α SMA	1:500	AB5694 (Abcam, UK)

IHC – Immunohistochemistry staining

Table 3.3 List of secondary antibodies used in this study

Name	Reactivity	Dilution	Catalogue No.
Goat anti-rabbit AF488	Rabbit	1:500	A11008 (ThermoFischer Scientific, USA)
Goat anti-rabbit AF555	Rabbit	1:500	A21422 (ThermoFischer Scientific, USA)

3.3.3 Brightfield and confocal microscopy

Histological stained (H&E and PSR) tissue samples were automatically scanned and digitised into 24-bit images with an AxioScan.Z1 slide scanner (Zeiss, Germany) equipped with a 20X 0.8NA Plan-Apochromat air objective and a Hitachi HV-F202SCL. Scaling per pixel for these images is 0.22 μ m by 0.22 μ m. For the visualisation of immunohistochemical stained tissue samples, Z-stacks of 10 optical sections were acquired using an upright confocal microscope TCS SP8 (Leica, Germany) with a 40X 1.2NA Plan-Apochromat or a 63X 1.32NA oil objective. Fluorophores were excited sequentially using a 405 nm, 488 nm and 532 nm wavelength laser. The images were acquired at 8-bit greyscale at 1024 x 1024 pixels with a Z-series frame average of 2 or 3. Maximum projections were calculated using Leica Application Suite X (Leica, Germany) for each of the Z-stacks recorded. All images were captured with identical parameters and camera settings to allow for direct comparison of staining intensity.

3.4 Results

3.4.1 Development of *ex vivo* adhesions required the presence of abrasion, clotted blood, and contact.

To explore a mouse peritoneal strip *ex vivo* adhesions model for studying the role of Cx43 during adhesion formation, the culture conditions required for the reproducible creation of *ex vivo* adhesions was first determined. Culturing of rat peritoneal strips in which clotted blood is placed between abraded peritoneal surfaces was previously demonstrated to produce a single adhesive band (Saed et al., 2016). One of the major modifications made to this published method in this study was the omission of sutures to prevent the unfolding of tissue strips. Instead, the half-folded mouse peritoneal strips were placed in a sterile tissue histology cassette. To refine this modification, the importance of contact between the mouse peritoneal surfaces in the development of *ex vivo* adhesions was explored in a pilot. Contact pressure between the strips was achieved by padding the strips with sterile cotton gauze after placing them in the sterile tissue histology cassette.

In the pilot run, peritoneal strips with clotted blood was placed between abraded peritoneal surfaces were cultured in the presence and absence of the gauze (n=7, each). Adhesions were found in all of the strips cultured in the presence of the gauze. In contrast, only 57% (4 out of 7) of the strips that were cultured in the absence of gauze resulted in adhesions. This concluded that contact pressure between the peritoneal surfaces was important for the development of *ex vivo* adhesions. All the strips mentioned later in this study were also cultured in the presence of the gauze.

Next, four conditions (n=5, each) were explored to determine which was most ideal for producing the most reliable adhesions in a mouse *ex vivo* adhesion model (summarised in **Table 3.1**). In conditions with no abrasions, no adhesions were observed in strips that were cultured in the presence (condition 3) or absence (condition 4) of clotted blood. In contrast, in conditions with abrasion, adhesions were only detected if the abraded peritoneal strips were cultured with clotted blood placed in between the peritoneal surfaces (condition 1) (**Fig. 3.2**). Under this ideal model condition, at least one adhesion band was always detected per strip macroscopically, although the total number of adhesion bands varied among peritoneal strips. It was found that the detected bands showed variation in strength and position along the abraded portion (**Fig. 3.2**). Overall, these observations suggests that the combination of abrasion trauma, presence of clotted blood, and contact between peritoneal surface, is essential for adhesion development in this mouse *ex vivo* adhesion model.

Preliminary microscopic assessment of adhesion formation in the abraded peritoneal strips cultured with clotted blood (condition 1, n=5), which were conditions that were found to be key for adhesion development, was performed using hematoxylin and eosin (H&E) histological stains. Eosinophilic matrices were observed between certain regions of juxtaposing mesothelium along the strip (**Fig. 3.3ci and ii**). Further assessment found these matrices could be positively stained with Picro-Sirius Red (PSR), indicating collagen deposition in these matrices (**Fig. 3.3di and ii**). In addition, surrounding mesothelia peripheral to the matrices expressed high levels of α SMA, and suggest mesothelial-to-mesenchymal transition and sub-mesothelial fibroblast activation (Sandoval et al., 2016) (**Fig. 3.3ei and ii**). These

microscopic evidences confirm the presence of adhesions and corroborate initial macroscopic observations where adhesions are only observed in certain regions. Interestingly, Cx43 protein levels in mesothelial cells and sub-mesothelial cells appeared to be higher in the regions where adhesions were observed (**Fig. 3.3fi and ii**).

Figure 3.2 Examples of *ex vivo* adhesions formed in peritoneal strips cultured under condition 1

Peritoneal strips cultured under different conditions were inspected for the presence of adhesion bands at 72 h post-induction of *ex vivo* adhesion formation. Cultures of peritoneal strips under condition 1 (n=5, 2 experimental repeats), in which clotted blood was placed between abraded surfaces, resulted in the reproducible development of adhesion bands (green arrowheads). Examples of peritoneal strips that were observed to have one (**A**) or two (**B**) thin filmy adhesion bands that may be broken by gentle traction. (**C**) Example of a peritoneal strip observed to have a thick adhesion band requiring blunt dissection to be broken.

Figure 3.3 Microscopic assessment of peritoneal strips that were cultured with a placement of blood clot between abraded surfaces.

(A) Schematic diagram differentiating the different layers of the cultured peritoneal strip. Boxed region illustrates an example of region of interest that focuses on the juxtaposing mesothelium and underlying muscle. (B) Representative stitched image of H&E-stained peritoneal strip. Five random regions (as indicated by black boxes) along a representative strip incubated under condition 1 (n=5) were selected and assessed with different histological (H&E and PSR) or immunohistochemical stains (α SMA and Cx43). (C) Some eosinophilic matrix (dotted lines) was only observed in certain regions (i and ii) between juxtaposing mesothelium along the strip. These regions were likely to be areas where adhesions have developed as indicated from positive staining of PSR (D) and α SMA

(green) (E). These regions were also found to have higher Cx43 protein levels (red) (F). Magnification used, 40X. Scale bar at 50 μ m.

3.4.2 Comparison of peritoneal strips cultured under different conditions

Histological differences between peritoneal strips cultured under the different conditions (n=5, each) were further investigated (**Fig. 3.4**). In condition 1, where peritoneal strips had placement of clotted blood between abraded surfaces, areas of adhesions were identified in certain regions of juxtaposed mesothelium as described earlier (**Fig. 3.4ai and ii**). In addition, the mesothelium appeared to be continuous along the strip, suggesting a certain degree of healing from abrasion trauma (**Fig. 3.4a**). However, the thickness of the mesothelium was varied across regions along the strip (**Fig. 3.4aiii**). In contrast with condition 2 (**Fig. 3.4b**), where peritoneal strips were abraded without the placement of clotted blood, there were regions of mesothelium still discontinuous or had poor integrity, even though there were also regions of thickened mesothelium along the strip. In condition 3 (**Fig. 3.4c**), where clotted blood was placed in the strip without abrasion, the mesothelium was continuous but remained mostly thin, with the exception of a few regions where the mesothelium appeared slightly thicker. Finally, in condition 4 (**Fig. 3.4d**), where both the placement of clotted blood and abrasion were absent, a thin continuous mesothelium was observed across the entire strip. These observations appear to suggest that condition 1 was the most ideal in generating *ex vivo* adhesions.

Figure 3.4 H&E assessment of cultured peritoneal strips cultured under different conditions

Three representative regions were selected for each peritoneal strip cultured under different conditions (n=5, each). (A) Under **condition 1**, where abrasion and clotted blood were present, the mesothelium appeared to be mostly continuous. Some regions along the strip saw the thickening of mesothelium and in certain regions of juxtaposed mesothelium, areas of adhesions were observed (marked with dotted lines in **i** and **ii**). (B) Under **condition 2**, where abrasion was present but clotted blood was absent, some regions of mesothelium with poor integrity (**i**) or discontinuous mesothelium (**ii** and **iii**) were observed along the strip. (C) Under **condition 3**, where abrasion is absent but clotted blood was present, thin mesothelium was observed in most of the regions (**i** and **iii**) except a few regions where mesothelium was thicker (**ii**). (D) Under **condition 4**, where both abrasion and clotted blood were absent, thin continuous mesothelium was observed across the entire strip (**i** to **iii**). Scale bar at 50 µm.

3.4.3 Development of *ex vivo* adhesions over time

Cultures of peritoneal strips with placement of clotted blood between abraded surfaces were harvested at various time points (n=5, per time point) to get a snapshot of the development of *ex vivo* adhesions at different stages. In general, the discontinuous mesothelium caused by the abrasion trauma appeared to heal over time, becoming continuous by 72 hours.

The mesothelium at 0 and 6 hours were not significantly different, being thin and discontinuous along the entire strip (**Fig. 3.5a and b**). At 0 hours, fibroblasts did not appear to be activated, as indicated by little/no α SMA staining on the mesothelium, but more staining was detected at 6 hours. Cx43 protein levels in these regions appeared to follow this pattern as well. At 24 hours (**Fig. 3.5c**), the blood clot seemed to have started lysing. Although discontinuous mesothelium was still observed, there were certain regions of the mesothelium that appeared to become slightly thicker than others. These regions stained more intensely for PSR with little α SMA and Cx43 staining.

At 48 hours (**Fig. 3.5d**), discontinuous mesothelium was less frequently observed, and mesothelium also appeared thicker in general. These observations suggest some degree of healing may have taken place. Regions of mesothelium stained densely with PSR also appeared to express higher levels of α SMA and Cx43. At 72 hours (**Fig. 3.5e**), as mentioned earlier in Figure 3, adhesions were observed between some regions of juxtaposing mesothelium along the strip. These adhesions stained positively for PSR and were surrounded by mesothelium expressing high levels of α SMA and Cx43.

Figure 3.5 Microscopic assessment of peritoneal strips cultured under condition 1 over 72 hours
 Abraded peritoneal strips cultured in the presence of blood (condition 1), resulting in the reproducible development of adhesion bands, were harvested at different time points. The strips of each time point (n=5) were assessed with different histological (H&E (**A**) and PSR (**B**)) or immunohistochemical stains (α SMA, in green (**C**) and Cx43, in red (**D**)). One representative region along the strip was selected to provide a general overview of the development over time. Dotted lines indicate areas where adhesions were detected. Magnification used, 40X. Scale bar at 50 μ m.

3.5 Discussion

3.5.1. Condition for creating adhesions required the presence of abrasion, clotted blood, and contact and its relevance to the human system.

Previously, (Saed et al., 2016) described a rat peritoneal strip *ex vivo* adhesion model. In the described rat model, obvious visible adhesive bands can be detected within 24 hours when abraded peritoneal strips were cultured in the presence of clotted blood.

Here, I have now described a new mouse peritoneal strip *ex vivo* adhesions model, which has refined the experimental conditions and will produce adhesions reproducibly. The key features of this model are the absence of sutures, which were previously used to prevent the unfolding of tissue strips, and the use of a standardised volume of clotted blood. Using this model, cultures of peritoneal strips with placement of clotted blood between abraded surfaces were also found to reproducibly develop adhesion bands at 72 hours. This corroborates with previous findings by Saed et al., (2016), which found that abrasion trauma and the presence of clotted blood are both important ingredients for the development of adhesion. Building on this, additional microscopic assessment using different histological stains (H&E and PSR) and immunohistochemical stains (α SMA and Cx43), were also performed and were shown to support macroscopic findings.

In addition to these findings, this study also found that contact between the peritoneal surfaces was important for the development of *ex vivo* adhesions. In the mouse *ex vivo* model, peritoneal strips were padded with sterile cotton gauze to

increase of contact between the opposing parietal surfaces. This simulated the approximation of two injured surfaces commonly performed in post-surgical in vivo adhesion experimental models (Bakkum et al., 1995; Poehnert et al., 2015; Rout et al., 2002; Wei et al., 2016a). Adhesions were found in all of the strips cultured in the presence of the gauze. In contrast, only 4 out of 7 strips resulted in adhesions when cultured in the absence of the gauze. This phenomenon shares clinical relevance, in which the importance of minimising contact between the two opposing traumatised regions is being highlighted and is also the thought design of clinical barrier agents targeting adhesion formation (Beyene et al., 2015).

In both models, adhesive bands were found reproducibly in cultures of peritoneal strips in which clotted blood was placed between abraded surfaces (condition 1). The adhesions generated under this condition were clinically relevant to the human system as they were consistent with the clinical observations of “Type 1B adhesions” (Diamond, 1997). The clotted blood simulates bleeding that may occur during surgery while the abraded surfaces simulate tissue trauma resulting from manipulations during surgical procedures. In both studies, no adhesive bands were observed when abrasion injury is absent. These conditions may suggest that abrasion trauma appears to be the main contributor to the model. However, in this study, no adhesions were formed in abraded strips cultured in the absence of blood (condition 2). This suggested that blood is also an essential ingredient for the development of adhesion. Fibrin from the bleed facilitates the initial adhesion of two surfaces during early adhesiogenesis (Wiseman, 2000). This observation is also consistent with clinical teaching, where it is important in achieving haemostasis to minimise adhesion development (Gonzalez-Quintero & Cruz-Pachano, 2009).

Although no adhesions were detected in abraded mice peritoneal strips cultured in the absence of blood after 72 hours, adhesions were described to be detected as early as 48 hours in the rat model in strips cultured under the same conditions (Saed et al., 2016). This difference of outcomes with the mouse model appears to suggest that rat tissues may have a higher propensity to form adhesions. Differences observed between the mouse and rat models in the adhesions developed from abraded strips cultured in the presence of clotted blood (condition 1) may further support this. Under condition 1, the adhesive bands generated in the new mouse model were observed only after being in culture for 72 hours and the majority of these adhesions were thin and filmy, requiring gentle traction to disrupt. In contrast, the adhesive bands described to be generated by the rat model, were observed as early as 24 hours and noted to be obvious, “dense and opaque” with appreciable vascularity (Saed et al., 2016). This may be due to interspecies differences in fibrinolytic activity (Albrechtsen, 1957; Smokovitis & Astrup, 1980; Wiseman, 2000).

3.5.2. Refinements for the mouse model

Previously, there was no standardisation of the amount of clotted blood placed between the strips in the rat model (Saed et al., 2016). This introduces potential variations as the presence of blood facilitates the initial adhesion between the two abraded surfaces by providing the model with a source of fibrin and immune cells (Ozel et al., 2005). Therefore, its volume was standardised to 200 μ L in this study. It is useful to point out that larger volumes of clot did not appear to increase the quality of adhesion created (pilot, data not shown).

In contrast to the rat model, no sutures were used in the mouse model to hold the strips together to prevent the abraded portions from unfolding while in culture. Suturing of numerous strips was performed in pilot tests and was found to be a time-consuming procedure that became logistically tedious. It may also give rise to contamination issues due to laxity in aseptic technique due to suture fatigue. Furthermore, suturing introduces additional grasping damage to the tissue and the presence of suture material may introduce a foreign body tissue reaction. These factors may introduce additional variables to the model, making it challenging to produce consistent and reproducible adhesions.

Therefore, the model was refined with the use of a tissue histology cassette to prevent the abraded portions from unfolding while in culture. The cassette also helps to keep the strips in place during culture and can be pre-labeled, helping with identification of the culture mediums. Folded peritoneal strips were padded with sterile cotton gauze to increase contact between the opposing parietal surfaces. By increasing the positional stability, not only the location and area of adhesion creation is standardised, but also the baseline incidence and extent of adhesions is increased (Wiseman, 2000). The move to a mouse model also has another advantage for researchers as genetically modified transgenic strains and commercial immunological reagents and/or assay kits designed for murine systems are widely available. This makes the mice model an attractive option.

3.5.3. Microscopic assessments

Preliminary microscopic assessment of adhesion formation was performed on strips in which adhesive bands were found reproducibly. The majority of the adhesive

bands were thin and filmy, which lysed upon unfolding of the strip. In order to preserve the adhesions in their physiological state, strips designated for microscopic assessment were not inspected macroscopically. Instead, the intact strips were tissue processed and paraffin embedded *en bloc* for downstream histological assessment, in which 5 tissue sections were sampled from 5 positions that were each 500 μm apart on the peritoneal strip. It was challenging to pinpoint the exact location of the adhesion at 72 hours, let alone at earlier time points. The presence of adhesions was identified by the presence of eosinophilic matrices between juxtaposing mesothelium. Collagen deposition was inferred from the positive staining of PSR in these matrices (Gao et al., 2017). In addition, surrounding mesothelium peripheral to the matrices expressed high levels of αSMA . This may suggest mesothelial-to-mesenchymal transition (Sandoval et al., 2016) and sub-mesothelial fibroblast activation to the adhesion phenotype (Saed & Diamond, 2004b) may have occurred in these areas. The presence of these markers confirmed the presence of adhesions. Regions where adhesions were not observed did not stain positively for these markers. Use of these markers will be useful for further assessing the efficacy of test articles in the future. Interestingly, Cx43 protein levels were also observed to be higher in regions where adhesions were present and may serve as a first line of evidence that Cx43 is implicated during adhesion formation.

In general, H&E assessment also showed that the mesothelial layer increased over time, similar to that described in a model of post-surgical adhesions (Bakkum & Trimpos-Kemper, 1995). This thickening of mesothelium is also noted in patients as well (Arung et al., 2011; diZerega & Campeau, 2001; Mutsaers et al., 2015). Conversely, conditions (2 to 4) that did not result in adhesions had mesothelium that

remained mostly thin. Thin mesothelium was also observed at early time points. The discontinuous nature of the mesothelium due to abrasion was observed less over time, suggesting that some degree of healing occurred. Unlike normal inflammatory events, little or almost no infiltration of immune cells was observed during the early time points, due to the nature of the *ex vivo* model. This may have limited the magnitude of the adhesion created (Wiseman, 2000). Unfortunately, it is difficult to make further meaningful conclusions because it is challenging to locate the exact point of adhesion creation during early culture.

3.5.4. Model suitability for studying the role of Cx43 during adhesion formation

In terms of cost effectiveness, the rat model still triumphs the mouse model. Due to the larger size of the rat peritoneal cavity, approximately 6 strips may be obtained from each rat. In comparison, at least 3 strips may be obtained from each mouse. Nevertheless, this still allows multiple arms of a study to be tested on strips obtained from one animal. Furthermore, paired dose-response comparisons and intra-animal paired comparisons for different test articles can be accomplished with the availability of multiple strips (Saed et al., 2016). The refined mouse model would be good at sorting out potential interventional options. For an example, it may be used to discern differences in the degree and duration of Cx43 knockdown between application of various doses of Cx43asODN and/or release profiles of potential sustained-release formulations. These variations may be compared on multiple unabraded strips for quantifying Cx43 expression at various timepoints of interest. Of course, due to size limitations of the mouse peritoneum, testing of agents may be more limited when compared to the rat model. However, the researchers using the

mice model will have access to genetically modified transgenic strains, which makes the mice model still an attractive option.

A major limitation of the *ex vivo* models is the lack of circulation. One of the early inflammatory responses to the inflicted injury of the peritoneum is characterised with the infiltration of the immune cells to the site of injury (Bakkum & Trimbos-Kemper, 1995; Beyene et al., 2015). Following, both resident and infiltrating cells release a cocktail of inflammatory mediators (Beyene et al., 2015). Many of these mediators have been shown to play a major role in subsequent adhesion formation (Maciver et al., 2011). For instance, one of earliest cytokines present in the injured peritoneal cavity is IL-1, which causes human peritoneal mesothelial cells to proliferate (Stadlmann et al., 2006). However, due to the nature of the model, only tissue resident immune cells and those that are found in the clotted blood were present. Therefore, this places a limited contribution from the presence of inflammation to the model, even though *ex vivo* adhesions can be generated with this model. Considering this, interpretations from either model need to be made carefully.

The mouse model is capable of generating multiple adhesive bands reproducibly in a strip. Therefore, the incidence of adhesions may be reported when assessing the efficacy of test articles in the mouse model, as adhesion-free outcome is a clinical meaningful term (Wiseman, 2000). However, the majority of the created adhesions in the mouse model were thin and filmy. This may make the model too “permissive”, such that all test articles become effective. On the contrary, the adhesive bands generated by the rat model were described to be dense, opaque, and vascularised.

More work has to be performed to explore ways to increase quality and quantity adhesion from this model. Although the number of times of abrasion was standardised, it is difficult to optimally apply consistent pressure during scalpel abrasion over the area due to the flaccidity of the strip. This may account for the observed heterogeneity of adhesive bands created across the strip. Consequently, this makes it challenging to pinpoint the exact point of adhesions creation, making the study of early molecular changes difficult. In addition, this makes microscopic assessments of adhesions for screening the efficacy of potential interventional options for reducing *ex vivo* adhesions challenging.

While it is tempting, Cx43asODN was not explored in the *ex vivo* model to identify specific cell populations that would have played a role in the Cx43-mediated adhesion formation. Such endeavour, regardless of *ex vivo* or *in vivo* systems, would require the fulfilment of at least three criteria: 1) observations of pathological changes of Cx43 in the investigated model; 2) a successful knockdown of Cx43 in cell population(s); 3) a significant biological outcome upon the treatment of Cx43asODN treatment (e.g. decrease in adhesions quantity or quality that is observed post-treatment). For the first criteria, it is challenging as discussed earlier, to pinpoint the exact point of adhesion creation in the *ex vivo* model and it is therefore difficult to pinpoint areas of pathological changes of Cx43 with certainty. For the second criteria, while knockdown of Cx43 in cell populations may be performed on unabraded strips (without adhesion creation), one would still require to optimise dosage and window of assessment, which would still have to be repeated in an *in vivo* system due to the differences in pharmacokinetics and the lack of circulation. Furthermore, it is likely that an *ex vivo* system does not contain the full

immune repertoire, and there is possibility that cell populations playing important roles in Cx43-mediated adhesion would be absent or present in limited quantities. For the third criteria, the majority of the created adhesions in the *ex vivo* mouse model is thin and filmy and may render the model too “permissive” when assessed macroscopically, such that all test articles become effective. More sensitive microscopic assessments may be performed but it is, again, difficult to pinpoint the exact point of adhesion creation. In addition, to further ensure that results generated from the treatment of Cx43asODN from the *ex vivo* model are not false positives, similar exploration must be repeated in an *in vivo* model. Considering these reasons, Cx43asODN was not explored in the *ex vivo* model for the identification of cell populations that may play a role in the Cx43-mediated adhesion formation. Instead, *in vivo* models of adhesion formation may be better suited for this purpose.

3.6 Conclusion

In this chapter, a new mouse peritoneal strip *ex vivo* adhesions model is established, which has refined the experimental conditions and will produce adhesions reproducibly. The key features of this model are the absence of sutures, which were previously used to prevent the unfolding of tissue strips, and the use of a standardised volume of clotted blood. Using this model, cultures of peritoneal strips with placement of clotted blood between abraded surfaces were found to reproducibly develop adhesion bands at 72 hours. Formed adhesions were identified macroscopically and presence of adhesions was identified by the presence of eosinophilic matrices between juxtaposing mesothelium and from the positive staining of collagen fibres stained red with PSR in these matrices. In addition, surrounding mesothelium peripheral to the matrices expressed high levels of α SMA,

suggesting mesothelial-to-mesenchymal transition and sub-mesothelial fibroblast activation to the adhesion phenotype. This same region was found to be elevated in Cx43 protein levels, serving as a first line of evidence that Cx43 may be implicated during adhesion formation. Unfortunately, it is challenging to track the exact point of adhesion creation during early culture (i.e. first 24 hours), limiting this model to study only study later molecular events or for assessing efficacy of therapeutics. Furthermore, the model lacks circulation and the contribution of the immune component to the overall adhesion formation. In light of these limitations, we turn our attention to *in vivo* models of adhesion formation. Attempts to identify the model that is most ideal for studying the role of Cx43 during adhesion formation is detailed in the next chapter.

Chapter 4: In search of an *in vivo* model of post-surgical peritoneal adhesion formation

4.1 Introduction

A better understanding of the fundamental processes involved in the complex pathogenesis of adhesion formation is essential for identifying new therapeutic targets. Many post-surgical adhesion animal models have been developed and refined over the years in an attempt to achieve this goal. In general, most animal models of post-surgical adhesions share a similar design; a selected animal species is first prepared for aseptic surgery and the designated organ of interest is subjected to a chosen adhesiogenic stimulus. Following that stage, a test article may be administered to some animals depending on experimental design. Post-recovery periods are usually allowed up to about one week before the developed adhesions are assessed by quantitative and/or qualitative criteria using gross macroscopic and/or histologic methods (Wiseman, 2000).

With many post-surgical adhesion animal models to decide from, a frequently asked question is “Which animal model is the gold standard for studying adhesions?” Unfortunately, there is no universal acceptance of one model (Beyene et al., 2015), and which model is superior is often debated (Kraemer et al., 2014; Whang et al., 2011). Furthermore, poor description and variations in surgical techniques, differences in assessment timings of adhesion formation and their respective scoring systems (Beyene et al., 2015; Ozel et al., 2005) make it challenging to directly

compare results reported in the literature. These factors collectively have hampered progress in the field.

In this chapter, the most common models of experimental adhesion formation are explored with respect to which is the most ideal for studying peritoneal adhesion formation. In order to do this, a systematic literature search was first conducted, and then experimental work was performed to identify the post-surgical peritoneal adhesion model that is most suitable for studying the role of Cx43 in post-surgical peritoneal adhesion.

The PubMed-NCBI database was searched between 1976 and 2016 to collect information on the different types of peritoneal adhesion post-surgical models performed in literature. This was conducted using the following key words: “peritoneal/peritoneum adhesion models”, yielding a total of 522 entries, of which only full-text English articles were identified. The abstracts were reviewed by this author and only experimental studies containing live animal peritoneal adhesion models were considered (reviews, articles with poorly described methods or other forms of adhesion were excluded). This left 106 articles as the source material for this literature review (**Fig. 4.1**).

Figure 4.1 PRISMA flowchat of literature search strategy

4.1.1 Selection of species

A plethora of animals has been employed to study adhesions, ranging from small animals (mice, rats, and hamsters) to larger animals (rabbits, dogs, pigs, horses, primates) (Beyene et al., 2015; Wiseman, 2000). The selection of species is clearly dependent on a number of factors but primarily these are the purpose of the study and cost. With respect to “purpose”, many models share features of providing tissue and time points. One of the less appreciated considerations can be model geometrical constraint. For example, when Interceed, an adhesion barrier, was investigated in mice, both injury in the peritoneum and increased formation of adhesions was observed (Haney & Doty, 1992a, 1992b, 1996). In contrast, the barrier was demonstrated to be beneficial in humans (Wiseman et al., 1999). This contrast in performance observed is due to difference in volume and buffer capacity of the peritoneal fluid in mouse and man. Specifically, the volume and buffer capacity of the peritoneal fluid in mouse was inadequate to neutralize the acidity of the barrier material of 1-2 cm², which would be comparable to almost 70 times of that used in humans on a per weight basis (Wiseman, 2000). Furthermore, significant remnants

of barrier material were noted, suggesting that phagocytic and hydrolytic processes were inadequate to handle such an amount of material in the mouse. In a similar theme, pharmacokinetic and pharmacodynamics differences also exist between species, which are important considerations for testing therapeutics (reviewed in detail by Wiseman (2000)).

Ultimately, restrictions in costs and logistics favour the use of smaller animals in most settings. Interestingly, while a handful of studies employ the use of mice, more investigators appear to favour the use of rats to develop their techniques for inducing adhesions (Gaertner, Hagerman, Felemovicius, Bonsack, & Delaney, 2008; Kraemer et al., 2014; Rajab et al., 2010; Whang et al., 2011). This has been speculated to be due to the perceived difficulty of creating adhesions in mice (Marshall et al., 2016). Nevertheless, they offer some definite advantages in some areas over rats as for example, genetically modified transgenic strains and commercial immunological reagents and/or assay kits designed for murine systems are widely available. This makes mice an attractive cost-effective way to explore the importance of particular gene products in the pathogenesis of adhesion formation. Since the majority of studies employ their mice and rats, my discussion in these remainder of the literature review focuses mainly based on findings from these species and this information is summarised in **Table 4.1**.

Table 4.1 Overview of rats and mice peritoneal adhesions models

Anatomical Position	Adhesiogenic Stimulus	References
Single Injury		
1. Bowel (Caecum)	a. Abrasion by gauze	20 ^{rh} , 21 ^{rh} , 71 ^{mh} , 76 ^{rh} , 112 ^{rh} , 145 ^{rh} , 152 ^{rh} , 162 ^{rh} , 165 ^m , 170 ^{rh} , 225 ^{rh} , 258 ^{rh} , 259 ^{rh} , 287 ^{rh} , 288 ^{rh} , 322 ^{rh} , 342 rd
	b. Abrasion by scalpel	9 ^{ma} , 19 ^{rh} , 80 ^{rdh} , 160 ^{rh} , 335 ^{rh}
	c. Abrasion by brush	138 ^{rh}
	d. Abrasion by abrasive paper	320 ^r
	e. Crush injury	38 ^r , 199 ^r
	f. Electrocautery	
	g. Excision	
	h. Incision	224 ^r
2. Sidewall	a. Abrasion by gauze	
	b. Abrasion by scalpel	125 ^{rh} , 130 ^{mh}
	c. Abrasion by brush	
	d. Abrasion by abrasive paper	320 ^r
	e. Crush injury	
	f. Electrocautery	16 ^r
	g. Excision	13 ^r , 23 ^r , 93 ^{mi} , 113 ^{ma} , 119 ^{ri} , 160 ^r , 176 ^{ri} , 192 ^r , 228 ^r , 320 ^r
	h. Incision	13 ^r , 204 ^{mi}
3. Uterine horn	i. Ischaemic buttons	1 ^{ri} , 47 ^{ri} , 54 ^{ri} , 58 ^{ri} , 132 ^{mi} , 156 ^{ri} , 168 ^{ri} , 242 ^{ri} , 238 ^{ri} , 263 ^{mi} , 314 ^{ri} , 320 ^{ri} , 330 ^{ri}
		24 ^{ra} , 307 ^{ra}
	a. Abrasion by gauze	
	b. Abrasion by scalpel	112 ^{rh} , 311 ^{rh}
	c. Abrasion by brush	
	d. Electrocautery	48 ^r , 55 ^r , 79 ^r , 116 ^m , 147 ^r , 202 ^m , 279 ^r , 303 ^r
	e. Crush injury	
4. Foreign body	a. Talcum	206 ^r , 207 ^r , 295 ^r
	b. Sponge discs	184 ^m , 185 ^m
Combinatorial Injury		
1a + 2a		131 ^m , 137 ^{rh} , 316 ^{rdh} , 317 ^{rh} , 318 ^{rah} , 296 rd
1a + 2b		27 ^{rh} , 36 ^r , 337 ^{rah}
1a + 2g		27 ^{rh} , 50 ^{rah} , 126 ^{rh} , 232 ^{rah} , 233 ^{rah} , 234 ^{rh} , 230 ^{rh} , 331 ^{ra} , 341 ^{rh}
1a + 2i		34 ^m , 263 ^m
1b + 2b		99 ^{rdh} , 158 ^r
1b + 2g		159 ^r
1c + 2c		221 ^{rh} , 301 ^{rh}
1c + 2g		14 ^{rh} , 282 ^{rh}
1d + 2g		52 ^r , 213 ^r
1e + 2g		25 ^{ri}
1f + 2f		106 ^{ra}
2b + 3a		118 ^{rh} , 249 ^{rah}
2b + 3b		339 ^{rh}
2c + 3c		313 ^{rh}
2h + 2g		65 ^{ri} , 239 ^r
2g + 3		25 ^{ra} , 119 ^{ra}
2g + 3d		312 ^{ri}
2g + 3e		22 ^{ra}

Some rat (**r**) and mouse (**m**) adhesion models may include one or more additional variables, which is denoted by the following: **a**, approximation of anatomic sites; **d**, desiccation (air or alcohol); **h**, haemorrhage; **i**, ischaemia. *References:* (Aarons et al., 2007; Andrews et al., 2012; Api et al., 2015;

Ara et al., 2006; Arung, Drion, & Detry, 2016; Aysan et al., 2002; Aysan, Basak, Kinaci, Yanar, & Coskun, 2007; Aysan, Bektas, Ersoz, Sari, & Huq, 2010; Bakkum et al., 1995; Bakkum et al., 1996; Bakkum & Trimbos-Kemper, 1995; Bakkum, van Blitterswijk, Dalmeijer, & Trimbos, 1994; Balbinotto et al., 2010; Bayhan et al., 2016; Bianchi et al., 2016; Bolgen, Vargel, Korkusuz, Menciloglu, & Piskin, 2007; Cassidy et al., 2015; Cetin et al., 2004; Chaturvedi, Lomme, Hendriks, & van Goor, 2013; Cho, Oh, & Lee, 2010; Chu et al., 2011; Coelho Junior et al., 2015; Cohen et al., 2007; Dalgic et al., 2015; de Virgilio et al., 1990; Dinarvand et al., 2012; Duran, Kuscu, Zeyneloglu, Saygili, & Batioglu, 2002; Elkins et al., 1984; Franconi, Roux, Lefebvre-Lacoeuille, & Lemaire, 2015; Gong et al., 2013; Gorvy, Herrick, Shah, & Ferguson, 2005; Guo et al., 2008; Halverson et al., 1999; Haney & Doty, 1994; Hasdemir et al., 2017; Hellebrekers et al., 2000; Hoban et al., 1994; Holschneider et al., 1997; Holsti et al., 2004; Hong et al., 2015; Ignjatovic et al., 2010; Irkorucu et al., 2009; Keskin et al., 2013; Kraemer et al., 2012; Kucuk et al., 2007; Kuroyanagi et al., 2014; Lai et al., 2005; Lauder, Garcea, Strickland, & Maddern, 2011; Lim et al., 2010; Liu et al., 2006; Lucas, Warejcka, Young, & Lee, 1996; Mendes, Campos, Rocha, & Andrade, 2009a; Mendes et al., 2009b; Miyamoto et al., 2010; Nagelschmidt & Saad, 1997; Nassif et al., 2016; Nehez et al., 2006; Nikeghbalian et al., 2007; Nikeghbalian et al., 2015; Oh et al., 2005; Otcu, Ozturk, Aldemir, Kilinc, & Dokucu, 2003; Ozler et al., 2011; Panahi et al., 2012; Parsaei, Karimi, Asadi, & Rafieian-Kopaei, 2013; Peker et al., 2013; Poehnert et al., 2015, 2015; Poehnert et al., 2015; Rasti, Parvaresh, Tavajoh, & Talaei, 2007; Reed et al., 2007; Sahbaz, Ersan, & Aydin, 2014; Sahbaz, Isik, Aynioglu, Gungorduk, & Gun, 2014; Suzuki et al., 2015; Takagi et al., 2013; Tarhan, Barut, & Sezik, 2008; Tarhan, Barut, Sutcu, Akdeniz, & Akturk, 2006; Torre, Favre, Pini Prato, Brizzolara, & Martucciello, 2002; Townsend et al., 2011; Turkoglu et al., 2015; Uysal et al., 2012; van Westreenen et al., 1999; Vural et al., 1998; Wallwiener et al., 2010; Wallwiener et al., 2006; Wallwiener et al., 2012; Wei et al., 2016a; Wei et al., 2015; Wei et al., 2016b; Whang et al., 2011; Winny et al., 2016; Wu et al., 2014; Wu et al., 2015; Xu et al., 2002a; Yang et al., 2012; Yang et al., 2011; Yilmaz et al., 2012; Yilmazlar, Kaya, Gurpinar, & Emiroglu, 1996)

4.1.2 Selection of anatomic site

Three anatomic sites are frequently used to exact adhesiogenic trauma: the uterus, the bowel, and the abdominal sidewall (refer to **Table 4.1**). Sometimes, more than one site is used in a single model (Bayhan et al., 2016; Dalgic et al., 2015; Turkoglu et al., 2015; Wei et al., 2016a). For example, adhesions were induced using a combination of gauze abrasion of the caecum with parietal peritoneum excision (Bayhan et al., 2016). In general, these sites are chosen to facilitate inference to the human equivalent anatomic site. However, it may be useful to point out that variations in adhesion development may also be influenced by differences in various species posture after surgery. In particular, humans tend to spend a longer period reclined in a supine position during the immediate post-operative period whereas laboratory animals often recover faster into a prone position and then mobilise

earlier to adopt their usual four-legged orientation. Therefore, the time they spent in their respective postures during this critical period may significantly influence the outcomes of adhesiogenesis (Wiseman, 2000).

Among the sites, the bowel region appears to be the favourite (refer to **Table 4.1**), with the caecum commonly employed to model intestinal surgery. Although the small bowel (Andrews et al., 2012; Xu, Rivkind, Pappo, Pikarsky, & Levi-Schaffer, 2002b) is also sometimes employed, the caecum of rodents and rabbits are easier to identify and access upon entry. This provides a surface area that is straightforward for the induction of adhesions (Burns et al., 1996). In this regard, abrasion trauma is a common adhesiogenic stimulus inflicted upon this site (Andrews et al., 2012; Irkorucu et al., 2009; Karaca, Gözalan, Yoldaş, Bilgin, & Tezer, 2013; Koca et al., 2016; Whang et al., 2011), although some studies employ tissue crush injuries (Bozkurt, Yuzbasioglu, Bulbuloglu, Gul, & Kale, 2009; Nagelschmidt & Saad, 1997).

Modeling of abdominal or gynaecologic surgery is provided by adhesiogenic trauma to the next popular site; the abdominal sidewall (refer to **Table 4.1**). These regions allow estimation of surface areas occupied by adhesions to be more readily performed, although variations may exist due to wound contraction (Wiseman, 2000). The creation of a parietal peritoneum excisional defect (80; 176; 187) and “ischaemic buttons” are the most common adhesiogenic stimuli employed on this anatomic site. Of note, the abdominal incision also influences the models whereby, paramedian transrectus incisions appear to develop more adhesions as compared to subcostal or midline incisions in rats, parallel to clinical experience (Moreno et al.,

1996). Therefore, the type of abdominal incisions for accessing the viscera for the induction of the post-surgical peritoneal adhesion models should be carefully considered and performed.

The third common site is the uterus region, in which the uterine horns are commonly used to model the effects of adhesions after human pelvic surgery. Sometimes, the contralateral uterine horn may serve as a control (Arora, Jaroudi, Hamilton, & Dayel, 1994; Rice, Shanti, Moghissi, & Leach, 1993). However, one must be cautious when interpreting results in these models as crossover effects may occur, particularly in small animals (e.g. mice, rats, hamsters), in which inflammatory signals from the injured uterine horn may spread to the contralateral uterine horn. This concern may explain why although this site has been explored in mice (Haney & Doty, 1994), rats (Keskin et al., 2013), and hamsters (Best, Rittenhouse, & Sueldo, 1991), rabbits appear to be a preferred animal choice (Wiseman, 2000). Electrocautery is the main method through which trauma is inflicted on this site (Cetin et al., 2004; Coelho Junior et al., 2015; Duran et al., 2002; Haney & Doty, 1994; Keskin et al., 2013; Uysal et al., 2012) although scalpel abrasion is sometimes employed (Guo et al., 2008; Vural et al., 1998).

4.1.3 Selection of Adhesiogenic Stimulus

Numerous types of adhesiogenic stimulus have been employed for the induction of adhesions (refer to **Table 4.2**). The nature, degree, and sometimes the combination of the inflicted stimulus significantly impact the duration and magnitude of inflammation. This would in turn influence adhesion development and the eventual capacity to treat them. Also, these variations play a significant role in determining

if an adhesion model becomes “challenging” or “permissive”, facilitating the particular inquiry to be carried out. However, it is important to ensure that the model does not become overly “permissive” such that all agents become effective, nor excessively “challenging” such that no agents are capable of reducing adhesions. While there is no single “perfect” method to employ, the adopted method should be well described and be able to produce reliable, consistent, and reproducible data. In this section, the primary types of adhesiogenic stimuli are discussed.

Table 4.2 Types of stimuli employed to induce adhesions

Stimulus	Additional variables
Abrasion injury	Approximation
Ischaemia injury	Anastomosis
Chemical injury	Contamination-infection
Crush injury	Desiccation (air or alcohol)
Electrocautery	Foreign body/materials-tissue irritation
Excision, Incision	Bleeding/Haemorrhage
Thermal injury	
Radiation injury	

Table adapted from Wiseman et al. (2000).

4.1.3.1 Abrasion injury

The most common adhesiogenic stimulus was use of an abrasion injury (refer to **Table 4.1**). One of the popular anatomic sites where abrasion is inflicted is the caecum. Although numerous instruments have been used (e.g. gauze, scalpel, brush, abrasive paper), a dry sterile gauze appears to be a favourite choice. Although abrasion may be regarded as one of most challenging methods to standardise (Wiseman, 2000), some efforts have been taken to refine the method. For example, a consistent area for abrasion may be facilitated through the use of templates (Burns et al., 1995; Burns et al., 1996). Regardless of the choice of anatomic site and instrument, the general consensus is that the end point of abrasion is the induction of petechial haemorrhages (Aysan et al., 2002; Irkorucu et al., 2009; Karaca et al.,

2016; Koca et al., 2016; Lai et al., 2005; Xu et al., 2002a). Fibrin from the bleed facilitates the initial adhesion of two surfaces during early adhesiogenesis (Wiseman, 2000). On this note, some studies also include desiccation for up to five minutes after haemorrhage has occurred. This is suggested to be a potent combination for the creation of adhesions (Ozel et al., 2005). Variations also include the addition of chemical irritants (e.g. iodine and alcohol) onto the abraded portion (Liebman et al., 1993; Osada et al., 1999). In an effort to produce adhesions more consistently, abrasive injuries, typically to the caecum, have often been combined with a parietal peritoneal abrasion or peritoneal excisional defect, sometimes with/without approximation of the two sites with sutures (refer to **Table 4.1**). Also, in the small bowel loop model, a loop of the small bowel is abraded and followed by the approximation of the injured serosal regions with the help of sutures (Andrews et al., 2012).

4.1.3.2 Excision injury

Excision injury is another common adhesiogenic stimulus (refer to **Table 4.1**), in which the parietal peritoneum with the underlying muscle is removed, usually performed on the sidewall using scalpel or surgical scissors. Further aggravation of adhesiogenesis may be achieved when muscle injury is exacted. Similar-sized biopsy punches can help standardise the areas of excision. On this note, greater adhesions formation is associated with larger defects (Ozel et al., 2005). While this injury is often applied in isolation, it may also be combined with other forms of serosal injury with/without approximation of two sites (Bayhan et al., 2016; Poehnert et al., 2015). Suturing to repair the defects is also another variation in an attempt by some to produce adhesions more consistently (Falk et al., 1998;

Hellebrekers et al., 2000). It is suggested that secondary ischaemia and foreign body reaction is likely to be responsible for contributing to this process (Ellis, 1971; Kraemer et al., 2014).

4.1.3.3 Presence of foreign bodies/materials

Another effective adhesiogenic stimuli are foreign bodies/materials, which have the capacity of generating a granulomatous reaction when placed in the peritoneum. These materials range from powders that are used to coat surgical gloves (e.g. talcum, starch), to irrigation solutions, to gauzes and meshes (prolene, polyglycolate, biologicals, etc), to sutures (absorbable and non-absorbable). In general, most studies in this category usually employ the use of powders (Nikeghbalian et al., 2007; Nikeghbalian et al., 2015). A dose-dependent relationship is observed between the amount of powder administered and the final adhesion and granulation formation (Jagelman & Ellis, 1973; Jahoda, Olson, & Kovacs, 2003; van den Tol et al., 2001). It is noted that certain regions are more affected than others (Jahoda et al., 2003). Therefore, when embarking on such studies, establishing beforehand the tissue region to be removed, and keeping it consistent, will help to remove possible bias. It is also good practice to only harvest from regions away (i.e. a different abdominal quadrant) from where the solution of powder was injected (Jahoda et al., 2003). On this note, it is also crucial to use non-powdered surgical gloves for experimental studies to avoid unwanted reactions.

Sutures of different composition generally produce different foreign body tissue reactions, with absorbable types (e.g. polyglycolic and polyglactin) resulting in a milder reaction compared to non-absorbable types (e.g. Prolene, silk). Absorbable

sutures are generally used for adhesion creation (Holmdahl, al-Jabreen, & Risberg, 1994; Rice et al., 1993). However, it is believed that the generated ischaemia due to tight suturing is main contributor for the adhesion creation instead of tissue reaction (Ellis, 1971; Holmdahl et al., 1994; Kraemer et al., 2014).

4.1.3.4 Ischaemic injury

Ischaemia is the restriction of blood supply to tissues, which can result in tissue death. This has been reported through use of a method of devascularisation (Wiseman, Huang, Johns, Rodgers, & Dizerega, 1994), ligation (Snoj, Ar'Rajab, Ahren, Larsson, & Bengmark, 1993), temporary vascular obstruction (Tsimoyiannis, 1989), and the most popular, the creation of ischaemic buttons (Whang et al., 2011). An early study elegantly demonstrated the significant contribution to adhesion formation of ischaemia induced by suturing (Ellis, 1962). The combination of ischaemia injury and abrasion injury appeared to aggravate adhesion created in a rabbit uterine horn model (Wiseman et al., 1994), in which adhesions were formed between the uterus and sites that were distant from the primary injury. On the contrary, the observed adhesions were only confined to the uterus in another study employing the identical model with only abrasion insult, thus highlighting the impact of ischaemia injury (Wiseman, Gottlick, & Diamond, 1992).

The majority of the studies were found to have chosen to inflict an ischaemic insult through the creation of an ischaemic peritoneal button (**Table 4.1**). Ligating an out-pouching of the abdominal sidewall, usually with a suture, creates this button. Although this method was suggested to be the best model compared to other post-surgical peritoneal adhesion models (Whang et al., 2011), the variation of adhesion

induction in this model may range from 62.3% to 92% in published literature (Kraemer et al., 2014; Mommers, Hong, Jongen, & Bouvy, 2018; Rajab et al., 2010; Sheldon et al., 2012). Multiple factors such as, type of suture material, skill of surgeon, size of button, and the duration of procedure, and evaluation may explain this. Although differences of histological response due to the types of sutures used have been speculated to affect the adhesiogenic performance of ischaemic button (Kharwadkar, Naique, & Molitor, 2005; Sajid, Parampalli, Baig, & McFall, 2011), a comprehensive comparison has yet to be performed. Nevertheless, the presence of a stronger response is likely to make adhesion creation more robust. Refining the surgical technique (i.e. minimising the angle between suture and sidewall) on post-mortem animals may help to reduce incidence of suture slippage (Mommers et al., 2018). Ultimately, one major advantage of this model is that the traumatised area is well defined and easily identified, facilitating the study of both early and late molecular events of adhesionogenesis.

4.1.4 Selection of timing and macroscopic assessment

A variety of timings have been used to evaluate adhesion formation depending on research inquiry, ranging from as early as 12 hours to 24 weeks post-induction (Ozel et al., 2005). In general, the earlier time frame (i.e. less than 24 hours) is useful for understanding adhesion pathogenesis (Ryan, Grobety, & Majno, 1971) although assessment is more frequently performed at later time frames (i.e. 7 to 10 days) since there is general agreement that adhesion formation is completed by then (Beyene et al., 2015; diZerega & Campeau, 2001; Gaertner et al., 2008; Holmdahl et al., 1994).

At the relevant time points, the animals are euthanised and followed by the assessment of adhesion that is ideally performed by researchers blinded to the test groups. Assessment is usually initiated with a U-shaped incision that encompasses the original site for access to the viscera (Beyene et al., 2015; Ozel et al., 2005). The point of second surgical entry is largely decided based on where the injury was inflicted. As a general rule, the incision for evaluation should be as distant as possible from the original entry and away from areas where adhesions are likely to form. This is so that adhesions may be preserved in their physiological state, as some filmy adhesions are easy to break upon gentle traction. Therefore, entry and exploration should be performed thoroughly and meticulously.

Macroscopic determination of adhesions appears to be the main assessment for the majority of experimental studies (Ozel et al., 2005). However, a wide range of different scoring systems exists, making it challenging to compare performance between models (Beyene et al., 2015). The number of adhesions found during assessment may be counted in a simple quantitative approach and reported (on a per site or per animal basis); or assessed qualitatively based on the density, tenacity, and strength (Adhesion Scoring Group, 1994; Cassidy et al., 2015; Hoffmann et al., 2009; Nair et al., 1974). Alternatively, the assessments may consider only the traumatised area (usually for caecum abrasion models), estimating the extent of the area that is covered by adhesions. This may be later reported as a numerical percentage or spilt on a 5-point scale (Chaturvedi et al., 2013; Leach, Burns, Dawe, SmithBarbour, & Diamond, 1998).

To newcomers of the field, usage of macroscopic assessments to measure adhesion characteristics (e.g. adhesion strength) may seem to be a concern of reproducibility and reliability. One possible way to overcome this is through the usage of devices to measure adhesion characteristics quantitatively, such as using a digital force meter when measuring adhesion strength. This has been explored by Kraemer and colleagues (Kraemer et al., 2011). However, the authors noted a problem with this method is where to apply the digital force meter. The authors discussed that the strongest point of the adhesion may be located separately and is not necessarily the point of insertion (Kraemer et al., 2011). Moreover, since measurement with the digital force meter score requires excision of the adhesion, which cannot be determined clinically *in situ* during assessment of clinical adhesiolysis, representing a huge disadvantage for this method of assessment in clinical setting (Kraemer et al., 2011). As such, assessment of adhesion strength remains a common parameter performed macroscopically. This is possible due to the well-defined terms used when describing the different grades of adhesion strength.

The adhesion strength scoring matrix ranks the detected adhesions into 5 possible grades (**Table 4.3**). The lowest and highest grade can be easily awarded to adhesions due to straightforward morphological descriptions provided by the scoring matrix (e.g. grade 0 – complete absence of adhesions; grade 4 – adhesion of all viscera to the abdominal wall). Next, adhesion strength that falls in between grades 0 and 4, can be further distinguished into 3 grades depending on ease (or effort) with which they can be broken (e.g. grade 1 – gentle traction requiring little effort to break adhesion; grade 2 – blunt dissection requiring a blunt instrument to break adhesion; grade 3 – sharp dissection requiring a sharp instrument to break adhesion). Sharp

and blunt dissection are common surgical techniques that are used frequently by adhesion scoring systems described in literature. The key difference between the two is that sharp dissection requires an instrument with a sharp edge (e.g. scalpel blade), whereas blunt dissection may be easily performed using a blunt instrument (e.g. blunt forceps) to skilfully pull apart adjoining tissues. In certain cases where adhesions are attached to tissues of weaker integrity (e.g. adipose tissues, as opposed to a tissue of stronger integrity like the muscular abdominal sidewall), sharp dissection is employed as opposed to blunt dissection that may result in unwanted critical tissue damage. Finally, gentle traction can also be easily performed with a blunt instrument, but the main difference is that it does not require much effort or skill in separating the adjoining tissues (e.g. do not need to consider the plane of dissection). This highlights the importance of performing the necropsy carefully and gently as adhesions of such nature can be easily broken if manipulation of tissues during necropsy is not performed mindfully and roughly.

Ultimately, the adopted scoring should be well defined, so as to enable similar understanding among different researchers, and possess the necessary capacity to capture faithfully the complexity of the adhesion formation. Since macroscopic assessment may be prone to intra- and inter-observer variability, it is also critical that other techniques (i.e. immunohistochemical, biochemical assays) complement this approach.

Table 4.3 Scoring matrix for assessing strength of peritoneal adhesions

Variable	Score	Description
Strength (Hoffmann et al., 2009)	0	Complete absence of adhesions
	1	Gentle traction requiring little effort to break adhesion
	2	Blunt dissection requiring a blunt instrument to break adhesion
	3	Sharp dissection requiring a sharp instrument to break adhesion
	4	Adhesion of all viscera to the abdominal wall

4.1.5 Other considerations

As mentioned above, regardless of method employed, the adopted method should be well described and be able to achieve consistent, reliable and reproducible data. It is critical that surgeon is well versed in the surgical technique. This may be achieved by extensive practice on post-mortem animals before living animals are attempted to ensure that obtained adhesions are adequately severe and created consistently across experiments. Also, well-defined methods of assessing adhesions should be adopted and observers performing the evaluation should ideally be blinded. The number of potential variables in the adopted method should be reduced (e.g. the use of templates, standardising the number of times of abrasion and amount of tissue grasped, etc.) such that consistency and reliability of the model can be reproduced easily. These refinements would simplify and streamline the surgery, reducing surgery time and surgeon fatigue. In addition, the adopted model should also be performed during the same period of the day to mitigate potential difference due to influences from the circadian rhythm (Sadek, Macklon, Bruce, Cagampang, & Cheong, 2011).

4.1.6 Selection of model for studying the role of Cx43 during adhesion formation

As it is challenging to directly compare results from different published reports, the performance of various models can only be informed if explored together under the same timing of evaluation and appropriate scoring matrix. Therefore, popular methods of creating peritoneal adhesions were selected from the literature (refer to **Table 4.1**) for comparison and exploration. Methods requiring specialised instruments/equipment (e.g. electrocautery) were excluded. In addition, methods combining more than one type injury were also excluded to reduce the potential number of variations. This narrowed down the available models to: the parietal ischaemic button model, the parietal peritoneum excision model, the small bowel loop model, and the caecal abrasion model. The performance of the models was evaluated on Cx43^{fl/fl} transgenic mice so that the role of Cx43 may be studied in the future using genetically modified transgenic strains (in **Chapter 6**). On this note, only male mice were used to avoid the confounding effects of the estrous cycle on adhesion formation (Bouckaert, van Wersch, Schellekens, Evers, & Rolland, 1984). Assessment was performed on day 7 post-surgery since there is general consensus that adhesion formation is completed by then (diZerega & Campeau, 2001). Finally, the performance of the explored models was then assessed using established scoring matrices assessing the quantity and quality of generated adhesions (Adhesion Scoring Group, 1994; Nair et al., 1974) to determine which model is the most ideal for the study of Cx43 in adhesion formation.

4.2 Aims and Objectives

Work in this chapter aims to explore popular *in vivo* adhesion models to identify the most ideal model for studying the role of Cx43 during adhesion formation. This would be achieved in two steps. First, types of adhesion models available are identified through a systematic literature search. Next, popular methods of creating peritoneal adhesions in literature are explored for comparison; these methods include the parietal ischaemic button model, the parietal peritoneum excision model, the small bowel loop model, and the caecal abrasion model. To determine which of these models are the most ideal, an assessment of developed adhesions would be performed on day 7 post-surgery. This assessment comprises of established macroscopic scoring matrices for evaluating the quality and quantity of adhesions, and also other factors in other considerations (e.g. degree of model standardisation, identification and definition of site trauma). Collectively, these steps would facilitate the identification of the most ideal model for studying the role of Cx43 during adhesion formation.

4.3 Materials and Method

4.3.1 Experimental animals

All animal procedures were performed in accordance with the animal protocols approved by the Institutional Animal Care and Use Committee of the Animal Research Facility of Nanyang Technological University, protocol number A0372. Mice were bred under specific pathogen-free conditions under a 12-hour light-dark cycle. Standard rodent pellet diet and water were supplied *ad libitum*. Only age-matched (7 to 12 weeks) males were used for experiments.

4.3.2 Induction of adhesion formation

The major steps for the induction of adhesions and the different models of post-surgical peritoneal adhesions (n=3, each) are described in the following sections and summarised in **Figure 4.2**.

4.3.2.1 Surgery preparation and entry

Prior to the procedure, fur around the incision site was removed using an electric trimmer and depilatory cream before returning the mice to their cages and allowing them to rest for at least 24 hours. Briefly, mice were anaesthetised with isoflurane and placed on a warm heating mat throughout the surgery. All surgeons used non-powdered surgical gloves during the surgery and all instruments were sterilised. Next, the midline incision site was sterilised with 7.5% (v/v) providone-iodine solution (**Fig. 4.2a**). A midline laparotomy of 4 cm was performed to facilitate the exposure of the parietal peritoneum (**Fig. 4.2b**). Depending on experiment design, the following four adhesion-inducing procedures may be employed on the mice.

4.3.2.2 Ischaemic button model of post-surgical peritoneal adhesions

Ischaemic buttons (IB, n=3) were created to induce adhesion formation as previously described by Sandoval et al., (2016) with minor modifications. Each ischaemic button was created by placing a 4/0 non-absorbable silk suture at the base of ~3 mm of the sidewall parietal peritoneum, which was grasped with a needle holder. Two ischaemic buttons, spaced at least 1 cm apart, were created on the peritoneum wall of each side of the midline (**Fig. 4.2c**).

4.3.2.3 Parietal peritoneum excision model of post-surgical peritoneal adhesions

The parietal peritoneum excision (PPE, n=3) procedure was adapted with modifications (Falk et al., 1998). After the midline laparotomy, the skin was carefully separated from the musculo-peritoneal layer to facilitate excision made on the latter layer. A defect was then created using a 6 mm biopsy punch on the left side of the midline incision. The parietal peritoneum, together with the underlying layer of muscle, was removed when the defect was created with the punch (**Fig. 4.2d**).

4.3.2.4 Small bowel loop model of post-surgical peritoneal adhesions

The small bowel loop (SBL, n=3) procedure was adapted with modifications (Andrews et al., 2012). A loop of small bowel was selected in the mid jejunal region and folded back into a juxtaposed position. A size 10 scalpel abraded the serosal surfaces of the selected bowel, which were to be placed in contact, for 20 times in the same direction, over a length of 10 mm. Two 7/0 polyglyconate sutures were later placed at each end of the abraded portion, holding the bowel in apposition (**Fig. 4.2e**).

4.3.2.5 Caecal abrasion model of post-surgical peritoneal adhesions

The caecal abrasion (CA, n=3) procedure is adapted as described by Hemadeh and colleagues (Hemadeh et al., 1993). Following midline laparotomy, the caecum was located and exposed. The anterior portion was abraded with a size 10 scalpel until petechial subserosal hemorrhages developed (**Fig. 4.2f**).

4.3.2.6 Surgery closure and post-recovery regime

The laparotomy site was then closed in two layers, the inner layer with simple interrupted polyglycolic acid absorbable 6.0 sutures (**Fig. 4.2g**) and the outer layer using metallic wound clips (**Fig. 4.2h**). After closure, mice were regularly monitored until euthanasia at 7 days post-surgery by euthanasia with CO₂. For post-recovery, the mice received buprenorphine (10 mL/10 g of animal weight) into the loose skin over the interscapular region.

Figure 4.2 Major steps during various adhesion-inducing operations in mice.

(A) Disinfection of the incision site after shaving. This was followed by a (B) midline incision of the skin. (C) Ischaemic button (IB) model. Two IBs, spaced approximately 1 cm apart, were created on the peritoneum on each side of the midline incision. Each button (yellow arrowheads) was created

by placing a silk ligature at the base of 3 mm of parietal tissue that is grasped with a needle holder. **(D)** Small bowel loop (SBL) model. A loop of small bowel was selected at middle small bowel level to be folded back into a juxtaposed position. The serosal surfaces of the selected bowel were scalpel abraded and two absorbable sutures (yellow arrowheads) were later placed at each end of the abraded portion, holding the bowel in apposition. **(E)** Parietal peritoneum excision (PPE) model. An excision is made through the musculo-peritoneal layer using a 6 mm biopsy punch on each side of the midline incision. **(F)** Caecal abrasion (CA) model. Caecum is located and exposed, before abrading the anterior portion with a scalpel blade until petechial subserosal haemorrhages (yellow arrowheads) develop. **(G)** Wound closure by applying simple interrupted sutures to the peritoneal musculature and **(H)** metallic clips to the skin.

4.3.3 Macroscopic assessment of peritoneal adhesions

An inversed U-shaped incision was made across the peritoneal cavity for necropsy and the adhesions in each animal were evaluated according to established scoring matrices. These matrices assessed peritoneal adhesions based on qualities such as, extent, involvement, strength, and severity (refer to **Table 4.4**) (Adhesion Scoring Group, 1994; Cassidy et al., 2015; Hoffmann et al., 2009; Nair et al., 1974). Three observers blinded to the experimental groups scored each mouse and their scores were later averaged to give a single score per mouse. Inter-observer scores did not differ more than 2. Scores are represented as a percentage of maximum score possible for each parameter and the mean parameter scores for each of the groups were later calculated.

Table 4.4 Scoring matrix for assessing peritoneal adhesions

Variable	Score	Description
Involvement (Quantity) (Nair et al., 1974)	0	Complete absence of adhesions
	1	1 adhesion band between the organs or between one organ and abdominal wall
	2	2 adhesion bands between the organs or between one organ and abdominal wall
	3	>2 adhesion bands between the organs or between one organ and abdominal wall or adhesions of intestinal loops without any adhesion to the abdominal wall. Significant adhesion.
	4	Adhesion of all viscera to the abdominal wall
Severity (Quality) (Adhesion Scoring Group, 1994)	0	Complete absence of adhesions
	1	Filmy, avascular
	2	Dense, opaque and/or vascular
	3	Very thick cohesive attachment

4.4 Results

4.4.1 Mice subjected to SBL did not survive until intended date of inspection.

Popular methods of creating peritoneal adhesions were explored in a pilot experiment to determine the most ideal method for the future study of Cx43 in peritoneal adhesions. These methods included the ischaemic button (IB), parietal peritoneum excision (PPE), small bowel loop (SBL), and the caecal abrasion (CA) procedures of adhesion creation (n=3, each). In the IB method of creating adhesions, two IBs were made on each side of the midline incision; each IB was created through a suture ligation of an out-pouching of the abdominal sidewall (**Fig. 4.2c**). Another popular method employed to induce adhesions was the PPE method. This method involved making an excision through the musculo-peritoneal layer using a biopsy punch on each side of the midline incision (**Fig. 4.2d**). Next, the SBL method of inducing adhesions was also explored. This involved abrading serosal surfaces of the selected bowel with scalpel before folding the abraded surfaces into apposition with the help of absorbable sutures (**Fig. 4.2e**). Finally, the last method explored to

induce adhesion was the CA procedure, which involved abrading the exposed caecum with a scalpel until petechial subserosal haemorrhages developed (**Fig. 4.2f**). Mice were allowed to recover after their respective procedures and were regularly monitored over the experimental period of 7 days before the induced adhesions were assessed.

Regular monitoring over the experimental period noted that the mice subjected to the caecal abrasion (CA), ischaemic button (IB), and parietal peritoneum excision (PPE) procedures recovered well post-surgery (n=3, each). In contrast, mice subjected to the small bowel loop (SBL) procedure (n=3) appeared tense from post-surgery day 3 onwards and were hunched with patches of hair piloerected. Their health was observed to have deteriorated over the following days. On post-surgery day 6, the mice appeared to be puffy with eyes half closed or more and showed significantly reduced respiration rate and quality. In addition, the mice were noted to have suppressed activity and elicited a mild response to touch stimulus. Under institutional animal care guidelines, euthanasia was performed. Immediate necropsy of the peritoneal cavity was performed after euthanasia. Small bowel distension from the stomach to the SBL was noted (**Fig. 4.3**). The portion of intestinal tract between SBL and the caecum was observed to be relatively empty compared to the portion of the tract between the SBL and stomach, indicating that small bowel obstruction may have occurred.

Figure 4.3 Small bowel obstruction observed in small bowel loop model.

Representative photo of small bowel obstruction observed upon necropsy for mice subjected to the SBL model. Small intestine distention (red arrowheads) was observed, starting from the stomach to where the SBL was made (yellow arrowhead). Intestinal tract after the SBL appear to be largely empty towards the caecum, suggesting small bowel obstruction.

4.4.2 Adhesions were observed in CA, IB, and PPE adhesion models.

Mice subjected to the CA, IB and PPE procedures (n=3, each) recovered well post-surgery and were euthanised on day 7 for assessment of adhesions formed. The peritoneal cavity was thoroughly explored to detect for the presence adhesions. In all animals, apart from the site of the procedure, all observers noted there to be no other adhesions in the peritoneal cavity (including the midline laparotomy site) in all animals. An overview of the adhesion bands detected upon necropsy is represented in **Figure 4.4**. At least 1 adhesion band was found in each mouse in the IB group (**Fig. 4.5a**). In contrast, no adhesion bands were found in 1 mouse in both CA and PPE group.

To further determine the performance of each model, adhesions were scored macroscopically based on quality and quantity of adhesion during necropsy by three

observers (refer to **Table 4.4**). Both quantity and quality scores for each model is expressed as a percentage of their respective maximum possible score. No significant differences in terms of quantity and quality were found among the various models. The CA and PPE models had similar quantity and quality scores (mean observer score of 16.7% and 37.0% respectively, **Fig. 4.5a and 4.5b**). In contrast, the IB model was found to have slightly lower quantity and quality scores (mean observer score of 12.5% and 21.8% respectively, **Fig. 4.5a and b**). Overall, the three models were found to have relatively close reproducibility in developing adhesions in terms of quantity (standard deviations for CA, IB, PPE = 14.4, 6.30, 14.4 respectively, **Fig. 4.5a**). However, in terms of quality, the IB model was substantially more reproducible than the CA and PPE models (standard deviations for CA, IB, PPE = 39.0, 5.61, 39.0 respectively, **Fig. 4.5b**).

Figure 4.4 Overview of the adhesions observed in various adhesion models upon necropsy.

(A) In the CA model, example of a single, opaque adhesion band between the caecum and the abdominal wall. The band can be broken through blunt dissection. (B) Presence of a single, filmy, and avascular adhesion band between the caecum head and itself in the CA model. The band can be broken easily through gentle traction. (C) Example of an absence of adhesions in the CA model. (D) In the IB model, presence of two adhesion bands between the IBs and fat pad. Bands are observed to be thick and cohesive, requiring sharp dissection to break adhesion. (E) Example of a single, opaque adhesion band between IB and liver lobe in the IB model. The band can be broken through blunt dissection. (F) Example of the absence of adhesions in the IB model. (G) In the PPE model, an example of a single, dense adhesion band between the defect and the fat pad. The band can be broken through blunt dissection. (H) Presence of a single, thin adhesion band, between the defect and the fat pad in the PPE model. Gentle traction easily breaks the band. (I) Absence of adhesions observed in the PPE model. Green arrowheads highlight adhesion bands. Red arrowheads show absence of adhesion bands.

Figure 4.5 Comparing the performance of popular adhesion models.

Mice (n=3 for each model) were freshly scored based on quantity and quality (refer to Table 1), upon euthanasia by three observers. The mean observer scores for quantity (**A**) and quality (**B**) for each mouse of each model are represented as a percentage of the maximum score possible. M1 to 3 are individual mice. Scores for mice 1 to 3 for each model are colour-coded from darkest to lightest shades representatively to facilitate visualisation. **CA** - Cecal Abrasion model. **IB** - Ischaemic Button model. **PPE** - Parietal Peritoneum Excision model.

4.5 Discussion

Despite an on-going and compelling need, there is still paucity in the existence of an ideal solution for preventing or treating peritoneal adhesions. Reliable adhesion-producing animal models are key to the development of effective strategies against peritoneal adhesions. Despite years of research, there is still lack of consistency and universal acceptance of a gold standard animal model for developing peritoneal adhesion. In this chapter, popular methods of developing adhesions, which were determined from a literature search, were explored to determine the most ideal adhesion-producing animal model for the study of the role of Cx43 during adhesion formation.

It is recognised that it is a complex endeavour to compare the results of distinct areas, depths, and causes of trauma. Therefore, this pilot study strived to assess the animal models in a fairest way possible, using adhesion-scoring matrices that are suitable to be applied to the models explored. Also, other characteristics of the models are also taken into consideration. One characteristic of an ideal animal model is the ability to be reproduced by investigators in a standardised fashion. The adhesiogenic stimulus of choice should be able to be induced consistently (Kraemer et al., 2014). This would allow the reproducible development of adhesions, which may later be objectively assessed, minimising the number of animals used. Also, the traumatised site should be easily identifiable and well defined. This would allow consistency in measurements and facilitate the investigation of early and late molecular events. Assessment is usually performed day 7 post-surgery since there is general consensus that adhesion formation is completed by then (diZerega & Campeau, 2001). Therefore, an ideal model should also ensure the survival of the

animal throughout this period of adhesion development. Finally, the model should neither be too challenging nor too permissive such that no agent reduces adhesions or all agents become efficacious (Wiseman, 2000).

In this study, four adhesion-producing procedures (SBL, CA, IB, and PPE) were explored after a detailed literature search. Assessment was performed at a clinical level, where adhesions were evaluated macroscopically for quantity and quality. Macroscopic tissue examination is a widely used method for the determination of adhesions in experimental studies (Ozel et al., 2005). These examinations utilise scoring systems that share relevance to the clinical assessment of adhesions (Adhesion Scoring Group, 1994; Wiseman, 2000). Adhesion quantity is a straightforward method that can be achieved quickly and objectively *in situ*. Adhesion quality is another *in situ* method that can be quickly performed and is based on morphological terms. This scoring system also further adds a dimension to adhesion quantity (e.g. two filmy avascular bands as opposed to two dense and opaque bands). Furthermore, this scoring has high clinical relevance as it provides technical information for planned surgical adhesiolysis. On the other hand, microscopic assessments are not widely performed in experimental studies (Beyene et al., 2015). Although a histologic scoring system was proposed by Zühlke in 1990, only a handful of studies have adopted this since then (Beyene et al., 2015; Ozel et al., 2005; Zuhlke, Lorenz, Straub, & Savvas, 1990). In addition, comparing morphometric results of distinct traumatised areas and distinct trauma depths fairly would be a complex endeavour.

Although the SBL procedure has been previously reported to produce reproducible adhesions between the opposing bowel segments (Andrews et al., 2012), in our hands, mice subjected to this procedure did not survive until experiment termination. No obvious adhesions were observed macroscopically upon necropsy. Instead, mice subjected to SBL were observed to have small bowel obstruction, which was suspected to be one of the possible reasons for the health deterioration. Due to the nature of the technique, a certain level of technical expertise is required, which in absence may have increased the likelihood of small bowel obstruction. Furthermore, this procedure was the longest surgical duration compared to the other procedures. This may be challenging if the procedure is to be performed on large experimental groups of mice.

In contrast, the PPE procedure required the least technical expertise. The musculo-peritoneal layer was easily separated from the skin and the area of excision was easily controlled through the use of standard sized biopsy punches. Similarly, little technical expertise was required for the CA procedure, as the caecum was easily located upon laparotomy and the abrasion of the caecum was easily performed. The abrasion trauma increases vascular permeability, leading to the recruitment of leukocytes and contributes to the consequent formation of adhesions (Kraemer et al., 2014). However, for the later part of the CA procedure, it is challenging to achieve petechial haemorrhage in a controlled manner. Due to the flaccidity of the caecum, it is difficult to optimally apply consistent pressure during scalpel abrasion over the area. Furthermore, the number of strokes of scalpel abrasion necessary to induce bleeding is often variable. Therefore, inability to perform these technicalities consistently may present a confounding element to reproducible adhesion

development. In addition, it may prove challenging to identify the initial traumatised site in both models.

Finally, the IB model required a moderate amount of technical expertise due to the requirement to perform simple suturing. In this study, four buttons were made, with two evenly distributed at the top and bottom on each side of the midline laparotomy. Due to the limitation of animal numbers, it is uncertain if the location of the IB would affect the likelihood of adhesion development. Although this is recognised as a potential confounder which warrants further investigation, the IB procedure triumphs over the other procedures in terms of its reproducibility in developing adhesions of similar quantity and quality. Both CA and PPE models were found to be lacking in terms of reproducibility, in which at least 1 animal was found to be absent of adhesions). This choice is further supported by similar conclusions made from an independent group comparing the performance of popular models of peritoneal adhesion (Whang et al., 2011). Another advantage of the IB procedure over the other models is the presence of a defined traumatised site that is easily identifiable due to placement of the suture. This allows for standardisation for performing measurements and tissue harvesting, which facilitates the study of early and late molecular events.

4.6 Conclusions

In light of these considerations, the most consistent and reproducible technique appears to be the IB procedure. In this model, adhesions were only detected at the defined traumatised site. This would allow for the standardisation in adhesion measurements and tissue sampling. Furthermore, the defined trauma site in the model facilitates the tracking of adhesion development, which is important when studying early and late molecular events (i.e. extent of inflammation). Although one obvious weakness of this pilot study is the number of animals, the advantages in standardisation and reproducibility of the IB model both in our hands and as reported in the literature (Whang et al., 2011) make it an attractive technique. Overall, work in this study identifies the IB model to be the best for the study of the role of Cx43 during adhesion formation (**Table 4.5**), and the development of potential strategies against adhesion formation in the future. This is explored in the next chapter.

Table 4.5 Overview of the performance of adhesion models explored

Parameters	SBL	CA	PPE	IB
Degree of standardisation for surgery	Poor	Poor	Strong	Moderate
Mice survival until end of study	No	Yes	Yes	Yes
Ease of assessment	Poor	++	++	+++
Quantity of adhesion	?	+	+	+
Quality of adhesion	?	++	++	+
Consistency of adhesion	?	+	+	+++
Identification and definition of site of trauma	Moderate	Poor	Moderate	Strong

Models explored: **SBL** – Small bowel loop; **CA** – Caecal abrasion; **PPE** – parietal peritoneum excision; **IB** – ischaemic button.

Chapter 5: The role of connexin43 during post-surgical peritoneal adhesion formation

5.1 Introduction

In response to cell injury, function and tissue integrity is maintained by the cellular and molecular mechanisms of tissue repair. This process typically involves two phases. The first regeneration phase involves the replacement of injured cells with cells of the same type. The second fibroplasia phase involves the replacement of normal tissue by connective tissue (Cogliati et al., 2016). Perturbations to these fine-tuned processes, usually due to a prolonged aggressive stimulus, often leads to fibrosis that results in excessive deposition of ECM components, resulting in pathological tissue healing (Cogliati et al., 2016; Wynn, 2007). Myofibroblasts, which are fibroblasts that have been activated and transformed, are the primary sources of ECM and collagen (Wynn, 2007, 2008).

Myofibroblasts usually arise from the activation of resident fibroblasts but may also originate from epithelial or endothelial cells that may have undergone mesenchymal transition. Also, they may originate from circulating fibrocytes from the bone marrow. Regardless, the derivation of myofibroblasts requires a concerted effort that relies on the sequential regulation of cellular events, which includes the differentiation into myofibroblasts, their migration and proliferation, synthesis of ECM, tissue remodeling, and apoptosis (Wynn & Ramalingam, 2012). These processes, which are also shared in adhesion pathogenesis, are tightly regulated and efficiently controlled through the communicative cellular network (Cogliati et al.,

2016). Along these lines, gap junctions may play a pivotal role in the regulation of these processes by providing a conduit for intercellular communication (Cogliati et al., 2016; Naus & Giaume, 2016). In addition, hemichannels also provide a pathway for the release of paracrine messengers. Since little is known about the role of connexins in peritoneal adhesion formation, understanding their role in fibrosis-related diseases may shed some light of their contribution to adhesion formation as some processes are similar between both aetiologies. Some of the current knowledge is briefly discussed below and is divided based on anatomical location, for the sake of clarity.

5.1.1 Skin

In the mammalian skin, a number of connexins is expressed. Of note, dermal fibroblasts express Cx43 and Cx45 predominantly, and Cx40 in lower quantities (Cogliati et al., 2016). Treatment of skin wounds in rats with Cx43asODN has been shown to accelerate the wound healing (Qiu et al., 2003). This was shown to be a result of reduced inflammation, increased proliferation of keratinocytes, and enhanced migration of dermal fibroblasts (Mori et al., 2006). Of note, treatment with Cx43asODN resulted in an increase TGF- β 1 from dermal fibroblasts, which was accompanied with increases in dermal fibroblast proliferation and migration, earlier differentiation into myofibroblasts, and increases in hydroxyproline content (Mori et al., 2006). In the same context, inhibition of hemichannels and gap junctions in dermal fibroblasts also resulted in enhanced migration (Wright, Berends, Flint, & Martin, 2013; Wright et al., 2012). Along these lines, fibroblasts harvested from hypertrophic scars or keloids are revealed to have reduced Cx43 protein levels compared to normal skin (Lu, Gao, Ogawa, & Hyakusoku, 2007).

Conversely, significant elevation of Cx43 protein levels is detected in fibroblasts found in the wound edges of human chronic diabetic foot ulcers (Mendoza-Naranjo et al., 2012). Similar elevation of Cx43 is also observed in wound-edge dermal fibroblasts in streptozotocin-induced diabetic rats (Mendoza-Naranjo et al., 2012). In the same study, fibroblasts cultured in high glucose concentrations to stimulate diabetic conditions exhibited significantly higher Cx43 protein levels and gap junction intercommunication, and reduced migration rates. Following on, prevention of glucose-induced Cx43 up-regulation with Cx43shRNA-transduced fibroblasts increased migration significantly (Mendoza-Naranjo et al., 2012).

Collectively, these evidences appear to suggest that Cx43 is essential for fibroblast homeostasis, particularly in the context of maintaining the balance between proliferation, apoptosis, migration, as well as for regulating the production of ECM.

5.1.2 Liver

In the liver, Cx32 is evenly expressed in all regions while Cx26 is preferentially expressed in the acinar periportal region (Kojima, Sawada, Zhong, Oyamada, & Mori, 1994). On the other hand, non-parenchymal liver cells mainly express Cx43, which includes the sinusoidal endothelial cells, Kupffer cells, and stellate cells (Eugenín, González, Sánchez, Brañes, & Sáez, 2007; Fischer et al., 2005; Hernandez-Guerra et al., 2014). Among these cells, stellate cells play an important role in the deposition and removal of ECM components in the perisinusoidal regions or fibrous septa upon its differentiation into myofibroblasts. During this differentiation, Cx43 protein levels increases, re-distributing to the cell plasma

surface. This is suggested to contribute to GJIC. Suppression of GJIC in primary hepatic stellate cells with carbenoxolone, a broad-spectrum gap junction manipulator, impedes its activation and reduces collagen and DNA synthesis (Uyama et al., 2003). In a separate study, treatment of rat hepatic stellate cells with Cx43-specific siRNA also exhibited reduced cell proliferation (Lim, Maubach, & Zhuo, 2009). These studies suggest that the reduction of Cx43 is protective against fibrosis. In contrast, excessive fibrosis was detected in Cx43^{+/-} transgenic mice compared to wild type controls upon liver fibrosis induction using carbon tetrachloride, (Cogliati et al., 2011).

5.1.3 Heart

During cardiac fibrosis, cardiac fibroblasts are activated, causing their differentiation into myofibroblasts. Once differentiated, the myofibroblasts are shown to be able to establish Ca²⁺-dependent Cx43-mediated communication with the cardiomyocytes, which is suggested to influence the secretion of paracrine/autocrine messengers of myofibroblasts and their function in scar contraction and collagen formation (Chilton et al., 2007).

Asazuma-Nakamura and colleagues demonstrated that TGF- β 1-induced cardiac fibroblast differentiation into myofibroblasts could be potentiated or inhibited by the over-expression or inhibition of Cx43 respectively (Asazuma-Nakamura et al., 2009). This phenomenon was suggested to be due to the competitive binding between endogenous Smad2/3 and Cx43 to microtubules (Dai, Nakagami, Tanaka, Hitomi, & Takamatsu, 2007). Of note, connexin hemichannels Cx43 and Cx45 in cardiac fibroblasts are also involved in the extracellular release of ATP, which

activates the purinergic P2Y receptors, leading to the increased protein levels of α SMA, and increase in mRNA expression of PAI-1, and collagen (Lu, Soleymani, Madakshire, & Insel, 2012).

5.1.4 Kidney

Tubulointerstitial fibrosis is one of the hallmarks of chronic kidney disease (CKD) that is frequently diagnosed in diabetic patients (Cogliati et al., 2016). It is characterised by an mesenchymal transition of epithelial cells into myofibroblasts, leading to increases of ECM deposition, fibrotic scar formation and renal function deterioration (Hills & Squires, 2010). Indeed, high glucose concentrations-induced TGF- β 1 release from human proximal tubule cells are able to induce mesenchymal transition. This is characterised by gradual expression of mesenchymal markers (e.g. N-cadherin and vimentin) and loss of epithelial markers (e.g. E-cadherin) (Zeisberg & Kalluri, 2004; Zheng et al., 2009). Another hallmark of CKD is the excessive recruitment of monocytes. Of note, Cx43 expression is found to be up-regulated within the damaged compartments of the kidney during the development of CKD in both humans and rodents (Abed et al., 2014; Cogliati et al., 2016). Reduction of its expression in Cx43^{+/-} transgenic mice or through administration of Cx43asODN in CKD models demonstrated a significant reduction of cell adhesion markers, which was followed by reductions in monocyte infiltration and interstitial renal fibrosis (Abed et al., 2014). This evidence appears to suggest the importance of Cx43 in the development of this aetiology.

5.1.5 Connexin43 in post-surgical peritoneal adhesions

Findings from different fibrosis-related diseases appear to suggest that Cx43 is essential for fibroblast migration, proliferation, differentiation, and apoptosis, as well as for regulating the production of ECM (Asazuma-Nakamura et al., 2009; Lu et al., 2007; Mendoza-Naranjo et al., 2012; Mori et al., 2006; Qiu et al., 2003). These events are central to early adhesion pathogenesis and it is likely that Cx43 is involved in these cellular processes as well. During adhesion formation, the fibrin deposits facilitates the invasion and proliferation of fibroblasts (Arung et al., 2011). These recruited fibroblasts become activated and differentiate into myofibroblasts, and later deposit extracellular matrix, leading to the development of adhesions (Braun & Diamond, 2014). On this note, myofibroblasts may also arise from mesenchymal transition from endothelial and mesothelial/epithelial sources, and Cx43 has been shown to be able to regulate this process (Abed et al., 2014; Asazuma-Nakamura et al., 2009; Chilton et al., 2007; Uyama et al., 2003).

Apart from being shown to regulate fibroblast homeostasis, Cx43 has also been demonstrated to regulate inflammation and coagulation, which are also central processes during early adhesion pathogenesis. As mentioned earlier, reduction of its expression in Cx43^{+/-} transgenic mice or through administration of Cx43asODN in CKD models exhibited a significant reduction of cell adhesion markers, which was followed by reductions in monocyte infiltration and interstitial renal fibrosis (Abed et al., 2014). In addition, treatment of thrombin-stimulated endothelial cells with carbenoxolone, a broad-spectrum connexin inhibitor, or Cx43 siRNA-mediated knockdown, reduces thrombin-induced vascular hyperpermeability, a process that facilitates recruitment of leukocytes and the accumulation of fibrin-rich exudates

during adhesion formation (Lum & Malik, 1996; O'Donnell, Birukova, Beyer, & Birukov, 2014). In another example, TNF- α induced tissue factor expression in endothelial cells was attenuated with the intracellular transfer of monoclonal antibodies against Cx43 (Okamoto, Akita, Hayashi, Shimaoka, & Suzuki, 2014). Because Cx43 has a potential role in many facets of adhesion pathogenesis, the role of Cx43 in post-surgical peritoneal adhesion formation is explored in this chapter.

5.1.6 Connexin43 antisense oligodeoxynucleotides

Antisense oligodeoxynucleotides (asODN) are a powerful tool in the molecular biologist's arsenal to study the function of genes through the knock down of protein expression (Ma, Rede, Naqvi, & Cook, 2000). One of the main benefits of this method is their selective knockdown of the expression and hence, function of proteins. Apart from their capacity for rational design, the generation of these sequences is relatively cheaper compared to maintaining a colony of transgenic mice gene knockouts.

Under general conditions, RNA polymerase II is responsible for the transcription of genes from the start site in the DNA, giving rise to heteronuclear RNA products. Subsequently, these RNA products undergo processing to become mature mRNA, which is subsequently translated into proteins after being trafficked into the cytoplasm. For the asODN to exert its mechanism of inhibition, the sequences are designed to be reverse complementary (antisense) to their intended target sense RNA sequence, allowing the two strands to form a RNA-DNA duplex. These sequences are usually synthesised in lengths of 13 to 35 nucleotides long, since at least 12 basepairs are estimated to be required for the formation of a hybrid with a phosphodiester backbone that is stable at physiological conditions (Mahato, Cheng,

& Guntaka, 2005). In general, there are two main mechanisms that asODN could inhibit gene expression: through steric hinderance of the processing of pre-mRNA and/or mRNA sequences that designated for processing or translation; and through activation of RNaseH-mediated degradation of the mRNA (Rayburn & Zhang, 2008).

The Cx43asODN used in this study is a 30 base long unmodified phosphodiester asODN sequence (refer to **Appendix II, Supplementary Table 3**) that has previously been demonstrated to specifically knockdown Cx43 expression via the RNaseH mechanism (Law, Zhang, Stott, Becker, & Green, 2006). Treatment of this sequence in tissues has resulted in notable biological effects, particularly in chick embryogenesis (Becker et al., 1999), accelerating the healing of normal (Mori et al., 2006; Qiu et al., 2003) and diabetic wounds (Wang, Lincoln, Cook, & Becker, 2007), reducing inflammation and improving recovery from spinal cord injury (Cronin, Anderson, Cook, Green, & Becker, 2008). In these studies, Cx43asODN is usually delivered in a 30% (w/v) FDA-approved Pluronic® F-127 poloxamer gel that is a liquid at lower temperatures (4°C) and sets into a gel at physiological temperature. Apart from acting as a weak surfactant that facilitates the cellular uptake of asODN, it also thought to be a local depot for the supply of asODN to the intended site of treatment (Green, Law, Lin, & Becker, 2001; Law et al., 2006). Penetration of at least 10 µm into the corneal epithelium was shown after 1 hour using fluorescent-tagged ODNs and Pluronic® F-127 gel in this mode of delivery (Law et al., 2006). In addition, no toxic side effects have been reported to date, making it a safe and well-established technique to temporally and selectively knockdown Cx43 levels for the study of its function in various biological contexts.

The choice of control treatment used in this study was a similar treatment regime with DI water instead of Cx43 asODN delivered in the same respective concentrations of Pluronic® F-127. Since the amount of DI water is little, most of the treatment for control can be considered Pluronic gel, which is FDA-approved for usage in humans (Diniz et al., 2015). Many controls for asODN exist (as reviewed by (Stein, 2001)). This includes the use of positive controls whereby downregulation is expected to show dose-dependency; and negative controls (i.e. scrambled asODN) is used to rule out non-sequence-related effects. Although the administration of a scrambled, non-functional asODN would be a useful negative control, ‘there is no evidence that scrambled oligonucleotides provide superior controls’ (Stein, 2001). In this work, a vehicle control is chosen instead to be used as negative control since scrambled controls, used at the same concentration, may be capable of inducing non-specific effects (Anselmet et al., 2002; Branch, 1998; Stein, 2001). On this note, scrambled controls are also seldom used in human clinical trials in the history of FDA-approved oligonucleotide therapies. While positive and negative controls are ideal, these options can quickly become expensive. In the light of limited funding for this work, the choice to use the vehicle as a control is also a more economical approach for gathering the first line of evidence of biological significance before the experiments may be repeated in the future with more rigorous controls upon the secure of more funding.

5.2 Aims and Objectives

Work in this chapter aims to explore the role of Cx43 in post-surgical peritoneal adhesion formation. This would be achieved by first determining whether or not Cx43 is implicated during adhesion formation. Potential pathological changes to

Cx43 protein levels during early phase adhesion formation (6 and 24 hours post-surgery) are assessed via immunohistochemistry. Next, the significance of the contribution of Cx43 levels during early adhesion formation would be investigated by treating an experimental group of mice with Cx43 antisense oligodeoxynucleotides delivered with two formulations of Pluronic® F-127 immediately post-creation of IBs. In the first formulation, 25 µL of 300 µM Cx43asODN is delivered in 30% (w/v) Pluronic® F-127 to each of the 4 ischaemic buttons. At this concentration, the gel sets onto the button upon application, allowing prolonged release until the breakdown of the gel, ensuring the button receives the treatment. This is further complemented with the subsequent formulation that is given immediately after the first. This subsequent formulation ensures further coverage of the peritoneal cavity through the delivery of additional 400 µL of 300 µM of Cx43 asODN in 20% (w/v) Pluronic® F-127 to the peritoneal cavity. Significance of the Cx43asODN-mediated knockdown of Cx43 protein levels would be determined through evaluations at both early and late phase (7 days post-surgery) of adhesion formation. Effects on late phase adhesion formation would be evaluated using macroscopic assessment based on established scoring matrices that would be later verified with microscopic assessment and further validated with measurement of the overall plasminogen-activating activity in the peritoneal cavity. Effects on early phase would be assessed in the context of leukocyte recruitment and TGF-β1 production to shed light on potential mechanisms of which Cx43 may have affected adhesion formation.

5.3 Materials and Methods

5.3.1 In vivo Work

5.3.1.1 *Experimental animals*

All animal procedures were performed in accordance with the animal protocols approved by the Institutional Animal Care and Use Committee of the Animal Research Facility of Nanyang Technological University, protocol number A0372. Mice were bred under specific pathogen-free conditions under a 12-hour light-dark cycle. Standard rodent pellet diet and water were supplied *ad libitum*. Only age-matched (7 to 12 weeks) males were used for experiments.

5.3.1.2 *Ischaemic button model of post-surgical peritoneal adhesions*

Ischaemic buttons (IB) were created to induce adhesion formation as previously described by Sandoval et al., (2016) with minor modifications. Prior to the procedure, fur around the incision site was removed using an electric trimmer and depilatory cream before returning the mice to their cages and allowing them to rest for at least 24 hours. Briefly, mice were anaesthetised with isoflurane and placed on a warm heating mat throughout the surgery. All surgeons used non-powdered surgical gloves during the surgery and all instruments were sterilised. Next, the midline incision site was sterilised with 7.5% (v/v) providone-iodine solution (**Fig. 5.1a**). A midline laparotomy of 4 cm was performed to facilitate the exposure of the parietal peritoneum (**Fig. 5.1b**). Each ischaemic button was created by placing a 4/0 non-absorbable silk suture at the base of ~3 mm of the sidewall parietal peritoneum, which was grasped with a needle holder. Two ischaemic buttons, spaced at least 1

cm apart, were created on the peritoneum wall of each side of the midline (**Fig. 5.1c**). Depending on study design, vehicle or treatment were added (detailed in the next subsection) before surgery closure. The laparotomy site was then closed in two layers, the inner layer with simple interrupted polyglycolic acid absorbable 6.0 sutures (**Fig. 5.1d**) and the outer layer using metallic wound clips (**Fig. 5.1e**). After closure, mice were regularly monitored until euthanasia with CO₂ at 6 and 24 hours post-surgery for early phase adhesion formation assessment and at 7 days post-surgery for late phase formation assessment. For post-recovery, the mice received buprenorphine (0.1 mL per 10 g of animal weight) into the loose skin over the interscapular region.

Figure 5.1 Major steps during various adhesion-inducing operations in mice.

(A) Disinfection of the incision site after shaving. This was followed by a (B) midline incision of the skin. (C) Ischaemic button (IB) model. Two IBs, spaced approximately 1 cm apart, were created on the peritoneum on each side of the midline incision. Each button (yellow arrowheads) was created by placing a silk ligature at the base of 3 mm of parietal tissue that is grasped with a needle holder. (D) Wound closure by applying simple interrupted sutures to the peritoneal musculature and (E) metallic clips to the skin.

5.3.1.3 Studying the role of Cx43 during peritoneal adhesion formation

Mouse Cx43 antisense oligodeoxynucleotides (asODN) (refer to **Appendix II, Supplementary Table 3** for sequence) was used in an IB model of peritoneal adhesion to investigate the role of Cx43 in adhesion formation. Surgery was performed as described earlier. Prior to surgery closure, each mouse received a two-

formulation treatment only after all the buttons were created for that mouse. In the first formulation, 25 μ L of 300 μ M of Cx43 asODN in 30% Pluronic[®] F-127 in DI H₂O was delivered to each of the 4 buttons. To ensure further coverage of the peritoneal cavity, an additional 400 μ L of 300 μ M of Cx43 asODN in 20% Pluronic[®] F-127 in DI H₂O was delivered immediately to the left and right flank of the peritoneal cavity. For control, a vehicle group received the same treatment regime with DI water instead of Cx43 asODN delivered in the same respective concentrations of Pluronic[®] F-127. After the respective treatments were administered, surgery closure and post-recovery regime was performed as described earlier. Mice were regularly monitored until euthanasia at various time points of interest (e.g. 6, 24 hours, and 7 days post-surgery).

5.3.1.4 Peritoneal lavage

Prior to necropsy, 3 mL of cold phosphate-buffered saline (PBS) was injected into the peritoneal cavity and the abdomen was gently agitated for 1 min. Approximately 2 mL of the fluid was recovered. Samples with blood contamination were omitted. Cellular debris was removed by microcentrifugation at 9000 g for 15 min at 4°C, before freezing the supernatant in 250 μ L aliquots at -80°C.

5.3.1.5 Peritoneal fluid tPA and PAI-1 total antigen ELISA

Thawed aliquots of the peritoneal lavage were analysed for the total antigen concentration of tPA and PAI-1 present, according to manufacturer's instructions (Molecular Innovations, USA). Samples that had coefficient of variation values more than 15% were excluded. Detected concentrations were normalised using total

protein content determined from a bicinchoninic acid assay performed on the same aliquot (ThermoFisher Scientific, USA).

5.3.1.6 Macroscopic assessment of peritoneal adhesions

An inversed U-shaped incision was made across the peritoneal cavity for necropsy and the adhesions in each animal were evaluated according to established scoring matrices. These matrices assessed peritoneal adhesions based on qualities such as, extent, involvement, severity, and strength (refer to **Table 5.1**) (Adhesion Scoring Group, 1994; Cassidy et al., 2015; Hoffmann et al., 2009; Nair et al., 1974). Three observers blinded to the experimental groups scored each mouse and their scores were later averaged to give a single score per mouse. Inter-observer scores did not differ more than 2. Scores are represented as a percentage of maximum score possible for each parameter (e.g. if 2 out of 4 buttons were found to have adhesions, a score of 50% will be awarded to the mouse for the extent of adhesion. After which, the mean parameter scores for each of the groups were calculated.

Table 5.1 Scoring matrix for assessing peritoneal adhesions

Variable	Score	Description
Extent (Cassidy et al., 2015)	0	Complete absence of adhesions
	1	Only 1 button has an adhesion band
	2	2 buttons have adhesion bands
	3	3 buttons have adhesion bands
	4	4 buttons have adhesion bands
Involvement (Quantity) (Nair et al., 1974)	0	Complete absence of adhesions
	1	1 adhesion band between the organs or between one organ and abdominal wall
	2	2 adhesion bands between the organs or between one organ and abdominal wall
	3	>2 adhesion bands between the organs or between one organ and abdominal wall or adhesions of intestinal loops without any adhesion to the abdominal wall. Significant adhesion.
	4	Adhesion of all viscera to the abdominal wall
Severity (Quality) (Adhesion Scoring Group, 1994)	0	Complete absence of adhesions
	1	Filmy, avascular
	2	Dense, opaque and/or vascular
	3	Very thick cohesive attachment
Strength (Hoffmann et al., 2009)	0	Complete absence of adhesions
	1	Gentle traction requiring little effort to break adhesion
	2	Blunt dissection requiring a blunt instrument to break adhesion
	3	Sharp dissection requiring a sharp instrument to break adhesion
	4	Adhesion of all viscera to the abdominal wall

5.3.2 Histology

5.3.2.1 Tissue processing and microtome sectioning

Tissue samples are collected and stored in 4% (w/v) paraformaldehyde at 4°C overnight. On the following day, the tissues were transferred into 70% (v/v) ethanol and stored at 4°C overnight. After, tissues were processed using the HistoCore PEARL (Leica, Germany) tissue processor and embedded in paraffin with the HistoCore Arcadia C (Leica, Germany). Following, tissue samples were sectioned at 5 µm thickness using a Leica RM2245 microtome (Leica, Germany) and mounted

onto Polysine[®] microscope slides and dried at 40°C for at least an hour before staining.

5.3.2.2 Haematoxylin and eosin (H&E) staining

Slides were loaded into Leica Autostainer XL (Leica, Germany) and a set programme was run (refer to **Appendix II, Supplementary Table 4**). At the end of the programme, glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at room temperature (RT).

5.3.2.3 Picro-Sirius red (PSR) staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and DI water. Slides were then immersed in Picro-Sirius red (Abcam, United Kingdom) for 1 hour. After, slides were transferred into two changes of acetic acid followed by two changes of 100% (v/v) ethanol. Slides were cleared again with two changes of Clearene (Leica, Germany) before glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at RT.

5.3.2.4 Immunofluorescence staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and PBS. Depending on the antibody, either heat-induced or proteolytic-induced antigen retrieval methods may be performed (refer to below sections). If no antigen retrieval was required, slides were then immersed in permeabilisation/blocking buffer (0.3% (v/v) Triton-X100, 1M Lysine in PBS) for 1 hour at RT before incubation in antibody buffer (1% (w/v) BSA in PBS) for 45 minutes. After incubation, the boundaries of

the tissue samples were dried and a hydrophobic perimeter was drawn around the sample with a hydrophobic pen (Agilent Technologies, United States). Tissue samples were then immunolabeled with primary antibody diluted in blocking solution (refer to **Table 5.2** for dilutions and incubation conditions regarding primary antibodies) and incubated overnight at 4°C. Controls without primary antibodies were performed similarly with antibody buffer in the absence of any primary antibody. After primary antibody incubation, tissues were washed twice, 3 minutes each, using washing buffer (0.05% (v/v) Tween-20 in PBS) before 1 hour incubation at RT with the appropriate secondary antibody (refer to **Table 5.3** for dilutions regarding secondary antibodies). After incubation, tissues were washed twice, 3 minutes each, with washing buffer before staining with DAPI (Life Technologies, USA) diluted at 1 in 10000 with antibody buffer. Finally, slides were washed twice, 3 minutes each, in PBS prior to applying Citiflour™ AF1 (Electron Microscopy Sciences, London) anti-fade mounting medium before mounting coverslips and sealing with nail varnish.

5.3.2.5 Heat-induced antigen retrieval (HIAR)

After rehydration, slides were submerged in Tris-EDTA antigen retrieval solution (10 mM Tris Base, 1 mM EDTA, 0.05% (v/v) Tween 20, pH 9.0) and incubated at 60°C for 30 minutes. Slides were rested in the same solution for 5 minutes at RT. After, slides were immersed in antibody buffer and the remainder of standard staining protocol was performed as described earlier.

Table 5.2 List of primary antibodies used in this study

Name	Protocol	Reactivity	Dilution	Catalogue No.
Rabbit polyclonal anti-Cx43	IHC	Cx43	1:1000	C6219 (Sigma, USA)
Rabbit polyclonal anti- α SMA	IHC	α SMA	1:500	AB5694 (Abcam, UK)
Rabbit monoclonal anti-TGF- β 1	IHC + HIAR	TGF- β 1	1:400	AB215715 (Abcam, UK)

IHC – Immunohistochemistry staining, **HIAR** – Heat induced antigen retrieval.

Table 5.3 List of secondary antibodies used in this study

Name	Reactivity	Dilution	Catalogue No.
Goat anti-rabbit AF488	Rabbit	1:500	A11008 (ThermoFischer Scientific, USA)
Goat anti-rabbit AF555	Rabbit	1:500	A21422 (ThermoFischer Scientific, USA)

5.3.3 Brightfield and confocal microscopy

Histological stained (H&E and PSR) tissue samples were automatically scanned and digitised into 24-bit images with an AxioScan.Z1 slide scanner (Zeiss, Germany) equipped with a 20X 0.8NA Plan-Apochromat air objective and a Hitachi HV-F202SCL. Scaling per pixel for these images is 0.22 μ m by 0.22 μ m. For the visualisation of immunohistochemical stained tissue samples, Z-stacks of 10 optical sections were acquired using an upright confocal microscope TCS SP8 (Leica, Germany) with a 40X 1.2NA Plan-Apochromat. Z-stacks were taken by first finding the centre position before proceeding to take slices above and below this centre point. The thickness of each Z is 0.699 μ m. Fluorophores were excited sequentially using a 405 nm, 488 nm and 532 nm wavelength laser. The images were acquired at 8-bit greyscale at 1024 x 1024 pixels with a Z-series frame average of 2 or 3. Maximum projections were calculated using Leica Application Suite X (Leica,

Germany) for each of the Z-stacks recorded. All images were captured with identical parameters and camera settings to allow for direct comparison of staining intensity.

5.3.4 Microscopic Assessment

5.3.4.1 Inclusion and exclusion criteria

Tissues samples were first subjected to a simple inclusion/exclusion criterion to identify samples with features that allowed fair comparison with one another. The inclusion criterion included only tissues sections with a visible button (refer to **Fig. 5.2a**) or with a visible button accompanied with sutures only at the base of the button (refer to **Fig. 5.2b**). The exclusion criterion rejected tissues sections with no visible button (refer to **Fig. 5.2c**) or with a visible button accompanied with sutures found at other locations other than at the base of the button (refer to **Fig. 5.2d**).

Figure 5.2 Inclusion and exclusion criteria

The inclusion criterion included only tissues sections with a visible button (A) or with a visible button accompanied with sutures only at the base of the button (B). The exclusion criterion rejected tissues sections with no visible button (C) or with a visible button accompanied with sutures found at other locations other than at the base of the button (D). **IB** – Ischaemic Button region, **S** – Suture region.

5.3.4.2 Extent of inflammation

For scoring the extent of inflammation, each IB sample was first divided into two regions, referred to as the “Within IB” and “Outside IB” regions (refer to **Fig. 5.3**). The “Within IB” region was defined as the area spanning 500 μm away from the periphery of the button. The “Outside IB” region was defined as the area encompassing 1500 μm away from both left and right flanks of the perimeter of the “Within IB” region. In each region, seven region-of-interests (ROIs), 62,500 μm^2 each, were evenly and objectively distributed along the mesothelial edge of the tissue. Each ROI was checked to ensure that more than half of the sampled area was covered by tissue containing both the mesothelium and the underlying muscle layer. Next, three observers blinded to the experimental groups scored the extent of inflammation in each ROI on a 5-point scale based on a scoring matrix described in **Figure 5.4**. Scores from the three observers were then averaged. Scores between observers did not differ more than 1.

Figure 5.3 Designating ROIs for scoring the extent of inflammation

Seven ROIs (black squares) were taken each from “Within IB” and “Outside IB” regions. The “Within IB” region was defined as the area spanning 500 μm away from the periphery of the button (green box). The “Outside IB” region was defined as the area encompassing 1500 μm away from both left and right flanks of the perimeter of the “Within IB” region (blue box). **IB** – Ischaemic Button region, **S** – Suture region.

Figure 5.4 Scoring matrix for grading extent of inflammation

Regions-of-interests were graded for the extent of inflammation based the above scoring matrix. Three representative images accompany the description of each grade, detailed in the following. **Grade 0**, little/no recruitment of leukocytes to the edge or muscle area; **Grade 1**, some recruitment of leukocytes to the edge but little/none to the muscle area; **Grade 2**, moderate recruitment of leukocytes to the edge and/or to the muscle area (no focal accumulation of leukocyte was observed); **Grade 3**, large recruitment of leukocyte to the edge and/or to muscle area (some focal accumulation of leukocytes was observed); **Grade 4**, extensive recruitment of leukocyte, which is limited to a particular region.

5.3.4.3 Fibrosis extent

During adhesion formation, fibrin deposits that persist allow the invasion and proliferation of fibroblasts, leading to the deposition of extracellular matrix proteins (e.g. collagen). As the adhesion develops and matures, the matrix becomes increasingly filled with more extracellular matrix proteins over time, eventually transforming the matrix into a more permanent fibrous connection. At an earlier stage (e.g. 7 days as opposed to 21 days) during adhesion development, the entire matrix may not necessarily be “filled” up with collagen, or other matrix proteins. The assessment of the extent of the fibrotic area (or matrix) is a good measure of area of the matured adhesion. In order to facilitate this analysis, several regions of

the IB sample were further identified. The entire fibrotic (EF) region was identified as the overall region that included the immediate fibrotic (IF) region and the regions of adhesion-inducing determinants, such as ischaemic button and suture (IBS) (refer to **Fig. 5.5**). The EF region was defined by region occupied by the abnormal encroaching of extracellular matrix that is observed across the entire ischaemic button tissue region and is bordered at the base of the ischaemic button by thickened mesothelium. The IBS region includes the ischaemic button region and the suture material region. The ischaemic button can be defined by the “ball of muscle” (due to the out-pouching) and can be distinguished from the matrix due to its pinkish characteristic. The suture material regions are determined by the presence of the suture material (greyish brown). The IF region is of the main interest and can be calculated by subtracting the area of the IBS region from the calculated area of the EF region. Note that these terms are also used in the following sections when determining fibrosis severity and protein levels of α SMA. Two ROIs were drawn on digital stitched images of tissue sections stained with H&E. One was drawn around the EF region and the other, around the IBS region. Areas of these regions were measured using image analysis software Zen Lite 2.3 (Zeiss, Germany) and used to calculate the area of IF region by subtracting the IBS region from EF region. Since the size of IBS region may influence the size of the IF region, the calculated area of the IF region is normalised to the area of IBS region to ensure fair comparison between the samples.

Figure 5.5 Defining regions around the ischaemic button

Representative digital stitched image of a section of the IB sample stained with H&E. The entire fibrotic (EF) region was identified as the total fibrotic region (area encompassed by red dashed lines). This is made up of the immediate fibrotic (IF) region and the regions of adhesion-inducing determinants such as the ischemic button and suture (IBS) (black dashed lines). The immediate fibrotic region (IF) may be determined by drawing ROIs to exclude the FR region from the IBS region or by the subtracting the FR region from IBS region. **B** – Ischaemic Button region, **S** – Suture region.

5.3.4.4 Fibrosis severity

ROIs were first drawn around the IF regions of the digital stitched images of tissue sections stained with PSR using ImageJ analysis software (version 1.51w; NIH) (**Fig. 5.6b**). These ROIs were saved for later use and documentation. Next, the amounts of collagen deposition were estimated using the greyscale threshold method described by Schipke et al., (2017) using ImageJ analysis software. Briefly, conversion of the acquired images to the RGB stacks generated greyscale images for red, green, and blue channels. Manual threshold was performed on the greyscale image corresponding to the green channel to detect collagen. Specific threshold values (80 to 255) were determined using at least 10 sections to ensure values were representative and then kept constant throughout image analysis. Next, an output mask is generated through manual thresholding at determined values (**Fig. 5.6c**).

The corresponding ROIs were loaded onto the respective output masks and the detected area was measured using the measure tool (**Fig. 5.6d**) and later normalised to the IBS area.

Figure 5.6 Example of image analysis workflow for PSR stained images.

(A) Example of PSR stained image to be analysed. (B) ROIs were first drawn around the IF regions of the digital stitched images of tissue sections stained with PSR using ImageJ. Next, the stained image to be analysed were converted to the RGB stacks, which generates greyscale images for red, green, and blue channels. Manual threshold was performed on the greyscale image corresponding to the green channel to detect collagen, which would produce an output mask. (C) Example of output mask after manual threshold (80 to 255). Specific threshold values were kept constant throughout analysis. (D) Example of the loading of corresponding ROIs for the respective sections so that the measure tool can be used to detect the area of interest in ImageJ.

5.3.4.5 Protein levels of α SMA

ROIs were drawn around the IF region in the digital stitched images of tissue sections immunostained with α SMA. Areas of blood vessels in which strong α SMA staining can be found, were excluded from the ROIs. Protein levels of α SMA in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the α SMA levels were determined by quantifying the area in the binary mask that is generated after performing manual threshold. Specific threshold values (80 to 255) were kept constant throughout image analysis. The detected area was measured using the measure tool and later normalised to the IBS area.

5.3.4.6 Protein levels of Cx43 (ischaemic buttons)

Tissue sections were immunostained with Cx43 and five fields-of-view were each acquired from regions within and away from the ischaemic button. From the acquired images, ROIs were drawn around the mesothelial and muscle regions. Protein levels of Cx43 in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the Cx43 levels were determined by quantifying the area in the binary mask generated after performing manual threshold. Specific threshold values were kept constant throughout image analysis (100 to 255). The measured Cx43 pixel area was then normalised to the pixel area of the ROI measured.

5.3.4.7 Protein levels of TGF- β 1 (ischaemic buttons)

Tissue sections were immunostained with TGF- β 1 and five fields-of-view were each acquired from regions within the ischaemic button. From the acquired images, ROIs were drawn around the muscle regions. Protein levels of TGF- β 1 in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the TGF- β 1 levels were determined by quantifying the area in the binary mask generated after performing manual threshold. Specific threshold values were kept constant throughout image analysis (90 to 255). The measured TGF- β 1 pixel area was then normalised to the pixel area of the ROI measured.

5.3.5 Statistical Analysis

All data is presented as mean value $\pm$ standard deviation. In general, all statistical comparisons made between vehicle and Cx43asODN treatment groups were made using a Mann-Whitney *U* test, unless mentioned otherwise. A two-way analysis of variance (ANOVA) followed by a post-hoc Tukey's multiple comparisons test was used for analysing the protein levels of Cx43 or the extent of inflammation in regions within and outside the IB regions at 6 and 24 hours post-surgery between vehicle and Cx43asODN treatment groups. All statistical comparisons were made using GraphPad Prism version 6.0f for Mac, (GraphPadTM Software, San Diego, USA).

5.4 Results

5.4.1 Adhesions were observed as early as 6 hours post-surgery during early phase adhesion formation

Previously, the ischaemic button (IB) model was determined to be the most ideal model for studying the role of Cx43 in post-surgical peritoneal adhesion formation. The ischaemic button (IB) model was further characterised after determining that it was the most ideal model for studying the role of Cx43 in post-surgical peritoneal adhesion formation in the previous chapter. Briefly, this model involves creating two IBs on each side of the midline incision. Each IB was created through a suture ligation of an out-pouching of the abdominal sidewall. Three windows of assessments, covering early and late phases of adhesiogenesis, were chosen for characterisation. Since an early window (i.e. less than 24 hours) is useful for understanding adhesion pathogenesis (Ryan et al., 1971), two time points (6 and 24 hours post-surgery, n=5 each) were chosen to provide coverage of early events occurring during the early phase of adhesiogenesis. A later window of assessment was performed at 7 days post-surgery (n=9). This was chosen to evaluate intervention efficacy since there is general agreement that adhesion formation is completed by 7 to 10 days post-surgery (Beyene et al., 2015; Gaertner et al., 2008; Holmdahl et al., 1994).

Peritoneal adhesions were detected as early as 6 hours post-surgery. By scoring the number of IBs found to have attached adhesive band, the mean extent of adhesions (refer to **Table 5.1**) at this time point was assessed and found to be $15 \pm 13.7\%$. These rare adhesions were immature in nature, being filmy and avascular (**Fig. 5.7a**).

In addition, the adhesions also broke easily with gentle traction. At 24 hours post-surgery, the mean extent of adhesions was detected to be two-folds more ($30 \pm 20.9\%$) of those detected at 6 hours post-surgery. These adhesions were slightly more mature, as they were observed to be slightly denser and less filmy than those detected at 6 hours post-surgery (**Fig. 5.7b**). However, these adhesions were still avascular and could be broken with gentle traction. In contrast, adhesions detected at 7 days post-surgery were markedly more mature than those detected during the early phase of adhesiogenesis. The majority of these attachments was denser, stronger, and generally required blunt (**Fig. 5.7d and e**) or sharp dissection (**Fig. 5.7f and g**) for breaking the adhesions. These late phase adhesions were also characterised by a more opaque and vascular appearance. As expected, these late phase adhesions were more extensive than early phase adhesions, with a mean extent score of $55.56 \pm 27.3\%$ (**Fig. 5.7c**). Interestingly, there appeared to be a difference in extent of adhesions ($p=0.0596$) in the cranial and caudal abdominal regions. Detected adhesions were mainly distributed to the caudal abdominal regions (65%), with most being attached to the epididymal fat pad (**Fig. 5.7f, g, and i**). Since the location of the IB appeared to influence the quantity of adhesion created, future analyses were performed on the IBs in the caudal abdominal region unless mentioned otherwise.

Figure 5.7 Types of adhesions observed at early and late phases of adhesion development

The panel shows representative images of adhesions (highlighted with green arrowheads) observed during early phases (6 and 24 hours, $n=5$ each) and late phase (7 days ($n=9$)) of adhesion development. (A) A filmy, avascular adhesion between the IB and the epididymal fat pad, detected at 6 hours post-surgery. These filmy adhesions were avascular and required gentle traction to break. (B) At 24 hours post-surgery, a denser, less filmy, avascular adhesion between the IB and the mesenteric fat was detected. (C) Extent of adhesions observed at early and late phase of adhesion development. (D-G) Adhesions detected at 7 days were markedly more matured. (D and E) Examples of opaque and vascular adhesive bands requiring blunt dissection to break, detected between the IB and (D) a liver lobe or (E) a small bowel loop at 7 days. (F and G) Examples of dense, opaque, and vascular adhesive bands requiring sharp dissection to break, detected between the IB and epididymal fat pad. Adhesive band detected in (G) was thicker and a more cohesive attachment compared to (F). (H) Majority of the adhesions at 7 days appeared to be mainly distributed in the lower abdominal region. (I) Fat pad distribution in the left abdominal region. Data is presented as mean value $\pm$ standard deviation. *RP* – Retroperitoneal fat pad; *E* – Epididymal fat pad

5.4.2 Significant upregulation of Cx43 at the IB during early adhesion formation.

Next, to investigate if there were any pathological changes to Cx43 protein levels occurring during early phase adhesiogenesis, tissue samples from the IB model and control sham-operated mice from 0, 6, and 24 hours post-surgery were immunostained for Cx43 (**Fig. 5.8a**). Cx43 protein levels in sham-operated mice from 0 hour post-surgery provided reference for levels at basal physiological state. Image analyses of the immunostained tissues were performed separately in the muscle and mesothelial regions.

At 6 hours post-surgery in the muscle regions within and outside IB, Cx43 protein levels increased 6.3- and 8.1- fold ($p \leq 0.01$) respectively from the basal physiological state (**Fig. 5.8b**). In contrast, a more dramatic increase in Cx43 protein levels was observed in the mesothelial regions. In these regions, at least a 21-fold increase from basal physiological state was detected within ($p \leq 0.05$) and outside IB ($p \leq 0.01$) (**Fig. 5.8c**). At 24 hours post-surgery, a slight 1.1-fold decrease in Cx43 protein levels from 6 hours post-surgery was observed in both muscle and mesothelial regions outside the IB. Despite this decrease, Cx43 protein levels remained significantly higher compared to basal physiological levels ($p \leq 0.05$). While no appreciable changes in Cx43 protein levels were observed in the muscle regions within the IB, a further increase of 1.4-fold in Cx43 protein levels in the mesothelial region within the IB was observed. When compared to basal physiological levels, this appeared to be the most significant increase of Cx43 levels ($p \leq 0.0001$). Overall, these results suggested that pathological increases of Cx43 protein levels were evident during

early phase adhesion formation, taking place mainly in the mesothelial regions within the IB.

Figure 5.8 Cx43 was significantly upregulated at the IB regions during early adhesion formation

(A) The panel shows representative immunofluorescence images of regions within and outside the IB samples at 0, 6, and 24 hours post-surgery (n=4, 5, 5, respectively). Longitudinal sections of IB sections were immunolabeled for Cx43 (red), and counterstained with DAPI (blue). IB samples at 0 hours (top two panels) revealed basal levels of Cx43 protein levels. At 6 hours (middle two panels), Cx43 levels appeared to be upregulated in both the mesothelial and muscle regions and were maintained for at least 24 hours post-surgery (bottom two panels). White dashed lines separate the mesothelial (upper) and muscle (lower) regions. Magnification used, 40X. Scale bar = 50 μ m. (B) Image analysis in the muscle regions outside the IB revealed overall higher Cx43 levels at 6- and 24-hours post-surgery. In contrast, no appreciable changes in Cx43 levels were observed in the muscle regions within the IB at 6 and 24 hours post-surgery. (C) Image analysis revealed significantly higher Cx43 levels in the mesothelial regions both within and outside the IB compared to basal physiological levels in 0 hour sham-operated controls. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a two-way ANOVA followed by a post-hoc Tukey's multiple comparisons test (* $p \leq 0.05$, ** $p \leq 0.01$, **** $p \leq 0.0001$).

5.4.3 Significant increase in extent of inflammation at the IB during early adhesion formation.

To investigate the extent of inflammation during early phase adhesiogenesis, tissue samples from the IB model and control sham-operated mice from 0, 6, and 24 hours post-surgery were stained with hematoxylin and eosin (H&E) histological stains (**Fig. 5.9a**). Ischaemic button regions with a higher degree of recruited leukocytes and microabscess received a higher score for the extent of inflammation by observers blinded to the experimental groups. Differences between the observer scores were less than 1, suggesting shared consensus among observers (**Fig. 5.9b**).

At 6 hours post-surgery, moderate recruitment of leukocytes was detected at muscle regions proximal to the periphery and mesothelial regions outside the IB (1.83 ± 0.34), while a larger recruitment of leukocytes was observed within the IB (2.21 ± 0.36) (**Fig. 5.9c**). At 24 hours post-surgery, the leukocytes recruited to regions outside the IB appeared to have slightly resolved (1.66 ± 0.32) while a more significant decrease of leukocytes was observed within the IB (1.46 ± 0.25) ($p \leq 0.05$).

(Fig. 5.9c). Overall, the extent of inflammation in the IB samples at 6 and 24 hours post-surgery was significantly higher than basal physiological state (0.05 ± 0.03) ($p \leq 0.0001$) (Fig. 5.9c).

Figure 5.9 Extent of inflammation was significantly increased during early adhesion formation
 (A) Representative ROIs from digital stitched images of H&E-stained IB samples at 0, 6 and 24 hours post-surgery (n=4, 5, 5, respectively). At 6 hours post-surgery outside the IB region, moderate recruitment of leukocytes was observed mainly to the mesothelial and muscle regions closer to the

periphery. At 24 hours post-surgery in the same region, almost similar magnitude of recruitment was detected, but generally fewer leukocytes were noted at muscle regions further from periphery. At 6 hours post-surgery within the IB region, a larger recruitment of leukocytes was detected when compared to outside the IB region. This magnitude of recruitment was noted to have reduced at 24 hours post-surgery. Scale bar = 50 μ m. (B) Mean scores for each experimental group by each observer was graphed and revealed inter-observer variability to be less than 1. (C) The extent of inflammation within the IB region at 6 hours post-surgery appeared to be significantly reduced at 24 hours post-surgery. The overall extent of inflammation in regions of the IB at both 6 and 24 hours post-surgery was still significant compared to 0 hour sham-operated controls. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a two-way ANOVA followed by a post-hoc Tukey's multiple comparisons test (* $p \leq 0.05$, **** $p \leq 0.0001$).

5.4.4 Treatment with Cx43asODN significantly reduced pathological increases of Cx43 in the mesothelial region within the IB.

After observing a pathological increase of Cx43 in the mesothelial region of the IB during early adhesion formation, Cx43asODN treatment was explored to investigate the role of Cx43 during adhesion formation. A pilot run was performed with Cx43asODN tagged with AlexaFluor488[®] fluorophore to inform on the extent of Cx43asODN distribution within the IB sample at 6 and 24 hours post-surgery. Microscopic assessment of IB samples at 6 hours post-surgery found the majority of the AlexaFluor488[®]-tagged Cx43asODN as deposits within the mesothelial regions within the IB region (**Fig. 5.10a**). The AlexaFluor488[®]-tagged Cx43asODN was also detected in the peripheral muscle region and was noted to have penetrated at least 180 μ m into the muscle layer distal from the mesothelium. It may be also important to point out that a tinge of green from the fluorophore was noticed in the bladders during necropsy, suggesting a degree of clearance of the fluorophore. At 24 hours post-surgery, deposits of the AlexaFluor488[®]-tagged Cx43 asODN in the mesothelial was less frequently observed and mainly localised to the regions proximal to the mesothelial edge. In addition, Cx43 protein levels appeared to be generally lower at 24 hours than 6 hours post-surgery. These findings suggested that

the main site of action of the Cx43asODN was at mesothelial regions and the accompanying muscle proximal to these regions within the IB.

After determining the extent of distribution of Cx43asODN, a larger study of similar design was subsequently conducted, in which regular Cx43asODN-treated IB samples were harvested at 6 and 24 hours post-surgery (n=6 each). At 6 hours post-surgery, a slight but insignificant decrease in the Cx43 protein levels was observed in the muscle and mesothelial regions within and outside the IB when compared to their vehicle controls (**Fig. 5.10b**). At 24 hours post-surgery, a more prominent decrease of Cx43 levels was observed. At muscle regions within and away from the IB, at least a 3-fold decrease in Cx43 protein levels was observed ($p \leq 0.05$). At mesothelial regions away from the IB, a 2.6-fold decrease was noted ($p = 0.0519$). The most significant decrease was a 4.2-fold decrease in Cx43 protein levels ($p \leq 0.01$) observed in the mesothelial regions within the IB. In summary, these observations suggest that treatment with Cx43asODN was able to significantly reduce pathological increases of Cx43 protein levels within the IB.

Figure 5.10 Cx43asODN significantly reduced Cx43 protein levels in the regions within the IB at 24 hours post-surgery.

(A) The panel shows representative immunofluorescence images revealing the distribution of AlexaFluor488[®]-tagged Cx43asODN and Cx43 protein levels at regions within the treated IBs at 6

and 24 hours post-surgery (n=3, each). AlexaFluor488®-tagged Cx43asODN (green, highlighted with green arrowheads) were visualised on longitudinal sections of IB sections immunolabeled for Cx43 (red, highlighted with red arrows), and counterstained with DAPI (blue). At 6 hours post-surgery (left 3 panels), AlexaFluor488®-tagged Cx43asODN were generally detected at the mesothelial regions and were also noted to have penetrated at least 180 µm into the muscle layer distal from the mesothelium (top left panel). At 24 hours post-surgery (right 3 panels), the presence of AlexaFluor488®-tagged Cx43asODN was less frequently observed and was mostly detected proximal to the mesothelial edge. Cx43 protein levels in these Cx43asODN-treated samples appeared to be generally lower at 24 hours post-surgery. White dashed lines separate the mesothelial (upper) and muscle (lower) regions. Magnification used, 40X. Scale bar = 50 µm. **(B)** Image analysis revealed significantly lower Cx43 protein levels in the mesothelial and muscle regions within the IB in the Cx43asODN-treated group (n=6) compared to the vehicle-treated group (n=5). Data is presented as mean value ± standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test. (*p≤0.05, **p≤0.01)

5.4.5 Treatment with Cx43asODN significantly reduced adhesion strength and severity at 7 days post-surgery.

After determining that Cx43asODN treatment significantly knocked down the early pathological increases of Cx43 protein levels during early adhesion formation, its effect on adhesion formation at 7 days post-surgery was evaluated. Macroscopic evaluation of the adhesions was conducted in a blinded manner by 3 observers using established scoring matrices (refer to **Table 5.1**) (Adhesion Scoring Group, 1994; Hoffmann et al., 2009; Nair et al., 1974). First, the adhesion involvement was assessed (refer to **Table 5.1**), in which a higher score for involvement was assigned if more adhesive bands were found attached to the button (Nair et al., 1974). Treatment with Cx43asODN did not appear to affect the adhesion involvement (p=0.295) (**Fig. 5.11a**). Next, the detected adhesions were assessed for their severity (refer to **Table 5.1**). Based on severity scoring matrix (Adhesion Scoring Group, 1994), adhesive bands that are opaque, denser, and more vascular, would receive a higher score. The majority of the adhesions in the vehicle-treated group (53.0 ± 29.7) were more opaque and denser compared to the adhesions in the Cx43asODN-treated

group (26.1 ± 21.1) (**Fig. 5.11b**). Finally, the detected adhesions were evaluated for their strength (refer to **Table 5.1**). Adhesive bands with lower tenacity, requiring less effort to break, were awarded lower scores (Hoffmann et al., 2009). In the vehicle-treated group, almost all of the adhesions required blunt dissection to break (43.5 ± 24.0). In contrast, almost all of the adhesions in the Cx43asODN-treated group required little effort to break the adhesion, with most breaking under gentle or little traction (20.0 ± 15.7) (**Fig. 5.11c**). Overall, macroscopic assessment of the detected adhesions suggested that treatment with Cx43asODN appeared to significantly reduce adhesion severity ($p \leq 0.05$) and strength at 7 days post-surgery ($p \leq 0.05$).

Figure 5.11 Treatment with Cx43asODN significantly reduced severity and strength of adhesions developed at 7 days post-surgery.

(A) Examples of peritoneal adhesions detected in the lower abdominal region at 7 days post-surgery. These adhesions (highlighted with green arrowheads) were assessed for qualities such as, involvement (B), severity (C), and strength (D), and scored by 3 observers blinded to the Cx43asODN- (n=10) and vehicle- (n=9) treated groups. Both groups did not receive significantly different scores for involvement. The Cx43asODN-treated groups received significantly lower severity and strength scores for the adhesions detected compared to the vehicle-treated group. The mean observer scores for each IB in the lower abdominal region were averaged for each mouse and

graphed as a percentage of maximum score possible for each parameter. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test (* $p \leq 0.05$).

5.4.6 Treatment with Cx43asODN significantly reduced the fibrotic extent and severity of adhesions.

To further validate the macroscopic observations, the IB samples were stained with H&E and PSR for microscopic evaluation of the extent (**Fig. 5.12**) and severity of fibrosis (**Fig. 5.13**) respectively. Only the IBs in the caudal abdominal region were analysed per animal since the location of the IB appeared to influence the quantity of adhesion created as discussed in the earlier. To ensure fair comparison between IB samples in an unbiased manner, the paraffin-embedded IB samples were first subjected to an inclusion/exclusion criterion to identify samples with features that allowed fair comparison. The inclusion criterion included only tissue sections with a visible button (refer to **Fig. 5.2a**) or with a visible button accompanied with sutures only at the base of the button (refer to **Fig. 5.2b**). The exclusion criterion rejected tissue sections with no visible button (refer to **Fig. 5.2c**) or those with a visible button that was accompanied with sutures found at the other locations other than at the base of the button (refer to **Fig. 5.2d**). Microscopic evaluations were then performed on the sections from treatment ($n=5$) and vehicle groups ($n=7$) that allowed fair comparison.

To determine the extent of fibrosis, immediate fibrotic regions in H&E stained IB samples were measured through image analysis. The fibrosis extent in the Cx43asODN-treated group (0.14 ± 0.07) was significantly smaller compared to the vehicle-treated group (0.49 ± 0.2) ($p \leq 0.01$), suggesting that the Cx43asODN

treatment reduced the extent of fibrosis (**Fig. 5.12c**). Following, the severity of fibrosis within the immediate fibrotic regions was further accessed by measuring the amount of collagen fibres stained red with PSR. Image analysis revealed that the amount of collagen deposition within the immediate fibrotic regions of the Cx43asODN-treated group (0.55 ± 0.45) were significantly reduced compared to the vehicle-treated group (1.63 ± 1.5) ($p \leq 0.05$). This suggested that the fibrosis severity was significantly reduced with the treatment of Cx43asODN (**Fig. 5.13c**). Together, these observations of significant reduction in fibrotic extent and severity corroborate previous macroscopic observations of reduced strength and severity of the adhesions.

Figure 5.12 Treatment with Cx43asODN significantly reduced fibrosis extent of the adhesions developed at 7 days post-surgery.

Representative digital stitched images of H&E-stained IB samples of the vehicle- (A, n=7) and Cx43asODN-treated (B, n=5) groups at post-surgery day 7, and their respective magnified regions (i, ii). Scale bars 50 and 500 μ m. (C) Image analysis revealed that fibrotic extent of the adhesions of the Cx43asODN-treated group was significantly reduced when compared to the vehicle-treated group. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test (** $p \leq 0.01$). Dashed lines highlight boundary of the immediate fibrotic region. **IB** – ischaemic button; **s** – suture; **m** – muscle; **IF** – immediate fibrotic region.

Figure 5.13 Treatment with Cx43asODN significantly reduced fibrosis severity of the adhesions developed at 7 days post-surgery.

Representative digital stitched images of PSR-stained IB samples of the vehicle- (**A**, $n=7$) and Cx43asODN-treated (**B**, $n=5$) groups at post-surgery day 7, and their respective magnified regions (*i*, *ii*). Scale bars 50 and 500 μ m. (**C**) Image analysis revealed that fibrotic severity of the adhesions of the Cx43asODN-treated group was significantly reduced compared to the vehicle-treated group. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney U test ($*p \leq 0.05$). Dashed lines highlight boundary of the immediate fibrotic region. *IB* – ischaemic button; *s* – suture; *m* – muscle; *IF* – immediate fibrotic region.

5.4.7 Effect of Cx43asODN treatment on the plasminogen-activating activity at 7 days post-surgery.

Next, the protein levels of tissue plasminogen activator (tPA) and plasminogen-activator inhibitor (PAI)-1 in the peritoneal cavity were investigated. This was

achieved by measuring their respective levels in the peritoneal fluid at 7 days post-surgery using commercial ELISA detection kits (**Fig. 5.14**). Following peritoneal injury, tPA plays an essential role in limiting adhesion formation. It does so by converting the inactive substrate plasminogen into its active form plasmin, a serine protease that dissolves fibrin (Beyene et al., 2015). In the peritoneal cavity, 95% of the plasminogen-activating activity (PAA) is contributed by tPA (Holmdahl et al., 1996). However, the ability of tPA in mounting an effective fibrinolytic response also depends on the presence of PAIs. By antagonising and forming inactive complexes with tPA, PAI-1 hampers the PAA (Beyene et al., 2015). Under pathological conditions, the PAA is markedly decreased, leading to the development of adhesions (**Fig. 1.1**) (Arung et al., 2011; Braun & Diamond, 2014). Treatment with Cx43asODN did not appear to affect the protein level of tPA in the peritoneal fluid ($p=0.263$) (**Fig. 5.14a**). In contrast, the protein level of PAI-1 in the peritoneal fluid of the Cx43asODN treated-group was significantly lower than the vehicle-treated group ($p\leq 0.01$) (**Fig. 5.14b**). This appears to suggest higher PAA activity in the Cx43asODN-treated group.

Figure 5.14 Effect of Cx43asODN treatment on tissue plasminogen activator (tPA) and plasminogen activator inhibitor-1 (PAI) proteins in the peritoneal fluid at day 7 post-surgery. (A) tPA protein levels at day 7 post-surgery were unchanged by Cx43asODN treatment ($n=7$). (B) Treatment with Cx43asODN significantly reduced PAI-1 protein levels at day 7 post-surgery when compared to vehicle-treated group. Statistical comparisons were made using a Mann-Whitney U test (** $p\leq 0.01$).

5.4.8 Treatment with Cx43asODN significantly reduced the degree of fibroblast activation at 7 days post-surgery

Under pathological conditions, a decreased PAA may lead to increased fibrin deposition, which will facilitate the invasion of fibroblasts and production of extracellular matrix, leading to the development of adhesions (Arung et al., 2011; Braun & Diamond, 2014). Adhesion fibroblasts, which are myofibroblasts isolated from adhesion tissues, are activated fibroblasts expressing high level of α SMA compared to normal peritoneal fibroblast (Saed & Diamond, 2004a; Saed, Zhao, Diamond, & Abu-Soud, 2006). Under pathological conditions leading to adhesion development, these adhesion fibroblasts are not cleared by apoptosis, persisting in the matrix, and scarring ensues (Fletcher, Jiang, Diamond, Abu-Soud, & Saed, 2008). Immunofluorescence of α SMA found its levels mostly in the regions proximal to the suture material and in the peripheral regions of the IB, suggesting the presence of adhesion fibroblasts in these regions (**Fig. 5.15**). Image analysis of the protein levels of α SMA in the immediate fibrotic regions revealed significantly lower levels of α SMA in the Cx43asODN-treated group compared to the vehicle-treated group ($p \leq 0.05$) (**Fig. 5.15c**). This suggested a lower degree of fibroblast activation upon Cx43asODN treatment.

C Effect of Cx43asODN on α -SMA expression

Figure 5.15 Treatment with Cx43asODN significantly reduced α SMA protein levels within the fibrotic region at 7 days post-surgery.

Representative digital stitched immunofluorescence images of IB samples of the vehicle- (A, n=7) and Cx43asODN-treated (B, n=5) groups at 7 days post-surgery, and their respective magnified regions (*i*, *ii*). Scale bars 50 and 500 μm. IB sections were immunolabeled for α SMA (green), and counterstained with DAPI (blue). (C) Image analysis revealed significantly reduced α SMA protein levels within the immediate fibrotic region in the Cx43asODN-treated group compared to the vehicle-treated group. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test (* $p \leq 0.05$). Dashed lines highlight the immediate fibrotic region. *IB* – ischaemic button; *s* – suture; *m* – muscle; *IF* – immediate fibrotic region.

5.4.9 Treatment with Cx43asODN reduced the extent of adhesion formation during early adhesion formation.

After determining that Cx43asODN intervention was effective by performing the different assessment on the developed adhesions at day 7 post-surgery, the effect of Cx43asODN on earlier events in adhesion formation was investigated. Adhesions were assessed at 6 and 24 hours post-surgery in the Cx43 asODN- and vehicle-treated group. Since developed adhesions were mostly immature during early adhesion formation (**Fig. 5.7a and b**), adhesions were only macroscopically evaluated for extent by scoring the number of IBs found to have attached adhesive band (Cassidy et al., 2015) (**Fig. 5.16**). At 6 hours post-surgery, the mean extent of adhesion of the vehicle-treated group was found to be $15 \pm 13.7\%$. In contrast, no peritoneal adhesions were detected in the Cx43 asODN-treated group ($p=0.06$). At 24 hours post-surgery, the mean extent of adhesions in the Cx43asODN-treated group ($12.5 \pm 20.9\%$) was slightly reduced compared to the vehicle-treated group ($30 \pm 20.9\%$) ($p=0.275$).

Figure 5.16 Cx43asODN treatment appeared to reduce the extent of adhesions during early adhesion formation

Peritoneal adhesions detected in the entire abdominal region were assessed and scored for extent of adhesion formation at 6 and 24 hours post-surgery by 3 observers blinded to the vehicle- ($n=5$) and Cx43asODN-treated ($n=6$) groups. At 6 hours, the vehicle-treated group had a mean extent score of 15% while no adhesions were observed in the Cx43asODN-treated group. At 24 hours, the vehicle-treated group had a mean extent score of 30% while Cx43asODN-treated group had about more than

two fold less adhesion compared to the vehicle group. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons between the treated groups and their respective time-matched control were made using a Mann-Whitney *U* test.

5.4.10 Effect of Cx43asODN treatment on the extent of inflammation during early adhesion formation.

The effect of Cx43asODN treatment on the extent of inflammation during early phase adhesiogenesis was further investigated. IB regions with a higher degree of recruited leukocytes and microabscess received a higher score for the extent of inflammation by observers blinded to the experimental groups. At both 6 and 24 hours post-surgery, Cx43asODN treatment appeared to slightly reduce the extent of inflammation outside the IB regions ($p=0.528$, $p=0.222$, respectively) when compared to its vehicle control (**Fig. 5.17b and d**). In contrast, a more significant reduction in the extent of inflammation was observed at regions within the IB at both 6 and 24 hours post-surgery ($p\leq 0.01$) (**Fig. 5.17c and e**). Taking into consideration that the most significant knockdown of Cx43 protein levels was also observed in this region (**Fig. 5.10b**), this might also suggest that the main site of action of Cx43asODN was within the IB region.

Next, transforming growth factor beta (TGF- β 1), proposed to be the main profibrotic mediator of adhesion formation (Chegini, 2008), was investigated within this region. Immunofluorescence staining of TGF- β 1 showed its levels mostly located in the muscle regions within the IB (**Fig. 5.18a**). Image analysis of the protein levels of TGF- β 1 in the immediate fibrotic regions revealed slightly lower levels in the Cx43asODN-treated group compared to the vehicle-treated group at 6 ($p=0.424$) and 24 hours ($p=0.31$) (**Fig. 5.18b and c**).

Figure 5.17 Cx43 asODN treatment significantly reduced the extent of inflammation within the IB at 6 and 24 hours

(A) Representative ROIs from digital stitched images of H&E-stained IB samples at 6 and 24 hours post-surgery. At regions outside of the IB at 6 and 24 hours, no notable changes in the extent of inflammation were observed although a clear reduction was observed in regions within the IB of the Cx43 asODN-treated group at both time points. Scale bar = 50 μ m. No differences were found between the vehicle-treated (n=5) and Cx43 asODN-treated group (n=6) for scores ascribed to the extent of inflammation of the regions outside of IB at 6 (B) and 24 hours (D). A significant decrease in the extent of inflammation in regions within the IB was found in the treated-groups from their respective time-matched vehicle-treated controls at 6 hours (C) and 24 hours (E). Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test (** $p \leq 0.01$).

Figure 5.18 Treatment with Cx43asODN appeared to reduce TGF-β1 protein levels within the IB region during early phase of adhesion formation.

(A) The panel shows representative immunofluorescence images revealing the distribution of TGF-β1 protein levels at regions within the IB at 6 and 24 hours post-surgery. Longitudinal sections of IB sections were immunolabeled for TGF-β1 (red, highlighted with yellow arrowheads), and counterstained with DAPI (blue). TGF-β1 protein levels observed in the muscle regions in the vehicle treated-group (n=5) at 6- and 24-hours appeared to be slightly reduced in the Cx43asODN treated group (n=6) at both time points. White dashed lines separate the mesothelial (upper) and muscle (lower) regions. Magnification used, 40X. Scale bar = 50 μm. (B) Image analysis revealed slightly lower TGF-β1 protein levels in the muscle regions within the IB in the Cx43 asODN-treated group compared to the vehicle-treated group. Data is presented as mean value ± standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test. (*p≤0.05)

5.5 Discussion

5.5.1 Brief comparison of the *ex vivo* mouse peritoneal strip model and *in vivo* IB model

In terms of clinical relevance, both *in vivo* and *ex vivo* systems were capable of generating adhesions consistent with the clinical observations of “Type 1B adhesions” were noted. This subtype is classified as “de novo adhesions that are produced at sites where a surgical procedure was performed and previously clear of adhesions” (Diamond & Nezhat, 1993). In addition, histology of the strips from the *ex vivo* samples revealed thickening of mesothelium, which is also observed in the *in vivo* IB model. This finding is also recently corroborated by others (Tsai et al., 2018). This observation is clinically relevant as mesothelium thickening is also observed in patients as well (Arung et al., 2011; diZerega & Campeau, 2001; Mutsaers et al., 2015). While Cx43 appears to be implicated in the *ex vivo* model as well, this was not fully explored because it was decided that the lack of means to locate the exact point of adhesion creation made it a challenge to make meaningful conclusions.

Despite some similarities between the *ex vivo* and *in vivo* models, majority of the adhesions observed in the *ex vivo* model were thin and filmy and could be broken through gentle traction. This contrasts with most of the adhesions found in the *in vivo* model, which are denser and required blunt dissection to break. This may suggest that adhesions formed in the *ex vivo* model are not at its “truest” potential, and essential “ingredients” for achieving this may be lacking in the *ex vivo* system. Some examples of this are the lack of circulation or the lack of the full immune

repertoire, which are inherent characteristics of most *ex vivo* systems. Naturally, *in vivo* systems remain closest to its native physiological environment and would be a preferred choice for understanding pathogenesis, which is critical for identifying new therapeutic targets.

5.5.2 Adhesions formed by the IB model and their distribution

Evaluation of adhesion formation has been previously performed at a variety of timings depending on the research inquiry, ranging from as early as 12 hours to 24th week post-induction (Ozel et al., 2005). In this study some adhesions were already detected in the IB model as early as 6 hours although their detection was infrequent. The detected adhesions were naturally immature and could be broken with gentle traction. The extent of adhesions was noted to increase over time, with its extent doubled by 24 hours post-surgery. Adhesions detected at this time point were still immature and could still be broken with little effort. Adhesions found in the earlier phase of adhesion formation were generally transparent, filmy, and avascular. In contrast, the majority of these adhesions observed in the later phase of adhesion formation were opaque, dense, vascular, and required at least blunt dissection to break. Also, the extent of adhesion formation at 7 days post-surgery was almost two-fold more than that detected at 24 hour post-surgery. Interestingly, the location of the developed adhesions appeared to be influenced by the location of the IB. Two-thirds of the total adhesions at 7 days post-surgery were being formed in the caudal abdominal region with the majority of the adhesions found adhered to the epididymal fat pad. The increased incidence of adhesions in this region is suspected to be due to prominence of epididymal fat pads in the caudal abdominal region (**Fig. 5.19**). While it may be possible that the proximity of the epididymal fat pad to the

IBs may have allowed the fat pad to “latch” onto the button or suture, differences in fibrinolytic activity among the organs may also have contributed to the increased incidence. Early work revealed the fibrinolytic activity to be heterogenous among whole tissues in humans and animals (Albrechtsen, 1957). Almost 4 times greater fibrinolytic activity was documented from surgical biopsies of serosal tissues from the omentum when compared to the gall bladder (Merlo, Fausone, & Castagna, 1983). However, the understanding of the fibrinolytic activity of tissues under physiological and pathological conditions is still limited and a detailed comparison of the influence of the location of the IB on the development of adhesions has yet to be performed in literature and may warrant further investigations. Since the location of the IB appeared to influence the quantity of adhesion created, analyses were only performed on the IBs in the caudal abdominal region.

Figure 5.19 Epididymal fat pad distribution in the caudal abdominal region

(A) Injection of PBS into peritoneal cavity suspended the epididymal fat pad, aiding the visualisation of the anatomical position of the epididymal fat pad. (B) Exposure of the inner peritoneal cavity to further aid visualisation. The abdomen is divided into a cranial and caudal portion by the umbilicus. The epididymal fat pads are located at the caudal region of the abdomen. Localisation of the epididymal fat pad at the caudal abdominal regions are highlighted with yellow arrowheads labeled *E*. *RP* –Retroperitoneal fat pad; *E* – Epididymal fat pad

5.5.3 Increases of Cx43 protein levels during early adhesion formation

During early adhesion formation, pathological increases of Cx43 protein levels were evident during early phase adhesion formation, (summarised in **Fig. 5.20**). At 6 hours post-surgery, higher increases of Cx43 levels were observed in both mesothelial and muscle regions outside the IB. At 24 hours post-surgery, these increases in Cx43 levels subsequently appeared to be slightly dampened. In contrast, Cx43 levels in the mesothelial regions within the IB appeared to increase over time. The most significant increase of Cx43 levels was observed at 24 hours post-surgery, after increasing another 1.4-folds from 6 hours post-surgery. Unfortunately, one limitation of the study was that no cell-specific markers were used to tease out the different cell types in which increases of Cx43 levels were specifically observed. Based on position and morphology of the cells in the Cx43 immunolabeled sections and the accompanying H&E sister sections, it is hypothesised that increases of Cx43 were observed mainly in the mesothelial cells and infiltrating leukocytes (**Fig. 5.21**).

Cx43 has been shown to act as a major gap junction protein in human pleural (Pelin, Hirvonen, & Linnainmaa, 1994) and peritoneal mesothelial cells (Mikula-Pietrasik et al., 2017; Ogawa et al., 2005; Tang et al., 2013). However, previous studies dealing with Cx43 in these cell types were mainly investigating its involvement in cancer and not in adhesion formation (Mikula-Pietrasik et al., 2017; Ogawa et al., 2005; Tang et al., 2013). Although limited literature is available to inform whether increases in Cx43 expression is likely to have an adverse effect on peritoneal mesothelial cells, various reports show that alterations of Cx43 expression is associated with human diseases (reviewed in (Trosko & Ruch, 1998; Willebrords, 2016)). Also, increases in Cx43 protein levels on neutrophils and

monocytes/macrophages during inflammatory conditions have been documented (Al-Ghadban, Kaissi, Homaidan, Naim, & El-Sabban, 2016; Branes, Contreras, & Saez, 2002; Eugenin, Branes, Berman, & Saez, 2003; Eugenin et al., 2007). For example, increases in protein levels of Cx40 and Cx43 were observed in neutrophils stimulated with TNF- α in endothelial cell-conditioned media (Branes et al., 2002). Along these lines, ATP may be released from Cx43 hemichannels from neutrophils and monocytes under inflammatory conditions (Calder et al., 2015). The accumulation of ATP may then activate purinergic receptors present on endothelial cells, triggering a lateral propagation of Ca²⁺-dependent signaling across the endothelium, which later results in the expression of leukocyte adhesion molecules (i.e. P-selectin) (Zernecke et al., 2006). This may facilitate the recruitment of leukocytes during early adhesion pathogenesis. Also, taking into consideration that these cell types share a common gap junction and the increase of Cx43 appeared to coincide with significant recruitment of leukocytes to the region, it is tempting to speculate that these cells may share gap junction intercellular communication, which may have a crucial role during adhesion formation.

Figure 5.20. Schematic diagram of Cx43 protein levels during early adhesion formation.

Relative protein levels of Cx43 in the mesothelial (circles) and muscle (squares) regions within and outside the IB are illustrated using shapes. The sizes of the shapes are representative of their fold

change from basal physiological levels (**Fig. 5.7**). Highest Cx43 protein levels were detected in the mesothelial regions within the IB at 24 hours post-surgery. Statistically significant changes in Cx43 protein levels at 24 hours post-surgery due to Cx43asODN treatment are denoted with downward pointing arrows: single arrow, ($p \leq 0.05$); two arrows, ($p \leq 0.01$). Cx43 protein level at 6 hours was not significantly reduced with treatment of Cx43asODN.

Figure 5.21 Neutrophils in the mesothelial regions

The panel shows representative immunofluorescence images of regions outside the IB samples at 6 hours post-surgery. Longitudinal sections of IB sections were immunolabeled for Cx43 (red), and counterstained with DAPI (blue). A single slice (top right) from a stack of slices that have been processed to be max projected (top left) is shown to facilitate identification of neutrophils (bottom left) and their accompanying Cx43 protein levels (bottom right). Neutrophils are identified by appearance of their multi-lobed nuclei (position highlighted with white arrowheads). Scale bar = 50 μm .

5.5.4 Role of Cx43 during adhesion formation

After observing a pathological increase of Cx43 in the mesothelial region of the IB during early adhesion formation, Cx43asODN treatment was explored to investigate the role of Cx43 during adhesion formation. Through macroscopic assessment of the adhesions at 7 days post-surgery, Cx43asODN was found to significantly reduce adhesion strength and severity. Following microscopic assessment, it was found that there was a smaller immediate fibrotic area that is accompanied by lesser deposition of collagen and a reduced degree of fibroblast activation. In addition, the protein level of PAI-1 in the peritoneal fluid of the Cx43asODN treated-group was significantly lower than the vehicle-treated group suggesting higher PAA activity at day 7 post-surgery. At an earlier window of assessment, Cx43asODN significantly reduced the degree of leukocyte recruitment and microabscesses observed. While no appreciable changes in cytokine levels were seen immediately at 6 hours post-surgery, TGF- β 1 levels were significantly reduced at 24 hours post-surgery. In the following subsections, the potential contribution of Cx43 to adhesion formation is discussed.

5.5.5.1 Role of Cx43 in inflammation and cell migration

Following peritoneal injury, the release of an initial cocktail of cytokines and chemokines sets the stage for the biochemical and cellular cascades that follow. While released cytokines kick starts the repair process, chemokines promote the recruitment of leukocytes to the site of injury. The early responders are predominantly neutrophils, with a gradual shift to macrophages after 24 hours (Maciver et al., 2011). The recruited leukocytes initiate another robust cytokine cascade and are suggested to play a crucial role in the outcome of adhesion formation

(Beyene et al., 2015) (Maciver et al., 2011). This was demonstrated when the reduction of neutrophils through the use of cyclophosphamide was shown to reduce the number and severity of adhesions formed (Vural et al., 1999). In another study, abrogating the interaction between chemokine (CCL1) and its receptor (CCR8) interaction in peritoneal macrophages hampered the migration of peritoneal macrophages, resulting in a significant reduction of peritoneal adhesions (Hoshino et al., 2007).

In this study, significant reduction in number of leukocytes recruited within the IB region at 6 and 24 hours post-surgery was observed in the Cx43asODN treated group. It is noteworthy to point out that this phenomenon has also been observed in other experimental disease models (Abed et al., 2014; Saredine et al., 2009). In a model of acute lung injury using intratracheal instillation of LPS, almost 50% fewer recruited neutrophils were observed in the bronchoalveolar lavages of Cx43^{+/-} mice or mice treated with connexin mimetic peptide, Gap26 (Saredine et al., 2009). Also, in both models of hypertension-induced and tubulointerstitial inflammation models of chronic kidney disease, infiltration of monocytes were significantly reduced with decreased Cx43 mRNA expression in Cx43^{+/-} mice or through Cx43asODN administration (Abed et al., 2014). Although these observations were made in different disease models, they corroborate with findings made in this study, further supporting the emerging role of Cx43 in the recruitment of leukocytes. Future investigations using *in vitro* systems or conditional-Cx43 transgenic mice will help determine which cell-derived Cx43 (i.e. mesothelial, neutrophil, macrophage) is a stronger contributor to leukocytes recruitment. This may elucidate

a method to fine-tune the recruitment of leukocytes to the site of injury, presenting a potential therapeutic avenue.

Considering recruited leukocytes release a robust cytokine cascade (Beyene et al., 2015), Cx43asODN-mediated reduction of leukocyte recruitment may have indirectly reduced the cytokines levels. Conversely, modulation of Cx43 has also been shown to exert direct effects on cytokine synthesis. In one example, treatment of lymphocytes with connexin mimetic peptides, Gap26 and Gap27, was demonstrated to temporally inhibit the synthesis of cytokines (Oviedo-Orta, Gasque, & Evans, 2001). This suggests that there may be a potential scenario where the manipulation of Cx43 expression may have directly contributed to changes in cytokines levels during early phase adhesion formation. Although no appreciable changes in cytokine levels were seen immediately at 6 hours post-surgery, TGF- β 1 levels were significantly reduced in the muscle regions within the IB at 24 hours post-surgery in the Cx43asODN-treated group.

The differentiation of fibroblasts into myofibroblasts in response to profibrotic cytokines is characterised by the acquisition of a *de novo* expression of α SMA (Desmouliere et al., 1993). Proposed to be a main profibrotic mediator (Chegini, 2008), TGF- β 1 has been shown to initiate and promote the expression of α SMA, playing an important role in regulating the myofibroblastic phenoconversion (Asazuma-Nakamura et al., 2009; Desmouliere et al., 1993; Rosenkranz, 2004; Tomasek, McRae, Owens, & Haaksma, 2005). Work by Dai and colleagues demonstrated Cx43 positively mediates the activity of TGF- β through the competitive binding to microtubules between endogenous Smads and Cx43 (Dai et

al., 2007). Later work by the same group demonstrated that over-expression or inhibition of Cx43 respectively potentiated or inhibited TGF- β 1-induced myofibroblastic phenocconversion (Asazuma-Nakamura et al., 2009). Based on distribution of the AlexaFluor488[®]-tagged Cx43asODN at 6- and 24-hours, it appears the Cx43asODN lasted for at least 6 hours, which is long enough to exert its action on invading fibroblasts and consequently inhibit/reduce their acquisition of *de novo* expression of α SMA. This may help to explain the reduction of α SMA expression and collagen deposition observed in the intermediate fibrotic region of the Cx43asODN-treated group. Considering this, future studies may attempt multiple treatments with Cx43asODN or explore the effects of next-generation modified asODN (i.e. GapmeRs) that possess a longer half-life.

The presence of TGF- β 1 during muscle injury has been documented (Goetsch & Niesler, 2016; Kim & Lee, 2017; Smith, Stauber, Waters, Alway, & Stauber, 2007). In skeletal muscles, TGF- β 1 is also regarded as one of the most profibrogenic factors (Delaney et al., 2017). It is capable of regulating the synthesis, degradation and the remodeling of the extracellular matrix, making it an important regulator of fibrosis development (Delaney et al., 2017; Kim & Lee, 2017). In one example, strong upregulation of TGF- β 1 and accompanying fibrosis was observed in skeletal muscles in response to damage and denervation (Zimowska et al., 2009). Treatment of TGF- β 1 and T β R1 antibodies to impair TGF- β 1 activity saw marked improvements in the regeneration of muscle that is accompanied with reduced fibrosis (Zimowska et al., 2009). In this study, TGF- β 1 protein levels at 24 hours post-surgery was significantly reduced in the muscle regions within the IB the Cx43asODN-treated group (**Fig. 5.17c**). Although fibrosis in the muscle regions

within the IB was not investigated in this study, α SMA protein levels also appears to be lowered in this region in the Cx43asODN treated group (**Fig. 26b**). This may be further investigated by studying the fibrotic responses in *in vitro* cultures of Cx43asODN-treated skeletal muscle that has undergone ischaemia/reperfusion damage through the use of a hypoxia chamber.

5.5.5.2 Role of Cx43 in plasminogen-activator activity

The formation of peritoneal adhesions depends on the equilibrium between fibrin production and fibrinolysis following peritoneal injury (Arung et al., 2011). The complete degradation of fibrin facilitates normal peritoneal healing. However, the incomplete degradation of fibrin may serve as a scaffolding matrix for invading fibroblasts and growth of new blood vessels, paving the way for adhesion formation (Arung et al., 2011). Plasminogen activators play a vital role in the fibrinolytic sequence. They convert inactive plasminogen substrates to its active form, plasmin, which then dissolves fibrin. In the peritoneal cavity, 95% of the plasminogen-activating activity (PAA) is contributed by tPA (Holmdahl et al., 1996). However, the activity of plasminogen activators also depends on the presence of PAIs. PAI-1, the main fibrinolytic inhibitor, hampers the fibrinolytic response by antagonising and forming inactive complexes with tPA (Maciver et al., 2011). Under pathological conditions, PAA is markedly decreased, leading to the development of adhesions (Arung et al., 2011; Braun & Diamond, 2014).

Work investigating the association between Cx43 and tPA levels is limited. Earlier *in vivo* investigation into the vascular effects of rotigaptide, a synthetic peptide capable of potentiating Cx43 communication, revealed that potentiation of Cx43

communication has no effect on tPA release in the forearm arterial circulation (Lang et al., 2008). In this study, the treatment with Cx43asODN did not appear to affect the protein level of tPA in the peritoneal fluid. While these observations may suggest little association between Cx43 and tPA, atorvastatin-treated human vascular smooth muscle cells were observed to decrease in Cx43 protein levels, which was accompanied with the increase of mRNA expression of tPA, suggesting the effect of Cx43 on tPA expression may be cell-type dependent (Dubuis et al., 2013). In the same study, the decrease of Cx43 protein levels was also accompanied by a decrease of PAI-1 protein levels. Along these lines, manipulation of Cx43 activity, via nonspecific gap junction inhibitors (carbenoxolone and probenecid) or siRNA knockdown, was suggested to decrease PAI-1 mRNA expression in rat cardiac fibroblasts by inhibiting ATP release that is involved in profibrotic signaling (Lu et al., 2012). The decrease in ATP levels was also accompanied by decreases in α SMA protein levels and collagen accumulation (Lu et al., 2012). Overall, this indicates that Cx43-mediated reduction of PAI-1 expression is a result of basal nucleotide signaling and may help explain the significant reduction of PAI-1 levels in Cx43asODN-treated group.

5.6 Conclusions

In this chapter, we set out to explore the role of Cx43 in an ischaemic button model of adhesion formation. Adhesions were generated by this model by creating an ischaemic button through a ligation of an out-pouching of the abdominal sidewall with a suture. Assessment performed during early adhesion formation found pathological increase of Cx43 protein levels in the mesothelial regions within the ischaemic button at 6 and 24 hours post-surgery that was accompanied by

observations of increases in leukocyte recruitment. The significance of this pathological increase was further pursued with the use of Cx43asODN, which was shown to penetrate up to 180 μ m into the muscle layer and significantly suppressed pathological increases Cx43 protein levels in the mesothelial regions within the ischaemic button. The effect of this early suppression of Cx43 levels was seen in subsequent macroscopic assessment of adhesion strength and severity, which was further validated by notable reductions in microscopic assessment of fibrosis extent and severity, the degree of α SMA protein levels. Also, levels of PAI-1 was also found to be significantly reduced. Collectively, this evidence suggested that down-regulation of Cx43 protein levels plays an important role in adhesion formation. Further investigations suggested that Cx43 mediates adhesion formation by regulating leukocyte recruitment and TGF- β 1 protein levels during early adhesion formation. This indicates that Cx43 may be a potential therapeutic target for post-surgical peritoneal adhesions.

Chapter 6: Exploring the role of endothelial-connexin43 in peritoneal adhesion formation

6.1 Introduction

Post-surgical peritoneal adhesions are considered the consequence of a pathological process of peritoneal healing to the peritoneal lining, usually arises from abdominal surgery (Arung et al., 2011; diZerega & Campeau, 2001; Mutsaers et al., 2015). Almost immediately upon injury, an inflammatory response is elicited at the peritoneal surface that is characterised by the release of a cocktail of cytokines and chemokines, recruitment of inflammatory cells, and the activation of the coagulation cascade (Beyene et al., 2015; Schnuriger et al., 2011).

Recent reviews suggest that endothelial cells have an essential role in the modulating cross talk between inflammation and blood coagulation events (Levi & van der Poll, 2010; O'Brien, 2012; Okamoto & Suzuki, 2017; Soon et al., 2016). A number of connexins have been detected in endothelial cells, particularly Cx32, Cx37, Cx40, and Cx43 (Okamoto & Suzuki, 2017; Soon et al., 2016). Through these Cx channels, endothelial cells can communicate with surrounding vascular components (i.e. adjacent endothelial cells, smooth muscle cells, leukocytes) (Okamoto & Suzuki, 2017; Soon et al., 2016). Depending on their position on the vascular tree and the health of the vessel in which they reside, their relative expression levels, size of gap junction plaques, and the selective permeability of channel may vary (Soon et al., 2016). In addition, changes in endothelial function in response to the pro-inflammatory or pro-coagulant stimuli that are present during adhesion formation

are accompanied by pathological alterations in endothelial Cx expression and gap junction function (O'Donnell et al., 2014; Okamoto et al., 2014). Conversely, accumulating evidence demonstrates that manipulation of the endothelial Cx levels is also able to regulate the spiraling inflammatory response (Abed et al., 2014; Okamoto et al., 2014; Zhang et al., 2015). Understanding how endothelial connexins play such a pivotal role in supporting endothelial homeostasis may open up new strategies for designing connexin-based therapeutics.

6.1.1 Endothelial connexins and inflammation

Endothelial connexin expression and gap junction functions are altered by various inflammatory stimuli (Okamoto & Suzuki, 2017). For example, 24 hour stimulation with pro-inflammatory cytokine, TNF- α , reduces endothelial Cx37 and Cx40 protein levels (Okamoto et al., 2011; van Rijen, van Kempen, Postma, & Jongsma, 1998). While this was accompanied by a reduction in gap junction activity, no changes were detected in Cx43 protein levels (Okamoto et al., 2011; van Rijen et al., 1998). On the contrary, LPS stimulation of endothelial cells appears to increase Cx43 protein levels. This has been demonstrated in LPS-treated human pulmonary artery endothelial cells, which was also shown to increase GJIC (O'Donnell et al., 2014). In addition, similar increases in Cx43 mRNA expression was observed in primary vascular endothelial cells cultured from the rat pulmonary veins (Zhang et al., 2015). In both studies, increases in endothelial-Cx43 were accompanied by increases in vascular permeability, which were both abrogated upon transfection with Cx43 siRNA (O'Donnell et al., 2014; Zhang et al., 2015). Along these lines, an increase of vascular permeability that was accompanied by up-regulation of endothelial-Cx43 in various organs in septic rats and this was also reduced by

carbenoxolone, a broad-spectrum connexin inhibitor. Although under pathophysiological conditions there exists a cocktail of pro-inflammatory mediators, these findings suggest that the manipulation of aberrant Cx expression and gap junction function (i.e. pathological increases in Cx43 expression) have the capacity to also influence the inflammatory responses of endothelial cells.

6.1.2 Endothelial connexins and coagulation

The stimulation of pro-inflammatory signals also elicits the blood coagulation cascade, by increasing tissue factor expression on the surface of endothelial cells and also the release of von Willebrand factor (Antoniak, 2018; Mackman, 2009). The elicited coagulation cascade initiates the mobilisation of coagulation factors that ultimately results in the generation of thrombin, which is responsible for the conversion of fibrinogen to fibrin (Engelmann & Massberg, 2013).

Conversely, thrombin and other factors of the coagulation cascade can also regulate inflammation (Coughlin, 2005; Popovic et al., 2012). Thrombin in particular is able to induce the expression of leukocyte adhesion molecules on endothelial cells (i.e. P-selectin) and the production of interleukin-6 and 8, leading to the chemotaxis of neutrophils and monocytes (Bizios, Lai, Fenton, & Malik, 1986; Kaplanski et al., 1998; Sugama et al., 1992). Furthermore, thrombin is reported to disrupt endothelial cell-cell junctions, resulting in an increase in vascular permeability (Lum & Malik, 1996; O'Donnell et al., 2014). Accumulating studies suggest thrombin up-regulates endothelial connexin expression and activity. In one example, thrombin-treated human pulmonary artery endothelial cells increased in vascular permeability, which was accompanied with the up-regulation of Cx43 mRNA expression and gap

junction function (O'Donnell et al., 2014). Subsequent investigations demonstrated that vascular permeabilisation and myosin light chain phosphorylation in thrombin-activated endothelial cells were both abrogated by treatment with carbenoxolone, a broad-spectrum connexin inhibitor, and Cx43 siRNA-mediated knockdown (O'Donnell et al., 2014).

Along these lines, manipulation of endothelial Cx expression and gap junction function can also regulate other coagulation responses. Tissue factor mRNA expression induced in endothelial cells upon TNF- α stimulation may be potentiated or attenuated with the intracellular transfer of monoclonal antibodies against Cx32 or Cx43 respectively (Okamoto et al., 2014). In addition, in a co-culture system, blockade of Cx43 reduced tissue factor mRNA expression that was induced in non-stimulated acceptor HUVECs from adjacent TNF- α -stimulated effector HUVECs (Okamoto et al., 2014). This suggests that Cx43 supports pro-coagulant signaling in stimulated endothelial cells.

6.1.3 Endothelial connexins and leukocyte adhesion

The adhesion of circulating leukocytes to inflamed endothelial cells and subsequent transmigration to the site of injury are important events in inflammation (Muller, 2013). Under pro-inflammatory conditions, endothelial cells regulate these processes through the expression of leukocyte adhesion molecules and chemokines (Okamoto & Suzuki, 2017). Studies investigating the endothelial Cx40 and Cx43 expression during the course of inflammation in the lungs suggest an opposite regulation of their expressions during inflammatory conditions (Fernandez-Cobo, Gingalewski, & De Maio, 1998; Rignault, Haefliger, Waeber, Liaudet, & Feihl,

2007; Sargedine et al., 2009; Zhang et al., 2015). Briefly, neutrophil transmigration across the endothelial barrier appears to be promoted by a high Cx43 to Cx40 ratio, while the opposite appears to hamper adhesion (Scheckenbach, Crespin, Kwak, & Chanson, 2011).

On this note, CD73 plays a pivotal role in supporting connexin regulation over these inflammatory processes (Scheckenbach et al., 2011). Under physiological conditions, endothelial cells constitutively express CD73, a 5'-ecto-ATPase that converts extracellular ATP into AMP that is anti-inflammatory (Zernecke et al., 2006). Under inflammatory conditions, ATP may be released from Cx43 hemichannels from neutrophils and monocytes (Calder et al., 2015). Also, under inflammatory conditions, endothelial cells have a reduced Cx40 protein levels that is accompanied by a reduction in CD73 protein levels (Chadjichristos et al., 2010; Rignault et al., 2007). This reduction in CD73 levels hampers the use of ATP as a substrate. Consequently, the accumulating ATP activates purinergic receptors present on endothelial cells, triggering a lateral propagation of Ca^{2+} -dependent signaling across the endothelium, which later results in the expression of leukocyte adhesion molecules (i.e. P-selectin) (Zernecke et al., 2006). Using an endothelial-Cx43 knockout further revealed that Cx43 is crucial in mediating this Ca^{2+} -dependent signaling (Parthasarathi et al., 2006). This further corroborates the notion that Cx43 supports pro-inflammatory signaling in stimulated endothelial cells by also facilitating the initial adhesions events of leukocyte recruitment (summarised in **Fig. 6.1**) (Scheckenbach et al., 2011).

Figure 6.1 Cx43 supports pro-inflammatory signaling in stimulated endothelial cells by facilitating the initial adhesions events of leukocyte recruitment.

(1) Under inflammatory conditions, ATP may be released from neutrophil-Cx43 hemichannels. (2) Also, under the same conditions, endothelial cells have an increase in Cx43 protein levels and a decrease in Cx40 protein levels. The decrease in Cx40 protein levels is accompanied by a reduction in CD73 protein levels, which hampers the use of ATP as a substrate. (3) Accumulating pool of ATP activates the purinergic receptors present on endothelial cells, triggering a lateral propagation of Ca^{2+} -dependent signaling across the endothelium that is mediated by Cx43. (4) The lateral propagation of Ca^{2+} -dependent signaling later results in the expression of leukocyte adhesion molecules (i.e. P-selectin).

6.1.4 Endothelial-Cx43 transgenic mice

These emerging evidences discussed above suggest that endothelial-Cx43 is a central regulator of these inflammatory processes. Furthermore, there is a gap of knowledge regarding its role during adhesion formation in the existing literature. In light of this, an *a priori* decision was made to explore the role of endothelial-Cx43 in peritoneal post-surgical adhesion formation using transgenic mice. Understanding the role of cell-specific connexins under pathological conditions has been made possible with the generation of numerous transgenic mice models (Chadjichristos et al., 2010; Westphalen et al., 2014). For this purpose, the Cre/lox site-specific

recombination system is useful for allowing the investigator to specifically control of the expression and activity of genes of interest in the desired cells through the conjugation of a tissue specific promoter to the Cre-recombinase (Forde et al., 2002; Gu, Zou, & Rajewsky, 1993; Metzger, Clifford, Chiba, & Chambon, 1995).

The *Tie2* promoter is frequently used in literature to manipulate gene expression in endothelial cells (Forde et al., 2002; Osei-Owusu et al., 2012). However, the expression of *Tie2* has been reported in both haematopoietic and endothelium cells (Dumont et al., 1994; Kisanuki et al., 2001; Takakura et al., 1998). Due to the transient *Tie2* expression in developing hematopoietic cells, this leads to the manipulation of the gene-of-interest (i.e. Cx43) in the haematopoietic lineages other than the endothelial cells (Tang, Harrington, Yang, Friesel, & Liaw, 2010). Therefore, Cx43 transgenic mice that are under the influence of a constitutive *Tie2* promoter, like those used by Parthasarathi and Véliz, would have Cx43 being manipulated in both endothelial and haematopoietic lineages (Koval et al., 2011; Parthasarathi et al., 2006; Veliz, Gonzalez, Duling, Saez, & Boric, 2008). This problem of non-specificity is mitigated in my study through the use of Cx43 transgenic mice that are under the influence of a tamoxifen-inducible *Tie2* promoter (Forde et al., 2002). Tamoxifen-induced recombination in the haematopoietic cell populations showed less than 0.2% recombination in B and T cell populations compared to over 90% recombination in constitutive *Tie2* Cre mice (Forde et al., 2002). No recombination was reported in other cell types of the haematopoietic lineage (i.e. macrophages). Tang and colleagues also further showed that circulating embryonic hematopoietic cells no longer had an active *Tie2* promoter (Tang et al., 2010).

In this study, tamoxifen-inducible cell-specific Cx43 transgenic mice were treated with tamoxifen to induce Cre recombination, and thus reducing endothelial-Cx43 protein levels in the process. This reduction allows for the investigation of the role of endothelial-Cx43 in adhesion formation and the potential to open up new strategies for fine-tuning connexin-based therapeutics for post-surgical peritoneal adhesions.

6.2 Aims and objectives

Work in this chapter explores the contribution of endothelial-Cx43 to adhesion formation. Effects of tamoxifen-induced endothelial-Cx43 knockdown on late phase adhesion formation were studied using tamoxifen-inducible endothelial-Cx43 specific transgenic mice in ischaemic button model of adhesion formation. After determining notable knockdown of endothelial-Cx43, its significance would be determined through evaluations at late phase of adhesion formation. Effects on late phase adhesion formation would be evaluated using macroscopic assessment based on established scoring matrices that and would be later verified with microscopic assessment and further validated with measurement of the overall plasminogen-activating activity in the peritoneal cavity.

6.3 Materials and Methods

6.3.1 In vivo Work

6.3.1.1 Experimental animals

All animal procedures were performed in accordance with the animal protocols approved by the Institutional Animal Care and Use Committee of the Animal Research Facility of Nanyang Technological University, protocol number A0372. Tie2-CreER^{T2} transgenic mice and Cx43^{fl/fl} transgenic mice were purchased from The European Mouse Mutant Archive and The Jackson Laboratory respectively. Mice were bred under specific pathogen-free conditions under a 12-hour light-dark cycle and were supplied standard rodent pellet diet and water *ad libitum*. For the generation of experimental tamoxifen-inducible endothelial cell-specific Cx43 transgenic mice (Tie2-CreER^{T2+/-}Cx43^{fl/fl}), Tie2-CreER^{T2} and Cx43^{fl/fl} transgenic mice were mated in-house and maintained on the C57BL/6 background. Briefly, the breeding strategy used to achieve this starts with the mating homozygous parents of both strains to obtain heterozygous genotype Tie2-CreER^{T2+/-}Cx43^{fl/-}, which was identified through genotyping. Next, mating of this genotype with the Cx43^{fl/fl} genotype allowed the generation of the Tie2-CreER^{T2+/-}Cx43^{fl/fl} genotype, which were identified through genotyping and used as experimental mice. Conversely, Tie2-CreER^{T2-/-}Cx43^{fl/fl} mice were used as their controls. The primers used for genotyping the mice are listed in **Appendix II, Supplementary Table 2**. For induction of Cre-mediated DNA recombination (EC-Cx43^{-/-}), 6 weeks old Cx43^{fl/fl} mice carrying the Tie2-CreER^{T2} transgene received daily oral gavage of 4 mg tamoxifen (Sigma-Aldrich, USA) for 5 days (n=9)(Anastassiadis et al., 2010; Forde

et al., 2002). Age matched tamoxifen-treated mice Cx43^{fl/fl} (EC-Cx43^{fl/fl}) served as a control group (n=6). All tamoxifen-treated mice were rested at least a week prior to the surgery. Only age-matched (7 to 12 weeks) males were used for experiments.

6.3.1.2 Ischaemic button model of post-surgical peritoneal adhesions

Ischaemic buttons (IB) were created to induce adhesion formation as previously described by Sandoval et al., (2016) with minor modifications. Prior to the procedure, fur around the incision site was removed using an electric trimmer and depilatory cream before returning the mice to their cages and allowing them to rest for at least 24 hours. Briefly, mice were anaesthetised with isoflurane and placed on a warm heating mat throughout the surgery. All surgeons used non-powdered surgical gloves during the surgery and all instruments were sterilised. Next, the midline incision site was sterilised with 7.5% (v/v) providone-iodine solution (**Fig. 6.2a**). A midline laparotomy of 4 cm was performed to facilitate the exposure of the parietal peritoneum (**Fig. 6.2b**). Each ischaemic button was created by placing a 4/0 non-absorbable silk suture at the base of ~3 mm of the sidewall parietal peritoneum, which was grasped with a needle holder. Two ischaemic buttons, spaced at least 1 cm apart, were created on the peritoneum wall of each side of the midline (**Fig. 6.2c**). The laparotomy site was then closed in two layers, the inner layer with simple interrupted polyglycolic acid absorbable 6.0 sutures (**Fig. 6.2d**) and the outer layer using metallic wound clips (**Fig. 6.2e**). After closure, mice were regularly monitored until euthanasia with CO₂ at 7 days post-surgery for late phase formation assessment. For post-recovery, the mice received buprenorphine (0.1 mL per 10 g of animal weight) into the loose skin over the interscapular region.

Figure 6.2 Major steps during various adhesion-inducing operations in mice.

(A) Disinfection of the incision site after shaving. This was followed by a (B) midline incision of the skin. (C) Ischaemic button (IB) model. Two IBs, spaced approximately 1 cm apart, were created on the peritoneum on each side of the midline incision. Each button (yellow arrowheads) was created by placing a silk ligature at the base of 3 mm of parietal tissue that is grasped with a needle holder. (D) Wound closure by applying simple interrupted sutures to the peritoneal musculature and (E) metallic clips to the skin.

6.3.1.3 Peritoneal lavage

Prior to necropsy, 3 mL of cold phosphate-buffered saline (PBS) was injected into the peritoneal cavity and the abdomen was gently agitated for 1 min. Approximately 2 mL of the fluid was recovered. Samples with blood contamination were omitted. Cellular debris was removed by microcentrifugation at 9000 g for 15 min at 4°C, before freezing the supernatant in 250 µL aliquots at -80°C.

6.3.1.4 Peritoneal fluid tPA and PAI-1 total antigen ELISA

Thawed aliquots of the peritoneal lavage were analysed for the total antigen concentration of tPA and PAI-1 present, according to manufacturer's instructions (Molecular Innovations, USA). Samples that had coefficient of variation values more than 15% were excluded. Detected concentrations were normalised using total protein content determined from a bicinchoninic acid assay performed on the same aliquot (ThermoFisher Scientific, USA).

6.3.1.5 Macroscopic assessment of peritoneal adhesions

An inversed U-shaped incision was made across the peritoneal cavity for necropsy and the adhesions in each animal were evaluated according to established scoring matrices. These matrices assessed peritoneal adhesions based on qualities such as, extent, involvement, severity, and strength (refer to **Table 6.1**) (Adhesion Scoring Group, 1994; Cassidy et al., 2015; Hoffmann et al., 2009; Nair et al., 1974). Three observers blinded to the experimental groups scored each mouse and their scores were later averaged to give a single score per mouse. Inter-observer scores did not differ more than 2. Scores are represented as a percentage of maximum score possible for each parameter. After which, the mean parameter scores for each of the groups were calculated.

Table 6.1 Scoring matrix for assessing peritoneal adhesions

Variable	Score	Description
Extent (Cassidy et al., 2015)	0	Complete absence of adhesions
	1	Only 1 button has an adhesion band
	2	2 buttons have adhesion bands
	3	3 buttons have adhesion bands
	4	4 buttons have adhesion bands
Involvement (Quantity) (Nair et al., 1974)	0	Complete absence of adhesions
	1	1 adhesion band between the organs or between one organ and abdominal wall
	2	2 adhesion bands between the organs or between one organ and abdominal wall
	3	>2 adhesion bands between the organs or between one organ and abdominal wall or adhesions of intestinal loops without any adhesion to the abdominal wall. Significant adhesion.
	4	Adhesion of all viscera to the abdominal wall
Severity (Quality) (Adhesion Scoring Group, 1994)	0	Complete absence of adhesions
	1	Filmy, avascular
	2	Dense, opaque and/or vascular
	3	Very thick cohesive attachment
Strength (Hoffmann et al., 2009)	0	Complete absence of adhesions
	1	Gentle traction requiring little effort to break adhesion
	2	Blunt dissection requiring a blunt instrument to break adhesion
	3	Sharp dissection requiring a sharp instrument to break adhesion
	4	Adhesion of all viscera to the abdominal wall

6.3.2 Histology

6.3.2.1 Tissue processing and microtome sectioning

Tissue samples are collected and stored in 4% (w/v) paraformaldehyde at 4°C overnight. On the following day, the tissues were transferred into 70% (v/v) ethanol and stored at 4°C overnight. After, tissues were processed using the HistoCore PEARL (Leica, Germany) tissue processor and embedded in paraffin with the HistoCore Arcadia C (Leica, Germany). Following, tissue samples were sectioned at 5 µm thickness using a Leica RM2245 microtome (Leica, Germany) and mounted

onto Polysine[®] microscope slides and dried at 40°C for at least an hour before staining.

6.3.2.2 Haematoxylin and eosin (H&E) staining

Slides were loaded into Leica Autostainer XL (Leica, Germany) and a set programme was run (refer to **Appendix II, Supplementary Table 4**). At the end of the programme, glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at room temperature (RT).

6.3.2.3 Picro-Sirius red (PSR) staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and DI water. Slides were then immersed in Picro-Sirius red (Abcam, United Kingdom) for 1 hour. After, slides were transferred into two changes of acetic acid followed by two changes of 100% (v/v) ethanol. Slides were cleared again with two changes of Clearene (Leica, Germany) before glass coverslips were mounted onto the slides using Organo mounting medium (Sigma, USA) and dried at RT.

6.3.2.4 Immunofluorescence staining

Briefly, slides were cleared with two changes of Clearene, before hydrating successively in 100%, 95%, and 70% (v/v) ethanol gradients and PBS. Depending on the antibody, either heat-induced or proteolytic-induced antigen retrieval methods may be performed (refer to below sections). If no antigen retrieval was required, slides were then immersed in permeabilisation/blocking buffer (0.3% (v/v) Triton-X100, 1M Lysine in PBS) for 1 hour at RT before incubation in antibody

buffer (1% (w/v) BSA in PBS) for 45 minutes. After incubation, the boundaries of the tissue samples were dried and a hydrophobic perimeter was drawn around the sample with a hydrophobic pen (Agilent Technologies, United States). Tissue samples were then immunolabeled with primary antibody diluted in blocking solution (refer to **Table 6.2** for dilutions and incubation conditions regarding primary antibodies) and incubated overnight at 4°C. Controls without primary antibodies were performed similarly with antibody buffer in the absence of any primary antibody. After primary antibody incubation, tissues were washed twice, 3 minutes each, using washing buffer (0.05% (v/v) Tween-20 in PBS) before 1 hour incubation at RT with the appropriate secondary antibody (refer to **Table 6.3** for dilutions regarding secondary antibodies). After incubation, tissues were washed twice, 3 minutes each, with washing buffer before staining with DAPI (Life Technologies, USA) diluted at 1 in 10000 with antibody buffer. Finally, slides were washed twice, 3 minutes each, in PBS prior to applying Citiflour™ AF1 (Electron Microscopy Sciences, London) anti-fade mounting medium before mounting coverslips and sealing with nail varnish.

6.3.2.5 Proteolytic-induced antigen retrieval (PIAR)

A 20X proteinase K stock solution (400 ug/ml) was prepared by dissolving 8 mg of proteinase K (Bioline, Singapore) in 50% (v/v) TE buffer (50 mM Tris Base, 1 mM EDTA, 0.5% (v/v) Triton X-100, pH 8.0), and 50% (v/v) glycerol. This stock solution was then diluted to a 1X working solution with TE Buffer. After rehydration, slides were submerged in the working solution and incubated for 5 minutes at 37°C. Following, slides were rested in the same solution for 5 minutes at RT before being washed in twice, 5 minutes each, with a proteinase K inactivating

wash buffer (1 mM PMSF in PBS). After, slides were immersed in antibody buffer and the standard staining protocol was performed. Slides were washed with proteinase K inactivating wash buffer instead of PBS prior to applying the anti-fade mounting medium. The remainder of standard staining protocol was performed as described earlier.

Table 6.2 List of primary antibodies used in this study

Name	Protocol	Reactivity	Dilution	Catalogue No.
Rabbit polyclonal anti-Cx43	IHC	Cx43	1:1000	C6219 (Sigma, USA)
Rabbit polyclonal anti- α SMA	IHC	α SMA	1:500	AB5694 (Abcam, UK)
Rat polyclonal anti-CD31	IHC + PIAR	CD31	1:25	553370 (BD Pharmingen, USA)

IHC – Immunohistochemistry staining, **HIER** – Heat induced antigen retrieval, **PIER** – Proteolytic-induced antigen retrieval.

Table 6.3 List of secondary antibodies used in this study

Name	Reactivity	Dilution	Catalogue No.
Goat anti-rat AF568	Rat	1:500	A11077 (ThermoFischer Scientific, USA)
Goat anti-rabbit AF488	Rabbit	1:500	A11008 (ThermoFischer Scientific, USA)
Goat anti-rabbit AF555	Rabbit	1:500	A21422 (ThermoFischer Scientific, USA)

6.3.3 Brightfield and confocal microscopy

Histological stained (H&E and PSR) tissue samples were automatically scanned and digitised into 24-bit images with an AxioScan.Z1 slide scanner (Zeiss, Germany) equipped with a 20X 0.8NA Plan-Apochromat air objective and a Hitachi HV-F202SCL. Scaling per pixel for these images is 0.22 μ m by 0.22 μ m. For the

visualisation of immunohistochemical stained tissue samples, Z-stacks of 10 optical sections were acquired using an upright confocal microscope TCS SP8 (Leica, Germany) with a 40X 1.2NA Plan-Apochromat or a 63X 1.32NA oil objective. Z-stacks were taken by first finding the centre position before proceeding to take slices above and below this centre point. The thickness of each Z is 0.699 μm . Fluorophores were excited sequentially using a 405 nm, 488 nm and 532 nm wavelength laser. The images were acquired at 8-bit greyscale at 1024 x 1024 pixels with a Z-series frame average of 2 or 3. Maximum projections were calculated using Leica Application Suite X (Leica, Germany) for each of the Z-stacks recorded. All images were captured with identical parameters and camera settings to allow for direct comparison of staining intensity.

6.3.4 Microscopic Assessment

6.3.4.1 Inclusion and exclusion criteria

Tissues samples were first subjected to a simple inclusion/exclusion criterion to identify samples with features that allowed fair comparison with one another. The inclusion criterion included only tissues sections with a visible button (refer to **Fig. 6.3a**) or with a visible button accompanied with sutures only at the base of the button (refer to **Fig. 6.3b**). The exclusion criterion rejected tissues sections with no visible button (refer to **Fig. 6.3c**) or with a visible button accompanied with sutures found at other locations other than at the base of the button (refer to **Fig. 6.3d**).

Figure 6.3 Inclusion and exclusion criteria

The inclusion criterion included only tissues sections with a visible button (A) or with a visible button accompanied with sutures only at the base of the button (B). The exclusion criterion rejected tissues sections with no visible button (C) or with a visible button accompanied with sutures found at other locations other than at the base of the button (D). **IB** – Ischaemic Button region, **S** – Suture region.

6.3.4.2 Fibrosis extent

During adhesion formation, fibrin deposits that persist allow the invasion and proliferation of fibroblasts, leading to the deposition of extracellular matrix proteins (e.g. collagen). As the adhesion develops and matures, the matrix becomes increasingly filled with more extracellular matrix proteins over time, eventually transforming the matrix into a more permanent fibrous connection. At an earlier stage (e.g. 7 days as opposed to 21 days) during adhesion development, the entire matrix may not necessarily be “filled” up with collagen, or other matrix proteins. The assessment of the extent of the fibrotic area (or matrix) is a good measure of area of the matured adhesion. In order to facilitate this analysis, several regions of the IB sample were further identified. The entire fibrotic (EF) region was identified as the overall region that included the immediate fibrotic (IF) region and the regions of adhesion-inducing determinants, such as ischaemic button and suture (IBS) (refer to **Fig. 6.4**). The EF region was defined by region occupied by the abnormal

encroaching of extracellular matrix that is observed across the entire ischaemic button tissue region and is bordered at the base of the ischaemic button by thickened mesothelium. The IBS region includes the ischaemic button region and the suture material region. The ischaemic button can be defined by the “ball of muscle” (due to the out-pouching) and can be distinguished from the matrix due to its pinkish characteristic. The suture material regions are determined by the presence of the suture material (greyish brown). The IF region is of the main interest and can be calculated by subtracting the area of the IBS region from the calculated area of the EF region. Note that these terms are also used in the following sections when determining fibrosis severity and protein levels of α SMA. Two ROIs were drawn on digital stitched images of tissue sections stained with H&E. One was drawn around the EF region and the other, around the IBS region. Areas of these regions were measured using image analysis software Zen Lite 2.3 (Zeiss, Germany) and used to calculate the area of IF region by subtracting the IBS region from EF region. Since the size of IBS region may influence the size of the IF region, the calculated area of the IF region is normalised to the area of IBS region to ensure fair comparison between the samples.

Figure 6.4 Defining regions around the ischaemic button

Representative digital stitched image of a section of the IB sample stained with H&E. The entire fibrotic (EF) region was identified as the total fibrotic region (area encompassed by red dashed lines). This is made up of the immediate fibrotic (IF) region and the regions of adhesion-inducing determinants such as the ischemic button and suture (IBS) (black dashed lines). The immediate fibrotic region (IF) may be determined by drawing ROIs to exclude the FR region from the IBS region or by the subtracting the FR region from IBS region. **B** – Ischaemic Button region, **S** – Suture region.

6.3.4.3 Fibrosis severity

ROIs were first drawn around the IF regions of the digital stitched images of tissue sections stained with PSR using ImageJ analysis software (version 1.51w; NIH) (**Fig. 6.5b**). These ROIs were saved for later use and documentation. Next, the amounts of collagen deposition were estimated using the greyscale threshold method described by Schipke et al., (2017) using ImageJ analysis software. Briefly, conversion of the acquired images to the RGB stacks generated greyscale images for red, green, and blue channels. Manual threshold was performed on the greyscale image corresponding to the green channel to detect collagen. Specific threshold values (80 to 255) were determined using at least 10 sections to ensure values were representative and then kept constant throughout image analysis. Next, an output mask is generated through manual thresholding at determined values (**Fig. 6.5c**).

The corresponding ROIs were loaded onto the respective output masks and the detected area was measured using the measure tool (**Fig. 6.5d**) and later normalised to the IBS area.

Figure 6.5 Example of image analysis workflow for PSR stained images.

(A) Example of PSR stained image to be analysed. (B) ROIs were first drawn around the IF regions of the digital stitched images of tissue sections stained with PSR using ImageJ. Next, the stained image to be analysed were converted to the RGB stacks, which generates greyscale images for red, green, and blue channels. Manual threshold was performed on the greyscale image corresponding to the green channel to detect collagen, which would produce an output mask. (C) Example of output mask after manual threshold (80 to 255). Specific threshold values were kept constant throughout analysis. (D) Example of the loading of corresponding ROIs for the respective sections so that the measure tool can be used to detect the area of interest in ImageJ.

6.3.4.4 Protein levels of α SMA

ROIs were drawn around the IF region in the digital stitched images of tissue sections immunostained with α SMA. Areas of blood vessels in which strong α SMA staining can be found, were excluded from the ROIs. Protein levels of α SMA in these regions were determined using in-house quantification algorithms using ImageJ analysis software (version 1.51w; NIH). Briefly, the α SMA levels were determined by quantifying the area in the binary mask that is generated after performing manual threshold. Specific threshold values (80 to 255) were kept constant throughout image analysis. The detected area was measured using the measure tool and later normalised to the IBS area.

6.3.5 Statistical Analysis

All data is presented as mean value $\pm$ standard deviation. In general, all statistical comparisons made between EC-Cx43^{fl/fl} and EC-Cx43^{-/-} groups were made using a Mann-Whitney *U* test, unless mentioned otherwise. Statistical comparisons made between EC-Cx43^{fl/fl} and EC-Cx43^{-/-} for the analysis of endothelial-Cx43 knockdown in the blood vessels was performed using a Student's *t*-test. All statistical comparisons were made using GraphPad Prism version 6.0f for Mac, (GraphPad™ Software, San Diego, USA).

6.4 Results

6.4.1 Endothelial-Cx43 protein levels in tamoxifen-inducible endothelial cell-specific Cx43 transgenic mice were significantly reduced upon tamoxifen treatment.

In order to investigate the role of endothelial-Cx43 in adhesion formation, tamoxifen-inducible endothelial cell-specific Cx43 transgenic mice (Tie2-CreER^{T2}Cx43^{fl/fl}) were given tamoxifen to induce Cre-mediated DNA recombination (now termed EC-Cx43^{-/-}). To assess the efficiency of recombination, dorsal skins harvested from the EC-Cx43^{-/-} group and its tamoxifen-treated Cre-absent controls (EC-Cx43^{fl/fl}) were co-immunolabeled for Cx43 and endothelial cell marker, CD31, to confirm the distribution of Cx43 in the blood vessel regions (**Fig. 6.6**). However, image analysis could not be performed on this set of immunostained sections because proteolytic-enzymatic antigen retrieval is required for the immunolabeling of CD31, which in the process may introduce artifacts that hamper

accurate image analysis. Therefore, image analysis of endothelial-Cx43 protein levels of these mice and their tamoxifen-treated Cre absent controls (EC-Cx43^{fl/fl}) was performed separately on a set of Cx43 immunolabeled sections that did not receive any form of antigen retrieval (**Fig. 6.7a**). Image analysis performed on this set of sections revealed that recombination efficiency was 57% ($p \leq 0.01$) (**Fig. 6.7b**). This suggested significant knockdown of endothelial-Cx43 in the EC-Cx43^{-/-} group and indicated its suitability to be used to study the role of endothelial-Cx43 in adhesion formation.

Figure 6.6 Endothelial-Cx43 protein levels in the tamoxifen-treated endothelial-specific Cx43 transgenic mice EC-Cx43^{-/-} was reduced upon tamoxifen treatment.

(A) The panel shows representative overlay and accompanying single channel immunofluorescence images of Cx43 protein levels in the blood vessel regions of skin samples taken from the dorsal region of the tamoxifen-treated mice. Endothelial-Cx43 (green, highlighted with green arrows) were visualised on longitudinally sectioned skin samples immunolabeled for Cx43 (green, highlighted with green arrows), endothelial cell marker CD31 (red, highlighted with red arrows), and counterstained with DAPI (blue). Endothelial-Cx43 protein levels in EC-Cx43^{-/-} mice appeared to be lower when compared EC-Cx43^{fl/fl} mice. White dashed lines represent the blood vessel regions that

are informed through the presence of positive CD31 staining. Note that antigen retrieval is needed for the immunolabeling of CD31 and may have interefered with Cx43, as they look “dustier” compared to immunostaining that were not subjected to antigen retrieval in **Figure 6.7**. Therefore, image analysis for Cx43 was not performed on this batch of immunostained sections. Magnification used, 63X. Scale bar = 25 μ m.

Figure 6.7 Endothelial-Cx43 protein levels in the tamoxifen-treated EC-Cx43^{-/-} transgenic mice was significantly reduced upon tamoxifen treatment.

(A) The panel shows images of Cx43 protein levels in the blood vessel regions of skin samples taken from the dorsal region of the tamoxifen-treated endothelial specific (EC-Cx43^{-/-}) transgenic mice.

Endothelial-Cx43 were visualised on longitudinally sectioned skin samples immunolabeled for Cx43 (red, highlighted with white arrows) and counterstained with DAPI (blue). Endothelial-Cx43 protein levels in EC-Cx43^{-/-} mice were lower compared to EC-Cx43^{fl/fl} mice. White dashed lines represent the blood vessel regions. Magnification used, 63X. Scale bar = 25 μ m. **(B)** Image analysis revealed significantly lower endothelial-Cx43 protein levels in EC-Cx43^{-/-} mice (n=8) compared EC-Cx43^{fl/fl} mice (n=9). Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using an unpaired Student's t-test with Welch's corrections (**p $\leq$ 0.01).

6.4.2 Knockdown of endothelial-Cx43 protein levels did not reduce adhesion formation.

After determining that endothelial-Cx43 protein levels was significantly reduced in the EC-Cx43^{-/-} group, its effect on adhesion formation was assessed at 7 days post-surgery in an ischaemic button (IB) model of adhesion formation. Macroscopic evaluation of the adhesions in the peritoneal cavity did not reveal notable differences in developed adhesions between the tamoxifen-treated groups (**Fig. 6.8a**). It is important to point out that at least 90% of the detected adhesions in EC-Cx43^{fl/fl} group were found distributed to the cranial abdominal region at necropsy (**Fig. 6.8b**). This is unusual considering that adhesions were previously found to be mainly distributed to the caudal abdominal region (65%) in the non-tamoxifen treated mice in **Chapter 5 (Fig. 5.7h)**. Adhesion distribution to the caudal abdominal region was attributed to the prominence of the epididymal fat pad in the same region. Considering this, it is hypothesised that tamoxifen treatment is likely to be responsible for the observed difference in distribution of adhesions between the two studies (**Chapter 5, Fig. 5.7h; Chapter 6, Fig. 5.7b**), by affecting epididymal fat. This prompted a retrospective literature search, which found that administration of tamoxifen (1mg/20g body weight) is sufficient to significantly reduce epididymal fat pad weight (Liu, L, et al., 2015). Considering that the presence of epididymal fat pad appeared to increase the extent of adhesions in the caudal abdominal region

(**Chapter 5, Fig. 5.7h**) and that the reduction of epididymal fat pad upon tamoxifen treatment may have reduced the extent of adhesion in the same region (**Chapter 6, Fig. 6.8b**), analyses in this study were only performed on the IBs in the cranial abdominal region.

Macroscopic evaluation of the adhesions detected in the cranial abdominal region was conducted in a blinded manner by 3 observers using established scoring matrices (**Table 6.1**) (Adhesion Scoring Group, 1994; Hoffmann et al., 2009; Nair et al., 1974). First, the adhesions involvement was assessed, in which a lower score for involvement was assigned if fewer adhesive bands were found attached to the button (**Table 6.1**) (Nair et al., 1974). The knockdown of endothelial-Cx43 did not appear to affect the development of adhesions ($p=0.214$) (**Fig. 6.8c**). The detected adhesions were then evaluated for their strength (Hoffmann et al., 2009). Adhesive bands with lower tensile strength, requiring less effort to break, received lower scores (**Table 6.1**). Most of the adhesions in the EC-Cx43^{fl/fl} group required blunt dissection to break (30.6 ± 18.8) while those found in the EC-Cx43^{-/-} group required slightly less effort to break (17.3 ± 20.5) (**Fig. 6.8a**). Adhesions in both groups were also subsequently evaluated for severity (Adhesion Scoring Group, 1994). Based on the score matrix, adhesive bands that were more translucent and less vascular adhesive bands would receive a lower score (**Table 6.1**). Slightly opaquer and sometimes denser adhesions were detected in the EC-Cx43^{fl/fl} group (28.2 ± 16.7) when compared to the EC-Cx43^{-/-} group (15.4 ± 18.2) (**Fig. 6.8a**). Statistical comparison of the macroscopic assessments of the detected adhesions revealed that knockdown of endothelial-Cx43 did not significantly reduce adhesion strength ($p=0.317$) and severity ($p=0.285$) at 7 days post-surgery (**Fig. 6.8d and e**).

Figure 6.8 Knockdown of endothelial-Cx43 protein levels did not appear to affect adhesion developed at 7 days post-surgery.

(A) Examples of peritoneal adhesions detected in the abdominal region at 7 days post-surgery (highlighted by green arrowheads). (B) Majority of the adhesions of the EC-Cx43^{fl/fl} mice at 7 days appeared to be mainly distributed in the cranial abdominal region. Developed adhesions in the cranial abdominal region at 7 days post-surgery in the EC-Cx43^{fl/fl} (n=6) and EC-Cx43^{-/-} (n=9) groups were assessed for qualities such as, involvement (C), severity (D), and strength (E), and scored by 3 observers blinded to the groups. The mean observer scores for each IB in the cranial abdominal region were averaged for each mouse and graphed as a percentage of maximum score possible for each parameter. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons between the groups were made using a Mann-Whitney *U* test did not find any significant difference.

6.4.3 The role of endothelial-Cx43 on adhesion formation.

To further verify the macroscopic observations, subsequent microscopic evaluation of the extent (Fig. 6.9) and severity (Fig. 6.10) of fibrosis in the IB samples was performed by using H&E and PSR histological stains respectively. To determine the extent of fibrosis, immediate fibrotic regions were measured through image analysis. The extent of fibrosis in the EC-Cx43^{-/-} group (0.23 ± 0.11) was similar to that of the EC-Cx43^{fl/fl} group (0.27 ± 0.2) ($p > 0.05$). Subsequently, measuring the amount of

collagen fibres stained red with PSR further accessed the severity of fibrosis within the immediate fibrotic regions. Comparison of measurements of collagen fibres stained red with PSR within these regions also did not reveal any significant differences between the two groups ($p=0.452$). Protein levels of α SMA, a marker for myofibroblasts, appear to be higher in the EC-Cx43^{-/-} group compared to the EC-Cx43^{fl/fl} group when assessed qualitatively. However, when assessed in this manner, the detected area of α SMA signal is not normalised to the IBS area, which may be misleading. Furthermore, areas of blood vessels contain strong α SMA signal may also give a false sense of high α SMA signal in the immediate fibrotic region. Therefore, quantitative measurement of α SMA in the immediate fibrotic regions of IB samples using image analysis methods to exclude areas of blood vessels was also performed. Image analysis found that protein levels of α SMA from the EC-Cx43^{-/-} group (0.17 ± 0.13) was comparable to that determined in the EC-Cx43^{fl/fl} group (0.23 ± 0.19) ($p=0.452$) (**Fig. 6.11**). Together, findings from the microscopic assessment corroborated the macroscopic observations made of the strength and severity of the adhesions that the down-regulation of endothelial-Cx43 had little effect. To further validate this conclusion, the protein levels of tPA and its inhibitor, PAI-1, in the peritoneal cavity were investigated to compare the overall plasminogen-activating activity in both groups. The knockdown of endothelial-Cx43 protein levels did not appear to affect the protein level of tPA in the peritoneal fluid. Similarly, no significant changes in protein level of PAI-1 in the peritoneal fluid were noted between the EC-Cx43^{-/-} groups and the EC-Cx43^{fl/fl} group ($p=0.370$) (**Fig. 6.12**). These observations suggested similar PAA activity in both groups, further confirming both macroscopic and microscopic assessments made of the adhesions in both groups.

Figure 6.9 Knockdown of endothelial-Cx43 protein levels did not appear to affect fibrosis extent of the adhesions developed at 7 days post-surgery.

Representative digital stitched images of H&E-stained IB samples of the EC-Cx43^{-/-} mice (**A**, n=6) and EC-Cx43^{fl/fl} mice (**B**, n=9) at post-surgery day 7, and their respective magnified regions (*i*, *ii*). (**C**) Image analysis did not reveal any significant changes in fibrotic extent of the adhesions in the EC-Cx43^{-/-} mice when compared to the EC-Cx43^{fl/fl} mice. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test. Dashed lines highlight boundary of the immediate fibrotic region. **IB** – ischaemic button; **s** – suture; **m** – muscle; **IF** – immediate fibrotic region.

Figure 6.10 Knockdown of endothelial-Cx43 protein levels did not appear to affect fibrosis severity of the adhesions developed at 7 days post-surgery.

Representative digital stitched images of PSR-stained IB samples of the EC-Cx43^{-/-} mice (A, n=6) and EC-Cx43^{fl/fl} mice (B, n=9) at post-surgery day 7, and their respective magnified regions (i, ii). (C) Image analysis did not reveal any significant changes in fibrotic severity of the adhesions in the EC-Cx43^{-/-} mice when compared to the EC-Cx43^{fl/fl} mice. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test. Dashed lines highlight boundary of the immediate fibrotic region. **IB** – ischaemic button; **s** – suture; **m** – muscle; **IF** – immediate fibrotic region.

Figure 6.11 Knockdown of endothelial-Cx43 protein levels did not appear to affect α SMA protein levels within the fibrotic region at 7 days post-surgery.

Representative digital stitched immunofluorescence images of images of IB samples of the EC-Cx43^{-/-} mice (**A**, n=6) and EC-Cx43^{fl/fl} mice (**B**, n=9) at post-surgery day 7, and their respective magnified regions (**i**, **ii**). IB sections were immunolabeled for α SMA (green), and counterstained with DAPI (blue). (**C**) Image analysis did not reveal any significant changes in α SMA protein levels within the immediate fibrotic region of the EC-Cx43^{-/-} mice when compared to the EC-Cx43^{fl/fl} mice. Data is presented as mean value $\pm$ standard deviation. Statistical comparisons were made using a Mann-Whitney *U* test. Dashed lines highlight boundary of the immediate fibrotic region. **IB** – ischaemic button; **s** – suture; **m** – muscle; **IF** – immediate fibrotic region.

Figure 6.12 Effect of endothelial-Cx43 knockdown on tissue plasminogen activator (tPA) and plasminogen activator inhibitor-1 (PAI) proteins in the peritoneal fluid at day 7 post-surgery. Knockdown of endothelial-Cx43 protein levels did not appear to affect (A) tPA protein levels or (B) PAI-1 protein levels at day 7 post-surgery. Statistical comparisons between EC-Cx43^{-/-} mice (A, n=1; B, n=8) and EC-Cx43^{fl/fl} mice (A, n=6; B, n=9) were made using a Mann-Whitney *U* test did not find any significant difference between the two groups.

6.4 Discussion

6.4.1 Potential effects of Tamoxifen on the model

Understanding the role of cell-specific connexins under pathological conditions has been made possible with the generation of numerous transgenic mice models (Chadjichristos et al., 2010; Westphalen et al., 2014). For this purpose, the Cre/lox site-specific recombination system is useful for the generation of conditional transgenic mice, specifically controlling the expression and activity of genes of interest in the desired cells (Gu et al., 1993). In addition, the Cre recombination event may be further controlled temporally. This is achieved through the use of fusion proteins that encompass Cre recombinase containing steroid hormone ligand-binding domains (Forde et al., 2002; Gu et al., 1993; Metzger et al., 1995). In this study, tamoxifen-inducible cell-specific Cx43 transgenic mice were treated with tamoxifen to induce Cre recombination. This was performed so that endothelial-

Cx43 protein levels may be reduced for the investigation of the role of endothelial-Cx43 in adhesion formation.

As mentioned earlier, a retrospective literature search found that administration of tamoxifen (1mg/20g body weight) in male Cre-absent transgenic mice is able to reduce body fat mass by 34%, which was later found to also significantly reduce epididymal adipose tissue ($P \leq 0.01$) (Liu et al., 2015). In the same study, tamoxifen was shown to increase apoptosis and autophagy markers in the sampled epididymal adipose tissue. Building on this, the investigators suggested that reactive oxygen species and oxidative stress were driving these processes and later showed increase in heme oxygenase 1, a sensitive marker of cellular oxidative stress, upon tamoxifen treatment (Liu et al., 2015; Poss & Tonegawa, 1997). Although, epididymal fat pad was not measured, it is likely that it was reduced by tamoxifen treatment. Coupled with the observation that only 10% of the detected adhesions in tamoxifen-treated EC-Cx43^{fl/fl} group were found distributed to the caudal abdominal region at necropsy, this observation supports the belief that epididymal fat pads facilitate adhesion formation in IB models of adhesion formation (Mommers et al., 2018). In light of this, only adhesions in the cranial abdominal region were analysed in this study.

The effect of tamoxifen treatment on adhesion formation remains controversial. It was shown to reduce adhesion formation in a cecal abrasion rat model assessed at 30 days post-operation (Karaca et al., 2013). However, no significant differences were observed between placebo and tamoxifen treated rats at day 3, 5 and 7 post-surgery in rat model studying anastomic healing (McNamara, Walsh, Kay, &

Bouchier-Hayes, 2003). Unfortunately, differences in models employed, tamoxifen dosage, and the timing of assessments make it difficult to compare both studies. This is further mired by the lack of experimental description; for the former study, it was not mentioned how long tamoxifen treatment was carried out for (Karaca et al., 2013) and no full-text could be retrieved for the latter study (McNamara et al., 2003). Regardless, these studies highlight the importance of having a washout period before experimental studies are conducted to reduce potential “off-target” tamoxifen effects and also having tamoxifen-treated controls when using tamoxifen for inducing Cre-mediated recombination. Both measures were adopted in this study so that experimental outcomes between in both experimental groups would be meaningful.

6.4.2 Endothelial Cx43 does not appear to be important for adhesion formation

In this study, macroscopic assessment of the detected adhesions between the EC-Cx43^{fl/fl} and EC-Cx43^{-/-} groups revealed no significant changes in adhesion strength and severity at 7 days post-surgery. These findings were corroborated by microscopic assessment of the IB samples and evaluation of the overall plasminogen-activating activity. Due to time and logistical constraints, assessment was not performed to compare the early phase of adhesion formation between the groups.

Prior work in **Chapter 5** had shown significant reduction in number of leukocytes recruited within the IB region at 6 and 24 hours post-surgery in the Cx43asODN-treated group. The same group was also shown to have significantly reduced

adhesion severity and strength at day 7 post-surgery, supporting previous findings that leukocyte recruitment is important for the development of adhesions (Hoshino et al., 2007; Vural et al., 1999). These observations also suggested Cx43 might have an important role in leukocyte recruitment. As mentioned earlier, Scheckenbach and colleagues have proposed that Cx43-dependent calcium signaling between endothelial cells supports leukocyte adhesion in the presence of a pro-inflammatory stimulus (Scheckenbach et al., 2011). This is based on observations that lateral propagation of calcium wave signaling along the endothelium of an endothelial Cx43 knockout mouse was significantly reduced (Parthasarathi et al., 2006). This reduction was thought to hamper endothelial P-selectin exocytosis, which mediates neutrophil rolling during initial recruitment events. This may explain why Véliz and colleagues found that the challenge of the endothelial-Cx43 knockout mouse with TNF- α hardly yielded any increase of leukocyte adhesion and transmigration events (Véliz et al., 2008).

Alternatively, the lack of increase in leukocyte adhesion and transmigration events in these endothelial Cx43 knockout mouse may also suggest that the leukocytes are refractory towards TNF- α (Veliz et al., 2008). It is important to point out that, although not statistically significant, the baseline leukocyte adhesion and transmigration events already appeared to be higher in the saline-injected endothelial Cx43 knockout mouse compared to wild-type saline-injected controls in this study (Veliz et al., 2008). This may indicate abnormal leukocyte behaviour. On a side note, it is important to note that the transgenic mice used by Parthasarathi and Véliz are manipulating Cx43 expression under the influence of a constitutive *Tie2* promoter, which may have caveats as discussed earlier in the introduction (**Section**

6.1.4). Hence, these observations of abnormal leukocyte behaviour at baseline (i.e. without pro-inflammatory stimulation) may be partially contributed to the manipulation of Cx43 on other haematopoietic cells due to the constitutive *Tie2* promoter, as earlier discussed. In light of this, these findings from Parthasarathi and Veliz warrant careful interpretation.

Recent work investigating gap junction-mediated regulation on endothelial cellular stiffness may help shed more light on leukocyte adhesion and transmigration events in the tamoxifen-inducible transgenic Cx43 mice used in this study (Okamoto & Suzuki, 2017). The degree of endothelial cellular stiffness may influence the regulation of leukocyte adhesion, spreading, and migration (Huvencers, Daemen, & Hordijk, 2015; Sun, Wu, & Yuan, 2011). Stiffer endothelial cells are able to increase the cell spreading of neutrophils by serving as an appropriate substrate for adhered neutrophils (Stroka & Aranda-Espinoza, 2011). Okamoto and colleagues demonstrated significant increase in endothelial stiffness in response to pro-inflammatory stimulus, TNF- α (Okamoto & Suzuki, 2017). It is important to point out that the same group also showed earlier that TNF- α treatment of endothelial cells does not significantly change the protein levels of Cx43 (Okamoto et al., 2011). The authors also transiently increase endothelial stiffness by reducing Cx43 activity using carbenoxolone, a broad-spectrum gap junction inhibitor, or a Cx43 monoclonal antibody (Okamoto & Suzuki, 2017). Following on, further increases of endothelial stiffness, which persisted for at least 24 hours, was shown upon the simultaneous application of both carbenoxolone and TNF- α compared to a single application of carbenoxolone or TNF- α (Okamoto & Suzuki, 2017).

Based on these findings, it is likely that neutrophil adhesion, spreading, and migration events in the EC-Cx43^{-/-} group may be more pronounced during the early phase of adhesion formation considering that the knockdown of endothelial-Cx43 in the group and TNF- α is being released during early adhesion formation (Beyene et al., 2015) (Stroka & Aranda-Espinoza, 2011).

In contrast, monocyte recruitment events might not follow the same manner as neutrophils in the EC-Cx43^{-/-} group. In a study to determine which cell-derived Cx43 is a stronger contributor to monocyte adhesion, freshly extracted monocytes and activated endothelial monolayer were treated with Gap26, a Cx43 connexin mimetic peptide (Abed et al., 2014). Gap26-treated monocytes saw significantly less monocyte adhesion to the endothelial monolayer, compared to Gap26-treated activated endothelial monolayer (Abed et al., 2014). This suggested that monocyte-Cx43 is a stronger contributor to adhesion events. Inferring from this finding, it suggests that monocyte recruitment events are likely to be reduced in the Cx43asODN-treated group in **Chapter 5** since Cx43asODN treatment is not specific to a particular cell type. Conversely, it suggests monocyte recruitment events are not affected in the EC-Cx43^{-/-} group in this study since Cre-mediated recombination of Cx43 occurred only in the endothelial cells.

In summary, the EC-Cx43^{-/-} group is inferred to have more pronounced neutrophil adhesion, spreading, and migration events while monocyte recruitment events are not affected. These hypothesised characteristics of the EC-Cx43^{-/-} group in leukocyte recruitment contribute little to adhesion formation considering no significant differences in adhesion outcomes were detected between the EC-Cx43^{-/-} and EC-

Cx43^{fl/fl} groups. To help elucidate this, future work should consider the staining of markers specific to these leukocyte populations in both experimental groups during early adhesion formation.

6.6 Conclusions

Work in this chapter set out to explore the contribution of endothelial-Cx43 to adhesion formation. Effects of tamoxifen-induced endothelial-Cx43 knockdown on late phase adhesion formation were studied using tamoxifen-inducible endothelial-Cx43 specific transgenic mice in ischaemic button model of adhesion formation. No notable differences in adhesion outcomes were observed between EC-Cx43^{-/-} and EC-Cx43^{fl/fl} groups, it indicates that endothelial-Cx43 knockdown does not play a significant role to adhesion formation. Also, it is hypothesised that increased neutrophil adhesion, spreading, and migration events due to endothelial-Cx43 knockdown contribute little to subsequent adhesion formation. However, more work using specific leukocyte markers would be required to elucidate this. Overall, the results from this chapter imply that other cell-derived Cx43 take on a more significant role in the events of adhesion pathogenesis. Further investigation into the contribution of Cx43 from other major cell types involved in this series of events will open up new strategies for fine-tuning connexin-based therapeutics for post-surgical peritoneal adhesions.

Chapter 7: Summary and Conclusion

Almost like a surgical footprint, 93% of patients undergoing intra-abdominal operations have been reported to develop adhesions (Ellis, 1997). Their presence may manifest a series of clinical issues, such as SBO, secondary infertility, chronic abdominal pain, adhesiolysis and related complications (ten Broek et al., 2016). Also, it increases the difficulty of repeated operations and length of recovery (Beck et al., 2000; Coleman et al., 2000). In 2001, the total cost of hospital and surgical expenditures for the management of adhesion-related complications was estimated to be at least \$5 billion in the USA. Considering that post-surgical peritoneal adhesions still represent a challenging clinical problem of significant magnitude and economic burden, much more research is warranted. Little is understood about the role connexins in post-surgical peritoneal adhesions formation, specifically Cx43.

This thesis hypothesises that Cx43 has an important role during early adhesion pathogenesis, and that a single treatment with Cx43asODN during this period will transiently reduce Cx43 protein levels, resulting in better adhesion formation outcomes due to fewer recruited leukocytes to the injury site, reduced activated fibroblasts, and promotion of plasminogen-activating activity. Therefore, this project aims to fill these research gaps by approaching it systematically using *ex vivo* and *in vivo* adhesion models.

In **Chapter 3**, a new mouse peritoneal strip *ex vivo* adhesions model was established. Reproducible adhesion bands were macroscopically observed after 72 hours in

cultures of peritoneal strips in which a standardised volume of clotted blood was placed between abraded surfaces. The presence of adhesions was identified in microscopic assessment through the presence of eosinophilic matrices between juxtaposing mesothelium and from the positive staining of collagen fibres stained red with PSR in these matrices. High levels of α SMA were also observed in the surrounding mesothelium peripheral to the matrices expressed, suggesting mesothelial-to-mesenchymal transition and sub-mesothelial fibroblast activation to the myofibroblasts. Elevation in Cx43 protein levels was also observed within the same region, serving as a first line of evidence that Cx43 may be implicated during adhesion formation. Unfortunately, tracking of the exact point of adhesion creation during early culture (i.e. first 24 hours) proved to be difficult, limiting this model for only the study of later molecular events or for assessing efficacy of therapeutics. In addition, there was a lack of circulation and the contribution of the immune component to the overall adhesion formation in this model. These limitations prompted the switch of focus to *in vivo* models of adhesion formation for the study of Cx43 role in adhesion formation.

Work in **Chapter 4** attempted to identify the most ideal model of adhesion formation for this inquiry. Available models were identified through a systematic literature search and the findings were discussed with respect to the selection of species, anatomical sites, adhesiogenic stimuli, and macroscopic assessment. Next, relevant models were explored and the ischaemic button (IB) model was concluded to be the most ideal model. This conclusion was based on established macroscopic assessment scoring for the quality and quantity of adhesions, and also evaluated based other considerations (e.g. degree of model standardisation, identification and

definition of site trauma). Adhesions were generated by this model by creating an IB through a ligation of an out-pouching of the abdominal sidewall with a suture. Subsequent adhesions generated were only detected at the defined traumatised site, allowing for the standardisation in adhesion measurements and tissue sampling. Importantly, the defined trauma site in the model facilitates the tracking of adhesion development, which is essential for studying early and late molecular events (i.e. extent of inflammation).

The role of Cx43 in post-surgical peritoneal adhesion formation was explored in an ischaemic button model of adhesion formation in **Chapter 5**. Pathological increase of Cx43 protein levels was observed in the mesothelial regions within the ischaemic button at 6 and 24 hours post-surgery. The increase was also accompanied by observations of increases in leukocyte recruitment. The significance of this pathological increase was investigated with the use of Cx43asODN. Treatment with Cx43asODN significantly suppressed pathological increases in Cx43 protein levels in the mesothelial regions within the ischaemic button.

Conclusion from later investigations suggested that up-regulation of Cx43 plays an important role in adhesion formation. Treatment with Cx43asODN was found to reduce macroscopic scores of adhesion strength and severity at 7 days post-surgery. This was further validated by notable reductions in microscopic assessment of fibrosis extent and severity, and the degree of α SMA protein levels. Also, levels of PAI-1 was also found to be significantly reduced. Further investigations into earlier phase of adhesion formation suggested that Cx43 mediates adhesion formation by regulating leukocyte recruitment and TGF- β 1 protein levels during early adhesion

formation. Attempts to further dissect the contribution of Cx43 are detailed in **Chapter 6**. The role of endothelial-Cx43 in adhesion formation was explored using tamoxifen-inducible endothelial-Cx43 specific transgenic mice in the IB model of adhesion formation. The investigation revealed that endothelial-Cx43 knockdown does not play a significant role to adhesion formation.

Future extensions of this work would include exploring other dosing regimens, such as prophylactic dosing or multiple dosing post-surgery. Also, considering most of the fibrotic reaction occurred around the sutures in this model, it would be interesting to explore the use of Cx43 asODN coated sutures. More importantly, future work would mainly focus on staining different markers to identify populations that are involved Cx43-mediated adhesion formation. Not only would this help with the isolation of populations for *in vitro* studies or generation of new transgenic mice models, it would also pave the way for the identification of existing ligands or design of ligands that interacts selectively with the cell surface receptor of the cell type of interest. Conjugation of the selected ligand to the asODN or its nanocarriers may improve its selective delivery to its sites of action.

Overall, this study suggests Cx43 plays a significant role in the events of adhesion pathogenesis by regulating early leukocyte recruitment and TGF- β 1 levels. Findings also indicate this regulatory role is likely to be mediated by Cx43 derived from cells other than endothelial cells. Further investigation into the contribution of Cx43 from other major cell types involved in this series of events will open up new strategies for fine-tuning connexin-based therapeutics for post-surgical peritoneal adhesions.

Publications

Soon, A. S., **Chua, J. W.**, & Becker, D. L. (2016). **Connexins in endothelial barrier function - novel therapeutic targets countering vascular hyperpermeability.** *Thromb Haemost*, 116(5), 852-867. doi:10.1160/TH16-03-0210

Appendix

APPENDIX I: SUPPLEMENTARY MATERIALS AND METHODS

Supplementary Table 1. *Ex vivo* media and tissue collection reagent components

Component	Amount	Catalogue Number, Brand
<i>Ex Vivo Media</i>¹		
Glucose	3 g	G7201 (Sigma, USA)
L-Methionine	1.5 mg	M5308 (Sigma, USA)
L-Leucine	5.2 mg	L8912 (Sigma, USA)
L-Lysine-Monohydrochloride	72.5 mg	L8662 (Sigma, USA)
Sodium Bicarbonate	2.2 g	S6297 (Sigma, USA)
MEM powder	9.43 g	61100061 (ThermoFischer Scientific, USA)
MEM vitamin solution	10 mL	11120052 (ThermoFischer Scientific, USA)
Non-essential a.a solution	10 mL	11140050 (ThermoFischer Scientific, USA)
Insulin	2.387 mL	12585014 (ThermoFischer Scientific, USA)
Tissue Collection Reagents²		
Solution A		
Fetal Bovine Serum	15 g	SV30160.03 (HyClone, USA)
Solution B		
Penicillin Streptomycin	20 mL	1804438 (ThermoFischer Scientific, USA)
Amphotericin B	8 mL	15290018 (ThermoFischer Scientific, USA)
Solution C		
Penicillin Streptomycin	20 mL	1804438 (ThermoFischer Scientific, USA)
Amphotericin B	4 mL	15290018 (ThermoFischer Scientific, USA)

¹ Media brought up to 1 L with deionised water.

² Reagent brought up to 500 mL with 1X phosphate buffered saline.

Supplementary Table 2. Primers used for genotyping

Transgenic mice	Sequence	Size
Tie2-CreER^{T2}	5' – CGA GTG ATG AGG TTC GCA AG – 3'	320 bp
	5' – TGA GTG AAC GAA CCT GGT CG – 3'	
Cx43^{fl/fl}	5' – CTT TGA CTC TGA TTA CAG AGC TTA A – 3'	580 bp
	5' – GTC TCA CTG TTA CTT AAC AGC TTG A – 3'	

Supplementary Table 3. Sequence of mouse Cx43 antisense oligodeoxynucleotides

Name	Sequence
Cx43 asODN	5' – GTA ATT GCG GCA GGA GGA ATT GTT TCT GTC-3'

Supplementary Table 4. Set Programme for H&E staining

Step	Station	Time	Media	Step	Station	Time	Media
1	1	5 min	Clearene	12	Wash 3	2 min	Deionised water
2	2	5 min	Clearene	13	10	5 s	Scott's Water
3	3	2 min	100% Ethanol	14	Wash 4	2 min	Deionised water
4	4	2 min	100% Ethanol	15	11	2 min	70% Ethanol
5	5	2 min	95% Ethanol	16	12	30 s	Eosin
6	6	2 min	95% Ethanol	17	13	2 min	70% Ethanol
7	7	2 min	70% Ethanol	18	14	2 min	95% Ethanol
8	Wash 1	2 min	Deionised water	19	15	2 min	95% Ethanol
9	8	1 min	Hematoxylin	20	16	2 min	100% Ethanol
10	Wash 2	3 min	Deionised water	21	17	2 min	100% Ethanol
11	9	3 s	Acid Alcohol	22	18	2 min	Clearene

References

1. Aarons, C. B., Cohen, P. A., Gower, A., Reed, K. L., Leeman, S. E., Stucchi, A. F., & Becker, J. M. (2007). Statins (HMG-CoA reductase inhibitors) decrease postoperative adhesions by increasing peritoneal fibrinolytic activity. *Ann Surg*, 245(2), 176-184. doi:10.1097/01.sla.0000236627.07927.7c
2. Abed, A., Toubas, J., Kavvadas, P., Authier, F., Cathelin, D., Alfieri, C., . . . Chadjichristos, C. E. (2014). Targeting connexin 43 protects against the progression of experimental chronic kidney disease in mice. *Kidney Int*, 86(4), 768-779. doi:10.1038/ki.2014.108
3. Adhesion Scoring Group. (1994). Improvement of interobserver reproducibility of adhesion scoring systems. Adhesion Scoring Group. *Fertil Steril*, 62(5), 984-988.
4. Ahmad, G., O'Flynn, H., Hindocha, A., & Watson, A. (2015). Barrier agents for adhesion prevention after gynaecological surgery. *Cochrane Database Syst Rev*(4), CD000475. doi:10.1002/14651858.CD000475.pub3
5. Al-Ghadban, S., Kaissi, S., Homaidan, F. R., Naim, H. Y., & El-Sabban, M. E. (2016). Cross-talk between intestinal epithelial cells and immune cells in inflammatory bowel disease. *Scientific Reports*, 6, 29783. doi:10.1038/srep29783 <https://www.nature.com/articles/srep29783#supplementary-information>
6. Al-Jabri, S., & Tulandi, T. (2011). Management and prevention of pelvic adhesions. *Semin Reprod Med*, 29(2), 130-137. doi:10.1055/s-0031-1272475
7. Albrechtsen, O. K. (1957). The fibrinolytic activity of animal tissues. *Acta Physiol Scand*, 39(2-3), 284-290. doi:10.1111/j.1748-1716.1957.tb01429.x
8. Anastassiadis, K., Glaser, S., Kranz, A., Berhardt, K., & Stewart, A. F. (2010). A practical summary of site-specific recombination, conditional mutagenesis, and tamoxifen induction of CreERT2. *Methods Enzymol*, 477, 109-123. doi:10.1016/S0076-6879(10)77007-5
9. Andrews, S., Ford, D., & Martin, P. (2012). Knockdown of osteopontin reduces the inflammatory response and subsequent size of postsurgical adhesions in a murine model. *Am J Pathol*, 181(4), 1165-1172. doi:10.1016/j.ajpath.2012.06.027
10. Anselmet, A., Mayat, E., Wietek, S., Layer, P. G., Payraastre, B., & Massoulie, J. (2002). Non-antisense cellular responses to oligonucleotides. *FEBS Lett*, 510(3), 175-180.
11. Antoniak, S. (2018). The coagulation system in host defense. *Res Pract Thromb Haemost*, 2(3), 549-557. doi:10.1002/rth2.12109
12. Aoki, S., Udo, K., Morimoto, H., Ikeda, S., Takezawa, T., Uchihashi, K., . . . Toda, S. (2013). Adipose tissue behavior is distinctly regulated by neighboring cells and fluid flow stress: a possible role of adipose tissue in peritoneal fibrosis. *J Artif Organs*, 16(3), 322-331. doi:10.1007/s10047-013-0702-8
13. Api, M., Cikman, M. S., Boza, A., Rabus, M. B., Onenerk, M., & Aker, F. V. (2015). Peritoneal closure over barbed suture to prevent adhesions: a randomized controlled trial in an animal model. *J Minim Invasive Gynecol*, 22(4), 619-625. doi:10.1016/j.jmig.2015.01.013
14. Ara, C., Karabulut, A. B., Kirimlioglu, H., Yilmaz, M., Kirimlioglu, V., & Yilmaz, S. (2006). Protective effect of aminoguanidine against oxidative stress in an experimental peritoneal adhesion model in rats. *Cell Biochem Funct*, 24(5), 443-448. doi:10.1002/cbf.1245
15. Arora, M., Jaroudi, K. A., Hamilton, C. J., & Dayel, F. (1994). Controlled comparison of interceed and amniotic membrane graft in the prevention of postoperative adhesions in the rabbit uterine horn model. *Eur J Obstet Gynecol Reprod Biol*, 55(3), 179-182.
16. Arung, W., Drion, P., & Detry, O. (2016). Sepramesh and postoperative peritoneal adhesions in a rat model. *Acta Chir Belg*, 116(6), 357-361. doi:10.1080/00015458.2016.1181322
17. Arung, W., Meurisse, M., & Detry, O. (2011). Pathophysiology and prevention of postoperative peritoneal adhesions. *World J Gastroenterol*, 17(41), 4545-4553. doi:10.3748/wjg.v17.i41.4545
18. Asazuma-Nakamura, Y., Dai, P., Harada, Y., Jiang, Y., Hamaoka, K., & Takamatsu, T. (2009). Cx43 contributes to TGF- β signaling to regulate differentiation of cardiac fibroblasts into myofibroblasts. *Experimental Cell Research*, 315(7), 1190-1199. doi:<https://doi.org/10.1016/j.yexcr.2008.12.021>

19. Aysan, E., Ayar, E., Aren, A., & Cifter, C. (2002). The role of intra-peritoneal honey administration in preventing post-operative peritoneal adhesions. *Eur J Obstet Gynecol Reprod Biol*, 104(2), 152-155.
20. Aysan, E., Basak, F., Kinaci, E., Yanar, H., & Coskun, H. (2007). Experimental adhesion model: effect of viscosities of fluids put in the peritoneal cavity on preventing peritoneal adhesions. *Exp Anim*, 56(5), 349-354.
21. Aysan, E., Bektas, H., Ersoz, F., Sari, S., & Huq, G. (2010). Effects of contractubex on the prevention of postoperative peritoneal adhesion. *J Surg Res*, 164(2), 193-197. doi:10.1016/j.jss.2010.05.045
22. Bakkum, E. A., Dalmeijer, R. A., Verdel, M. J., Hermans, J., van Blitterswijk, C. A., & Trimbos, J. B. (1995). Quantitative analysis of the inflammatory reaction surrounding sutures commonly used in operative procedures and the relation to postsurgical adhesion formation. *Biomaterials*, 16(17), 1283-1289.
23. Bakkum, E. A., Emeis, J. J., Dalmeijer, R. A., van Blitterswijk, C. A., Trimbos, J. B., & Trimbos-Kemper, T. C. (1996). Long-term analysis of peritoneal plasminogen activator activity and adhesion formation after surgical trauma in the rat model. *Fertil Steril*, 66(6), 1018-1022.
24. Bakkum, E. A., & Trimbos-Kemper, T. C. (1995). Natural course of postsurgical adhesions. *Microsurgery*, 16(9), 650-654.
25. Bakkum, E. A., van Blitterswijk, C. A., Dalmeijer, R. A., & Trimbos, J. B. (1994). A semiquantitative rat model for intraperitoneal postoperative adhesion formation. *Gynecol Obstet Invest*, 37(2), 99-105. doi:10.1159/000292534
26. Balbinotto, R. P., Trindade, M. R., Muller, A. L., Nunes, A. G., Da Silva, R., Meyer, F. S., & Cerski, C. T. (2010). Experimental model of the formation of pelvic adhesions by videolaparoscopic in female rabbits. *Acta Cir Bras*, 25(1), 34-36.
27. Bayhan, Z., Zeren, S., Kocak, F. E., Kocak, C., Akcilar, R., Kargi, E., . . . Akcilar, A. (2016). Antiadhesive and anti-inflammatory effects of pirfenidone in postoperative intra-abdominal adhesion in an experimental rat model. *J Surg Res*, 201(2), 348-355. doi:10.1016/j.jss.2015.11.033
28. Beck, D. E., Cohen, Z., Fleshman, J. W., Kaufman, H. S., van Goor, H., Wolff, B. G., & Adhesion Study Group Steering, C. (2003). A prospective, randomized, multicenter, controlled study of the safety of Seprafilm adhesion barrier in abdominopelvic surgery of the intestine. *Dis Colon Rectum*, 46(10), 1310-1319. doi:10.1097/01.DCR.0000089117.62296.E4
29. Beck, D. E., Ferguson, M. A., Opelka, F. G., Fleshman, J. W., Gervaz, P., & Wexner, S. D. (2000). Effect of previous surgery on abdominal opening time. *Dis Colon Rectum*, 43(12), 1749-1753.
30. Becker, D. L., McGonnell, I., Makarenkova, H. P., Patel, K., Tickle, C., Lorimer, J., & Green, C. R. (1999). Roles for alpha 1 connexin in morphogenesis of chick embryos revealed using a novel antisense approach. *Dev Genet*, 24(1-2), 33-42. doi:10.1002/(sici)1520-6408(1999)24:1/2<33::aid-dvg5>3.0.co;2-f
31. Becker, J. M., & Stucchi, A. F. (2004). Intra-abdominal Adhesion Prevention: Are We Getting Any Closer? *Ann Surg*, 240(2), 202-204. doi:10.1097/01.sla.0000133118.38686.d0
32. Best, C. L., Rittenhouse, D., & Sueldo, C. E. (1991). A comparison of TC7 and 32% dextran 70 for prevention of postoperative adhesions in hamsters. *Obstet Gynecol*, 78(5 Pt 1), 858-860.
33. Beyene, R. T., Kavalukas, S. L., & Barbul, A. (2015). Intra-abdominal adhesions: Anatomy, physiology, pathophysiology, and treatment. *Curr Probl Surg*, 52(7), 271-319. doi:10.1067/j.cpsurg.2015.05.001
34. Bianchi, E., Boekelheide, K., Sigman, M., Lamb, D. J., Hall, S. J., & Hwang, K. (2016). Ghrelin ameliorates adhesions in a postsurgical mouse model. *J Surg Res*, 201(1), 226-234. doi:10.1016/j.jss.2015.10.044
35. Bizios, R., Lai, L., Fenton, J. W., 2nd, & Malik, A. B. (1986). Thrombin-induced chemotaxis and aggregation of neutrophils. *J Cell Physiol*, 128(3), 485-490. doi:10.1002/jcp.1041280318
36. Bolgen, N., Vargel, I., Korkusuz, P., Menciloglu, Y. Z., & Piskin, E. (2007). In vivo performance of antibiotic embedded electrospun PCL membranes for prevention of abdominal adhesions. *J Biomed Mater Res B Appl Biomater*, 81(2), 530-543. doi:10.1002/jbm.b.30694

37. Bouckaert, P. X., van Wersch, J. W., Schellekens, L. A., Evers, J. L., & Rolland, R. (1984). Haemostatic and fibrinolytic properties of peritoneal fluid in the menstrual cycle. *Br J Obstet Gynaecol*, 91(3), 256-259.
38. Bozkurt, S., Yuzbasioglu, M. F., Bulbuloglu, E., Gul, M., & Kale, I. T. (2009). Prevention of postoperative peritoneal adhesions by administration of estrogen. *J Invest Surg*, 22(4), 263-267.
39. Branch, A. D. (1998). A good antisense molecule is hard to find. *Trends Biochem Sci*, 23(2), 45-50.
40. Branes, M. C., Contreras, J. E., & Saez, J. C. (2002). Activation of human polymorphonuclear cells induces formation of functional gap junctions and expression of connexins. *Med Sci Monit*, 8(8), Br313-323.
41. Braun, K. M., & Diamond, M. P. (2014). The biology of adhesion formation in the peritoneal cavity. *Semin Pediatr Surg*, 23(6), 336-343. doi:10.1053/j.sempedsurg.2014.06.004
42. Brochhausen, C., Schmitt, V. H., Rajab, T. K., Planck, C. N. E., Krämer, B., Wallwiener, M., . . . Kirkpatrick, C. J. (2011). Intraperitoneal adhesions—an ongoing challenge between biomedical engineering and the life sciences. *Journal of Biomedical Materials Research Part A*, 98A(1), 143-156. doi:doi:10.1002/jbm.a.33083
43. Burns, E. M., Currie, A., Bottle, A., Aylin, P., Darzi, A., & Faiz, O. (2013). Minimal-access colorectal surgery is associated with fewer adhesion-related admissions than open surgery. *Br J Surg*, 100(1), 152-159. doi:10.1002/bjs.8964
44. Burns, J. W., Skinner, K., Colt, J., Sheidlin, A., Bronson, R., Yaacobi, Y., & Goldberg, E. P. (1995). Prevention of tissue injury and postsurgical adhesions by precoating tissues with hyaluronic acid solutions. *J Surg Res*, 59(6), 644-652. doi:10.1006/jsre.1995.1218
45. Burns, J. W., Skinner, K., Colt, M. J., Burgess, L., Rose, R., & Diamond, M. P. (1996). A hyaluronate based gel for the prevention of postsurgical adhesions: evaluation in two animal species. *Fertil Steril*, 66(5), 814-821.
46. Calder, B. W., Rhett, J. M., Bainbridge, H., Fann, S. A., Gourdie, R. G., & Yost, M. J. (2015). Inhibition of Connexin 43 Hemichannel-Mediated ATP Release Attenuates Early Inflammation During the Foreign Body Response. *Tissue Eng Part A*, 21(11-12), 1752-1762. doi:10.1089/ten.tea.2014.0651
47. Cassidy, M. R., Sherburne, A. C., Heydrick, S. J., & Stucchi, A. F. (2015). Combined intraoperative administration of a histone deacetylase inhibitor and a neurokinin-1 receptor antagonist synergistically reduces intra-abdominal adhesion formation in a rat model. *Surgery*, 157(3), 581-589. doi:10.1016/j.surg.2014.09.031
48. Cetin, M., Duran, B., Demirkoprulu, N., Guvenal, T., Erden, O., & Cetin, A. (2004). Effects of diazeniumdiolates (NONOates) and methylene blue on the reduction of postoperative adhesion in rats. *Gynecol Obstet Invest*, 57(4), 186-190. doi:10.1159/000076477
49. Chadjichristos, C. E., Scheckenbach, K. E., van Veen, T. A., Richani Saredidine, M. Z., de Wit, C., Yang, Z., . . . Kwak, B. R. (2010). Endothelial-specific deletion of connexin40 promotes atherosclerosis by increasing CD73-dependent leukocyte adhesion. *Circulation*, 121(1), 123-131. doi:10.1161/CIRCULATIONAHA.109.867176
50. Chaturvedi, A. A., Lomme, R. M., Hendriks, T., & van Goor, H. (2013). Prevention of postsurgical adhesions using an ultrapure alginate-based gel. *Br J Surg*, 100(7), 904-910. doi:10.1002/bjs.9131
51. Chegini, N. (2008). TGF-beta system: the principal profibrotic mediator of peritoneal adhesion formation. *Semin Reprod Med*, 26(4), 298-312. doi:10.1055/s-0028-1082388
52. Chilton, L., Giles, W. R., & Smith, G. L. (2007). Evidence of intercellular coupling between co-cultured adult rabbit ventricular myocytes and myofibroblasts. *J Physiol*, 583(Pt 1), 225-236. doi:10.1113/jphysiol.2007.135038
53. Cho, W. J., Oh, S. H., & Lee, J. H. (2010). Alginate film as a novel post-surgical tissue adhesion barrier. *J Biomater Sci Polym Ed*, 21(6-7), 701-713. doi:10.1163/156856209x435835
54. Chu, D. I., Lim, R., Heydrick, S., Gainsbury, M. L., Abdou, R., D'Addese, L., . . . Becker, J. M. (2011). N-acetyl-l-cysteine decreases intra-abdominal adhesion formation through the upregulation of peritoneal fibrinolytic activity and antioxidant defenses. *Surgery*, 149(6), 801-812. doi:10.1016/j.surg.2011.02.015
55. Coelho Junior, E. R., Costa, L. O., Alencar, A. V., Barbosa, A. P., Pinto, F. C., & Aguiar, J. L. (2015). Prevention of peritoneal adhesion using a bacterial cellulose hydrogel, in experimental study. *Acta Cir Bras*, 30(3), 194-198. doi:10.1590/s0102-865020150030000005

56. Cogliati, B., Da Silva, T. C., Aloia, T. P., Chaible, L. M., Real-Lima, M. A., Sanches, D. S., . . . Dagli, M. L. (2011). Morphological and molecular pathology of CCL4-induced hepatic fibrosis in connexin43-deficient mice. *Microsc Res Tech*, 74(5), 421-429. doi:10.1002/jemt.20926
57. Cogliati, B., Menecier, G., Willebrords, J., Da Silva, T. C., Maes, M., Pereira, I. V. A., . . . Vinken, M. (2016). Connexins, Pannexins, and Their Channels in Fibroproliferative Diseases. *J Membr Biol*, 249(3), 199-213. doi:10.1007/s00232-016-9881-6
58. Cohen, P. A., Gower, A. C., Stucchi, A. F., Leeman, S. E., Becker, J. M., & Reed, K. L. (2007). A neurokinin-1 receptor antagonist that reduces intraabdominal adhesion formation increases peritoneal matrix metalloproteinase activity. *Wound Repair Regen*, 15(6), 800-808. doi:10.1111/j.1524-475X.2007.00291.x
59. Coleman, M. G., McLain, A. D., & Moran, B. J. (2000). Impact of previous surgery on time taken for incision and division of adhesions during laparotomy. *Dis Colon Rectum*, 43(9), 1297-1299.
60. Confino, E., Friberg, J., Vermesh, M., Thomas, W., & Gleicher, N. (1988). Effects of progesterone on postoperative adhesion formation in hysterectomized rabbits. *Int J Fertil*, 33(2), 139-142.
61. Coughlin, S. R. (2005). Protease-activated receptors in hemostasis, thrombosis and vascular biology. *J Thromb Haemost*, 3(8), 1800-1814. doi:10.1111/j.1538-7836.2005.01377.x
62. Cronin, M., Anderson, P. N., Cook, J. E., Green, C. R., & Becker, D. L. (2008). Blocking connexin43 expression reduces inflammation and improves functional recovery after spinal cord injury. *Mol Cell Neurosci*, 39(2), 152-160. doi:10.1016/j.mcn.2008.06.005
63. Dai, P., Nakagami, T., Tanaka, H., Hitomi, T., & Takamatsu, T. (2007). Cx43 mediates TGF-beta signaling through competitive Smads binding to microtubules. *Mol Biol Cell*, 18(6), 2264-2273. doi:10.1091/mbc.e06-12-1064
64. Dalgic, T., Oymaci, E., Bostanci, E. B., Cakir, T., Kece, C., Erguder, I., & Akoglu, M. (2015). Effects of carbon dioxide pneumoperitoneum on postoperative adhesion formation and oxidative stress in a rat cecal abrasion model. *Int J Surg*, 21, 57-62. doi:10.1016/j.ijsu.2015.06.082
65. de Virgilio, C., Dubrow, T., Sheppard, B. B., MacDonald, W. D., Nelson, R. J., Lesavoy, M. A., & Robertson, J. M. (1990). Fibrin glue inhibits intra-abdominal adhesion formation. *Arch Surg*, 125(10), 1378-1381; discussion 1381-1372.
66. Delaney, K., Kasprzycka, P., Ciemerych, M. A., & Zimowska, M. (2017). The role of TGF-beta1 during skeletal muscle regeneration. *Cell Biol Int*, 41(7), 706-715. doi:10.1002/cbin.10725
67. Desmouliere, A., Geinoz, A., Gabbiani, F., & Gabbiani, G. (1993). Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *J Cell Biol*, 122(1), 103-111.
68. Diamond, M. P. (1997). Animal Adhesion Models: Design, Variables, and Relevance. In G. S. diZerega, A. H. DeCherney, M. P. Diamond, R. C. Dunn, E. P. Goldberg, A. F. Haney, J. A. Rock, T. Tulandi, & K. E. Rodgers (Eds.), *Pelvic Surgery: Adhesion Formation and Prevention* (pp. 65-70). New York, NY: Springer New York.
69. Diamond, M. P., El-Hammady, E., Munkarah, A., Bieber, E. J., & Saed, G. (2005). Modulation of the expression of vascular endothelial growth factor in human fibroblasts. *Fertil Steril*, 83(2), 405-409. doi:10.1016/j.fertnstert.2004.06.073
70. Diamond, M. P., & Nezhat, F. (1993). Adhesions after resection of ovarian endometriomas. *Fertil Steril*, 59(4), 934-935; author reply 935-936.
71. Dinarvand, P., Hashemi, S. M., Seyedjafari, E., Shabani, I., Mohammadi-Sangcheshmeh, A., Farhadian, S., & Soleimani, M. (2012). Function of poly (lactic-co-glycolic acid) nanofiber in reduction of adhesion bands. *J Surg Res*, 172(1), e1-9. doi:10.1016/j.jss.2011.10.014
72. Diniz, I. M., Chen, C., Xu, X., Ansari, S., Zadeh, H. H., Marques, M. M., . . . Moshaverinia, A. (2015). Pluronic F-127 hydrogel as a promising scaffold for encapsulation of dental-derived mesenchymal stem cells. *J Mater Sci Mater Med*, 26(3), 153. doi:10.1007/s10856-015-5493-4
73. diZerega, G. S. (1997). Biochemical events in peritoneal tissue repair. *Eur J Surg Suppl*(577), 10-16.

74. DiZerega, G. S. (2000). Use of Adhesion Prevention Barriers in Pelvic Reconstructive and Gynecologic Surgery. In G. S. diZerega (Ed.), *Peritoneal Surgery* (pp. 379-399). New York, NY: Springer New York.
75. diZerega, G. S., & Campeau, J. D. (2001). Peritoneal repair and post-surgical adhesion formation. *Hum Reprod Update*, 7(6), 547-555.
76. Du, X. H., Liu, J. Q., Xin, K., & Liu, G. H. (2015). Dexamethasone and sodium carboxymethyl cellulose prevent postoperative intraperitoneal adhesions in rats. *Braz J Med Biol Res*, 48(4), 344-348. doi:10.1590/1414-431x20144211
77. Dubuis, C., May, L., Alonso, F., Luca, L., Mylonaki, I., Meda, P., . . . Haeffliger, J. A. (2013). Atorvastatin-loaded hydrogel affects the smooth muscle cells of human veins. *J Pharmacol Exp Ther*, 347(3), 574-581. doi:10.1124/jpet.113.208769
78. Dumont, D. J., Gradwohl, G., Fong, G. H., Puri, M. C., Gertsenstein, M., Auerbach, A., & Breitman, M. L. (1994). Dominant-negative and targeted null mutations in the endothelial receptor tyrosine kinase, tek, reveal a critical role in vasculogenesis of the embryo. *Genes Dev*, 8(16), 1897-1909.
79. Duran, H. E., Kuscu, E., Zeyneloglu, H. B., Saygili, E., & Batioglu, S. (2002). Lipiodol versus methylene blue for prevention of postsurgical adhesion formation in a rat model. *Eur J Obstet Gynecol Reprod Biol*, 102(1), 80-82.
80. Elkins, T. E., Ling, F. W., Ahokas, R. A., Abdella, T. N., Homsey, C. A., & Malinak, L. R. (1984). Adhesion prevention by solutions of sodium carboxymethylcellulose in the rat. II. *Fertil Steril*, 41(6), 929-932.
81. Elkins, T. E., Stovall, T. G., Warren, J., Ling, F. W., & Meyer, N. L. (1987). A histologic evaluation of peritoneal injury and repair: implications for adhesion formation. *Obstet Gynecol*, 70(2), 225-228.
82. Ellis, H. (1962). The aetiology of post-operative abdominal adhesions. An experimental study. *Br J Surg*, 50, 10-16.
83. Ellis, H. (1971). The cause and prevention of postoperative intraperitoneal adhesions. *Surg Gynecol Obstet*, 133(3), 497-511.
84. Ellis, H. (1997). The clinical significance of adhesions: focus on intestinal obstruction. *Eur J Surg Suppl*(577), 5-9.
85. Ellis, H. (1998). The magnitude of adhesion related problems. *Ann Chir Gynaecol*, 87(1), 9-11.
86. Ellis, H., Moran, B. J., Thompson, J. N., Parker, M. C., Wilson, M. S., Menzies, D., . . . Crowe, A. M. (1999). Adhesion-related hospital readmissions after abdominal and pelvic surgery: a retrospective cohort study. *Lancet*, 353(9163), 1476-1480. doi:10.1016/S0140-6736(98)09337-4
87. Engelmann, B., & Massberg, S. (2013). Thrombosis as an intravascular effector of innate immunity. *Nat Rev Immunol*, 13(1), 34-45. doi:10.1038/nri3345
88. Eugenin, E. A., Branes, M. C., Berman, J. W., & Saez, J. C. (2003). TNF-alpha plus IFN-gamma induce connexin43 expression and formation of gap junctions between human monocytes/macrophages that enhance physiological responses. *J Immunol*, 170(3), 1320-1328.
89. Eugénin, E. A., González, H. E., Sánchez, H. A., Brañes, M. C., & Sáez, J. C. (2007). Inflammatory conditions induce gap junctional communication between rat Kupffer cells both in vivo and in vitro. *Cellular immunology*, 247(2), 103-110. doi:10.1016/j.cellimm.2007.08.001
90. Evans, D. M., McAree, K., Guyton, D. P., Hawkins, N., & Stakleff, K. (1993). Dose dependency and wound healing aspects of the use of tissue plasminogen activator in the prevention of intra-abdominal adhesions. *Am J Surg*, 165(2), 229-232.
91. Evans, W., H., De Vuyst, E., & Leybaert, L. (2006). The gap junction cellular internet: connexin hemichannels enter the signalling limelight. *Biochem J*, 397(Pt 1), 1-14. doi:10.1042/bj20060175
92. Evans, W. H., & Martin, P. E. (2002). Gap junctions: structure and function (Review). *Mol Membr Biol*, 19(2), 121-136. doi:10.1080/09687680210139839
93. Falk, K., Holmdahl, L., Halvarsson, M., Larsson, K., Lindman, B., & Bengmark, S. (1998). Polymers that reduce intraperitoneal adhesion formation. *Br J Surg*, 85(8), 1153-1156. doi:10.1046/j.1365-2168.1998.00784.x
94. Fazio, V. W., Cohen, Z., Fleshman, J. W., van Goor, H., Bauer, J. J., Wolff, B. G., . . . Gohl, J. (2006). Reduction in adhesive small-bowel obstruction by Seprafilm adhesion barrier after intestinal resection. *Dis Colon Rectum*, 49(1), 1-11. doi:10.1007/s10350-005-0268-5

95. Fernandez-Cobo, M., Gingalewski, C., & De Maio, A. (1998). Expression of the connexin 43 gene is increased in the kidneys and the lungs of rats injected with bacterial lipopolysaccharide. *Shock*, 10(2), 97-102.
96. Fischer, R., Reinehr, R., Lu, T. P., Schonicke, A., Warskulat, U., Dienes, H. P., & Haussinger, D. (2005). Intercellular communication via gap junctions in activated rat hepatic stellate cells. *Gastroenterology*, 128(2), 433-448.
97. Fletcher, N. M., Jiang, Z. L., Diamond, M. P., Abu-Soud, H. M., & Saed, G. M. (2008). Hypoxia-generated superoxide induces the development of the adhesion phenotype. *Free Radic Biol Med*, 45(4), 530-536. doi:10.1016/j.freeradbiomed.2008.05.002
98. Forde, A., Constien, R., Grone, H. J., Hammerling, G., & Arnold, B. (2002). Temporal Cre-mediated recombination exclusively in endothelial cells using Tie2 regulatory elements. *Genesis*, 33(4), 191-197. doi:10.1002/gene.10117
99. Franconi, F., Roux, J., Lefebvre-Lacoeuille, C., & Lemaire, L. (2015). Imaging visceral adhesion to polymeric mesh using pneumoperitoneal-MRI in an experimental rat model. *Surg Endosc*, 29(6), 1567-1573. doi:10.1007/s00464-014-3843-9
100. Fukasawa, M., Girgis, W., & diZerega, G. S. (1991). Inhibition of postsurgical adhesions in a standardized rabbit model: II. Intraperitoneal treatment with heparin. *Int J Fertil*, 36(5), 296-301.
101. Gaertner, F., & Massberg, S. (2016). Blood coagulation in immunothrombosis-At the frontline of intravascular immunity. *Semin Immunol*, 28(6), 561-569. doi:10.1016/j.smim.2016.10.010
102. Gaertner, W. B., Hagerman, G. F., Felemovicius, I., Bonsack, M. E., & Delaney, J. P. (2008). Two experimental models for generating abdominal adhesions. *J Surg Res*, 146(2), 241-245. doi:10.1016/j.jss.2007.08.012
103. Gao, Q., Wei, G., Wu, Y., Yao, N., Zhou, C., Wang, K., . . . Li, X. (2017). Paeoniflorin prevents postoperative peritoneal adhesion formation in an experimental rat model. *Oncotarget*, 8(55), 93899-93911. doi:10.18632/oncotarget.21333
104. Goetsch, K. P., & Niesler, C. U. (2016). The extracellular matrix regulates the effect of decorin and transforming growth factor beta-2 (TGF-beta2) on myoblast migration. *Biochem Biophys Res Commun*, 479(2), 351-357. doi:10.1016/j.bbrc.2016.09.079
105. Gomel, V., Urman, B., & Gurgan, T. (1996). Pathophysiology of adhesion formation and strategies for prevention. *J Reprod Med*, 41(1), 35-41.
106. Gong, C. Y., Wu, Q. J., Liao, J. F., Qi, T. T., Yang, B., Wang, Y. J., . . . Qian, Z. Y. (2013). Prevention of postsurgical cauterization-induced peritoneal adhesions by biodegradable and thermosensitive micelles. *J Biomed Nanotechnol*, 9(12), 1984-1995.
107. Gonzalez-Quintero, V. H., & Cruz-Pachano, F. E. (2009). Preventing adhesions in obstetric and gynecologic surgical procedures. *Rev Obstet Gynecol*, 2(1), 38-45.
108. Goodenough, D. A., Goliger, J. A., & Paul, D. L. (1996). Connexins, connexons, and intercellular communication. *Annu Rev Biochem*, 65, 475-502. doi:10.1146/annurev.bi.65.070196.002355
109. Gorvy, D. A., Herrick, S. E., Shah, M., & Ferguson, M. W. (2005). Experimental manipulation of transforming growth factor-beta isoforms significantly affects adhesion formation in a murine surgical model. *Am J Pathol*, 167(4), 1005-1019.
110. Green, C. R., Law, L. Y., Lin, J. S., & Becker, D. L. (2001). Spatiotemporal Depletion of Connexins Using Antisense Oligonucleotides. In R. Bruzzone & C. Giaume (Eds.), *Connexin Methods and Protocols* (pp. 175-185). Totowa, NJ: Humana Press.
111. Gu, H., Zou, Y. R., & Rajewsky, K. (1993). Independent control of immunoglobulin switch recombination at individual switch regions evidenced through Cre-loxP-mediated gene targeting. *Cell*, 73(6), 1155-1164.
112. Guo, Q., Li, Q. F., Liu, H. J., Li, R., Wu, C. T., & Wang, L. S. (2008). Sphingosine kinase 1 gene transfer reduces postoperative peritoneal adhesion in an experimental model. *Br J Surg*, 95(2), 252-258. doi:10.1002/bjs.5890
113. Halverson, A. L., Barrett, W. L., Bhanot, P., Phillips, J. E., Iglesias, A. R., Jacobs, L. K., & Sackier, J. M. (1999). Intraabdominal adhesion formation after preperitoneal dissection in the murine model. *Surg Endosc*, 13(1), 14-16.
114. Haney, A. F., & Doty, E. (1992a). Comparison of the peritoneal cells elicited by oxidized regenerated cellulose (Interceed) and expanded polytetrafluoroethylene (Gore-Tex Surgical Membrane) in a murine model. *Am J Obstet Gynecol*, 166(4), 1137-1146; discussion 1146-1139.

115. Haney, A. F., & Doty, E. (1992b). Murine peritoneal injury and de novo adhesion formation caused by oxidized-regenerated cellulose (Interceed [TC7]) but not expanded polytetrafluoroethylene (Gore-Tex Surgical Membrane). *Fertil Steril*, 57(1), 202-208.
116. Haney, A. F., & Doty, E. (1994). The formation of coalescing peritoneal adhesions requires injury to both contacting peritoneal surfaces. *Fertil Steril*, 61(4), 767-775.
117. Haney, A. F., & Doty, E. (1996). The peritoneal response to expanded polytetrafluoroethylene and oxidized regenerated cellulose surgical adhesion barriers. *Artif Cells Blood Substit Immobil Biotechnol*, 24(2), 121-141.
118. Hasdemir, P. S., Ozkut, M., Guvenal, T., Uner, M. A., Calik, E., Koltan, S. O., . . . Ozbilgin, K. (2017). Effect of Pirfenidone on Vascular Proliferation, Inflammation and Fibrosis in an Abdominal Adhesion Rat Model. *J Invest Surg*, 30(1), 26-32. doi:10.1080/08941939.2016.1215578
119. Hellebrekers, B. W., Trimbo-Kemper, G. C., Bakkum, E. A., Trimbo, J. B., Declerck, P. J., Kooistra, T., & Emeis, J. J. (2000). Short-term effect of surgical trauma on rat peritoneal fibrinolytic activity and its role in adhesion formation. *Thromb Haemost*, 84(5), 876-881.
120. Hemadeh, O., Chilukuri, S., Bonet, V., Hussein, S., & Chaudry, I. H. (1993). Prevention of peritoneal adhesions by administration of sodium carboxymethyl cellulose and oral vitamin E. *Surgery*, 114(5), 907-910.
121. Hernandez-Guerra, M., Gonzalez-Mendez, Y., de Ganzo, Z. A., Salido, E., Garcia-Pagan, J. C., Abrante, B., . . . Quintero, E. (2014). Role of gap junctions modulating hepatic vascular tone in cirrhosis. *Liver Int*, 34(6), 859-868. doi:10.1111/liv.12446
122. Hershlag, A., Otterness, I. G., Bliven, M. L., Diamond, M. P., & Polan, M. L. (1991). The effect of interleukin-1 on adhesion formation in the rat. *Am J Obstet Gynecol*, 165(3), 771-774.
123. Hervé, J.-C., Derangeon, M., Sarrouilhe, D., Giepmans, B. N. G., & Bourmeyster, N. (2012). Gap junctional channels are parts of multiprotein complexes. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1818(8), 1844-1865. doi:<https://doi.org/10.1016/j.bbamem.2011.12.009>
124. Hills, C. E., & Squires, P. E. (2010). TGF-beta1-induced epithelial-to-mesenchymal transition and therapeutic intervention in diabetic nephropathy. *Am J Nephrol*, 31(1), 68-74. doi:10.1159/000256659
125. Hoban, L. D., Pierce, M., Quance, J., Hayward, I., McKee, A., Gowda, D. C., . . . Williams, T. (1994). Use of polypentapeptides of elastin to prevent postoperative adhesions: efficacy in a contaminated peritoneal model. *J Surg Res*, 56(2), 179-183. doi:10.1006/jsre.1994.1029
126. Hoffmann, N. E., Siddiqui, S. A., Agarwal, S., McKellar, S. H., Kurtz, H. J., Gettman, M. T., & Ereth, M. H. (2009). Choice of hemostatic agent influences adhesion formation in a rat cecal adhesion model. *J Surg Res*, 155(1), 77-81. doi:10.1016/j.jss.2008.08.008
127. Holmdahl, L. (2000). The Plasmin System, a Marker of the Propensity to Develop Adhesions. In G. S. diZerega (Ed.), *Peritoneal Surgery* (pp. 117-131). New York, NY: Springer New York.
128. Holmdahl, L., al-Jabreen, M., & Risberg, B. (1994). Experimental models for quantitative studies on adhesion formation in rats and rabbits. *Eur Surg Res*, 26(4), 248-256. doi:10.1159/000129342
129. Holmdahl, L., Eriksson, E., al-Jabreen, M., & Risberg, B. (1996). Fibrinolysis in human peritoneum during operation. *Surgery*, 119(6), 701-705.
130. Holschneider, C. H., Cristoforoni, P. M., Ghosh, K., Punyasavatsut, M., Abed, E., & Montz, F. J. (1997). Endogenous versus exogenous IL-10 in postoperative intraperitoneal adhesion formation in a murine model. *J Surg Res*, 70(2), 138-143. doi:10.1006/jsre.1997.5124
131. Holsti, M. A., Chitnis, T., Panzo, R. J., Bronson, R. T., Yagita, H., Sayegh, M. H., & Tzianabos, A. O. (2004). Regulation of postsurgical fibrosis by the programmed death-1 inhibitory pathway. *J Immunol*, 172(9), 5774-5781.
132. Hong, G. S., Schwandt, T., Stein, K., Schneiker, B., Kummer, M. P., Heneka, M. T., . . . Wehner, S. (2015). Effects of macrophage-dependent peroxisome proliferator-activated receptor gamma signalling on adhesion formation after abdominal surgery in an experimental model. *Br J Surg*, 102(12), 1506-1516. doi:10.1002/bjs.9907
133. Hoshino, A., Kawamura, Y. I., Yasuhara, M., Toyama-Sorimachi, N., Yamamoto, K., Matsukawa, A., . . . Dohi, T. (2007). Inhibition of CCL1-CCR8 interaction prevents aggregation of macrophages and development of peritoneal adhesions. *J Immunol*, 178(8), 5296-5304.

134. Hubbard, T. B., Khan, M. Z., Carag, V. R., Albites, V. E., & Hricko, G. M. (1967). The pathology of peritoneal repair: its relation to the formation of adhesions. *Ann Surg*, 165(6), 908-916.
135. Huvencuers, S., Daemen, M. J., & Hordijk, P. L. (2015). Between Rho(k) and a hard place: the relation between vessel wall stiffness, endothelial contractility, and cardiovascular disease. *Circ Res*, 116(5), 895-908. doi:10.1161/circresaha.116.305720
136. Ichinose, A., Takio, K., & Fujikawa, K. (1986). Localization of the binding site of tissue-type plasminogen activator to fibrin. *J Clin Invest*, 78(1), 163-169. doi:10.1172/JCI112546
137. Ignjatovic, D., Aasland, K., Pettersen, M., Sund, S., Chen, Y., Spasojevic, M., & Nesgaard, J. M. (2010). Intra-abdominal administration of bevacizumab diminishes intra-peritoneal adhesions. *Am J Surg*, 200(2), 270-275. doi:10.1016/j.amjsurg.2009.08.038
138. Irkocucu, O., Ferahkose, Z., Memis, L., Ekin, O., & Akin, M. (2009). Reduction of postsurgical adhesions in a rat model: a comparative study. *Clinics (Sao Paulo)*, 64(2), 143-148.
139. Ivarsson, M. L., Bergstrom, M., Eriksson, E., Risberg, B., & Holmdahl, L. (1998). Tissue markers as predictors of postoperative adhesions. *Br J Surg*, 85(11), 1549-1554.
140. Jagelman, D. G., & Ellis, H. (1973). Starch and intraperitoneal adhesion formation. *Br J Surg*, 60(2), 111-114.
141. Jahoda, A. E., Olson, M. K., & Kovacs, E. J. (2003). Murine model of peritoneal adhesion formation. *Methods Mol Med*, 78, 141-148. doi:10.1385/1-59259-332-1:141
142. Kaidi, A. A., Gurchumelidze, T., Nazzari, M., Figert, P., Vanterpool, C., & Silva, Y. (1995). Tumor necrosis factor- α : a marker for peritoneal adhesion formation. *J Surg Res*, 58(5), 516-518. doi:10.1006/jsre.1995.1081
143. Kaminski, P., Gajewska, M., Wielgos, M., Szymusik, I., Ziolkowska, K., & Bartkowiak, R. (2006). The usefulness of laparoscopy and hysteroscopy in the diagnostics and treatment of infertility. *Neuro Endocrinol Lett*, 27(6), 813-817.
144. Kaplanski, G., Marin, V., Fabrigoule, M., Boulay, V., Benoliel, A. M., Bongrand, P., . . . Farnarier, C. (1998). Thrombin-activated human endothelial cells support monocyte adhesion in vitro following expression of intercellular adhesion molecule-1 (ICAM-1; CD54) and vascular cell adhesion molecule-1 (VCAM-1; CD106). *Blood*, 92(4), 1259-1267.
145. Karaca, G., Aydin, O., Pehlivanli, F., Kocael, A., Pekcici, R., Duymus, E., . . . Guler, O. (2016). Effect of ankaferd blood stopper in experimental peritoneal adhesion model. *Ann Surg Treat Res*, 90(4), 213-217. doi:10.4174/ast.2016.90.4.213
146. Karaca, T., Gözalan, A. U., Yoldaş, Ö., Bilgin, B. Ç., & Tezer, A. (2013). Effects of tamoxifen citrate on postoperative intra-abdominal adhesion in a rat model. *International Journal of Surgery*, 11(1), 68-72. doi:<https://doi.org/10.1016/j.ijsu.2012.11.015>
147. Keskin, H. L., Sirin, Y. S., Keles, H., Turgut, O., Ide, T., & Avsar, A. F. (2013). The aromatase inhibitor letrozole reduces adhesion formation after intraperitoneal surgery in a rat uterine horn model. *Eur J Obstet Gynecol Reprod Biol*, 167(2), 199-204. doi:10.1016/j.ejogrb.2012.12.002
148. Kharwadkar, N., Naik, S., & Molitor, P. J. (2005). Prospective randomized trial comparing absorbable and non-absorbable sutures in open carpal tunnel release. *J Hand Surg Br*, 30(1), 92-95. doi:10.1016/j.jhsb.2004.10.009
149. Kim, J., & Lee, J. (2017). Role of transforming growth factor- β in muscle damage and regeneration: focused on eccentric muscle contraction. *J Exerc Rehabil*, 13(6), 621-626. doi:10.12965/jer.1735072.536
150. Kirdak, T., Uysal, E., & Korun, N. (2008). [Assessment of effectiveness of different doses of methylprednisolone on intraabdominal adhesion prevention]. *Ulus Travma Acil Cerrahi Derg*, 14(3), 188-191.
151. Kisanuki, Y. Y., Hammer, R. E., Miyazaki, J., Williams, S. C., Richardson, J. A., & Yanagisawa, M. (2001). Tie2-Cre transgenic mice: a new model for endothelial cell-lineage analysis in vivo. *Dev Biol*, 230(2), 230-242. doi:10.1006/dbio.2000.0106
152. Koca, Y. S., Tarhan, O. R., Kaya, S., & Gokce Ceylan, B. (2016). Effects of saline lavage temperature on peritoneal fibrinolysis and adhesion formation. *Ulus Travma Acil Cerrahi Derg*, 22(1), 1-6. doi:10.5505/tjtes.2015.95881
153. Kojima, T., Sawada, N., Zhong, Y., Oyamada, M., & Mori, M. (1994). Sequential changes in intercellular junctions between hepatocytes during the course of acute liver injury and restoration after thioacetamide treatment. *Virchows Arch*, 425(4), 407-412.
154. Koval, M., Billaud, M., Straub, A. C., Johnstone, S. R., Zarbock, A., Duling, B. R., & Isakson, B. E. (2011). Spontaneous lung dysfunction and fibrosis in mice lacking connexin

- 40 and endothelial cell connexin 43. *Am J Pathol*, 178(6), 2536-2546. doi:10.1016/j.ajpath.2011.02.045
155. Kraemer, B., Wallwiener, C., Rajab, T. K., Brochhausen, C., Wallwiener, M., & Rothmund, R. (2014). Standardised models for inducing experimental peritoneal adhesions in female rats. *Biomed Res Int*, 2014, 435056. doi:10.1155/2014/435056
 156. Kraemer, B., Wallwiener, M., Wallwiener, C. W., Juhasz-Boess, I., Hartkopf, A., Wallwiener, D., & Rajab, T. K. (2012). Differential mRNA expression of TACR1 after ischemic peritoneal trauma: a pilot animal study. *Arch Gynecol Obstet*, 285(3), 717-720. doi:10.1007/s00404-011-2041-4
 157. Krause, T. J., Katz, D., Wheeler, C. J., Ebner, S., & McKinnon, R. D. (1999). Increased levels of surgical adhesions in TGFbeta1 heterozygous mice. *J Invest Surg*, 12(1), 31-38.
 158. Kucuk, H. F., Kaptanoglu, L., Kurt, N., Uzun, H., Eser, M., Bingul, S., . . . Akyol, H. (2007). The role of simvastatin on postoperative peritoneal adhesion formation in an animal model. *Eur Surg Res*, 39(2), 98-102. doi:10.1159/000099156
 159. Kuroyanagi, M., Yamamoto, A., Shimizu, N., Toi, A., Inomata, T., Takeda, A., & Kuroyanagi, Y. (2014). Development of anti-adhesive spongy sheet composed of hyaluronic acid and collagen containing epidermal growth factor. *J Biomater Sci Polym Ed*, 25(12), 1253-1265. doi:10.1080/09205063.2014.926579
 160. Lai, P. H., Chang, Y., Liang, H. C., Chen, S. C., Wei, H. J., & Sung, H. W. (2005). Peritoneal regeneration induced by an acellular bovine pericardial patch in the repair of abdominal wall defects. *J Surg Res*, 127(2), 85-92. doi:10.1016/j.jss.2005.03.016
 161. Lang, N. N., Myles, R. C., Burton, F. L., Hall, D. P., Chin, Y. Z., Boon, N. A., & Newby, D. E. (2008). The vascular effects of rotigaptide in vivo in man. *Biochem Pharmacol*, 76(10), 1194-1200. doi:10.1016/j.bcp.2008.08.022
 162. Lauder, C. I., Garcea, G., Strickland, A., & Maddern, G. J. (2011). Use of a modified chitosan-dextran gel to prevent peritoneal adhesions in a rat model. *J Surg Res*, 171(2), 877-882. doi:10.1016/j.jss.2010.06.028
 163. Law, L. Y., Zhang, W. V., Stott, N. S., Becker, D. L., & Green, C. R. (2006). In Vitro Optimization of Antisense Oligodeoxynucleotide Design: An Example Using the Connexin Gene Family. *J Biomol Tech*, 17(4), 270-282.
 164. Leach, R. E. M. D., Burns, J. W. P. D., Dawe, E. J. D. V. M., SmithBarbour, M. D. B. S., & Diamond, M. P. M. D. (1998). Reduction of Postsurgical Adhesion Formation in the Rabbit Uterine Horn Model with Use of Hyaluronate/ Carboxymethylcellulose Gel¹. *Fertil Steril*, 69(3), 415-418. doi:10.1016/S0015-0282(97)00573-6
 165. Lee, M. T., Lee, C. C., Wang, H. M., Chou, T. H., Wu, M. C., Hsueh, K. L., & Chen, S. C. (2016). Hypothermia Increases Tissue Plasminogen Activator Expression and Decreases Post-Operative Intra-Abdominal Adhesion. *PLoS One*, 11(9), e0160627. doi:10.1371/journal.pone.0160627
 166. Levi, M., & van der Poll, T. (2010). Inflammation and coagulation. *Crit Care Med*, 38(2 Suppl), S26-34. doi:10.1097/CCM.0b013e3181c98d21
 167. Liebman, S. M., Langer, J. C., Marshall, J. S., & Collins, S. M. (1993). Role of mast cells in peritoneal adhesion formation. *Am J Surg*, 165(1), 127-130.
 168. Lim, M. C., Maubach, G., & Zhuo, L. (2009). TGF-beta1 down-regulates connexin 43 expression and gap junction intercellular communication in rat hepatic stellate cells. *Eur J Cell Biol*, 88(12), 719-730. doi:10.1016/j.ejcb.2009.08.003
 169. Lim, R., Stucchi, A. F., Morrill, J. M., Reed, K. L., Lynch, R., & Becker, J. M. (2010). The efficacy of a hyaluronate-carboxymethylcellulose bioresorbable membrane that reduces postoperative adhesions is increased by the intra-operative co-administration of a neurokinin 1 receptor antagonist in a rat model. *Surgery*, 148(5), 991-999. doi:10.1016/j.surg.2010.01.024
 170. Liu, H. J., Wu, C. T., Duan, H. F., Wu, B., Lu, Z. Z., & Wang, L. (2006). Adenoviral-mediated gene expression of hepatocyte growth factor prevents postoperative peritoneal adhesion in a rat model. *Surgery*, 140(3), 441-447. doi:10.1016/j.surg.2005.12.014
 171. Liu, L., Zou, P., Zheng, L., Linarelli, L. E., Amarell, S., Passaro, A., . . . Cheng, Z. (2015). Tamoxifen reduces fat mass by boosting reactive oxygen species. *Cell Death Dis*, 6, e1586. doi:10.1038/cddis.2014.553
 172. López-Cabrera, M. (2014). Mesenchymal Conversion of Mesothelial Cells Is a Key Event in the Pathophysiology of the Peritoneum during Peritoneal Dialysis. *Advances in Medicine*, 2014, 17. doi:10.1155/2014/473134

173. Lower, A. M., Hawthorn, R. J., Clark, D., Boyd, J. H., Finlayson, A. R., Knight, A. D., . . . Clinical Research, G. (2004). Adhesion-related readmissions following gynaecological laparoscopy or laparotomy in Scotland: an epidemiological study of 24 046 patients. *Hum Reprod*, 19(8), 1877-1885. doi:10.1093/humrep/deh321
174. Lu, D., Soleymani, S., Madakshire, R., & Insel, P. A. (2012). ATP released from cardiac fibroblasts via connexin hemichannels activates profibrotic P2Y2 receptors. *FASEB J*, 26(6), 2580-2591. doi:10.1096/fj.12-204677
175. Lu, F., Gao, J., Ogawa, R., & Hyakusoku, H. (2007). Variations in gap junctional intercellular communication and connexin expression in fibroblasts derived from keloid and hypertrophic scars. *Plast Reconstr Surg*, 119(3), 844-851. doi:10.1097/01.prs.0000255539.99698.f4
176. Lucas, P. A., Warejcka, D. J., Young, H. E., & Lee, B. Y. (1996). Formation of abdominal adhesions is inhibited by antibodies to transforming growth factor-beta1. *J Surg Res*, 65(2), 135-138. doi:10.1006/jsre.1996.0355
177. Lum, H., & Malik, A. B. (1996). Mechanisms of increased endothelial permeability. *Can J Physiol Pharmacol*, 74(7), 787-800.
178. Ma, D. D. F., Rede, T., Naqvi, N. A., & Cook, P. D. (2000). Synthetic oligonucleotides as therapeutics: The coming of age *Biotechnology Annual Review* (Vol. 5, pp. 155-196): Elsevier.
179. Maciver, A. H., McCall, M., & James Shapiro, A. M. (2011). Intra-abdominal adhesions: cellular mechanisms and strategies for prevention. *Int J Surg*, 9(8), 589-594. doi:10.1016/j.ijsu.2011.08.008
180. Mackman, N. (2009). The role of tissue factor and factor VIIa in hemostasis. *Anesth Analg*, 108(5), 1447-1452. doi:10.1213/ane.0b013e31819bceb1
181. Mahato, R. I., Cheng, K., & Guntaka, R. V. (2005). Modulation of gene expression by antisense and antigene oligodeoxynucleotides and small interfering RNA. *Expert Opin Drug Deliv*, 2(1), 3-28. doi:10.1517/17425247.2.1.3
182. Marshall, C. D., Hu, M. S., Leavitt, T., Barnes, L. A., Cheung, A. T., Malhotra, S., . . . Longaker, M. T. (2016). Creation of Abdominal Adhesions in Mice. *J Vis Exp*(114). doi:10.3791/54450
183. McNamara, D. A., Walsh, T. N., Kay, E., & Bouchier-Hayes, D. J. (2003). Neoadjuvant antiangiogenic therapy with tamoxifen does not impair gastrointestinal anastomotic repair in the rat. *Colorectal Dis*, 5(4), 335-341.
184. Mendes, J. B., Campos, P. P., Rocha, M. A., & Andrade, S. P. (2009a). Cilostazol and pentoxifylline decrease angiogenesis, inflammation, and fibrosis in sponge-induced intraperitoneal adhesion in mice. *Life Sci*, 84(15-16), 537-543. doi:10.1016/j.lfs.2009.01.023
185. Mendes, J. B., Rocha, M. A., Araujo, F. A., Moura, S. A., Ferreira, M. A., & Andrade, S. P. (2009b). Differential effects of rolipram on chronic subcutaneous inflammatory angiogenesis and on peritoneal adhesion in mice. *Microvasc Res*, 78(3), 265-271. doi:10.1016/j.mvr.2009.08.008
186. Mendoza-Naranjo, A., Cormie, P., Serrano, A. E., Wang, C. M., Thrasivoulou, C., Sutcliffe, J. E., . . . Becker, D. L. (2012). Overexpression of the gap junction protein Cx43 as found in diabetic foot ulcers can retard fibroblast migration. *Cell Biol Int*, 36(7), 661-667. doi:10.1042/cbi20110628
187. Menzies, D., & Ellis, H. (1990). Intestinal obstruction from adhesions--how big is the problem? *Annals of The Royal College of Surgeons of England*, 72(1), 60-63.
188. Merlo, G., Fausone, G., & Castagna, B. (1983). Fibrinolytic activity of mesothelial lining of the displaced peritoneum. *Am J Med Sci*, 286(2), 12-14.
189. Metzger, D., Clifford, J., Chiba, H., & Chambon, P. (1995). Conditional site-specific recombination in mammalian cells using a ligand-dependent chimeric Cre recombinase. *Proceedings of the National Academy of Sciences of the United States of America*, 92(15), 6991-6995.
190. Mikula-Pietrasik, J., Uruski, P., Szubert, S., Szpurek, D., Sajdak, S., Tykarski, A., & Ksiazek, K. (2017). Malignant ascites determine the transmesothelial invasion of ovarian cancer cells. *Int J Biochem Cell Biol*, 92, 6-13. doi:10.1016/j.biocel.2017.09.002
191. Milingos, S., Kallipolitis, G., Loutradis, D., Liapi, A., Mavrommatis, K., Drakakis, P., . . . Michalas, S. (2000). Adhesions: laparoscopic surgery versus laparotomy. *Ann N Y Acad Sci*, 900, 272-285.

192. Miyamoto, T., Tamura, M., Kabashima, N., Serino, R., Shibata, T., Furuno, Y., . . . Otsuji, Y. (2010). An integrin-activating peptide, PHSRN, ameliorates inhibitory effects of conventional peritoneal dialysis fluids on peritoneal wound healing. *Nephrol Dial Transplant*, 25(4), 1109-1119. doi:10.1093/ndt/gfp601
193. Mommers, E. H., Hong, L., Jongen, A., & Bouvy, N. D. (2018). Baseline performance of the ischaemic button model for induction of adhesions in laboratory rats. *Lab Anim*, 23677218773116. doi:10.1177/0023677218773116
194. Moreno, A., Aguayo, J. L., Zambudio, G., Ramirez, P., Canteras, M., & Parrilla, P. (1996). Influence of abdominal incision on the formation of postoperative peritoneal adhesions: an experimental study in rats. *Eur J Surg*, 162(3), 181-185.
195. Mori, R., Power, K. T., Wang, C. M., Martin, P., & Becker, D. L. (2006). Acute downregulation of connexin43 at wound sites leads to a reduced inflammatory response, enhanced keratinocyte proliferation and wound fibroblast migration. *J Cell Sci*, 119(Pt 24), 5193-5203. doi:10.1242/jcs.03320
196. Moris, D., Chakedis, J., Rahnama-Azar, A. A., Wilson, A., Hennessy, M. M., Athanasiou, A., . . . Pawlik, T. M. (2017). Postoperative Abdominal Adhesions: Clinical Significance and Advances in Prevention and Management. *J Gastrointest Surg*, 21(10), 1713-1722. doi:10.1007/s11605-017-3488-9
197. Muller, W. A. (2013). Getting leukocytes to the site of inflammation. *Vet Pathol*, 50(1), 7-22. doi:10.1177/0300985812469883
198. Mutsaers, S. E., Birnie, K., Lansley, S., Herrick, S. E., Lim, C. B., & Prêle, C. M. (2015). Mesothelial cells in tissue repair and fibrosis. *Front Pharmacol*, 6. doi:10.3389/fphar.2015.00113
199. Nagelschmidt, M., & Saad, S. (1997). Influence of polyethylene glycol 4000 and dextran 70 on adhesion formation in rats. *J Surg Res*, 67(2), 113-118. doi:10.1006/jsre.1996.4973
200. Nair, S. K., Bhat, I. K., & Aurora, A. L. (1974). Role of proteolytic enzyme in the prevention of postoperative intraperitoneal adhesions. *Arch Surg*, 108(6), 849-853.
201. Nappi, C., Di Spiezo Sardo, A., Greco, E., Guida, M., Bettocchi, S., & Bifulco, G. (2007). Prevention of adhesions in gynaecological endoscopy. *Hum Reprod Update*, 13(4), 379-394. doi:10.1093/humupd/dml061
202. Nassif, J., Abbasi, S. A., Kechli, M. K., Boutary, S. S., Ghulmiyyah, L., Khalifeh, I., . . . Nassar, A. H. (2016). Effect of the Mode of Application of Cryopreserved Human Amniotic Membrane on Adhesion Formation after Abdomino-Pelvic Surgery in a Mouse Model. *Front Med (Lausanne)*, 3, 10. doi:10.3389/fmed.2016.00010
203. Naus, C. C., & Giaume, C. (2016). Bridging the gap to therapeutic strategies based on connexin/pannexin biology. *J Transl Med*, 14(1), 330. doi:10.1186/s12967-016-1089-0
204. Nehez, L., Tingstedt, B., Vodros, D., Axelsson, J., Lindman, B., & Andersson, R. (2006). Novel treatment in peritoneal adhesion prevention: protection by polypeptides. *Scand J Gastroenterol*, 41(9), 1110-1117. doi:10.1080/00365520600554550
205. Nielsen, M. S., Axelsen, L. N., Sorgen, P. L., Verma, V., Delmar, M., & Holstein-Rathlou, N. H. (2012). Gap junctions. *Compr Physiol*, 2(3), 1981-2035. doi:10.1002/cphy.c110051
206. Nikeghbalian, S., Atefi, S., Kazemi, K., Roshan, N., Tanideh, N., & Jalaian, H. (2007). Effect of oral D-penicillamine vs. colchicine on experimentally induced peritoneal adhesions in rats. *Fertil Steril*, 88(4 Suppl), 1187-1189. doi:10.1016/j.fertnstert.2007.01.109
207. Nikeghbalian, S., Vafaei, H., Moradian, F., Kazemi, K., Tanideh, N., Shayan, L., & Nikeghbalian, Z. (2015). Administration of Intravenous Infliximab for Prevention of Peritoneal Adhesions Formation in Rats. *Bull Emerg Trauma*, 3(3), 97-103.
208. Norrman, B., Wallen, P., & Ranby, M. (1985). Fibrinolysis mediated by tissue plasminogen activator. Disclosure of a kinetic transition. *Eur J Biochem*, 149(1), 193-200.
209. O'Brien, M. (2012). The reciprocal relationship between inflammation and coagulation. *Top Companion Anim Med*, 27(2), 46-52. doi:10.1053/j.tcam.2012.06.003
210. O'Donnell, J. J., 3rd, Birukova, A. A., Beyer, E. C., & Birukov, K. G. (2014). Gap junction protein connexin43 exacerbates lung vascular permeability. *PLoS One*, 9(6), e100931. doi:10.1371/journal.pone.0100931
211. O'Leary, D. P., & Coakley, J. B. (1992). The influence of suturing and sepsis on the development of postoperative peritoneal adhesions. *Ann R Coll Surg Engl*, 74(2), 134-137.
212. Ogawa, T., Hayashi, T., Tokunou, M., Nakachi, K., Trosko, J. E., Chang, C. C., & Yorioka, N. (2005). Suberoylanilide hydroxamic acid enhances gap junctional intercellular communication via acetylation of histone containing connexin 43 gene locus. *Cancer Res*, 65(21), 9771-9778. doi:10.1158/0008-5472.CAN-05-0227

213. Oh, S. H., Kim, J. K., Song, K. S., Noh, S. M., Ghil, S. H., Yuk, S. H., & Lee, J. H. (2005). Prevention of postsurgical tissue adhesion by anti-inflammatory drug-loaded pluronic mixtures with sol-gel transition behavior. *J Biomed Mater Res A*, 72(3), 306-316. doi:10.1002/jbm.a.30239
214. Okabayashi, K., Ashrafian, H., Zacharakis, E., Hasegawa, H., Kitagawa, Y., Athanasiou, T., & Darzi, A. (2014). Adhesions after abdominal surgery: a systematic review of the incidence, distribution and severity. *Surg Today*, 44(3), 405-420. doi:10.1007/s00595-013-0591-8
215. Okamoto, T., Akita, N., Hayashi, T., Shimaoka, M., & Suzuki, K. (2014). Endothelial connexin 32 regulates tissue factor expression induced by inflammatory stimulation and direct cell-cell interaction with activated cells. *Atherosclerosis*, 236(2), 430-437. doi:10.1016/j.atherosclerosis.2014.07.025
216. Okamoto, T., Akiyama, M., Takeda, M., Akita, N., Yoshida, K., Hayashi, T., & Suzuki, K. (2011). Connexin32 protects against vascular inflammation by modulating inflammatory cytokine expression by endothelial cells. *Exp Cell Res*, 317(3), 348-355. doi:10.1016/j.yexcr.2010.10.018
217. Okamoto, T., & Suzuki, K. (2017). The Role of Gap Junction-Mediated Endothelial Cell–Cell Interaction in the Crosstalk between Inflammation and Blood Coagulation. *Int J Mol Sci*, 18(11). doi:10.3390/ijms18112254
218. Ordoñez, J. L., Domínguez, J., Evrard, V., & Koninckx, P. R. (1997). The effect of training and duration of surgery on adhesion formation in the rabbit model. *Human Reproduction*, 12(12), 2654-2657. doi:10.1093/humrep/12.12.2654
219. Osada, H., Minai, M., Tsunoda, I., Fujii, T. K., Tsubata, K., & Satoh, K. (1999). The effect of hyaluronic acid-carboxymethylcellulose in reducing adhesion reformation in rabbits. *J Int Med Res*, 27(6), 292-296. doi:10.1177/030006059902700605
220. Osei-Owusu, P., Sabharwal, R., Kaltenbronn, K. M., Rhee, M. H., Chapleau, M. W., Dietrich, H. H., & Blumer, K. J. (2012). Regulator of G protein signaling 2 deficiency causes endothelial dysfunction and impaired endothelium-derived hyperpolarizing factor-mediated relaxation by dysregulating Gi/o signaling. *J Biol Chem*, 287(15), 12541-12549. doi:10.1074/jbc.M111.332130
221. Otcu, S., Ozturk, H., Aldemir, M., Kilinc, N., & Dokucu, A. I. (2003). Influence of the platelet-activating factor receptor antagonist BB-882 on intra-abdominal adhesion formation in rats. *Eur Surg Res*, 35(1), 41-45. doi:10.1159/000067034
222. Oviedo-Orta, E., Gasque, P., & Evans, W. H. (2001). Immunoglobulin and cytokine expression in mixed lymphocyte cultures is reduced by disruption of gap junction intercellular communication. *FASEB J*, 15(3), 768-774. doi:10.1096/fj.00-0288com
223. Ozel, H., Avsar, F. M., Topaloglu, S., & Sahin, M. (2005). Induction and assessment methods used in experimental adhesion studies. *Wound Repair Regen*, 13(4), 358-364. doi:10.1111/j.1067-1927.2005.130402.x
224. Ozler, M., Ersoz, N., Ozerhan, I. H., Topal, T., Oter, S., & Korkmaz, A. (2011). The effect of alpha-lipoic acid in the prevention of peritoneal adhesions. *Turk J Gastroenterol*, 22(2), 190-194.
225. Panahi, F., Sadraie, S. H., Khoshmohabat, H., Shahram, E., Kaka, G., & Hosseinalipour, M. (2012). Macroscopic and pathological assessment of methylene blue and normal saline on postoperative adhesion formation in a rat cecum model. *Int J Surg*, 10(9), 537-541. doi:10.1016/j.ijssu.2012.08.009
226. Parker, M. C., Ellis, H., Moran, B. J., Thompson, J. N., Wilson, M. S., Menzies, D., . . . Crowe, A. M. (2001). Postoperative adhesions: ten-year follow-up of 12,584 patients undergoing lower abdominal surgery. *Dis Colon Rectum*, 44(6), 822-829; discussion 829-830.
227. Parker, M. C., Wilson, M. S., Menzies, D., Sunderland, G., Clark, D. N., Knight, A. D., . . . Clinical Adhesions Research, G. (2005). The SCAR-3 study: 5-year adhesion-related readmission risk following lower abdominal surgical procedures. *Colorectal Dis*, 7(6), 551-558. doi:10.1111/j.1463-1318.2005.00857.x
228. Parsaei, P., Karimi, M., Asadi, S. Y., & Rafieian-Kopaei, M. (2013). Bioactive components and preventive effect of green tea (*Camellia sinensis*) extract on post-laparotomy intra-abdominal adhesion in rats. *Int J Surg*, 11(9), 811-815. doi:10.1016/j.ijssu.2013.08.014
229. Parthasarathi, K., Ichimura, H., Monma, E., Lindert, J., Quadri, S., Issekutz, A., & Bhattacharya, J. (2006). Connexin 43 mediates spread of Ca²⁺-dependent proinflammatory responses in lung capillaries. *J Clin Invest*, 116(8), 2193-2200. doi:10.1172/JCI26605

230. Peker, K., Inal, A., Sayar, I., Sahin, M., Gullu, H., Inal, D. G., & Isik, A. (2013). Prevention of intraabdominal adhesions by local and systemic administration of immunosuppressive drugs. *Iran Red Crescent Med J*, 15(12), e14148. doi:10.5812/ircmj.14148
231. Pelin, K., Hirvonen, A., & Linnainmaa, K. (1994). Expression of cell adhesion molecules and connexins in gap junctional intercellular communication deficient human mesothelioma tumour cell lines and communication competent primary mesothelial cells. *Carcinogenesis*, 15(11), 2673-2675.
232. Poehnert, D., Abbas, M., Kreipe, H. H., Klemptner, J., & Winny, M. (2015). Evaluation of 4DryField(R) PH as Adhesion Prevention Barrier Tested in an Optimized Adhesion Model in Rats. *Eur Surg Res*, 55(4), 341-351. doi:10.1159/000441025
233. Poehnert, D., Abbas, M., Kreipe, H. H., Klemptner, J., & Winny, M. (2015). High reproducibility of adhesion formation in rat with meso-stitch approximation of injured cecum and abdominal wall. *Int J Med Sci*, 12(1), 1-6. doi:10.7150/ijms.8870
234. Poehnert, D., Abbas, M., Maegel, L., Sambale, F., Lavrentieva, A., Kreipe, H. H., . . . Winny, M. (2015). Evaluation of the biological tolerability of the starch-based medical device 4DryField(R) PH in vitro and in vivo a rat model. *J Biomater Appl*, 30(4), 463-471. doi:10.1177/0885328215592257
235. Popovic, M., Smiljanic, K., Dobutovic, B., Syrovets, T., Simmet, T., & Isenovic, E. R. (2012). Thrombin and vascular inflammation. *Mol Cell Biochem*, 359(1-2), 301-313. doi:10.1007/s11010-011-1024-x
236. Poss, K. D., & Tonegawa, S. (1997). Reduced stress defense in heme oxygenase 1-deficient cells. *Proc Natl Acad Sci U S A*, 94(20), 10925-10930.
237. Qiu, C., Coutinho, P., Frank, S., Franke, S., Law, L. Y., Martin, P., . . . Becker, D. L. (2003). Targeting connexin43 expression accelerates the rate of wound repair. *Curr Biol*, 13(19), 1697-1703.
238. Rajab, T. K., Wauschkunn, C. A., Smaxwil, L., Kraemer, B., Wallwiener, M., & Wallwiener, C. W. (2010). An improved model for the induction of experimental adhesions. *J Invest Surg*, 23(1), 35-39. doi:10.3109/08941930903469474
239. Rasti, M., Parvaresh, E., Tavajoh, S., & Talaei, M. (2007). The comparison of diphenhydramine HCl and Nedocromil sodium in prevention of abdominal postoperative adhesion formation in rat models: an experimental study. *Int J Surg*, 5(6), 384-387. doi:10.1016/j.ijsu.2007.04.013
240. Ravi, M., Paramesh, V., Kaviya, S. R., Anuradha, E., & Solomon, F. D. (2015). 3D cell culture systems: advantages and applications. *J Cell Physiol*, 230(1), 16-26. doi:10.1002/jcp.24683
241. Rayburn, E. R., & Zhang, R. (2008). Antisense, RNAi, and Gene Silencing Strategies for Therapy: Mission Possible or Impossible? *Drug Discov Today*, 13(11-12), 513-521. doi:10.1016/j.drudis.2008.03.014
242. Reed, K. L., Heydrick, S. J., Aarons, C. B., Prushik, S., Gower, A. C., Stucchi, A. F., & Becker, J. M. (2007). A neurokinin-1 receptor antagonist that reduces intra-abdominal adhesion formation decreases oxidative stress in the peritoneum. *Am J Physiol Gastrointest Liver Physiol*, 293(3), G544-551. doi:10.1152/ajpgi.00226.2007
243. Rice, V. M., Shanti, A., Moghissi, K. S., & Leach, R. E. (1993). A comparative evaluation of Poloxamer 407 and oxidized regenerated cellulose (Interceed [TC7]) to reduce postoperative adhesion formation in the rat uterine horn model. *Fertil Steril*, 59(4), 901-906.
244. Rignault, S., Haefliger, J. A., Waeber, B., Liaudet, L., & Feihl, F. (2007). Acute inflammation decreases the expression of connexin 40 in mouse lung. *Shock*, 28(1), 78-85. doi:10.1097/shk.0b013e3180310bd1
245. Risberg, B. (1997). Adhesions: preventive strategies. *Eur J Surg Suppl*(577), 32-39.
246. Robb, W. B., & Mariette, C. (2014). Strategies in the prevention of the formation of postoperative adhesions in digestive surgery: a systematic review of the literature. *Dis Colon Rectum*, 57(10), 1228-1240. doi:10.1097/DCR.0000000000000191
247. Robbins, G. F., Brunschwig, A., & Foote, F. W. (1949). Deperitonealization: clinical and experimental observations. *Ann Surg*, 130, 466-470.
248. Rosenkranz, S. (2004). TGF-beta1 and angiotensin networking in cardiac remodeling. *Cardiovasc Res*, 63(3), 423-432. doi:10.1016/j.cardiores.2004.04.030
249. Rout, U. K., Saed, G. M., & Diamond, M. P. (2002). Transforming growth factor-beta1 modulates expression of adhesion and cytoskeletal proteins in human peritoneal fibroblasts. *Fertil Steril*, 78(1), 154-161.

250. Ryan, G. B., Grobety, J., & Majno, G. (1971). Postoperative peritoneal adhesions. A study of the mechanisms. *Am J Pathol*, 65(1), 117-148.
251. Sadek, K., Macklon, N., Bruce, K., Cagampang, F., & Cheong, Y. (2011). Hypothesis: Role for the circadian Clock system and sleep in the pathogenesis of adhesions and chronic pelvic pain? *Med Hypotheses*, 76(3), 453-456. doi:10.1016/j.mehy.2010.11.020
252. Saed, G. M., Al-Hendy, A., Salama, S. A., & Diamond, M. P. (2008). Adenovirus-mediated expression of cyclooxygenase-2 antisense reverse abnormal genetic profile of human adhesion fibroblasts. *Fertil Steril*, 89(5 Suppl), 1455-1460. doi:10.1016/j.fertnstert.2007.04.061
253. Saed, G. M., & Diamond, M. P. (2004a). Differential expression of alpha smooth muscle cell actin in human fibroblasts isolated from intraperitoneal adhesions and normal peritoneal tissues. *Fertil Steril*, 82 Suppl 3, 1188-1192. doi:10.1016/j.fertnstert.2004.02.147
254. Saed, G. M., & Diamond, M. P. (2004b). Molecular characterization of postoperative adhesions: the adhesion phenotype. *J Am Assoc Gynecol Laparosc*, 11(3), 307-314.
255. Saed, G. M., Fletcher, N. M., & Diamond, M. P. (2016). The Creation of a Model for Ex Vivo Development of Postoperative Adhesions. *Reprod Sci*, 23(5), 610-612. doi:10.1177/1933719115607997
256. Saed, G. M., Zhang, W., & Diamond, M. P. (2001). Molecular characterization of fibroblasts isolated from human peritoneum and adhesions. *Fertil Steril*, 75(4), 763-768.
257. Saed, G. M., Zhao, M., Diamond, M. P., & Abu-Soud, H. M. (2006). Regulation of inducible nitric oxide synthase in post-operative adhesions. *Human Reproduction*, 21(6), 1605-1611. doi:10.1093/humrep/dei500
258. Sahbaz, A., Ersan, F., & Aydin, S. (2014). Effect of Nigella sativa oil on postoperative peritoneal adhesion formation. *J Obstet Gynaecol Res*, 40(2), 532-537. doi:10.1111/jog.12172
259. Sahbaz, A., Isik, H., Aynioglu, O., Gungorduk, K., & Gun, B. D. (2014). Effect of intraabdominal administration of Allium sativum (garlic) oil on postoperative peritoneal adhesion. *Eur J Obstet Gynecol Reprod Biol*, 177, 44-47. doi:10.1016/j.ejogrb.2014.03.018
260. Sajid, M. S., Parampalli, U., Baig, M. K., & McFall, M. R. (2011). A systematic review on the effectiveness of slowly-absorbable versus non-absorbable sutures for abdominal fascial closure following laparotomy. *Int J Surg*, 9(8), 615-625. doi:10.1016/j.ijsu.2011.09.006
261. Saltzman, A. K., Olson, T. A., Mohanraj, D., Carson, L. F., & Ramakrishnan, S. (1996). Prevention of postoperative adhesions by an antibody to vascular permeability factor/vascular endothelial growth factor in a murine model. *Am J Obstet Gynecol*, 174(5), 1502-1506.
262. Sammour, T., Kahokehr, A., Soop, M., & Hill, A. G. (2010). Peritoneal damage: the inflammatory response and clinical implications of the neuro-immuno-humoral axis. *World J Surg*, 34(4), 704-720. doi:10.1007/s00268-009-0382-y
263. Sandoval, P., Jimenez-Heffernan, J. A., Guerra-Azcona, G., Perez-Lozano, M. L., Rynne-Vidal, A., Albar-Vizcaino, P., . . . Lopez-Cabrera, M. (2016). Mesothelial-to-mesenchymal transition in the pathogenesis of post-surgical peritoneal adhesions. *J Pathol*, 239(1), 48-59. doi:10.1002/path.4695
264. Sarieddine, M. Z., Scheckenbach, K. E., Foglia, B., Maass, K., Garcia, I., Kwak, B. R., & Chanson, M. (2009). Connexin43 modulates neutrophil recruitment to the lung. *J Cell Mol Med*, 13(11-12), 4560-4570. doi:10.1111/j.1582-4934.2008.00654.x
265. Scheckenbach, K. E., Crespin, S., Kwak, B. R., & Chanson, M. (2011). Connexin channel-dependent signaling pathways in inflammation. *J Vasc Res*, 48(2), 91-103. doi:10.1159/000316942
266. Schnuriger, B., Barmparas, G., Branco, B. C., Lustenberger, T., Inaba, K., & Demetriades, D. (2011). Prevention of postoperative peritoneal adhesions: a review of the literature. *Am J Surg*, 201(1), 111-121. doi:10.1016/j.amjsurg.2010.02.008
267. Sheldon, H. K., Gainsbury, M. L., Cassidy, M. R., Chu, D. I., Stucchi, A. F., & Becker, J. M. (2012). A sprayable hyaluronate/carboxymethylcellulose adhesion barrier exhibits regional adhesion reduction efficacy and does not impair intestinal healing. *J Gastrointest Surg*, 16(2), 325-333. doi:10.1007/s11605-011-1709-1
268. Smith, C. A., Stauber, F., Waters, C., Alway, S. E., & Stauber, W. T. (2007). Transforming growth factor-beta following skeletal muscle strain injury in rats. *J Appl Physiol* (1985), 102(2), 755-761. doi:10.1152/jappphysiol.01503.2005
269. Smokovitis, A., & Astrup, T. (1980). The fibrinolytic response in injured animal tissues normally low in fibrinolytic activity. *Haemostasis*, 9(3), 175-187. doi:10.1159/000214356

270. Smolarek, S., Shalaby, M., Paolo Angelucci, G., Missori, G., Capuano, I., Franceschilli, L., . . . Sileri, P. (2016). Small-Bowel Obstruction Secondary to Adhesions After Open or Laparoscopic Colorectal Surgery. *JSLs*, 20(4). doi:10.4293/JSLs.2016.00073
271. Snoj, M., Ar'Rajab, A., Ahren, B., Larsson, K., & Bengmark, S. (1993). Phospholipase-resistant phosphatidylcholine reduces intra-abdominal adhesions induced by bacterial peritonitis. *Res Exp Med (Berl)*, 193(2), 117-122.
272. Soon, A. S., Chua, J. W., & Becker, D. L. (2016). Connexins in endothelial barrier function - novel therapeutic targets countering vascular hyperpermeability. *Thromb Haemost*, 116(5), 852-867. doi:10.1160/TH16-03-0210
273. Sortini, D., Feo, C. V., Maravegias, K., Carcoforo, P., Pozza, E., Liboni, A., & Sortini, A. (2006). Role of peritoneal lavage in adhesion formation and survival rate in rats: an experimental study. *J Invest Surg*, 19(5), 291-297. doi:10.1080/08941930600889409
274. Stadlmann, S., Pollheimer, J., Renner, K., Zeimet, A. G., Offner, F. A., & Amberger, A. (2006). Response of human peritoneal mesothelial cells to inflammatory injury is regulated by interleukin-1b and tumor necrosis factor-a. *Wound Repair and Regeneration*, 14(2), 187-194. doi:doi:10.1111/j.1743-6109.2006.00109.x
275. Stein, C. A. (2001). The experimental use of antisense oligonucleotides: a guide for the perplexed. *J Clin Invest*, 108(5), 641-644. doi:10.1172/jci13885
276. Stroka, K. M., & Aranda-Espinoza, H. (2011). Endothelial cell substrate stiffness influences neutrophil transmigration via myosin light chain kinase-dependent cell contraction. *Blood*, 118(6), 1632-1640. doi:10.1182/blood-2010-11-321125
277. Sugama, Y., Tiruppathi, C., offakidevi, K., Andersen, T. T., Fenton, J. W., 2nd, & Malik, A. B. (1992). Thrombin-induced expression of endothelial P-selectin and intercellular adhesion molecule-1: a mechanism for stabilizing neutrophil adhesion. *J Cell Biol*, 119(4), 935-944.
278. Sun, C., Wu, M. H., & Yuan, S. Y. (2011). Nonmuscle myosin light-chain kinase deficiency attenuates atherosclerosis in apolipoprotein E-deficient mice via reduced endothelial barrier dysfunction and monocyte migration. *Circulation*, 124(1), 48-57. doi:10.1161/CIRCULATIONAHA.110.988915
279. Suzuki, T., Kono, T., Bochimoto, H., Hira, Y., Watanabe, T., & Furukawa, H. (2015). An injured tissue affects the opposite intact peritoneum during postoperative adhesion formation. *Sci Rep*, 5, 7668. doi:10.1038/srep07668
280. Swank, D. J., & Jeekel, H. (2004). Laparoscopic adhesiolysis in patients with chronic abdominal pain. *Curr Opin Obstet Gynecol*, 16(4), 313-318.
281. Sylvia, S., Juergen, P., Kathrin, R., G., Z. A., A., O. F., & Albert, A. (2006). Response of human peritoneal mesothelial cells to inflammatory injury is regulated by interleukin-1b and tumor necrosis factor-a. *Wound Repair and Regeneration*, 14(2), 187-194. doi:doi:10.1111/j.1743-6109.2006.00109.x
282. Takagi, K., Araki, M., Fukuoka, H., Takeshita, H., Hidaka, S., Nanashima, A., . . . Nakajima, N. (2013). Novel powdered anti-adhesion material: preventing postoperative intra-abdominal adhesions in a rat model. *Int J Med Sci*, 10(4), 467-474. doi:10.7150/ijms.5607
283. Takakura, N., Huang, X. L., Naruse, T., Hamaguchi, I., Dumont, D. J., Yancopoulos, G. D., & Suda, T. (1998). Critical role of the TIE2 endothelial cell receptor in the development of definitive hematopoiesis. *Immunity*, 9(5), 677-686.
284. Takase-Sanchez, M. M., Brooks, H. M., Hale, D. S., & Heit, M. H. (2015). Obliterative Surgery for the Treatment of Pelvic Organ Prolapse: A Patient Survey on Reasons for Surgery Selection and Postoperative Decision Regret and Satisfaction. *Female Pelvic Med Reconstr Surg*, 21(6), 325-331. doi:10.1097/SPV.0000000000000198
285. Tang, B., Peng, Z. H., Yu, P. W., Yu, G., Qian, F., Zeng, D. Z., . . . Luo, H. X. (2013). Aberrant expression of Cx43 is associated with the peritoneal metastasis of gastric cancer and Cx43-mediated gap junction enhances gastric cancer cell diapedesis from peritoneal mesothelium. *PLoS One*, 8(9), e74527. doi:10.1371/journal.pone.0074527
286. Tang, Y., Harrington, A., Yang, X., Friesel, R. E., & Liaw, L. (2010). The contribution of the Tie2+ lineage to primitive and definitive hematopoietic cells. *Genesis*, 48(9), 563-567. doi:10.1002/dvg.20654
287. Tarhan, O. R., Barut, I., & Sezik, M. (2008). An evaluation of normal saline and taurolidine on intra-abdominal adhesion formation and peritoneal fibrinolysis. *J Surg Res*, 144(1), 151-157. doi:10.1016/j.jss.2007.09.006

288. Tarhan, O. R., Barut, I., Sutcu, R., Akdeniz, Y., & Akturk, O. (2006). Pentoxifylline, a methyl xanthine derivative, reduces peritoneal adhesions and increases peritoneal fibrinolysis in rats. *Tohoku J Exp Med*, 209(3), 249-255.
289. ten Broek, R. P., Bakkum, E. A., Laarhoven, C. J., & van Goor, H. (2016). Epidemiology and Prevention of Postsurgical Adhesions Revisited. *Ann Surg*, 263(1), 12-19. doi:10.1097/SLA.0000000000001286
290. ten Broek, R. P., Issa, Y., van Santbrink, E. J., Bouvy, N. D., Kruitwagen, R. F., Jeekel, J., . . . van Goor, H. (2013a). Burden of adhesions in abdominal and pelvic surgery: systematic review and met-analysis. *BMJ*, 347, f5588. doi:10.1136/bmj.f5588
291. ten Broek, R. P., Kok-Krant, N., Bakkum, E. A., Bleichrodt, R. P., & van Goor, H. (2013b). Different surgical techniques to reduce post-operative adhesion formation: a systematic review and meta-analysis. *Hum Reprod Update*, 19(1), 12-25. doi:10.1093/humupd/dms032
292. ten Broek, R. P. G., Stommel, M. W. J., Strik, C., van Laarhoven, C., Keus, F., & van Goor, H. (2014). Benefits and harms of adhesion barriers for abdominal surgery: a systematic review and meta-analysis. *Lancet*, 383(9911), 48-59. doi:10.1016/S0140-6736(13)61687-6
293. Tomasek, J. J., McRae, J., Owens, G. K., & Haaksma, C. J. (2005). Regulation of α -Smooth Muscle Actin Expression in Granulation Tissue Myofibroblasts Is Dependent on the Intronic CArG Element and the Transforming Growth Factor- β 1 Control Element. *The American Journal of Pathology*, 166(5), 1343-1351. doi:[https://doi.org/10.1016/S0002-9440\(10\)62353-X](https://doi.org/10.1016/S0002-9440(10)62353-X)
294. Topley, N., Brown, Z., Jorres, A., Westwick, J., Davies, M., Coles, G. A., & Williams, J. D. (1993). Human peritoneal mesothelial cells synthesize interleukin-8. Synergistic induction by interleukin-1 beta and tumor necrosis factor-alpha. *Am J Pathol*, 142(6), 1876-1886.
295. Torre, M., Favre, A., Pini Prato, A., Brizzolara, A., & Martucciello, G. (2002). Histologic study of peritoneal adhesions in children and in a rat model. *Pediatr Surg Int*, 18(8), 673-676. doi:10.1007/s00383-002-0872-6
296. Townsend, K. L., Race, A., Keane, M., Miller, W., Dishaw, L., Fisher, E. R., . . . Allen, M. J. (2011). A novel hydrogel-coated polyester mesh prevents postsurgical adhesions in a rat model. *J Surg Res*, 167(2), e117-124. doi:10.1016/j.jss.2009.09.024
297. Trosko, J. E., & Ruch, R. J. (1998). Cell-cell communication in carcinogenesis. *Front Biosci*, 3, d208-236.
298. Tsai, J. M., Sinha, R., Seita, J., Fernhoff, N., Christ, S., Koopmans, T., . . . Rinkevich, Y. (2018). Surgical adhesions in mice are derived from mesothelial cells and can be targeted by antibodies against mesothelial markers. *Science Translational Medicine*, 10(469), eaan6735. doi:10.1126/scitranslmed.aan6735
299. Tsimoyiannis, E. C. (1989). Role of feces, necrotic tissue, and various blocking agents in the prevention of adhesions. *Ann Surg*, 209(4), 502-503.
300. Turkcapar, A. G., Ozarslan, C., Erdem, E., Bumin, C., Erverdi, N., & Kutlay, J. (1995). The effectiveness of low molecular weight heparin on adhesion formation in experimental rat model. *Int Surg*, 80(1), 92-94.
301. Turkoglu, A., Gul, M., Yuksel, H. K., Alabalik, U., Ulger, B. V., Uslukaya, O., & Avci, Y. (2015). Effect of intraperitoneal curcumin instillation on postoperative peritoneal adhesions. *Med Princ Pract*, 24(2), 153-158. doi:10.1159/000369020
302. Uyama, N., Shimahara, Y., Okuyama, H., Kawada, N., Kamo, S., Ikeda, K., & Yamaoka, Y. (2003). Carbenoxolone inhibits DNA synthesis and collagen gene expression in rat hepatic stellate cells in culture. *J Hepatol*, 39(5), 749-755.
303. Uysal, B., Demirbag, S., Poyrazoglu, Y., Cayci, T., Yesildaglar, N., Guven, A., . . . Korkmaz, A. (2012). Medical ozone therapy decreases postoperative uterine adhesion formation in rats. *Arch Gynecol Obstet*, 286(5), 1201-1207. doi:10.1007/s00404-012-2435-y
304. van den Tol, M. P., Haverlag, R., van Rossen, M. E., Bonthuis, F., Marquet, R. L., & Jeekel, J. (2001). Glove powder promotes adhesion formation and facilitates tumour cell adhesion and growth. *Br J Surg*, 88(9), 1258-1263. doi:10.1046/j.0007-1323.2001.01846.x
305. Van Der Krabben, A. A., Dijkstra, F. R., Nieuwenhuijzen, M., Reijnen, M. M., Schaapveld, M., & Van Goor, H. (2000). Morbidity and mortality of inadvertent enterotomy during adhesiotomy. *Br J Surg*, 87(4), 467-471. doi:10.1046/j.1365-2168.2000.01394.x
306. van Rijen, H. V., van Kempen, M. J., Postma, S., & Jongsma, H. J. (1998). Tumour necrosis factor alpha alters the expression of connexin43, connexin40, and connexin37 in human umbilical vein endothelial cells. *Cytokine*, 10(4), 258-264. doi:10.1006/cyto.1997.0287

307. van Westreenen, M., van den Tol, P. M., Pronk, A., Marquet, R. L., Jeekel, J., & Leguit, P. (1999). Perioperative lavage promotes intraperitoneal adhesion in the rat. *Eur Surg Res*, 31(2), 196-201. doi:10.1159/000008639
308. Veliz, L. P., Gonzalez, F. G., Duling, B. R., Saez, J. C., & Boric, M. P. (2008). Functional role of gap junctions in cytokine-induced leukocyte adhesion to endothelium in vivo. *Am J Physiol Heart Circ Physiol*, 295(3), H1056-H1066. doi:10.1152/ajpheart.00266.2008
309. von Dembowski, T. (1889). Über die Ursachen der peritonealen Adhäsionen nach chirurgischen Eingriffen mit Rücksicht auf die Frage des Ileus nach Laparotomien. *Langenbecks Arch Chir*, 37, 745-748.
310. Vural, B., Canturk, N. Z., Esen, N., Solakoglu, S., Canturk, Z., Kirkali, G., & Sokmensuer, C. (1999). The role of neutrophils in the formation of peritoneal adhesions. *Hum Reprod*, 14(1), 49-54.
311. Vural, B., Mercan, R., Corakci, A., Ozeren, S., Keskin, N., Vural, S., . . . Erk, A. (1998). A trial of reducing adhesion formation in a uterine horn model. *Gynecol Obstet Invest*, 45(1), 58-61. doi:10.1159/000009912
312. Wallwiener, C. W., Kraemer, B., Wallwiener, M., Brochhausen, C., Isaacson, K. B., & Rajab, T. K. (2010). The extent of adhesion induction through electrocoagulation and suturing in an experimental rat study. *Fertil Steril*, 93(4), 1040-1044. doi:10.1016/j.fertnstert.2008.12.002
313. Wallwiener, M., Brucker, S., Hierlemann, H., Brochhausen, C., Solomayer, E., & Wallwiener, C. (2006). Innovative barriers for peritoneal adhesion prevention: liquid or solid? A rat uterine horn model. *Fertil Steril*, 86(4 Suppl), 1266-1276. doi:10.1016/j.fertnstert.2006.05.023
314. Wallwiener, M., Wallwiener, C. W., Molinas, R., Rajab, T. K., Brucker, S. Y., Kraemer, B., & Hartkopf, A. D. (2012). Intraabdominal adhesion formation is associated with differential mRNA expression of metabolic genes PDHb and SDHa. *Arch Gynecol Obstet*, 286(3), 683-686. doi:10.1007/s00404-012-2364-9
315. Wang, C. M., Lincoln, J., Cook, J. E., & Becker, D. L. (2007). Abnormal connexin expression underlies delayed wound healing in diabetic skin. *Diabetes*, 56(11), 2809-2817. doi:10.2337/db07-0613
316. Wei, G., Chen, X., Wang, G., Fan, L., Wang, K., & Li, X. (2016a). Effect of Resveratrol on the Prevention of Intra-Abdominal Adhesion Formation in a Rat Model. *Cell Physiol Biochem*, 39(1), 33-46. doi:10.1159/000445603
317. Wei, G., Chen, X., Wang, G., Jia, P., Xu, Q., Ping, G., . . . Li, X. (2015). Inhibition of cyclooxygenase-2 prevents intra-abdominal adhesions by decreasing activity of peritoneal fibroblasts. *Drug Des Devel Ther*, 9, 3083-3098. doi:10.2147/dddt.s80221
318. Wei, G., Zhou, C., Wang, G., Fan, L., Wang, K., & Li, X. (2016b). Keratinocyte Growth Factor Combined with a Sodium Hyaluronate Gel Inhibits Postoperative Intra-Abdominal Adhesions. *Int J Mol Sci*, 17(10). doi:10.3390/ijms17101611
319. Westphalen, K., Gusarova, G. A., Islam, M. N., Subramanian, M., Cohen, T. S., Prince, A. S., & Bhattacharya, J. (2014). Sessile alveolar macrophages communicate with alveolar epithelium to modulate immunity. *Nature*, 506(7489), 503-506. doi:10.1038/nature12902
320. Whang, S. H., Astudillo, J. A., Sporn, E., Bachman, S. L., Miedema, B. W., Davis, W., & Thaler, K. (2011). In search of the best peritoneal adhesion model: comparison of different techniques in a rat model. *J Surg Res*, 167(2), 245-250. doi:10.1016/j.jss.2009.06.020
321. Willebrords, J. (2016). Connexins and their channels in inflammation. *51(6)*, 413-439. doi:10.1080/10409238.2016.1204980
322. Winny, M., Grethe, L., Maegel, L., Jonigk, D., Lippmann, T., Klempnauer, J., & Poehnert, D. (2016). Impairment of the Peritoneal Surface as a Decisive Factor for Intestinal Adhesions in Intraperitoneal Onlay Mesh Surgery - Introducing a New Rat Model. *Int J Med Sci*, 13(2), 108-112. doi:10.7150/ijms.14056
323. Wiseman, D. M. (2000). Animal Adhesion Models: Design, Variables, and Relevance. In G. S. diZerega (Ed.), *Peritoneal Surgery* (pp. 459-476). New York, NY: Springer New York.
324. Wiseman, D. M. (2008). Disorders of adhesions or adhesion-related disorder: monolithic entities or part of something bigger--CAPPS? *Semin Reprod Med*, 26(4), 356-368. doi:10.1055/s-0028-1082394
325. Wiseman, D. M., Gottlick, L. E., & Diamond, M. P. (1992). Effect of thrombin-induced hemostasis on the efficacy of an absorbable adhesion barrier. *J Reprod Med*, 37(9), 766-770.

326. Wiseman, D. M., Huang, W. J., Johns, D. B., Rodgers, K. E., & Dizerega, G. S. (1994). Time-dependent effect of tolmetin sodium in a rabbit uterine adhesion model. *J Invest Surg*, 7(6), 527-532.
327. Wiseman, D. M., Trout, J. R., Franklin, R. R., & Diamond, M. P. (1999). Metaanalysis of the safety and efficacy of an adhesion barrier (Interceed TC7) in laparotomy. *J Reprod Med*, 44(4), 325-331.
328. Wright, C. S., Berends, R. F., Flint, D. J., & Martin, P. E. (2013). Cell motility in models of wounded human skin is improved by Gap27 despite raised glucose, insulin and IGFBP-5. *Exp Cell Res*, 319(4), 390-401. doi:10.1016/j.yexcr.2012.12.013
329. Wright, J. A., Richards, T., & Becker, D. L. (2012). Connexins and Diabetes. *Cardiology Research and Practice*, 2012, 8. doi:10.1155/2012/496904
330. Wu, Q., Li, L., Wang, N., Gao, X., Wang, B., Liu, X., . . . Gong, C. (2014). Biodegradable and thermosensitive micelles inhibit ischemia-induced postoperative peritoneal adhesion. *Int J Nanomedicine*, 9, 727-734. doi:10.2147/ijn.s55497
331. Wu, Q., Wang, N., He, T., Shang, J., Li, L., Song, L., . . . Gong, C. (2015). Thermosensitive hydrogel containing dexamethasone micelles for preventing postsurgical adhesion in a repeated-injury model. *Sci Rep*, 5, 13553. doi:10.1038/srep13553
332. Wynn, T. A. (2007). Common and unique mechanisms regulate fibrosis in various fibroproliferative diseases. *J Clin Invest*, 117(3), 524-529. doi:10.1172/jci31487
333. Wynn, T. A. (2008). Cellular and molecular mechanisms of fibrosis. *J Pathol*, 214(2), 199-210. doi:10.1002/path.2277
334. Wynn, T. A., & Ramalingam, T. R. (2012). Mechanisms of fibrosis: therapeutic translation for fibrotic disease. *Nat Med*, 18(7), 1028-1040. doi:10.1038/nm.2807
335. Xu, X., Pappo, O., Garbuzenko, E., Bischoff, S. C., Rivkind, A., & Levi-Schaffer, F. (2002a). Mast cell dynamics and involvement in the development of peritoneal adhesions in the rat. *Life Sci*, 70(8), 951-967.
336. Xu, X., Rivkind, A., Pappo, O., Pikarsky, A., & Levi-Schaffer, F. (2002b). Role of mast cells and myofibroblasts in human peritoneal adhesion formation. *Ann Surg*, 236(5), 593-601. doi:10.1097/01.SLA.0000033037.13104.C4
337. Yang, A. H., Chen, J. Y., & Lin, J. K. (2003). Myofibroblastic conversion of mesothelial cells. *Kidney Int*, 63(4), 1530-1539. doi:10.1046/j.1523-1755.2003.00861.x
338. Yang, B., Gong, C., Zhao, X., Zhou, S., Li, Z., Qi, X., . . . Qian, Z. (2012). Preventing postoperative abdominal adhesions in a rat model with PEG-PCL-PEG hydrogel. *Int J Nanomedicine*, 7, 547-557. doi:10.2147/ijn.s26141
339. Yang, B., Yang Gong, C., Yong Qian, Z., Zhao, X., Yu Li, Z., Tao Zhou, S., . . . Quan Wei, Y. (2011). Prevention of abdominal adhesion formation by thermosensitive PECE-hydrogel in a rat uterine horn model. *J Biomed Mater Res B Appl Biomater*, 96(1), 57-66. doi:10.1002/jbm.b.31739
340. Yilmaz, Y., Celik, I. H., Pampal, A., Demirel, G., Topal, F., Oguz, S. S., . . . Dilmen, U. (2012). Effects of different pulmonary surfactants in the prevention of postoperative intraabdominal adhesion formation. *J Pediatr Surg*, 47(8), 1560-1565. doi:10.1016/j.jpedsurg.2012.01.080
341. Yilmazlar, T., Kaya, E., Gurpinar, E., & Emiroglu, H. (1996). Efficacy of tenoxicam on intra-abdominal adhesion prevention in a rat model. *J Int Med Res*, 24(4), 352-357. doi:10.1177/030006059602400406
342. Yin, J., Meng, Q., Zhang, G., & Sun, Y. (2009). Differential methotrexate hepatotoxicity on rat hepatocytes in 2-D monolayer culture and 3-D gel entrapment culture. *Chem Biol Interact*, 180(3), 368-375. doi:10.1016/j.cbi.2009.04.004
343. Zeisberg, M., & Kalluri, R. (2004). The role of epithelial-to-mesenchymal transition in renal fibrosis. *J Mol Med (Berl)*, 82(3), 175-181. doi:10.1007/s00109-003-0517-9
344. Zernecke, A., Bidzhekov, K., Ozuyaman, B., Fraemohs, L., Liehn, E. A., Luscher-Firzlaff, J. M., . . . Weber, C. (2006). CD73/ecto-5'-nucleotidase protects against vascular inflammation and neointima formation. *Circulation*, 113(17), 2120-2127. doi:10.1161/circulationaha.105.595249
345. Zhang, J., Yang, G. M., Zhu, Y., Peng, X. Y., Li, T., & Liu, L. M. (2015). Role of connexin 43 in vascular hyperpermeability and relationship to Rock1-MLC20 pathway in septic rats. *Am J Physiol Lung Cell Mol Physiol*, 309(11), L1323-1332. doi:10.1152/ajplung.00016.2015
346. Zheng, G., Lyons, J. G., Tan, T. K., Wang, Y., Hsu, T. T., Min, D., . . . Harris, D. C. (2009). Disruption of E-cadherin by matrix metalloproteinase directly mediates epithelial-

- mesenchymal transition downstream of transforming growth factor-beta1 in renal tubular epithelial cells. *Am J Pathol*, 175(2), 580-591. doi:10.2353/ajpath.2009.080983
347. Zimowska, M., Duchesnay, A., Dragun, P., Oberbek, A., Moraczewski, J., & Martelly, I. (2009). Immunoneutralization of TGFbeta1 Improves Skeletal Muscle Regeneration: Effects on Myoblast Differentiation and Glycosaminoglycan Content. *Int J Cell Biol*, 2009, 659372. doi:10.1155/2009/659372
348. Zuhlke, H. V., Lorenz, E. M., Straub, E. M., & Savvas, V. (1990). [Pathophysiology and classification of adhesions]. *Langenbecks Arch Chir Suppl II Verh Dtsch Ges Chir*, 1009-1016.