

ORIGINAL RESEARCH COMMUNICATION

---

# Hydrogen Sulfide Is an Endogenous Regulator of Aging in *Caenorhabditis elegans*

Bedoor Qabazard,<sup>1,\*</sup> Ling Li,<sup>1,\*</sup> Jan Gruber,<sup>2,3</sup> Meng Teng Peh,<sup>4</sup> Li Fang Ng,<sup>2</sup> Srinivasan Dinesh Kumar,<sup>5</sup> Peter Rose,<sup>6</sup> Choon-Hong Tan,<sup>7</sup> Brian W. Dymock,<sup>8</sup> Feng Wei,<sup>8</sup> Suresh C. Swain,<sup>1</sup> Barry Halliwell,<sup>2</sup> Stephen R. Stürzenbaum,<sup>1</sup> and Philip K. Moore<sup>4</sup>

## Abstract

**Aims:** To investigate the role of endogenous hydrogen sulfide (H<sub>2</sub>S) in the control of aging and healthspan of *Caenorhabditis elegans*. **Results:** We show that the model organism, *C. elegans*, synthesizes H<sub>2</sub>S. Three H<sub>2</sub>S-synthesizing enzymes are present in *C. elegans*, namely cystathionine  $\gamma$  lyase (CSE), cystathionine  $\beta$  synthetase, and 3-mercaptopyruvate transferase (MPST or 3-MST). Genetic deficiency of *mpst-1* (3-MST orthologue 1), but not *cth-2* (CSE orthologue), reduced the lifespan of *C. elegans*. This effect was reversed by a pharmacological H<sub>2</sub>S donor (GYY4137). GYY4137 also reduced detrimental age-dependent changes in a range of physiological indices, including pharyngeal contraction and defecation. Treatment of *C. elegans* with GYY4137 increased the expression of several age-related, stress response, and antioxidant genes, whereas MitoSOX Red fluorescence, indicative of reactive oxygen species generation, was increased in *mpst-1* knockouts and decreased by GYY4137 treatment. GYY4137 additionally increased the lifespan in short-lived *mev-1* mutants with elevated oxidative stress and protected wild-type *C. elegans* against paraquat poisoning. The lifespan-prolonging and health-promoting effects of H<sub>2</sub>S in *C. elegans* are likely due to the antioxidant action of this highly cell-permeable gas. **Innovation:** The possibility that novel pharmacological agents based on the principle of H<sub>2</sub>S donation may be able to retard the onset of age-related disease by slowing the aging process warrants further study. **Conclusion:** Our results show that H<sub>2</sub>S is an endogenous regulator of oxidative damage, metabolism, and aging in *C. elegans* and provide new insight into the mechanisms, which control aging in this model organism. *Antioxid. Redox Signal.* 00, 000–000.

## Introduction

**H**YDROGEN SULFIDE (H<sub>2</sub>S) AFFECTS cell metabolism notably by inhibiting cytochrome *c* oxidase and thereby reducing mitochondrial adenosine triphosphate (ATP) production (4). It was originally thought that H<sub>2</sub>S would kill cells at low concentrations (by starving the cell of ATP), but it is now becoming increasingly clear that physiologically relevant concentrations are not toxic and that H<sub>2</sub>S actually exhibits cytoprotective effects in a wide range of cells and

tissues (11). For example, H<sub>2</sub>S protects the heart (12) against damage due to ischemia reperfusion.

Many of the cytoprotective effects of H<sub>2</sub>S are likely due to its ability to quench, or modulate, the formation of damaging free radical species. It has been known for many years that reactive oxygen species (ROS) trigger accumulating damage in aging cells thus causing loss of function or even death and that they may also act as signaling molecules (1). The fact that exogenous H<sub>2</sub>S gas prolongs the lifespan of *Caenorhabditis elegans* (18) suggests a link between this gas

---

<sup>1</sup>School of Biomedical Science, King's College London, London, United Kingdom.

Departments of <sup>2</sup>Biochemistry and <sup>4</sup>Pharmacology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore.

<sup>3</sup>Yale-NUS College, Science Division, Singapore, Singapore.

<sup>5</sup>Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore.

<sup>6</sup>School of Life Sciences, University of Lincoln, Lincoln, United Kingdom.

<sup>7</sup>Division of Chemistry and Biological Chemistry, Nanyang Technological University, Singapore, Singapore.

<sup>8</sup>Department of Pharmacy, National University of Singapore, Singapore, Singapore.

\*These authors contributed equally to this work.

### Innovation

While exogenous hydrogen sulfide (H<sub>2</sub>S) reportedly protects *Caenorhabditis elegans* against stress and increases longevity, the role of endogenous H<sub>2</sub>S, if any, is unknown. Here, we report that (i) *C. elegans* contains cystathionine  $\gamma$  lyase, cystathionine  $\beta$  synthetase, and 3-mercaptopyruvate transferase enzyme orthologues and synthesizes H<sub>2</sub>S, (ii) genetic deficiency of *mpst-1* reduces endogenous H<sub>2</sub>S production and negatively affects health and lifespan, and (iii) GYY4137, which releases low concentrations of H<sub>2</sub>S over an extended time period, prolongs the lifespan and retards age-dependent deterioration in physiological functions. GYY4137 seems well suited as a prototype for the development of a new class of drugs with beneficial effects on healthy aging.

and aging. However, whether *C. elegans* synthesizes H<sub>2</sub>S naturally and whether endogenous H<sub>2</sub>S affects the lifespan in this organism perhaps by an antioxidant action is not known.

In the present work, we have characterized endogenous H<sub>2</sub>S biosynthesis in *C. elegans* and investigated its effect and mechanism of action on lifespan and other markers of health in this organism.

### Results

#### Endogenous H<sub>2</sub>S synthesis in *C. elegans*

Mammals synthesize H<sub>2</sub>S using a combination of the following enzymes: cystathionine  $\gamma$  lyase (CSE), cystathionine  $\beta$  synthetase (CBS), and 3-mercaptopyruvate transferase (MPST; usually abbreviated in the literature to 3-MST). CSE and CBS convert L-cysteine to H<sub>2</sub>S in the presence of pyridoxal 5' phosphate (PPP), whereas 3-MST utilizes 3-mercaptopyruvate (3-MP) as substrate. All three of these enzymes are evolutionarily conserved in lower species with two CSE orthologues (*cth-1*, *cth-2*), two CBS orthologues (*cbs-1*, *cbs-2*), and seven 3-MST orthologues (*mpst1-7*) predicted in *C. elegans* (16).

In preliminary experiments, we demonstrated the presence of CSE, CBS, and 3-MST in adult *C. elegans* using immunofluorescence (Fig. 1A–F) and diaminobenzidine staining (Supplementary Fig. S1A–C; Supplementary Data are available online at [www.liebertpub.com/ars](http://www.liebertpub.com/ars)). By quantitative polymerase chain reaction (qPCR), we also confirmed the presence and relative expression levels of all the named genes predicted to encode for CSE, CBS, and 3-MST proteins (Supplementary Fig. S2). In separate experiments, we assessed the formation of H<sub>2</sub>S from exogenous L-cysteine (to assay the CSE and CBS activity) and 3-MP (to assay the 3-MST activity) in homogenates of adult *C. elegans* and found significant biosynthesis after incubation with either L-cysteine (1.49 ± 0.05 nmol/mg protein/min, *n* = 12) or 3-MP (1.09 ± 0.05 nmol/mg/min, *n* = 3). Finally, we used a sensitive rp-HPLC assay to identify endogenous L-cysteine (2.03 ± 0.47  $\mu$ M, *n* = 6), 3-MP (0.13 ± 0.03  $\mu$ M, *n* = 6), and H<sub>2</sub>S itself (3.97 ± 0.87  $\mu$ M, *n* = 6) in homogenates of adult *C. elegans*. From these results, we conclude that H<sub>2</sub>S, its synthesizing enzymes, and substrates thereof occur naturally in *C. elegans*.

#### 3-MST regulates the lifespan and health of *C. elegans*


Since exposure of *C. elegans* to H<sub>2</sub>S gas (50 ppm) prolongs the lifespan in this organism (18), we wondered whether naturally produced (*i.e.*, endogenous) H<sub>2</sub>S could exhibit a similar lifespan-prolonging effect in this organism.

To test this possibility, we examined the effect of genetic (RNAi) knockdown of CSE or 3-MST on lifespan. CBS was excluded from this part of the study since RNAi-mediated knockdown of the CBS orthologue reportedly impairs the development (time to adulthood >9 days) and caused abnormalities in the gut and pharynx (30). There are no previous descriptions of CSE or 3-MST knockdown in *C. elegans*. The knockdown efficiency of *cth-2* (ZK1127.10) and *mpst-1* (D2023) as determined by qPCR was similar, that is, 66.5% ± 0.6% and 64.2% ± 0.6%, respectively (*n* = 3). While lifespan was unchanged in *cth-2* RNAi animals (mean lifespan: 12.15 ± 0.16 days compared to (cf) 12.23 ± 0.16 days, *p* > 0.05), it was substantially reduced (36.5%) in *mpst-1* RNAi animals (mean lifespan: 7.76 ± 0.18 days, *p* < 0.05) (Fig. 2A), implying a possible role for 3-MST, but probably not CSE, in regulating the lifespan in this organism.

Further experiments were then carried out using *C. elegans* carrying an *mpst-1* null deletion allele (ok2040). Initial characterization of the *mpst-1* knockout phenotype revealed a reduction in brood size (4 days: 41.8 ± 2.7 c.f. 206.9 ± 4.9 animals, *p* < 0.05) but no significant difference in the rate of development to the adult stage between *mpst-1* knockouts and wild-type controls (Supplementary Fig. S3A). Loss of *mpst-1* and 3-MST protein was confirmed in these animals by qPCR (Supplementary Fig. S3B) and immunohistochemically (Supplementary Fig. S3C–F). The former also revealed significant, presumably compensatory, upregulation of five of the other six 3-MST orthologues (*i.e.*, *mpst 3-7*) in these *mpst-1* deletion mutants. Subsequent lifespan experiments confirmed the short-lived (15.4% reduced compared to wild-type controls) phenotype of *mpst-1* knockouts (Fig. 2B). Interestingly, partial knockdown of *mpst-1* was more effective than complete deletion of this gene in terms of shortening the lifespan in these animals. Again, this may be due to the compensatory upregulation of other 3-MST orthologues in *mpst-1* knockouts as noted above.

Since lifespan was detrimentally affected in *mpst-1* knockouts, we wondered whether there was a corresponding decline in the general health status of these animals. To assess this possibility, we compared the healthspan in *mpst-1* knockout and wild-type *C. elegans*. While healthspan deteriorated over time in both wild types and mutants, the rate of loss of normal healthy behavior was significantly accelerated in the *mpst-1* knockouts (Fig. 2C).

Finally, we attempted to correlate the changes in lifespan and healthspan found in *mpst-1* knockouts with the ability of these animals to synthesize H<sub>2</sub>S. The amount of H<sub>2</sub>S detected in homogenates of synchronized adult (day 5), *mpst-1* knockout *C. elegans* (38.99 ± 3.32 nmol/mg, *n* = 8), was ~57% less than that found in wild types (92.25 ± 19.80 nmol/mg, *n* = 8, *p* < 0.05), which shows that the deletion of *mpst-1* significantly reduces H<sub>2</sub>S production in these animals. The fact that H<sub>2</sub>S production was not completely abolished in *mpst-1* knockouts is probably due to compensatory upregulation of other 3-MST orthologues coupled with the continued presence in these animals of additional H<sub>2</sub>S-synthesizing enzymes, such as CSE and CBS.



**FIG. 1. Expression of hydrogen sulfide ( $H_2S$ )-synthesizing enzymes in *Caenorhabditis elegans*.** Transverse sections showing immunoreactive staining for cystathionine  $\gamma$  lyase (CSE) (A), cystathionine  $\beta$  synthetase (CBS) (C), and 3-mercaptopyruvate transferase (3-MST) (E) in *C. elegans*. 4',6-diamidino-2-phenylindole (DAPI) staining of nuclei is shown in (B, D, and F). Magnification is  $\times 20$ . Horizontal bar shows scale (100  $\mu m$ ).

#### Mimicking the effect of endogenous $H_2S$ using the slow-releasing donor, GYY4137


Since genetic deficiency of *mpst-1* reduced lifespan and decreased  $H_2S$  levels in *C. elegans*, we decided to investigate whether replenishing  $H_2S$  in *mpst-1* knockout animals was able to reverse the lifespan deficit.

We have previously reported that GYY4137 is a slow-releasing  $H_2S$  donor (15). In preliminary experiments, GYY4137 (100  $\mu M$ ) was incubated at 20°C in aqueous vehicle for up to 30 days to mimic, as far as possible, the treatment of *C. elegans* during a complete lifespan experiment. Under these conditions, the hydrolysis of GYY4137 was slow with increasing concentrations of  $H_2S$  ( $\sim 5$ –7  $\mu M$ ) generated each day up to a plateau at day 7 thereafter increasing only very slowly ( $< 0.5 \mu M$  per day) up to day 30 (Fig. 2D). The maximum daily exposure of *C. elegans* to  $H_2S$  during GYY4137 treatment in the first 7 days can therefore not be  $> 50\%$  higher than the concentration found naturally in homogenates (*i.e.*, 3.97  $\mu M$ ). Indeed, the true exposure of *C. elegans* to  $H_2S$  gas released from GYY4137 in this way is likely to be considerably less as some  $H_2S$  generated will not be absorbed and  $H_2S$  release from GYY4137 declines after day 7. Previous researchers have demonstrated the lifespan-enhancing effect of  $H_2S$  in *C. elegans* following exposure to  $H_2S$  gas (50 ppm) (18). This is equivalent to a dissolved  $H_2S$  concentration of 1.47 mM, which is more than 370 times greater than the presumed endogenous concentration of this gas in

*C. elegans*. For this reason, we chose not to use  $H_2S$  gas in the present experiments, but GYY4137, which most likely better mimics the natural exposure rate of *C. elegans* to this gas.

Treatment of *mpst-1* knockout animals with GYY4137 (Fig. 2B) completely restored, without overshooting, the shortened lifespan of *mpst-1* knockouts (Fig. 2B). This result suggests that replenishment with  $H_2S$ , under these conditions, rescued the *mpst-1* knockout phenotype and, importantly, also pinpoints  $H_2S$  as a key regulator of lifespan under this perturbation. Intriguingly, GYY4137 also prolonged lifespan in wild-type *C. elegans* (Fig. 2E) an effect not seen when time-expired (released  $> 85\%$  of its available  $H_2S$ ) compound was used (Fig. 2F). This effect of GYY4137 was similar to, or even greater than the lifespan-enhancing response reported for rapamycin (100  $\mu M$ , 19% increase; 26) or resveratrol (1 mM, 19% increase; 29) or those typically seen with antioxidant interventions (24) but less than that in *C. elegans* exposed to  $H_2S$  gas (50 ppm) in which lifespan was increased by 74% (18). The apparent discrepancy between our results using GYY4137 to donate  $H_2S$  and those of other researchers using  $H_2S$  gas may be explained by the very different concentrations of  $H_2S$  to which these animals are exposed by the two treatment regimens.

Since loss of endogenous  $H_2S$  following *mpst-1* knockout reduced healthspan in *C. elegans*, we also determined whether GYY4137 treatment had the reverse effect, that is, to improve the healthspan in wild-type animals. GYY4137 not only significantly increased overall healthspan (Fig. 3A) but also, at



**FIG. 2. Response of *C. elegans* to 3-MST deficiency and effect of GYY4137.** (A) Effect of *cth-2* and *mpst-1* RNAi knock-down on lifespan,  $n = 130\text{--}288$  animals per group,  $*p < 0.05$  compared to wild type. (B) Comparison of lifespan in wild-type and *mpst-1* (*ok2040*) null animals and rescue of *mpst-1* null animals by GYY4137 (GYY;  $100\ \mu\text{M}$ ),  $n = 340\text{--}516$  animals per group,  $*p < 0.05$ , *mpst-1* null versus wild type,  $+p < 0.05$  *mpst-1* null versus *mpst-1* null + GYY4137. (C) Healthspan comparison in wild-type and *mpst-1* knockout animals as assessed over 30 days,  $n = 108\text{--}149$  animals per group,  $*p < 0.05$ . (D) Spontaneous release of H<sub>2</sub>S from GYY4137 ( $100\ \mu\text{M}$ ) incubated for up to 30 days at 20°C. Results are mean  $\pm$  SE mean,  $n = 6\text{--}9$  (where not shown standard error bars lie within dimensions of symbol). (E) Effect of GYY4137 ( $100\ \mu\text{M}$ ) on lifespan of wild-type *C. elegans*,  $n = 114\text{--}123$ ,  $*p < 0.05$ . (F) Effect of time-expired GYY4137 ( $100\ \mu\text{M}$ , 10 days at room temperature) on lifespan of *C. elegans*,  $n = 67\text{--}86$  animals per group,  $p > 0.05$ .


least partially, reversed a number of other age-dependent physiological/behavioral changes, including the decline in body length and area (Fig. 3B, C) and the reduction in pharyngeal pumping rate (Fig. 3D), defecation (Fig. 3E), and head thrashing (Fig. 3F).

Thus, the treatment of *C. elegans* with continuous low concentrations of exogenous H<sub>2</sub>S, released from GYY4137, (i) rescued the phenotype (*i.e.*, normalized the lifespan) of *mpst-1* knockouts, (ii) increased the lifespan of wild-type *C. elegans*, and (iii) improved overall health, that is, reduced the age-related deterioration of physiological parameters in this organism.

#### Mechanism of action of H<sub>2</sub>S

Since knockout of *mpst-1* is likely to affect the expression of a wide range of genes due to compensatory changes and

many of these are presumably secondary to altered health and/or lifespan, we decided, as a first step to understanding the role of H<sub>2</sub>S in these processes, to examine the effect of GYY4137 on the transcriptome profile of wild-type *C. elegans*. To gauge intra-assay variation and to justify statistical analysis, three separate groups of animals were treated with either GYY4137 ( $100\ \mu\text{M}$ ) or vehicle (water). Microarray analysis showed a good degree of conformity between individual batches with significant ( $p < 0.05$ ) changes in the expression of 1045 genes between GYY4137- and vehicle-treated groups (Supplementary Fig. S4). Gene ontology analysis revealed significant differences in the expression of 18 aging- and 31 stress-related genes (Supplementary Tables S1 and S2). Notable age-related genes upregulated by GYY4137 include *dod-6* ( $1.4\times$ ), *dod-17* ( $1.6\times$ ), *dod-21* ( $8.7\times$ ), *dod-22* ( $4.6\times$ ),




**FIG. 3. Effect of GYY4137 on health and behavior of *C. elegans*.** Time course (days of adulthood, *i.e.*, post L4) on the effect of GYY4137 (GYY; numbers in parenthesis show concentration in  $\mu\text{M}$ ) on (A) healthspan,  $n=76-92$ , (B) body length,  $n=20$ , (C) body area,  $n=20$ , (D) pharyngeal pumping rate,  $n=10$ , (E) defaecation rate,  $n=10$ , and (F) head shake frequency studied on day 1 of adulthood of *C. elegans*,  $n=20$ . Results show mean  $\pm$  SEM,  $*p < 0.05$ .

*dod-23* ( $1.8\times$ ), and *dod-24* ( $1.9\times$ ) as well as *ins-7* (insulin-related,  $2.4\times$ ). *Daf-16* is a key gene involved in aging in *C. elegans* and its activation triggers numerous downstream genes collectively referred to as downstream of *daf-16* or *dod* genes (9). Stress response and antioxidant genes upregulated by GYY4137 included *hsp-17* (heat shock protein,  $1.6\times$ ), *catalase* ( $1.4\times$ ), and *glutathione S-transferase* ( $1.15-1.2\times$ ).

The role of mitochondrial ROS, ROS-mediated damage, endogenous antioxidant systems, and small-molecule antioxidants in the control of lifespan in *C. elegans* is the subject of much active debate and research (1, 5).  $\text{H}_2\text{S}$  is a powerful, highly membrane permeable reducing agent and metabolic modulator, and the possibility that its effect on health and lifespan may be mediated through an antioxidant mechanism is intriguing. We therefore decided to investigate this possibility in greater depth.

First, we showed that GYY4137 exhibited pronounced antioxidant activity in two standard biochemical assays *in vitro* (Supplementary Fig. S5A, B). In addition, GYY4137 increased the survival of paraquat-exposed wild-type *C. elegans* (Fig. 4A). Paraquat kills *C. elegans* by promoting ROS generation, an effect associated with the upregulation of the stress response gene, *pqm-1* ( $1.5\times$ ). In further experiments, we used three separate assays to monitor ROS production and associated damage in adult *C. elegans*. In the first assay, mitochondrial ROS generation, as determined using the fluorescent dye MitoSOX Red, was reduced in *C. elegans* treated with GYY4137 (Fig. 4B) but was increased (cf. wild type) in *mpst-1* null animals (Fig. 4C). In the second assay, mtDNA lesions due to oxidative damage were significantly higher in *mpst-1* knockouts than in untreated and GYY4137-treated wild-type controls (Fig. 4D). Using this assay, we also noted a trend toward lower mtDNA damage levels in GYY4137-treated wild types,

although this effect did not reach statistical significance ( $p=0.07$ , Fig. 4D). However, mtDNA damage in wild-type controls is extremely low, in the order of one lesion per mtDNA molecule and hence it is not surprising that damage levels are not suppressed even further by GYY4137 treatment. In the third assay, the protein carbonyl (PC) content, another marker of oxidative damage, was significantly reduced by GYY4137 in wild-type animals (Fig. 4E), while a nonsignificant ( $p=0.08$ ) trend to increased PC content was noted in *mpst-1* null animals (Fig. 4F). Thus, each of these biochemical assays supports the hypothesis that the degree of oxidative stress/ROS generation is correlated with  $\text{H}_2\text{S}$  concentration, that is, reduced by GYY4137 treatment and increased in *mpst-1* knockouts with diminished endogenous  $\text{H}_2\text{S}$ -synthesizing capacity. In addition, we used the dichlorofluorescein diacetate (DCF-DA) fluorescence assay to assess ROS production from *C. elegans* in these experiments. We note that the validity of this assay using lysed worms has been challenged since this causes the release of transition metal ions, such as iron, which can generate ROS by redox cycling. However, in the present experiments, we used whole animals that had not been disrupted and in which iron therefore remained sequestered. We are aware of the controversy regarding the actual ROS responsible for DCF-DA fluorescence and, more generally regarding the meaning of this fluorescence. However, we believe that when used in whole *C. elegans* and interpreted with the appropriate caveats and especially when combined with other assays, DCF-DA fluorescence can be a useful indicator of global ROS flux. Indeed, we have previously utilized this assay to show a clear age-dependent increase in this marker of global ROS flux (5). Using DCF-DA, we noted increased ROS production in *mpst-1* RNAi *C. elegans*, which was reduced by GYY4137 treatment both in



**FIG. 4.** H<sub>2</sub>S and markers of oxidative stress in *C. elegans*. (A) Effect of GYY4137 (GYY; 100  $\mu$ M) on paraquat-evoked (10 mM) toxicity. Data show median survival assessed daily over a 7-day period,  $n = 154$ – $187$ ,  $*p < 0.05$ . (B) Effect of GYY4137 (100  $\mu$ M) treatment on reactive oxygen species (ROS) production determined by MitoSOX Red fluorescence, data show mean  $\pm$  SEM of three independent experiments,  $*p < 0.05$ . (C) MitoSOX Red fluorescence indicative of ROS production in wild-type (control) and *mpst-1* knockout *C. elegans*, data show mean  $\pm$  SEM of three independent experiments,  $*p < 0.05$ . (D) Measurement of mtDNA damage in wild type (control), wild type treated with GYY4137 (100  $\mu$ M), and *mpst-1* knockout *C. elegans*, data show mean  $\pm$  SEM of three independent experiments,  $*p < 0.05$ . Measurement of protein carbonyl content (PCC) in GYY4137-treated (GYY; 100  $\mu$ M) wild-type controls (E) and in *mpst-1* knockout *C. elegans* (F), data show mean  $\pm$  SEM,  $n = 10$ – $13$ ,  $*p < 0.05$ . (G) Effect of GYY4137 (GYY; numbers in parenthesis show concentration in  $\mu$ M) on survival of *mev-1* (*kn1*) *C. elegans* mutants,  $n = 52$ – $58$ ,  $*p < 0.05$ .

these mutants and in wild-type controls (Supplementary Fig. S6).

Finally, we used the *C. elegans* mutant, *mev-1* (*kn1*), which is short lived due to its overproduction of ROS. GYY4137 significantly increased the lifespan in *mev-1* *C. elegans* again pointing to an antioxidant effect as a likely mechanism of action (Fig. 4G).

Taken together, these results strongly suggest that both endogenous and exogenous H<sub>2</sub>S prolongs the lifespan and healthspan in *C. elegans* by acting as antioxidant and/or modulator of ROS production.

## Discussion

Exposing *C. elegans* to H<sub>2</sub>S in air increases their lifespan, improves their tolerance to elevated temperature (18), and increases their resistance to killing by pseudomonas (2). While pharmacological treatment with H<sub>2</sub>S gas in this way clearly protects *C. elegans* against some forms of stress and increases their longevity, it is not known whether endogenous H<sub>2</sub>S, made by *C. elegans*, has similar activity. This is important as it would imply a hitherto unknown physiological role of this molecule and is therefore the focus of the present work. We describe herein four important findings, (i) *C. elegans* contains functional CSE, CBS, and 3-MST enzyme orthologues and substrates thereof and synthesize H<sub>2</sub>S, (ii) genetic deficiency of *mpst-1* negatively affects health and lifespan and reduces endogenous H<sub>2</sub>S production, (iii) GYY4137, which releases low, physiologically relevant, concentrations of H<sub>2</sub>S over a period of several weeks, prolongs lifespan, and retards

age-dependent deterioration in physiological functions, and (iv) the effect of H<sub>2</sub>S deficiency (in *mpst-1* knockouts) and excess (after GYY4137 treatment) are likely due to the modulation of ROS-mediated damage by H<sub>2</sub>S.

To date, a total of 11 orthologues of CSE, CBS, and 3-MST have been predicted to occur in the *C. elegans* genome (Wormbase; <http://wormbase.org/>). These enzymes play important roles in regulating sulfur metabolism in this organism as they do in mammals. Whether these enzymes, like their mammalian counterparts, also synthesize H<sub>2</sub>S, has not previously been shown. CSE and CBS have been extensively studied and the biological roles of H<sub>2</sub>S generated by their activity have been well characterized in mammals. In contrast, the biological relevance of 3-MST-derived H<sub>2</sub>S is not so clear. 3-MST catalyzes the transsulfuration of 3-MP to pyruvate during degradation of L-cysteine and, in this way, can also increase the intracellular concentration of thioredoxin and reduced glutathione (GSH), both of which contribute to cellular redox homeostasis (19). The fact that MitoSOX Red and mtDNA (marker of general ROS production/activity) were elevated in *mpst-1* knockouts suggests that these animals were in a state of oxidative stress. 3-MST also synthesizes H<sub>2</sub>S in mammalian liver, kidney, and brain *in vitro* (32), and since H<sub>2</sub>S is not only a strong reducing agent and scavenger of free radicals in mammalian cells (31) but also increases intracellular GSH levels (10), it seems likely that at least part of the protective effect of 3-MST in oxidative stress in *C. elegans* is due to its ability to generate H<sub>2</sub>S. We therefore propose that (i) 3-MST deficiency reduces the lifespan in *C. elegans* because of diminished formation of endogenous antioxidant H<sub>2</sub>S, (ii)

GY4137 restores normal biological aging to these mutants by replenishing the lost H<sub>2</sub>S gas, and (iii) GYY4137 prolongs the lifespan in wild-type *C. elegans* by a similar mechanism.

Several lines of experimental evidence support these proposals. Alongside changes in free radical generation in *mpst-1* mutants, which have been noted above, we also report here that GYY4137 (i) has a potent antioxidant activity in biochemical assays *in vitro*, (ii) alters the expression of a range of genes associated with both aging and oxidation and reduction, (iii) reduces markers of oxidative stress including Mito-SOX Red and DCF-DA fluorescence, mtDNA damage, and PCs, (iv) decreases paraquat lethality, which is believed to be secondary to ROS generation (8), and (v) enhances the lifespan of *mev-1* mutants, which exhibit premature death due to overproduction of superoxide anions due to depleted complex II enzyme succinate-CoQ oxidoreductase activity (6). Taken together, these data suggest that both endogenous and exogenous H<sub>2</sub>S oppose oxidative stress through both direct and indirect mechanisms in *C. elegans* and that this likely forms the basis for the effect of this gas on lifespan and health status in this organism.

In conclusion, small amounts of endogenous H<sub>2</sub>S, generated either from 3-MP by 3-MST enzyme activity or by spontaneous decomposition of GYY4137, prolongs the lifespan and delays the adverse health consequences of aging in *C. elegans* by an antioxidant effect. Whether endogenous H<sub>2</sub>S also prolongs the lifespan in mammals is a key question, which has yet to be directly addressed. However, circumstantial evidence, such as the identification of increased aortic CSE and CBS expression in aged rats (23) and reduced plasma H<sub>2</sub>S concentration in humans older than 50 years (3), suggests that H<sub>2</sub>S may play a part in physiological changes in aging mammals as well. Moreover, while the treatment with H<sub>2</sub>S gas is clearly not a viable H<sub>2</sub>S delivery strategy in the clinic, the water-soluble H<sub>2</sub>S-releasing drug, GYY4137, seems well suited, at least as a prototype, for the development of a new class of drugs with beneficial effects on healthy aging.

## Materials and Methods

### Use of *C. elegans*

All *C. elegans* strains were obtained from the *Caenorhabditis* Genetic Centre (University of Minnesota) and cultured on nematode growth medium (NGM, 3 g/L NaCl, 17 g/L bacto-agar, 2.5 g/L bacto-peptone, 5 mg/L cholesterol, 25 mM KPO<sub>4</sub> [pH 6], 1 mM MgSO<sub>4</sub>, and 1 mM CaCl<sub>2</sub>) seeded with *Escherichia coli* OP50 (OD<sub>595</sub> 3.4–3.6) at 20°C as described elsewhere (7). *C. elegans* strains used in this study were Bristol N2 (wild type), *mev-1(kn1)*, and *mpst-1(ok2040)*. The *mpst-1* mutant strain was backcrossed to the wild type four times and single worm genotyping carried out using standard procedures. Only homozygous mutant strains were used. For some experiments, RNAi-mediated silencing *via* double-stranded RNA (dsRNA) was used to knock down *C. elegans* genes that are orthologous to the human H<sub>2</sub>S-synthesizing enzymes, CSE and 3-MST. Genes targeted (and their enzyme product) were *cth-2* ZK1127.10 (CSE) and *mpst-1* D2023.5 (3-MST). *E. coli* strain HT115 (DE3) expressing dsRNA of the gene of interest (*cth-2* or *mpst-1* cloned into pPD129.36) was fed to *C. elegans* maintained on NGM plates containing isopropyl β-D-1-thiogalactopyranoside and ampicillin. RNAi controls were fed HT115 (DE3) bacteria containing the empty pPD129.36 vector.

### *C. elegans* synchronization and drug treatment

Adult wild-type and mutant animals were washed from NGM plates with M9 buffer (22 mM KH<sub>2</sub>PO<sub>4</sub>, 42.2 mM Na<sub>2</sub>HPO<sub>4</sub>, 85.5 mM NaCl, and 1 mM MgSO<sub>4</sub> · 7H<sub>2</sub>O) and centrifuged (2500g, 2 min). The resulting pellets were washed with M9 buffer and digested with hypochlorite solution (1.0%–1.3% w/v sodium hypochlorite, 500 mM sodium hydroxide). Samples were mixed by vigorous shaking (2.5 min), centrifuged (2500g, 1 min), and the supernatant was discarded. Thereafter, the pellets were resuspended in M9 buffer, centrifuged (2500g, 1 min), and the wash procedure was repeated four times after which 10 volumes of M9 buffer was added to the final egg pellets, and the resulting suspension gently rotated overnight at room temperature to allow hatching into the first larval stage (L1) animals. Synchronized L1 animals were cultured (48 h) in sealed tubes in the medium containing M9 buffer and *E. coli* OP50 (1:2 v/v) in the presence or absence of GYY4137 (100 μM) or water (vehicle control) at room temperature. During this time, animals developed to the fourth larval stage (L4).

### Determination of the effect of drugs on aging and life traits of *C. elegans*

The effect of drugs on lifespan of *C. elegans* was assessed at 20°C using L4 worms as detailed previously (7). Lifespan was monitored daily, and death was scored by failure of the animal to move in response to gentle prodding with a platinum wire. Drug-induced changes in body size (length and area) of *C. elegans* were observed using a digital camera (×1.5 magnification) attached to a microscope (Nikon UK Ltd.). The volumetric area within the perimeter of the nematodes was determined using Image-Pro Express v5.1 software (Media Cybernetics). Additional age-related parameters, including head thrashing, pharyngeal pumping, and defecation rates, were assessed by manual counting using a fully apochromatic corrected stereomicroscope (Leica M205 with FusionOptics™ with a zoom capacity of 20.5:1 and a resolution of 1050 lp/mm).

A further index of age-related deterioration in physiological capacity (*i.e.*, healthspan), based on measurement of the progressive decline in locomotor function with aging was used (20, 27). Briefly, *C. elegans* (up to 200 animals) were observed for locomotor activity at daily intervals and subdivided into three groups: class A animals were healthy, showed spontaneous movements, and were highly mobile; class B animals showed movement only after prodding; and class C nematode movement was restricted to the head and/or tail upon prodding with a platinum wire. Only A type nematodes were counted as healthy. Healthspan curves were determined by scoring the percentage of worms in the synchronized aging cohort that still remain in the healthy category (A-type) as a function of age. Data are expressed as % healthy (*i.e.*, category A).

### Survival assay in the presence of paraquat

Paraquat (1,1'-dimethyl-4,4'-bipyridinium dichloride) resistance was determined by transferring L4 worms previously treated (48 h) with either GYY4137 (100 μM) or water (vehicle) to plates containing paraquat (10 mM) with/without GYY4137 (100 μM) and scored as described above for lifespan determination.

### Quantitative real-time polymerase chain reaction

*C. elegans* were homogenized in the presence of 0.5-mm glass beads to maximize tissue/cell breakage and total RNA extracted according to the manufacturer's instructions using Tri-reagent (Sigma). Total RNA (0.5–1  $\mu$ g) was reverse-transcribed to cDNA using M-MLV reverse transcriptase enzyme (Promega), and real-time qPCR was performed with primers specific for each enzyme. A list of primers used is provided in Supplementary Table S3.

### Microarray analysis

*C. elegans* (minimum, 5000 animals) were incubated with GYY4137 (100  $\mu$ M) or water vehicle from L1 to L4. Three separate batches of animals were prepared for each treatment. *C. elegans* were homogenized, and total RNA (~20  $\mu$ g) was extracted according to the manufacturer's instructions using Trizol reagent (Invitrogen) and purified using an RNeasy kit followed by a DNase digestion kit (Qiagen). RNA was not amplified before labeling. Synthesis of double-stranded cDNA and biotin-labeled cRNA was performed according to the manufacturer's instruction (Affymetrix). Fragmented cRNA preparations were hybridized to *C. elegans* genome oligonucleotide arrays using Affymetrix hybridization Oven 640, washed, and scanned on a GeneChip Scanner 3000 (Affymetrix). Initial data extraction and normalization within each array was performed using Affymetrix GCOS software. Data intensities were log transformed and normalized within, and between, arrays using the quantile normalization method, and two-tail, pair-wise analysis or a two-way analysis of variance was employed using Spotfire Decision Site software package 7.2 v10.0 (Spotfire, Inc.) to extract the statistically significant data from each group of animals. Each treatment replicate was analyzed twice (technical repeats). Thereafter, the log<sub>2</sub> values (GYY4137-treated net intensity/normalized control net intensity) from each replicate were averaged. A gene was considered for analysis if the hybridization signal for the corresponding probe was deemed to be present on the array (significantly greater than background level,  $p < 0.05$ ) in at least two of three relevant arrays in the drug-treated experiment. A gene was considered induced/repressed if the difference in expression levels was significant ( $p < 0.05$ ) compared to the control using Student's *t* test and Significance Analysis of Microarrays analysis. Gene enrichment analysis was performed using DAVID 6.7 to obtain a ranking of functional categories based on co-occurrence with sets of genes in the gene list ( $p < 0.05$ ). Genes were annotated using Wormbase (<http://wormbase.org/>). The MIAME accession number for the microarray data reported in this article is GSE16975.

### Assay of antioxidant effect of GYY4137

The antioxidant effect of GYY4137 (and NaHS for comparison) *in vitro* was assessed using two standard assays. The 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS) assay was carried out as described elsewhere (25), and data are expressed as the Trolox equivalent antioxidant capacity. The reducing power of GYY4137 and NaHS was determined using the potassium ferricyanide–ferric chloride method (21). ROS formation in *C. elegans* was measured using MitoSOX Red mitochondrial superoxide indicator (Molecular Probes) as described elsewhere (13) with modifications. For this assay,

~150 wild-type animals were treated with GYY4137 (100  $\mu$ M) or vehicle in agar plates for 24 h before being manually transferred into each well of a black 96-well microtiter plate containing M9 buffer (100  $\mu$ l). Similar experiments were carried out using *mpst-1* knockouts. MitoSOX red reagent (100  $\mu$ l of 20  $\mu$ M solution) was added to each well, and superoxide-associated fluorescence was assessed kinetically (every 2 min for 5 h, room temperature) with a Synergy H1 multimode microplate reader (BioTek Instruments) using excitation and emission frequencies of 510 and 580 nm, respectively. All samples were normalized to bacterial control in separate wells on the same plate. In some experiments, ROS formation was also determined using DCF-DA as described previously (5). To measure sequence-specific mitochondrial DNA (mtDNA) damage, a minimum of 10,000 animals were washed three times on ice with S basal buffer (100.1 mM NaCl, 5.7 mM K<sub>2</sub>HPO<sub>4</sub>, 44.1 mM KH<sub>2</sub>PO<sub>4</sub>, 0.01 mM cholesterol) and then twice with isolation buffer (210 mM mannitol, 70 mM sucrose, 0.1 mM EDTA, 5 mM Tris-HCl, pH 7.4). Animals were collected by centrifugation and the resulting *C. elegans* pellets were homogenized in isolation buffer using a Teflon-glass homogenizer. Debris and nuclei were removed from the homogenate by centrifugation (600g, 4°C, 10 min), and the supernatant containing mitochondria was further centrifuged (7200g, 4°C, 10 min) to obtain a mitochondrial pellet, which was then resuspended in Tris-EDTA buffer (50 mM Tris-HCl, 0.1 mM EDTA, pH 7.4). DNA from crude mitochondria was purified using Prepman Ultra Sample Preparation Reagent (Applied Biosystems), according to the manufacturer's protocol. Quantitative PCR was performed using the GeneAMP XL PCR kit (Applied Biosystems) based on a protocol described elsewhere (17). mtDNA damage was assessed by comparing amplification efficiency in a 6.3 kb region of the mitochondrial genome (17) to a 71 bp region (forward primer: 5'-GAGCGTCATTTATTGGGAAGAAGA-3' and reverse primer: 5'-TGTGCTAATCCCATAAATGTAACCTT-3'). PC content was measured as described elsewhere (27). Briefly, about 200 wild-type or *mpst-1* knockout worms exposed to GYY4137 (100  $\mu$ M) on agar plates for 24 h were collected, washed to remove residual bacteria, and resuspended in phosphate-buffered saline containing tween 20 (0.1% v/v) and phenylmethylsulfonyl fluoride (1 mM). Samples were then sonicated, and protein concentration of the lysates was determined by the Dc-Protein Assay (Bio-Rad). Worm lysates (5  $\mu$ l, 2  $\mu$ g) were mixed with 12% v/v SDS (5  $\mu$ l) and 2,4-dinitrophenylhydrazine (DNPH, 20 mM, 10  $\mu$ l) for 15 min at room temperature and neutralized with Tris-Cl buffer (2 mM containing 30% v/v glycerol, 75  $\mu$ l). Slot blot was conducted (1 h, room temperature) using anti-DNPH antibody (1:150; Chemicon International) with secondary detection using horseradish peroxidase (HRP)-conjugated anti-rabbit IgG antibody (1:300). All chemicals and antibodies for this assay were from the Oxyblot Protein Oxidation Detection Kit (Chemicon International). Antibody-bound proteins were detected by chemiluminescence using a Chemidoc XRS imaging system (Bio-Rad).

### Immunohistochemical detection of H<sub>2</sub>S-synthesizing enzymes

*C. elegans* were fixed with 4% v/v paraformaldehyde in phosphate buffer at 4°C and washed with PBS. Whole mount or cryo-sections (10  $\mu$ m) of the worms were then washed with

PBS and blocked in 5% v/v normal rabbit or mouse serum for 1 h at room temperature. Sections were then incubated overnight with the following primary antibodies: rabbit anti-CBS (1:200; Abcam), rabbit anti-3-MST (1:200; Abcam), and mouse anti-CSE (1:250; Abcam) in PBS containing 0.1% v/v Triton X-100 (PBS-TX). Sections were washed in PBS and incubated in anti-rabbit IgG conjugated to FITC (1:200; Sigma) or anti-mouse IgG conjugated to Cy3 (1:200; Sigma) both diluted in PBS-TX for 1 h at room temperature and then counterstained with 4',6-diamidino-2-phenylindole (DAPI). 3-MST was also stained in *mpst-1* mutants. Control sections were also incubated as described above but without the primary antibodies and showed no significant staining. Photomages were captured in a fluorescence microscope equipped with a digital camera (Olympus BX51; Olympus).

#### Measurement of H<sub>2</sub>S release from GYY4137 and H<sub>2</sub>S-synthesizing activity

GYY4137 (100  $\mu$ M) was incubated in buffer solution in capped glass vials and incubated at 20°C. Measurements were conducted on days 0–7 and 30. The concentration of H<sub>2</sub>S was determined using the fluorescent probe, dansyl azide (0.6 mM) (22), with excitation and emission wavelengths of 344 and 533 nm, respectively. Enzymatic H<sub>2</sub>S synthesis by *C. elegans* homogenate was measured as described (14) with the following modifications: 430  $\mu$ l supernatant derived from a homogenate of at least 4000 adult (day 5) worms was incubated with PPP (2 mM) and substrate L-cysteine (10 mM) and sealed with a double parafilm layer to avoid leakage of H<sub>2</sub>S gas generated after incubation at 37°C for 30 min. In experiments to measure the 3-MST activity, homogenates were incubated with 3-MP (1 mM) instead of L-cysteine and PPP was omitted. Baseline controls contained trichloroacetic acid (TCA; 10% w/v, 250  $\mu$ l) prepared in parallel. At the end of the incubation period, zinc acetate (1% w/v, 250  $\mu$ l) was injected to trap the H<sub>2</sub>S followed by TCA (10% w/v, 250  $\mu$ l) to terminate the reaction. Subsequently, *N,N*-dimethyl-*p*-phenylenediamine sulfate dye (NNDPD; 20 mM) in 7.2 M HCl was added, followed by the addition of FeCl<sub>3</sub> (30 mM) in 1.2 M HCl. After centrifugation (14,000 rpm, 4 min, 4°C), absorbance (670 nm) of the resulting methylene blue in the supernatant was measured using a 96-well microplate reader (Tecan Systems) and compared against a standard curve of NaHS. Results are expressed as nmol H<sub>2</sub>S formed per mg protein per min. Protein concentration in each homogenate was estimated according to the manufacturer's instructions using the Bradford assay (Bio-Rad). The presence of H<sub>2</sub>S, cysteine, and 3-MP in homogenates of *C. elegans* was determined after derivatizing with monobromobimane, using a reverse-phase high-pressure liquid chromatography with fluorescence detection as described elsewhere (28). In brief, *C. elegans* (~1000 animals) were homogenized and 15  $\mu$ l homogenate incubated with Tris-HCl buffer (70  $\mu$ l, 100 mM, pH 9.5, 0.1 mM diethylene triamine pentaacetic acid) and monobromobimane solution (50  $\mu$ l, 10 mM in acetonitrile). The reaction was stopped after 30 min by adding ice-cold sulfosalicylic acid (50  $\mu$ l, 200 mM), and samples were vortexed and placed on ice before rp-hplc as described elsewhere (28). Detection of thiols was assessed by measurement of the fluorescent sulfide-dibimane derivatives (excitation at 390 nm/emission at 475 nm). The retention times of authentic cysteine, 3-MP, and H<sub>2</sub>S were 4.1  $\pm$  0.02, 14.5  $\pm$  0.01, and 24.4  $\pm$  0.01 min (all *n* = 3), respectively.

#### Statistical analysis

Data show mean  $\pm$  SEM. Multiple comparisons were made using one-way ANOVA followed by *post hoc* Tukey test. The difference between lifespan and healthspan survival curves was calculated using the log-rank (Mantel-Cox) test. Statistical significance was set at *p* < 0.05.

#### Acknowledgments

This work was supported by the King's College London (S.R.S.), the National University of Singapore (P.K.M.), and a grant (MOE2012-T2-2-003) from the Ministry of Education, Singapore (P.K.M., J.G., B.H.). B.Q. was supported by a State of Kuwait funded studentship.

#### Authors' Contribution

SRS, P.K.M., J.G., B.H., and L.L. designed the research; B.Q. and L.L. also performed research, analyzed data, and contributed to the writing of the article; M.T.P., L.F.N., S.D.K., P.R., and F.W. performed research; C-H.T. and B.W.D. contributed new reagents/analytic tools; SRS and SCS analyzed microarray data; J.G., SRS and P.K.M. analyzed data and wrote the article.

#### Author Disclosure Statement

The authors declare no conflicts of interest.

#### References

1. Back P, Braeckman BP, and Matthijssens F. ROS in aging *Caenorhabditis elegans*: damage or signaling? *Oxid Med Cell Longev*, 2012. [Epub ahead of print]; DOI: 10.1155/2012/608478.
2. Budde MW and Roth MB. The response of *Caenorhabditis elegans* to hydrogen sulfide and hydrogen cyanide. *Genetics* 189: 521–532, 2011.
3. Chen YH, Yao WZ, Geng B, Ding YL, Lu M, Zhao MW, and Tang CS. Endogenous hydrogen sulfide in patients with COPD. *Chest* 128: 3205–3211, 2005.
4. Cooper CE and Brown GC. The inhibition of mitochondrial cytochrome oxidase by the gases carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide: chemical mechanism and physiological significance. *J Bioenerg Biomembr* 40: 533–539, 2008.
5. Gruber J, Ng LF, Fong S, Wong YT, Koh SA, Chen CB, Shui G, Gheong FW, Schaffe S, Wenk MR, and Halliwell B. Mitochondrial changes in ageing *caenorhabditis elegans*—what do we learn from superoxide dismutase knockouts? *PLoS One* 6: e19444, 2011.
6. Hosokawa H, Ishii N, Ishida H, Ichimori K, Nakazawa H, and Suzuki K. Rapid accumulation of fluorescent material with aging in an oxygen-sensitive mutant *mev-1* of *Caenorhabditis elegans*. *Mech Ageing Dev* 74: 161–170, 1994.
7. Hughes S and Stürzenbaum SR. Single and double metallothionein knockout in the nematode *C. elegans* reveals cadmium dependent and independent toxic effects on life history traits. *Environ Pollut* 145: 395–340, 2007.
8. Kearney M, Matthijssens F, Sharpe M, Vanfleteren J, and Gems D. Superoxide dismutase mimetics elevate superoxide dismutase activity *in vivo* but do not retard aging in the nematode *Caenorhabditis elegans*. *Free Radic Biol Med* 37: 239–250, 2004.
9. Kenyon C. A pathway that links reproductive status to lifespan in *Caenorhabditis elegans*. *Ann N Y Acad Sci*. 1204: 156–162, 2010.

10. Kimura Y, Goto Y, and Kimura H. Hydrogen sulfide increases glutathione production and suppresses oxidative stress in mitochondria. *Antioxid Redox Signal* 12: 1–13, 2010.
11. Kimura H, Shibuya N, and Kimura Y. Hydrogen sulfide is a signaling molecule and a cytoprotectant. *Antioxid Redox Signal* 17: 45–57, 2012.
12. King AL and Lefer DJ. Cytoprotective actions of hydrogen sulfide in ischaemia-reperfusion injury. *Exp Physiol* 96: 840–846, 2011.
13. Kulich SM, Horbinski C, Patel M, and Chu CT. 6-Hydroxydopamine induces mitochondrial ERK activation. *Free Radic Biol Med* 43: 372–383, 2007.
14. Li L, Bhatia M, Zhu YZ, Zhu YC, Ramnath RD, Wang ZJ, Anuar FB, Whiteman M, Salto-Tellez M, and Moore PK. Hydrogen sulfide is a novel mediator of endotoxic shock. *FASEB J* 19: 1196–1198, 2005.
15. Li L, Whiteman M, Guan YY, Neo KL, Cheng Y, Lee SW, Zhao Y, Baskar R, Tan CH, and Moore PK. Characterisation of a novel, water soluble hydrogen sulfide releasing molecule (GYY4137): new insights into the biology of hydrogen sulphide. *Circulation* 117: 2351–2360, 2008.
16. Mathew ND, Schlipalius DI, and Ebert PR. Sulfurous gases as biological messengers and toxins: comparative genetics of their metabolism in model organisms. *J Toxicol* 2011: 394970, 2011.
17. Melov S, Lithgow GJ, Fischer DR, Tedesco PM, and Johnson TE. Increased frequency of deletions in the mitochondrial genome with age of *Caenorhabditis elegans*. *Nucleic Acids Res* 23: 1419–1425, 1995.
18. Miller DL and Roth MB. Hydrogen sulfide increases thermotolerance and lifespan in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* 104: 20618–206122, 2007.
19. Nagahara N and Katayama A. Post-translational regulation of mercaptopyruvate sulfurtransferase via a low redox potential cysteine sulfenate in the maintenance of redox homeostasis. *J Biol Chem* 280: 34569–34576, 2005.
20. Onken B and Driscoll M. Metformin induces a dietary restriction-like state and the oxidative stress response to extend *C. elegans* Healthspan via AMPK, LKB1, and SKN-1. *PLoS One* 5: e8758, 2010.
21. Oyaizu M. Studies on products of browning reaction: anti-oxidative activity of products of browning reaction prepared from glucosamine. *Jap J Nutr* 44: 307–315, 1986.
22. Peng H, Cheng Y, Dai C, King AL, Predmore BL, Lefer DJ, and Wang B. A fluorescent probe for fast and quantitative detection of hydrogen sulfide in blood. *Angew Chem Int Ed Engl* 50: 9672–9675, 2011.
23. Predmore BL, Alendy MJ, Ahmed KI, Leeuwenburgh C, and Julian D. The hydrogen sulfide signaling system: changes during aging and the benefits of caloric restriction. *Age (Dordr)* 32: 467–481, 2010.
24. Pun PB, Gruber J, Tang SY, Schaffer S, Ong RL, Fong S, Ng LF, Cheah I, and Halliwell B. Ageing in nematodes: do antioxidants extend lifespan in *Caenorhabditis elegans*? *Biogerontology* 11: 17–30, 2010.
25. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, and Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 26: 1231–1237, 1999.
26. Robida-Stubbs S, Glover-Cutter K, Lamming DW, Mizunuma M, Narasimhan SD, Neumann-Haefelin E, Sabatini DM, and Blackwell TK. TOR signaling and rapamycin influence longevity by regulating SKN-1/Nrf and DAF-16/FoxO. *Cell Metab* 15: 713–724, 2012.
27. Schaffer S, Gruber J, Ng LF, Fong S, Wong YT, Tang SY, and Halliwell B. The effect of dichloroacetate on health- and lifespan in *C. elegans*. *Biogerontology* 12: 195–209, 2011.
28. Shen X, Pattillo CB, Pardue S, Bir SC, Wang R, and Kevil CG. Measurement of plasma hydrogen sulfide *in vivo* and *in vitro*. *Free Radic Biol Med* 50: 1021–1031, 2011.
29. Viswanathan M, Kim SK, Berdichevsky A, and Guarente L. A role for SIR2.1. Regulation of ER stress response genes in determining *C. elegans* life span. *Dev Cell* 9: 605–615, 2005.
30. Vozdek R, Hnízda A, Krijt J, Kostrouchová M, and Kožich V. Novel structural arrangement of nematode cystathionine  $\beta$ -synthases: characterization of *Caenorhabditis elegans* CBS-1. *Biochem J* 443: 535–547, 2012.
31. Whiteman M, Armstrong JS, Chu SH, Jia-Ling S, Wong BS, Cheung NS, Halliwell B, and Moore PK. The novel neuro-modulator hydrogen sulfide: an endogenous peroxynitrite ‘scavenger’? *J Neurochem* 90: 765–768, 2004.
32. Wróbel M, Jurkowska H, Sliwa L, and Srebro Z. Sulfurtransferases and cyanide detoxification in mouse liver, kidney, and brain. *Toxicol Mech Methods* 14: 331–337, 2004.

Address correspondence to:  
 Dr. Stephen R. Stürzenbaum  
 School of Biomedical Science  
 King's College London  
 Franklin Wilkins Building  
 150, Stamford Street  
 London SE1 9NH  
 United Kingdom

E-mail: stephen.sturzenbaum@kcl.ac.uk.

Dr. Philip K. Moore  
 Department of Pharmacology  
 Yong Loo Lin School of Medicine  
 National University of Singapore  
 Singapore 117597  
 Singapore

E-mail: dprmpk@nus.edu.sg.

Date of first submission to ARS Central, May 25, 2013; date of final revised submission, September 10, 2013; date of acceptance, October 5, 2013.

#### Abbreviations Used

3-MP = 3-mercaptopyruvate
3-MST = 3-mercaptopyruvate transferase
CBS = cystathionine $\beta$ synthetase
CSE = cystathionine $\gamma$ lyase
DAB = diaminobenzidine
DCF = 2,7-dichlorofluorescein
DCF-DA = 2,7-dichlorofluorescein diacetate
GO = gene ontology
GSH = glutathione
H <sub>2</sub> S = hydrogen sulfide
NGM = nematode growth medium
PPP = pyridoxal 5' phosphate
ROS = reactive oxygen species
TCA = trichloroacetic acid