Delivery system designs for platinum prodrug in cancer treatment

Lim, Wei Qi

2019

Lim, W. Q. (2019). Delivery system designs for platinum prodrug in cancer treatment.
Doctoral thesis, Nanyang Technological University, Singapore.

<https://hdl.handle.net/10356/104277>

<https://doi.org/10.32657/10356/104277>

**NANYANG
TECHNOLOGICAL
UNIVERSITY**

SINGAPORE

**DELIVERY SYSTEM DESIGNS FOR PLATINUM
PRODRUG IN CANCER TREATMENT**

LIM WEI QI
Interdisciplinary Graduate School
NTU- Northwestern Institute for Nanomedicine

DELIVERY SYSTEM DESIGNS FOR PLATINUM PRODRUG IN CANCER TREATMENT

LIM WEI QI

**Interdisciplinary Graduate School
NTU- Northwestern Institute for Nanomedicine**

A thesis submitted to the Nanyang Technological University in
partial fulfillment of the requirement for the degree of
Doctor of Philosophy

2019

Statement of Originality

I hereby certify that the work embodied in this thesis is the result of original research, is free of plagiarised materials, and has not been submitted for a higher degree to any other University or Institution.

10 July 2019

.....

Date

.....

Lim Wei Qi

Supervisor Declaration Statement

I have reviewed the content and presentation style of this thesis and declare it is free of plagiarism and of sufficient grammatical clarity to be examined. To the best of my knowledge, the research and writing are those of the candidate except as acknowledged in the Author Attribution Statement. I confirm that the investigations were conducted in accord with the ethics policies and integrity standards of Nanyang Technological University and that the research data are presented honestly and without prejudice.

19 June 2019

.....

Date

.....

Professor Zhao Yanli

Authorship Attribution Statement

This thesis contains material from two papers published in the following peer-reviewed journal(s) where I was the first and/or corresponding author.

Chapter 2 is published as Lim, W. Q.; Phua, S. Z. F.; Chen, H.; Zhao, Y., An oxaliplatin(IV) prodrug-based supramolecular self-delivery nanocarrier for targeted colorectal cancer treatment. *Chem. Commun.* **2018**, 54 (90), 12762-12765. DOI: 10.1039/C8CC07858K

The contributions of the co-authors are as follows:

- Prof Zhao and Dr Chen Hongzhong provided the initial research direction.
- I drafted the manuscript, which was revised by Prof Zhao and Ms Phua Soo Zeng, Fiona.
- I performed all the material synthesis and characterizations, in-vitro experiments and data analysis, with the assistance of Ms Phua Soo Zeng.
- Ms Phua Soo Zeng, Fiona carried out compounds' characterization by high resolution mass spectroscopy.

Chapter 3 is published as Lim, W. Q.; Yang, G.; Phua, S. Z. F.; Chen, H.; Zhao, Y., Self-assembled Oxaliplatin(IV) Prodrug-Porphyrin Conjugate for Combinational Photodynamic Therapy and Chemotherapy. *ACS Appl. Mater. Interfaces* **2019**, 11 (18), 16391-16401. DOI: 10.1021/acsami.9b04557

The contributions of the co-authors are as follows:

- Prof Zhao suggested the project direction and edited the manuscript drafts.
- I drafted the manuscript which was revised by Ms Phua Soo Zeng, Fiona.
- I performed all the materials synthesis, purifications and characterizations. I also carried out the in-vitro experiments and conducted data analysis.

- Dr. Chen Hongzhong suggested the compounds synthesis procedures and provided starting materials for the synthesis.
- Dr Yang Guangbao planned the in-vivo experiments and built the animal models. I carried out the in vivo experiments. Ms Phua Soo Zeng Fiona helped with the histological experiments.

Chapter 4 is published as Lim, W. Q.; Phua, S. Z. F.; Zhao, Y., Redox-Responsive Polymeric Nanocomplex for Delivery of Cytotoxic Protein and Chemotherapeutics. *ACS Appl. Mater. Interfaces* **2019**, *11* (35), 31638-31648. DOI: 10.1021/acsami.9b09605

The contributions of the co-authors are as follows:

- Prof Zhao refined the project idea and edited the manuscript drafts.
- I drafted the manuscript which was revised by Ms Phua Soo Zeng, Fiona.
- I performed all the materials synthesis, purifications and characterizations. I also carried out the in-vitro experiments and conducted data analysis.

10 July 2019

.....
Date

.....
Lim Wei Qi

Acknowledgements

I would like to express my sincere gratitude to numerous people who have given me support and help during my four years of Ph.D. studies.

First and foremost, I am especially grateful to my supervisor, Professor Zhao Yanli for his constant guidance, invaluable support and the opportunity to pursue research. He placed his trust in me and gave me freedom to explore projects that I am interested in. He always provides invaluable advices for my research and suggestions to improve my academic writing. Under his supervision, I have gained a lot of skills such as research skills and critical thinking, which allow me to become more confident and resilient.

Also, the time, precious advices and helpful discussions which my thesis advisory committee members Professor Amy Paller and Assistant Professor Chen Gang had provided in my research is greatly appreciated. I am thankful to my mentors, Dr Luo Zhong and Dr Chen Hongzhong who patiently guided me and shared their research knowledge and experiences with me. I would also like to thank the past and present members of the Zhao's group; Dr Nguyen Kim Truc, Dr Qu Qiuyu, Dr Zhao Lingzhi, Dr Sreejith Sivaraapanicker, Dr Yang Guangbao, Dr Wang Dongdong, Dr Xiang Huijing, Dr Zhang Yuanyuan, Dr Li Menghuan, Dr Fengtao, Dr Tan Si Yu, Dr Ang Chung Yen, Dr Yang Chaolong, Dr Xing Pengyao, Dr Quan Hongping for their help in one way or another.

I am also fortunate to have wonderful friends and lab members- Dr Wang Yang, Ms Bindra Anivind Kaur, Dr Nakayama Masanari, Dr Phoebe Tham Huijun, Dr Zheng Chunchuan, Mr Eddy Wong Mun Fei, Mr Deblin Jana, Mr Ong Wee Kong, Mr Toh Wei Liang and Mr Victor Xu Hesheng who made my time in this research group memorable. Special mentions to Ms Phua Soo Zeng Fiona who not only provided valuable

suggestions and research assistance but gave me emotional support and words of encouragement in these four years. Also, to Mr Andreas Dickescheid, whose friendship and assistance in HPLC helped my research. Additionally, special thanks to Dr Wong Yee Shan for the administrative support and answering of all my queries.

Importantly, I want to express my sincere appreciations to my beloved family members, whose unwavering support and understanding allowed me to focus on my research. Their unconditional love and sacrifices helped me persevere through this long and rewarding journey.

Lastly, I would like to thank Interdisciplinary Graduate School and Nanyang Technological University for their financial support and academic opportunity to complete my research studies in a highly regarded academic environment.

Thank you all!

Table of Contents

Acknowledgements	I
Table of Contents	III
Summary	IV
Chapter 1 Introduction	6
1.1 Cancer and the different treatment modalities	6
1.2 Combination therapy	8
1.3 Nanomedicine	8
1.4 Supramolecular chemistry	18
1.5 Prodrugs	21
1.6 Platinum-based drugs	26
1.7 Platinum(IV) prodrugs-based nano-delivery system for cancer therapy	35
1.8 Research gaps and objectives	40
Chapter 2 An oxaliplatin(IV) prodrug-based supramolecular self-delivery nanocarrier for targeted colorectal cancer treatment	43
2.1 Introduction	43
2.2 Experimental Section	46
2.3 Results and Discussion	54
2.4 Conclusion	65
Chapter 3 Self-assembled Oxaliplatin(IV) Prodrug-Porphyrin Conjugate for Combinational Photodynamic Therapy and Chemotherapy	66
3.1 Introduction	66
3.2 Experimental Section	71
3.3 Results and Discussion	83
3.4 Conclusion	98
Chapter 4 Redox-responsive polymeric nanocomplex for delivery of cytotoxic protein & chemotherapeutics	99
4.1 Introduction	99
4.2 Experimental Section	103
4.3 Results and Discussion	116
4.4 Conclusion	138
Chapter 5 Conclusion and Future Outlook	139
List of publications	142
References	144

Summary

Platinum-based drugs are among the most powerful and widely used chemotherapy drugs. However, it is challenging to selectively deliver and accumulate these therapeutic agents in the targeted pathological regions and avoid off-target toxicity to healthy tissues. Recently, developing novel platinum prodrugs to overcome the aforementioned challenges and facilitate tumor-activated cytotoxicity has been a favored strategy. Particularly, delivering the drugs in nanostructures rather than their molecular form takes advantage of the enhanced permeability and retention effect for selective tumor accumulation and improves pharmacokinetics. Therefore, there is increasing attention on exploring the construction of platinum prodrug-based nanoparticles for effective cancer treatment. Adapting from natural phenomena of amphiphilic self-assembly, this dissertation highlights the different approaches to fabricate platinum(IV) prodrug self-assembled delivery systems and its combination with other therapeutic agents to enhance treatment efficacy *in vitro* and *in vivo*.

Firstly, Chapter 1 includes a literature review of nanomedicine, with the focus on platinum-based chemotherapy drugs. The current trends and limitations of nanotechnology in cancer treatment is first introduced, and the strategies for drug self-delivery systems is highlighted. In the second part of this chapter, an overview of the recent development in platinum drug delivery system is illustrated with recent literature examples. At the end of this chapter, we discuss the research gaps and present the outline of this dissertation.

Following which, in Chapter 2, we demonstrate how established amide coupling chemistry was harnessed to conjugate relevant guest moiety to a platinum(IV) prodrug scaffold. With complementary binding to a host molecule tethered with an active targeting ligand to form a supramolecular complex, the amphiphilic self-assembled

nanoparticles enhanced targeting specificity towards cancer cells and achieved controlled drug release in vitro.

To maximize treatment efficacy, combination therapy which promotes synergistic anticancer effects has been preferred over single chemotherapy. Hence, in Chapter 3, a supramolecular self-assembled nanoconjugate based on a redox-responsive oxaliplatin prodrug and a light-activatable photosensitizer for combined chemotherapy-photodynamic therapy is reported. Importantly, involvement of host-guest chemistry in the design enabled precise control over the therapeutics loading ratio, ensuring that the anticancer drugs combination acted synergistically in vitro and in vivo.

Lastly, we explored the use of a polymeric oxaliplatin prodrug to fabricate self-assembled nanocomplexes as therapeutic proteins delivery vehicles. Challenges of delivering therapeutic proteins intracellularly limit their clinical applications. In Chapter 4, we describe how therapeutic proteins and oxaliplatin could be modified to carry complementary electrostatic charges and mixed together to form nanostructures. The sophisticated delivery system released the therapeutic cargoes in response to reducing cellular environment and activated the caged protein and prodrug to induce efficient apoptosis.

Using nanotechnology and the elegance of self-assembly, these proposed delivery systems investigated in this dissertation provides an inspiring outlook for use of prodrug-based nanocarrier in cancer therapy.

Chapter 1

Introduction

1.1 Cancer and the different treatment modalities

Cancer is a broad term describing the abnormal and uncontrollable cell growth. The possibility of cancer to invade and metastasize to other parts of body makes it an intimidating disease. Depending on the type, location and stage of the disease, it could result in life-threatening situations. Despite advancements in technology, cancer mortality remains high and is estimated to result in approximately 17 million deaths annually by 2030.¹ The worrying outlook of increasing incidents of cancer cases has spurred substantial efforts in cancer research which results in better survival rates within the last few decades.

Existing treatments for cancer patients include surgery, radiotherapy, chemotherapy and immunotherapy.² At present, surgery and radiotherapy are commonly employed for early stage tumors. Nevertheless, the efficacy of these treatment options is still limited for cases when the cancer metastasized.³ While systemic whole-body treatment such as chemotherapy might be able to address the issue of metastasis, conventional chemotherapy drugs often cause undesired effects and poor treatment efficiency owing to its non-selective distribution of the anticancer drugs in the body. This means that beside the cancer cells, anticancer drugs could be internalized by the healthy tissues. This translates to greater dosage of drugs required to inhibit tumor growth, with undesirable side-effects on other healthy cells. Another form of systematic therapy, immunotherapy which elicits own body's immune response against cancer has seen better potential of higher treatment effectiveness with lower side effects.⁴⁻⁵ However, as a relatively new technique, the long term effect of immunotherapy has yet to be studied.

Other than traditional molecular chemotherapy drugs, many proteins and peptides also possess biological activities which could act as potential therapeutics.⁶ For instance, endostatin which is a peptide inhibitor of angiogenesis is now under clinical development and has shown potential as an anticancer drug. L-asparaginase which inhibits tumor growth by degrading asparagine that is overexpressed in tumor cells, is used in the treatment of leukemia.⁷ Driven by the advances in solid-phase peptide synthesis and proteomics, large scale production of clinical-grade peptides and proteins is easily obtained and further progress on discovery of protein anticancer drugs is underway. However, the poor physiological stability due to protease degradation, rapid renal elimination, possible trigger of immunogenicity and barriers to cell translocation hampers their translation as first-line anticancer agents. Numerous approaches such as polymer conjugation, modifications with tumor cell targeting ligands and encapsulation in microreservoir delivery systems have been proposed. Nonetheless, more efforts are required on effective delivery of such potent proteins and peptides to their target sites.

More recently, phototherapies including photodynamic therapy (PDT) and photothermal therapy (PTT) have gained attraction for their high tumor eradication efficiency, desirable spatial and temporal control and, negligible side effects to healthy tissues.⁸ PDT involves the delivery of photosensitizers to target sites and upon excitation of light of a suitable wavelength, generate cytotoxic reactive oxygen species (ROS) which will oxidize cellular organelles, resulting in cell ablation.⁹ However, the reliance of endogenous oxygen levels and poor aqueous solubility of the conventional photosensitizers often hinder its efficiency in clinical applications. PTT is another light activated treatment modality which employs near-infrared photoabsorbers to generate thermal energy for eradication of tumor.¹⁰ Although PTT has shown encouraging results for complete ablation of tumor with low recurrence rate, most PTT agents are inorganic

materials or noble metal nanostructures which could pose biodegradability issues and long-term toxicity concerns.¹¹

1.2 Combination therapy

Acknowledging the limitations of each conventional treatment modality, the concept of combination therapy which is the strategy of combining two or more therapeutic agents becomes the cornerstone of cancer therapy. With the heterogeneousness of tumor environment, unpredictable cellular mechanisms and pathways which potentiate issues of drug resistance and cancer metastasis, increasing evidence show that monotherapy is not efficacious for complete cancer treatment.¹² Comparatively, the approach of combining different treatment modalities that targets separate cellular survival pathways could provide better therapeutic efficacy compared to the monotherapy.¹³ However, challenges such as optimizing dosages, controlling and maintaining proper ratio of therapeutic agents and realizing suitable drug release profiles that influence the therapy outcome have yet to be overcome.

1.3 Nanomedicine

To achieve success for the treatment of cancer, transporting the therapeutic agents to exact pathological site is vital as the mechanism of actions of these agents mainly take effect within the cells. Other than the lack of drug selectivity, traditional therapeutic drug molecules suffer from rapid renal clearance and fluctuating drug blood levels. Moreover, the repeated administration of high doses of drugs to guarantee therapeutic outcomes could result in acquired drug resistance, compromising therapeutic efficacy. The concept of drug delivery system (DDS) was hence proposed to address those issues. Therapeutic agents are commonly loaded into DDS, which helps to improve the drug solubility, enhance blood circulation and increase accumulation at

the tumor cells to avoid toxicity to the normal cells. DDSs development has now advanced to the use of targeted nanocarriers.¹⁴⁻¹⁵

Figure 1. Schematic illustration of nanoparticles selectively accumulating at tumor site via the EPR effect. Reproduced with permission.¹⁶ Copyright 2014 Ivyspring International Publisher.

Nanomaterials are developed as DDS because of their abilities to exploit the enhanced permeability and retention (EPR) effect (Figure 1).¹⁷⁻¹⁸ For normal healthy vasculature, the endothelial cells have tight junctions which prevent extravasation of nano-sized particles into normal tissues. On the other hand, to acquire more nutrients and metabolites for its aberrant growth, solid tumor present an irregular and leaky vasculature with ineffective lymphatic drainage. The high permeability of the tumor tissues provides an opportunity for nanoparticulate of certain sizes to pass through and accumulate discriminately in tumor cells. Owing to the peculiar heterogeneity of the tumor vasculature and patient characteristics, there is no clear-cut size limit of the nanoparticles. Generally, nanoparticles with sizes 7-400 nm are recommended for the accumulation in tumor tissues via EPR effect.

1.3.1 Classification of nanomaterials

At present, an extensive list of materials has been used in the fabrication of nano-sized DDS (Figure 2) and are classified into few broad categories: organic, biological and inorganic-based nanoparticles. Organic nanoparticles include polymers,¹⁹ dendrimers,²⁰ micelles²¹ and liposomes.²² These classes of nanomaterials are biodegradable and biocompatible which greatly facilitate their advances for clinical translations. To date, most nanoparticle-based DDS permitted by U.S. Food and Drug Administration (FDA) utilize organic materials. One example is Doxil, which is the first FDA-approved nanoformulation that involves encapsulation of doxorubicin in PEGylated liposomes. The PEGylation helps to avoid clearance by the phagocytes and minimize interactions with plasma proteins, prolonging circulation time. The variation of synthesis methods and chemical functionalities alongside with facile post-modifications also make organic materials attractive choice for development of DDS. Drug loading is generally achieved by physical encapsulation of the drugs or conjugating the drug molecules to the carrier using cleavable covalent bonds.

Figure 2. Examples of different nanomaterials used to prepare drug delivery systems. Reproduced with permission.²³ Copyright 2015 Elsevier.

Another class of nanomaterial is derived from biological-based sources. Prominent examples of this category include exosomes, proteins and DNA-templated delivery systems.²⁴⁻²⁷ Offering high biocompatibility, low immunogenicity with excellent physiological stability, these bio-inspired nanocarriers are attractive alternatives to organic polymers for efficient drug and gene delivery. Interestingly, DNA-templated nanocarriers can be designed with unique structural complexities, allowing programmable properties and biological functions. Additionally, the plethora of natural bio-sources and advent of sophisticated biotechnology widen the scope of biomaterials-based nanocarriers that can be obtained. Further development of the applications of these biomaterials as DDS requires deeper understanding of fundamental concepts that remains elusive. For instance, the exact mechanisms of the interactions between exosomes and tumor cells or the self-assembly mechanisms of protein nanocages require further elucidation. More productive purification methods for scale-up synthesis and reproducibility of DNA nanocarriers and exosomes are also highly required. Such issues must be addressed for use in clinical applications.

Inorganic nanomaterials have also been of growing interest in drug delivery applications. Examples of inorganic nanoparticles comprise of quantum dots, silica-based nanoparticles and metal-based (e.g., gold, iron) nanoparticles²⁸. Unlike most organic nanomaterials which have no therapeutic functions, the unique inherent photonic, electric or magnetic properties of inorganic nanoparticles can be harness as therapeutic or imaging modalities. Shi et al. reported an oxygen-vacancy-enhanced relativity of a Gd³⁺-doped tungsten bronze (PEG-Na_xGdWO₃) nanorod with high-performance magnetic resonance imaging (MRI) ($r_1 = 80 \text{ mM}^{-1}\text{s}^{-1}$ at 0.7T) for

photothermal therapy.²⁹ Upon irradiation with a 980 nm laser, intravenous administration of their nanoparticles substantially suppressed tumor growth in mice. Furthermore, the tumor was completely eradicated by intratumor administration without recurrence during the study period of 95 days. The oxygen vacancies largely responsible for the high magnetic relativity and outstanding photothermal performance enabled PEG- Na_xGdWO_3 to be applied as efficient in-vivo MRI-guided tumor photothermal therapy. Despite the tremendous therapeutic potential of these inorganic nanomaterials, its long-term biosafety is not yet thoroughly understood since the biological effects could vary enormously with size, morphology and chemical compositions, thereby hindering their progress in clinical applications.

Figure 3. (A) Schematic illustration of porphyrinic metal-organic framework aMMTm nanoparticles for combination PDT and antiangiogenesis. Reproduced with permission.³⁰ Copyright 2019 Wiley-VCH. (B) Schematic illustration of Au-GI nanosystem for combinatorial photothermal, photodynamic and chemotherapy. Reproduced with permission.³¹ Copyright 2018 Wiley-VCH.

Lately, greater emphasis is placed on developing hybrid material DDS composing of both organic and inorganic components that could integrate the beneficial features of both components and possess distinct advantages over the individual

counterparts. Examples include nanosized metal-organic frameworks (MOFs) and coordination polymers. These are prepared using transition metal ions and polydentate bridging organic ligands to physically encapsulate drugs or have therapeutic properties as the carrier itself. Nie et al. exemplified the advantages of such multifunctional nanosystem by integrating a porphyrinic Zr-MOFs (aMMTm) decorated with a tumor cell membrane for the delivery of vascular endothelial growth factor receptor 2 inhibitor apatinib to achieve PDT and anti-angiogenesis therapy (Figure 3A).³⁰ The biomimic decoration mediated homotypic targeting, allowing selective accumulation in tumor while the use of MOFs-based photosensitizer effectively reduced self-quenching, improved loading amount and enhanced therapeutic outcomes. Similarly, Kim's group decorated gold nanoparticles with cytosine rich DNA i-motif and G-quadruplex hybrid (Au-GI) for combined anticancer therapy (Figure 3B).³¹ The G-quadruplex hybridized with the i-motif which can interact with photosensitizers and doxorubicin to load as cargoes. Inside the acidic tumor environment, the i-motif were constituted leading to aggregation of the gold nanoparticles, which efficiently generate heat for photothermal therapy under NIR irradiation.

Each of these class of material systems has its own advantages and limitations, which makes the selection of an appropriate material to be a complicated decision. Depending on the applications of the DDS, single or hybrid material-based DDS can be engineered to provide a system with desired functionalities.

1.3.2 Consideration factors for the design of nanocarrier

With the plethora of materials available for the fabrication of DDS, deciding on the appropriate choice of material is challenging. Moreover, there are key considerations in designing the DDS to maximize therapeutic efficacy. Generally, any DDS must be

biocompatible, stable under physiological conditions and allows for high drug loading capacity. Also, it should be able to achieve targeted site delivery and controllable drug release to minimize off-site toxicity.³²

To achieve targeted site delivery, conjugation of various targeting ligands of the nano-based DDS has been reported (Figure 4).³³ Targeting ligands such as antibiotics³⁴, aptamers³⁵, vitamins³⁶ and peptides³⁷ bind specifically to particular receptors overexpressed on the surfaces of cancer cells, and subsequently allowing the nanocarriers to be internalized via specific pathways.³⁸⁻³⁹

Figure 4. Illustration of the biophysicochemical properties of DDS. Reproduced with permission.¹⁴ Copyright 2012 Wiley-VCH.

Another major consideration for the design of efficient DDS is to ensure controllable release.⁴⁰⁻⁴¹ The DDS should transport the encapsulated drugs without drug leakage before cellular uptake and then release the drug in a controllable manner. To increase selectivity, internal stimuli such as pH, enzymes and redox agents or remote triggers including light and temperature can be harness to activate the release of the loaded drugs.⁴²

Figure 5. Different cleavable linkers used in fabrication of stimuli-responsive nanocarriers. The dashed line shows the bond that is cleaved when responded to the corresponding stimulus given in parentheses. Reproduced with permission.⁴³ Copyright 2012 Elsevier.

Internal stimuli make use of the intrinsic differences between the growth mechanism of cancer and normal cells to trigger a response in the DDS for drug release. As listed in Figure 5, cleavable bonds that are responsive to internal stimuli can be incorporated into the design of DDS or drug conjugation to temporarily maintain the stability of the encapsulated drugs in the formulation and achieve triggered release at desired tumor site. For example, the erratic growth of tumor results in a hypoxic environment and forces cancer cells to undergo anaerobic respiration which produces lactic acid and consequently much lower pH than normal tissues.⁴⁴ The difference in acidity of the intracellular environment of cancer and normal cells can be exploited to release drugs from DDS only in the tumors. One approach is to conjugate drugs onto the carrier material using pH-labile bonds or incorporate these bonds in the skeletal structure of the carrier materials. Commonly used pH-sensitive bonds include acetal, hydrazine, β -carboxylic acid amide linkages.⁴⁵⁻⁴⁶ Normally stable under neutral pH conditions, the pH-responsive DDS will degrade or dissociate in acidic environment,

alongside with pH-triggered drug release. Another strategy is to integrate protonatable groups (e.g., amino or carboxyl groups) into the DDS. Upon entry into the tumor microenvironment, where the pH is lower than their acid dissociation constant, these groups are protonated, leading to destabilization of the DDS and consequent release of the encapsulated drugs.

The significantly higher concentration of bio-reductants inside tumor cells than in the cellular exterior or in normal cells (≈ 10 mM vs ≈ 1 μ M) is another endogenous stimulus that could be taken advantage of for controlled drug release.⁴⁷ A commonly studied redox-labile bond is disulfide bond which can be incorporated as intra-carrier cross-linker. When the DDS are internalized into highly reducing cellular environment, the disulfide bond will be reduced, resulting in degradation of the DDS and release of drugs.

More recently, researchers have harnessed the elevated reactive oxygen species (ROS) levels in tumors for ROS-responsive drug release.⁴⁸ The enhanced ROS production in tumor cells may be a result of the abnormal metabolic activity, cellular proliferation and mitochondrial dysfunction which impaired ROS detoxification.⁴⁹ To exploit this biochemical feature for controlled release, ROS-sensitive DDS which incorporate thioketal, thioether or boronic ester bonds have been developed.⁵⁰

Still, there are great challenges in controlling the release and dosage of the drug by using internal stimuli. The heterogeneity of tumors and the endogenous variations between the tumor microenvironment and in normal cells that sometimes might be negligible could lead to ineffective responding of the sensitive bonds. In contrast, remote stimuli such as magnetic field and light could enable precise control of the timing of release within specific sites.⁵¹ For DDS with inbuilt magnetic-sensitive materials, application of an alternating magnetic field would result in vibration motion or produce

heat to cause disintegration or dissociation of the DDS.⁵² Light-triggered system involves materials that transfer absorbed light energy into heat or production of ROS that in turn be used as stimuli to promote drug release. Currently, most of the light sources used for the activation are ultraviolet (UV)- or near-infrared (NIR)- based.

With rational design of the nano-sized DDS, it offers advantages of high drug loading, site-targeted accumulation and spatial controlled drug release at the tumor site, which greatly improve therapeutic efficacy of the chemotherapy drugs while minimizing the non-specific toxicity to healthy tissues. Presently, the focus is to develop nanomaterials to deliver combination therapies against cancer. Multifunctional nanomaterials for mechanism-based combination therapies could help to greatly advance the development of optimal cancer treatment strategy.⁵³ Lin et al. exemplified the unique properties of inorganic materials by developing a novel nanocomposite (BPs@Au@Fe₃O₄) which comprised of iron oxide nanoparticles and gold nanoparticles assembled on black phosphorus sheets.⁵⁴ The combination of photothermal and photodynamic effect from the black phosphorus nanosheets together with the plasmonic photothermal effect from the gold nanoparticles mediated by a low-power NIR laser exhibited excellent tumor growth inhibition. In addition, the iron oxide nanoparticles acted as magnetic contrast agent, allowing monitoring of the treatment efficacy with magnetic resonance imaging. Nonetheless, no nanomedicine for combination therapy has been approved for clinical application yet. Challenges such as simplifying fabrication process with high batch-to-batch reproducibility, precise control over the dosage of drugs loaded and released from the nanocarrier and optimization of the nano-combination to achieve maximal therapeutic responses are yet to be overcome. Thus, it is of interest in this thesis to offer probable facile strategy of a DDS to deliver combination therapeutics for synergistic cancer therapy.

1.4 Supramolecular chemistry

In recent years, numerous DDS leveraging supramolecular chemistry have garnered interest. The concept of ‘Supramolecular chemistry’ was first put forth by Johannes Diderik van der Waals in 1873. Employing the non-covalent interactions between molecules (Figure 6) which are dynamic and tunable, new strategies to engineer DDS can be realized.⁵⁵

Figure 6. Examples of various non-covalent interactions between molecules.

Generally, the interplay of the various non-covalent interactions could direct the spontaneous association of molecules to construct ordered and stable systems. This phenomenon is also known as molecular self-assembly.⁵⁶⁻⁶² The inherent properties of the self-assembly process help to introduce complexity and/or functionality in the supramolecular system which would otherwise be difficult to achieve by classical covalent synthesis. Also, multiple synthesis steps and complicated purification steps could be avoided, making this approach more cost effective. Because of the facile preparation and modular properties of the self-assembled systems, supramolecular chemistry which provides encapsulation and targeted release mechanism is incorporated in the design of DDS. Examples of the supramolecular complexes include micelles, vesicles and liquid crystals.⁶³ In addition, hybrid nanocomposites can be prepared with a variety of materials and organization of the individual entities into hierarchical architectures, hence contributing to designs of multifunctional self-assembled DDS.

Among the various noncovalent interactions, host-guest interaction based on macrocycles which includes calixarenes, crown ethers, cyclodextrin, pillar[n]arenes and cucurbit[n]urils has been widely studied. For a typical host-guest inclusion complex, the macrocycles act as the host to provide a cavity that encapsulates guest molecules via noncovalent interactions.

Figure 7. Schematic illustration of the structure of β -cyclodextrin and its typical guest molecules. Reproduced with permission.⁶⁴ Copyright 2014 American Chemical Society.

One prominent host molecule, cyclodextrin (CDs) is widely employed in drug delivery. As a family of cyclic oligosaccharides, CDs are made of macrocyclic glucose subunits connected by α -1,4 glycosidic bonds. Naturally occurring α -, β -, and γ -CD are made of 6, 7 and 8 glucose subunits respectively. Possessing a toroidal shape, the primary and secondary hydroxyl groups are exposed on the exterior wall of the toroid, making it hydrophilic. On the other hand, the interior of the CDs is considerably less hydrophilic and act as suitable cavity to host hydrophobic molecules such as adamantane, cholesterol, azobenzene and ferrocene (Figure 7).⁶⁴ The earliest uses of these macrocycles in drug delivery were to act as excipients which improved the solubility and cell permeability of the encapsulated hydrophobic drugs.⁶⁵⁻⁶⁶

Figure 8. Schematic illustration of supramolecular nanoparticles formed from amine-modified β -CD and PEGylated adamantane for delivery of succinobucol. Reproduced with permission.⁶⁷ Copyright 2019 John Wiley and Sons.

The specific binding and relative high stability of a host-guest complex provide robust and reliable methodology for the construction of supramolecular systems. Importantly, the favorable biocompatibility of these host macrocycles and the reversibility of the host-guest complex formation makes them attractive candidates for fabrication of stimuli-responsive supramolecular systems. To illustrate such concept, Li et al. utilized protonatable aliphatic amine-modified CDs for pH-responsive drug delivery and release.⁶⁸ They constructed a supramolecular system formed between PEGylated adamantane and hydrophobic β -CD derivative with loading of water-insoluble succinobucol in the core for treatment of lung metastasis (Figure 8). In the acidic cellular environment, the amine groups on the CDs were protonated, disturbing the hydrophilic-hydrophobic balance of the nanoparticles and achieving pH-specific drug release. Other than molecular drugs, gene therapeutics like DNA can also be delivered with such strategy.⁶⁹⁻⁷⁰

1.5 Prodrugs

Classically, the term ‘prodrug’ relates to the biologically inactive drug derivatives in which the activity of the parent drug has been temporarily masked. At certain cellular locations, the prodrugs transform via enzymatic and/or chemical processes to the pharmacologically bioactive parent drug (Figure 9A).⁷¹ Other than ensuring site-specific activity, such modifications help to improve the water solubility, increase its chemical stability and circulation time, thus overcoming the shortcomings of most chemotherapy drugs.

Figure 9. (A) Schematic illustration of mechanistic action of prodrugs and the common functional groups on parent drug amenable to modifications. Reproduced with permission.⁷¹ Copyright 2008 Springer Nature. (B) Schematic illustration of H₂O₂-responsive DNA alkylating nitrogen mustard prodrug. Reproduced with permission.⁷² Copyright 2011 American Chemical Society. (C) Schematic illustration of NIR light-responsive paclitaxel prodrug. Reproduced with permission.⁷³ Copyright 2016 American Chemical Society.

Active site of drugs can be modified with enzymes-cleavable linkage. Due to the high specificity of enzymes, such prodrugs are will only be activated in tissues that overexpress the enzymes, allowing controlled release.⁷⁴⁻⁷⁵ In terms of chemically triggered release, prodrugs can be obtained via modifications with redox-responsive (Figure 9B) or light-responsive linker (Figure 9C) and be released in the presence of reducing agents or oxidative stress and photoirradiation respectively.⁷²⁻⁷³

1.5.1 Prodrug-based self-assembled delivery systems

With the deeper understanding of the non-covalent mechanisms behind self-assembled systems and the advantages of prodrugs, there is an increasing interest in developing prodrug-based nanoparticles for cancer therapy.⁷⁶ The most studied systems would be polymer-based drug-conjugated DDS.⁷⁷⁻⁸¹ Typically, hydrophobic drugs are conjugated to hydrophilic polymers. The amphiphilic balance between the drug and polymer allows self-assembly. During the self-assembly process, additional therapeutic agents could be introduced into the DDS.

Figure 10. (A) Schematic illustration representing the preparation of EGCG/herceptin nanocomplexes and its representative TEM image. Cell viability assay showed the nanocomplexes with higher cytotoxicity over free EGCG or proteins. Reproduced with permission.⁸² Copyright 2014 Springer Nature. (B) Schematic illustration showing formation of liposome-like CPT nanocapsules encapsulating DOX with enhanced cytotoxicity over the free drugs. Reproduced with permission.⁸³ Copyright 2010 American Chemical Society.

For example, (–)-Epigallocatechin-3-O-gallate (EGCG) that has been reported to exhibit anticancer effects was used to form a self-assembled micellar nanocomplex encapsulating anticancer drug herceptin for cancer therapy (Figure 10A).⁸² Administered at a non-toxic dosage, derivatized EGCG forms stable micellar complexes with herceptin and augment its cytotoxicity with better tumor selectivity and accumulation compared to the free proteins. In another work, Murdoch et al. used low molecular weight prodrug molecules to fabricate nanocarriers (Figure 10B).⁸³ The hydrophobic CPT was attached with a short ethylene glycol chain, facilitating self-

assembly to give stable liposome-like nanocapsules with high doxorubicin loading efficiency.

However, these traditional polymer-based drug-conjugated DDS often suffer from low drug loading since the molecular drugs are appended to high molecular weight polymers. Thus, construction of prodrug self-assembled systems where the low molecular weight prodrugs are rationally modified with inherent amphiphilic properties to facilitate self-assembly into nanoparticles under specific conditions (i.e. pH, concentration, ionic strength and kinetic factors) has garnered research interest.⁸⁴ With the drug acting as both the backbone of the carrier as well as the cargoes, such prodrug self-assembled systems are also known as carrier-free DDS. Such designs yield few advantages: (1) superior drug loading; (2) elimination of preparation steps for additional carriers; (3) minimize carrier-induced toxicity and immunogenicity, (4) prolong drug circulation and achieve selective accumulation due to EPR effect and (5) controlled drug release at cancerous tissues, reducing systemic toxicity.

Figure 11. (A) Schematic illustration of β -sheet peptide-CPT prodrug conjugated nanotubes with high drug loading. Reproduced with permission.⁸⁵ Copyright 2013 American Chemical Society. (B) Self-delivery prodrug system fabricated between RGD peptide and camptothecin for targeting integrin-overexpressing tumor. Reproduced with permission.⁸⁶ Copyright 2016 Springer Nature.

Since majority of the anticancer drugs are hydrophobic, short peptides with its excellent biocompatibility, bioactivity and tunable amphiphilicity have been found to be advantageous for constructing amphiphilic prodrugs. Recently, Cui et al. reported their work on a self-assembled β -sheet peptide-CPT conjugated nanotubes with high drug loading of 38% (Figure 11A).⁸⁵ Targeting peptides have also been successfully modified to paclitaxel which formed nanospheres via self-assembly and demonstrated specific tumor accumulation. Zhang's group fabricated a tumor-targeting prodrug self-delivery system, inclusive of the RGD peptide and CPT (Figure 11B) for better anticancer efficacy.⁸⁶ The integrin-targeting action of the peptide helped to enhance cellular internalization and in turn improved therapeutic efficacy of the chemotherapy drug.

Figure 12. (A) Schematic illustration of self-assembly of amphiphilic Ir-Cb nanoparticles and on-demand drug release. Reproduced with permission.⁸⁷ Copyright 2014 American Chemical Society. (B) Schematic illustration of preparation of amphiphilic pPhD nanoparticles for combination therapy. Reproduced with permission.⁸⁸ Copyright 2018 Springer Nature.

Molecular self-assembly of amphiphilic drug-drug conjugate based on chemical conjugation or non-covalent interaction were also designed for combination therapy.⁸⁹⁻⁹⁰ By conjugating hydrophobic chlorambucil with hydrophilic Ir via a hydrolysable ester linkage, Yan et al. pioneered the work on amphiphilic Ir-Cb conjugate self-assembled nanodrug, which can achieve synergistic cytotoxicity (Figure 12A).⁸⁷ In a work by Li's group, a charge-transformable pPhD nanoparticles constructed from pheophorbide a and doxorubicin for combination photodynamic therapy and chemotherapy was reported (Figure 12B).⁸⁸ The conjugation of hydrophobic pheophorbide a photosensitizer and hydrophilic doxorubicin via a pH-responsive hydrazine bond resulted in amphiphilic PhD monomer which first assembled into ultra-small micelle-like particles and further crosslinked by dual-aldehyde terminated polyethylene glycol (PEG) to give relatively stable nanoparticles. The pH responsiveness of pPhD nanoparticles allowed deep tumor penetration with improved synergistic in-vitro and in-vivo therapeutic outcome over the non-PEGylated and non-transformable nanoparticles.

1.6 Platinum-based drugs

Today, a broad spectrum of anticancer drugs is in clinical use as monotherapy or as cocktail combination therapy. Particularly, platinum-based drugs are widely used as chemotherapeutics, as it is used in estimated 50- 70% chemotherapy cocktails regime.⁹¹

The first platinum-based drug, Cisplatin was granted FDA approval in the 1970s as a chemotherapeutic and is used in treating various cancers such as bladder, sarcomas

and testicular cancers.⁹² Its administration in chemotherapy regime has greatly improved the survival rates of cancer patients; particularly having attained a tremendous improvement in testicular cancer mortality from 85% to 10%.⁹³

The mechanism of cell killing of cisplatin occurs via the cross linking of the platinum (Pt) core with the nuclear DNA. Studies show that when cisplatin enters cells, ligand exchange occurs between its labile chloride ligands and water.⁹⁴⁻⁹⁵ The resulting electrophilic Pt(II) complexes are able to react with nucleophilic purine bases of nucleic acid, specifically guanine at the N-7 position and form intrastrand cross-links, inhibiting DNA synthesis or repair, eventually resulting in apoptosis.⁹⁶

Despite its therapeutic success in treating testicular cancer, cisplatin has been found to induce side effects such as renal neurotoxicity, nausea and nephrotoxicity.⁹⁷ Furthermore, some cancers including breast, ovarian and colorectal cancers exhibited resistance to cisplatin therapy. These limitations spearheaded the quest to develop other novel platinum complexes with anti-proliferative properties (Figure 13).⁹⁸⁻⁹⁹

Figure 13. Examples of platinum-based anticancer drugs.

New platinum drugs are strategically designed by varying the labile leaving ligands and non-labile leaving ligands.¹⁰⁰⁻¹⁰² The lability of the leaving ligands will decide the reaction and aquation kinetics of the platinum complex. Ligands that are highly labile will in turn induce high toxicity as their rapid loss from the platinum

complex lead to unselective binding to off-target proteins and biological nucleophiles.¹⁰³

Hence, carboplatin which has relatively stable 1,1-cyclobutanedicarboxylate ligand (aquation rate constant $\sim 10^{-8} \text{ s}^{-1}$) as leaving group causes fewer side effects than cisplatin with its labile chloride ligands (aquation rate constant $\sim 10^{-5} \text{ s}^{-1}$).¹⁰⁴

The non-leaving ligands are mainly nitrogen donors as they form thermodynamically stable coordination bonds with platinum.¹⁰² Since the non-leaving amine ligands remain in the final Pt-DNA adducts, they can be modulated to alter the cytotoxicity and mechanism of action of the platinum drug. Oxaliplatin which belongs to the 1,2-diaminocyclohexane (DACH) ligand family exhibits vastly different spectra of activity from cisplatin which contains cis-diammine ligands.¹⁰⁵ The formation of sterically hindered and more hydrophobic DACH-Pt-DNA adducts inhibit DNA synthesis more effectively and with superior cytotoxicity than cis-diammine-Pt-DNA. Thus, oxaliplatin is potent in treatment of colorectal cancer which are inherently resistant to cisplatin.

As it can be seen, the conventional concept of platinum drug development involves the different combinations of labile leaving group and the non-labile leaving group before going on to test the pharmacodynamics and cytotoxicity of the resulting drug agent. Such an approach may be considered as “trial and error”, requiring a long timeframe to modify and achieve appropriate solubility, metabolic stability, membrane permeability, distribution and half-life. Moreover, there is often a lack of clear understanding of the exact molecular targets and mechanisms of action of the novel platinum agent, which prevents clinical translation and commercialization of the new drug.

Most of the novel platinum compounds only offer incremental toxicity over existing platinum-based drugs, which is unsurprising since majority are structural

derivatives of cisplatin. Clinically, it is a huge challenge to develop another potential platinum drug with distinct advantageous properties from cisplatin and oxaliplatin. Alternatively, the focus should be shifted from designing new platinum-based drugs to improving or enhancing the targeted accumulation and maximize the drug efficacy of existing platinum drugs.

1.6.1 Nanoparticulate platinum drugs-loaded delivery system

Preferably, the platinum drugs should only accumulate at the tumor sites and kill cancer cells without affecting the normal cells. However, the nonspecific systemic distribution and poor selectivity results in side effects whilst increasing dosage will eventually lead to drug resistance, hereby limits the application of the platinum drugs. To overcome the unfavorable situations, two main strategies are often implemented.¹⁰⁶

One strategy is the encapsulation of the platinum drugs in DDS to improve physiological stability and enhance tumor accumulation.¹⁰⁷⁻¹⁰⁹ Barthelemy and coworkers demonstrated that nucleoside-lipids-based nanoparticles could efficiently deliver cisplatin into various cancer cell lines to enhance the antitumor efficacy.¹¹⁰ Their novel encapsulation technique controlled the precipitation of cisplatin into aggregates followed by directed self-assembly by the nucleoside polar heads into high drug-loaded and stable nanoparticles. Their system reported an unprecedented in-vitro toxicity of up to 130-folds higher than free cisplatin. To overcome multidrug resistance resulting from hypoxic condition, Chen et al. developed a nano-carrier system which can release both Pt(II)-based drugs and O₂.¹¹¹ In their work, PLGS polymer was used to encapsulate Pt(II) drugs and catalase within the core to enhance cellular accumulation. The catalase produces O₂ from intracellular H₂O₂, resulting in increased internal pressure which ruptured the nanoparticle shell and released the encapsulated drugs.

Figure 14. Schematic illustration of solvent displacement preparation of PLGA-based nanoparticles ratiometrically containing cisplatin core and GMP core for synergistic anticancer therapy. Reproduced with permission.¹¹² Copyright 2014 John Wiley and Sons.

To achieve combination therapy, Huang et al. introduced a PLGA-based nanoparticle that ratiometrically loaded cisplatin and gemcitabine monophosphate (Figure 14).¹¹² To resolve the disparity in the physicochemical properties of the two drugs, they proposed the use of dioleoyl phosphatidic coated calcium phosphate cores to separately load the two drugs as hydrophobic cores, which were subsequently encapsulated into the same PLGA carrier via a solvent displacement method. This

strategy shown precise ratiometric control over drug loading and cellular accumulation, which is key to achieving synergistic anticancer therapy.

1.6.2 Platinum(IV) prodrugs

The other strategy for improved platinum drugs' efficacy is to develop inactive Pt(IV) prodrugs. They are prepared by introducing two axial ligands to the Pt(II) drug (Figure 15) which not only tune the redox potential of the platinum core but could help to change the hydrophobic/hydrophilicity balance or to conjugate a functional moiety.¹¹³

Figure 15. Composition of platinum(IV) prodrugs. Reproduced with permission.¹¹³ Copyright 2014 American Chemical Society.

The inherent properties of Pt(IV) prodrugs provide two main advantages.^{101, 114} Firstly, Pt(IV) prodrugs have to undergo reduction before aquation and subsequent reaction with biological nucleophiles. Reduction to its cytotoxic analog form was in account of high intracellular concentration of endogenous reducing agents, for example, ascorbic acid and glutathione in cancer cells. This reduces side effects to normal cells as activation of the active drug is occurred selectively in cancer cells.¹¹⁵ Secondly, Pt(IV) prodrugs have considerably slower ligand exchange rates and better inertness than its Pt(II) counterparts. The better stability of Pt(IV) prodrugs favors longer circulation

lifetime in blood, facilitates accumulation in tumor sites and also avoids undesirable interactions with proteins and other biomolecules during blood circulation.

Satraplatin, with two polar acetate axial ligands was the first reported Pt(IV) agent for oral administration. Its reduction half-life of 50 mins provided sufficient time for intestinal adsorption and exhibited better toxicity behavior over cisplatin in preclinical studies. Despite entering clinical trials, it was unable to obtain FDA approval as the overall survival rates of patients did not improve.¹¹⁶ Still, satraplatin demonstrated the advantages of enhanced lipophilicity and stability of Pt(IV) prodrugs over the unmodified Pt(II) agents.

The inclusion of biologically active agents as the axial ligands can produce dual-threat Pt(IV) agents.¹¹⁷⁻¹¹⁸ One prominent Pt (IV) prodrug that has been developed is the conjugation of ethacrynic acid to cisplatin to form ethacraplatin.¹¹⁹ As an effective inhibitor of glutathione-S-transferase (GST) isozymes, ethacrynic acid is often used in combination with cisplatin to treat cisplatin-resistant cell lines. Upon cellular uptake, ethacraplatin reduces in-vivo releasing the ethacrynic acid and a cytotoxic cisplatin, thus reversing cisplatin associated drug resistance. In the work by Dyson's group, ethacraplatin displayed better inhibitory effects of GST as compared to ethacrynic acid.¹¹⁹ Besides, a range of cell lines treated with ethacraplatin in a 24 hr incubation period resulted in higher cytotoxicity compared to treatment with cisplatin alone.

In addition to reduction-activated Pt(IV) prodrugs, photoactivated diazido Pt(IV) complexes were developed to overcome the inadequate selectivity and uncontrollable reduction activation.¹²⁰⁻¹²² These diazido Pt(IV) prodrugs harness the site-specific property of light as an external stimulus and get reduced to active Pt(II) drugs when excited with ultraviolet or visible light. Sadler et al. synthesized trans, trans, trans-[Pt(N₃)₂(OH)₂(NH₃)(py)] containing an octahedral Pt(IV) core with almost linear azido

ligands.¹²² The Pt(IV) complex displayed good dark stability and was inert to cellular reducing agent but became a potent toxin when photoactivated. Upon light irradiation, the prodrug was shown to be more cytotoxic toward cisplatin-resistant A2780cis cells. The strategy of using light to activate diazido Pt(IV) prodrugs could lessen the side effects of unselective platinum-based drugs by preferentially irradiating the site of interest with light. Furthermore, unlike conventional photodynamic therapy, light-activated diazido Pt(IV) could work well in hypoxic tumors as it does not require oxygen for its mode of action.

1.6.3 Preparation of platinum(IV) complexes

The synthesis routes for preparing Pt(IV) complexes are fairly established and complexes with different axial ligands can be obtained.¹¹³ The most common synthetic route occurs via two-electron oxidation of a Pt(II) precursor by excess hydrogen peroxide in water, which gives trans-dihydroxo Pt(IV) complexes.¹²³ The stereochemistry of the equatorial ligands in the obtained Pt(IV) complex is retained as in the initial Pt(II) species. The nucleophilic dihydroxo axial ligands in the Pt(IV) complexes can then react with other electrophiles such as acid anhydrides¹²⁴ or acid chlorides,¹²⁵ and isocyanates to give dicarboxylate and dicarbamate Pt(IV) complexes respectively (Scheme 1). Alternatively, coupling trans-dihydroxo Pt(IV) species to activated carboxylic acids with suitable coupling reagent such as TBTU also gives trans dicarboxylate complexes.¹²⁶

A useful group of trans Pt(IV) dicarboxylates can be obtained by ring-opening reactions of Pt(IV) dihydroxo complexes with cyclic anhydrides. Carboxylation with succinic, maleic, glutaric and phthalic anhydrides produces Pt(IV) carboxylates with terminal carboxyl functional group that can further undergo amide or ester coupling reactions (Scheme 1).¹²⁷⁻¹²⁹ Such reactions are conveniently used to tether targeting

groups to Pt(IV) complexes to minimize unselective cellular uptake¹³⁰⁻¹³³ or couple to a range of amines and alcohols to adjust their lipophilicities.¹³⁴⁻¹³⁶

Scheme 1. Possible reaction routes of symmetrical dihydroxo platinum(IV) complexes.

Recently, axial ligands with carbonyl, azide and thio-reactive maleimide functional groups are also reported.¹¹³ Carbonyl functional groups could be reacted with organic hydrazines and hydroxylamines to form hydrazones and oximes which are useful functionalities to fabricate novel Pt(IV) complexes. Kowol et al. prepared Pt(IV) complex with axial ligand containing thiol-reactive maleimide functional group. This facilitates conjugation to human serum albumin at its cysteine residue for assisted delivery into CT-26 tumors.¹³⁷

Besides symmetrical Pt(IV) complexes, preparation of mixed hydroxo-carboxylato complexes could allow the introduction of different functionalities and

greater molecular complexity.¹²⁴ Mixed trans hydroxo-carboxylato Pt(IV) complexes can be attained via peroxide oxidation of Pt(II) species with excess carboxylic acid.¹²⁶ Further functionalization with acid anhydrides could yield mixtures of dicarboxylate complexes. Otherwise, limiting the amount of acid anhydride reacting with dihydroxo Pt(IV) under mild conditions also result in the monocarboxylate product.^{113, 138} Similarly, the monocarboxylate species can be obtained by reacting dihydroxo Pt(IV) with limiting pre-activated carboxylic acids.

1.7 Platinum(IV) prodrugs-based nano-delivery system for cancer therapy

To facilitate more efficient delivery of platinum drugs, research on integrating the prodrug strategy and nanotechnology into a single system is emerging. By developing as prodrug-based nanocarriers, it combines the advantages of reduction-responsive release of the bioactive form of the Pt(IV) prodrugs and promotes tumour accumulation of the prodrugs via the EPR effect.

Figure 16. (A) Schematic illustration of cisplatin(IV)-encapsulated Pt(II)-based supramolecular cage delivery system.¹³⁹ Published by The Royal Society of Chemistry. (B) Schematic illustration of ethacraplatin-loaded nanomicelles with high GST inhibitory activity for treating cisplatin-resistant BEL7404-CP20 cells. Reproduced with permission.¹⁴⁰ Copyright 2017 Elsevier.

The simplest form of this concept is prodrug-encapsulated nanoparticles.¹⁴¹⁻¹⁴³ The lipophilic Pt(IV) prodrugs can be physically encapsulated in a wide range of inorganic or polymeric nanoparticles. The Lippard's group designed a novel DDS comprising of hexanuclear Pt(II) cage which encapsulated Pt(IV) prodrug molecules (Figure 16A).¹³⁹ This self-assembled system was shown to react with biological reductant ascorbic acid and release cytotoxic cisplatin. In-vitro experiments demonstrated the capability of the system to facilitate enhanced cellular accumulation

and prompted cell cycle arrests in A2780 cell line. Biologically active molecules that impair drug resistance mechanism can be tethered to the platinum core to treat drug-resistant cells. Liang et al. demonstrated the advantages of such systems with the use of DSPE-PEG nanomicelles to deliver ethacraplatin, a glutathione S-transferases (GST) inhibitor modified cisplatin prodrug (Figure 16B).¹⁴⁰ The nanosystem was able to inhibit the activity of GST and circumvent deactivation of cisplatin, achieving more than two-folds reduction in IC₅₀ value when cisplatin-resistant BEL7404-CP20 cell line was treated with their formulation as compared to free cisplatin. By encapsulating the ethacraplatin in micelles, it presented low systemic toxicity and circumvented the nephrotoxicity of cisplatin. One main disadvantage of the noncovalent encapsulation of prodrugs in DDS is the relatively low drug loading capacity due to the high proportion of carrier materials.

For higher drug loading efficiency, Li and co-workers reported a small molecular weight peptide-conjugate cisplatin(IV) prodrug that could undergo enzyme-triggered self-assembly into nanostructures.¹⁴⁴ Tumor overly expresses phosphatases which catalyze dephosphorylation of the short peptide substrate, allowing in-situ supramolecular self-assembly of the prodrug nanofiber that enhanced drug accumulation.

Similar to typical DDS, prodrug-based nanocarriers can adopt combination therapy approach to maximize synergistic chemotherapeutic efficacy against individual drug targets.¹⁴⁵ Kabanov's group co-loaded cisplatin(IV) prodrug and etoposide into amphiphilic block copolymer system with over 50% wt. drug loading.¹⁴⁶ The encapsulation of drugs combination in the micelles improved tumour accumulation of both drugs and demonstrated better antitumor activity over the free drugs combination in several animal models. Other therapeutics agents such as gene medicine, proteins or

photosensitizers could also be co-encapsulated with platinum prodrugs to overcome platinum drug resistance via combinational therapy.¹⁴⁷⁻¹⁵³ Imaging modalities could also be incorporated to monitor the therapeutic effects of the nanoparticle system.¹⁵⁴ Although these delivery strategies have improved cytotoxicity, their in-vivo behaviors remain elusive. Physical encapsulation of the drugs could have premature release issues and is also challenging to guarantee the integrity of the delivery system before reaching the target tumor.

Figure 17. (A) Schematic illustration of the preparation of PLA-cisplatin(IV) polymer by microfluidic technology, with low IC₅₀ values for treatment in vitro. Reproduced with permission.¹⁵⁵ Copyright 2010 National Academy of Sciences. (B) Schematic illustration of triply drugs-conjugated polymer nanoparticles via ring-opening metathesis polymerization.¹⁵⁶ Reproduced with permission. Copyright 2014 American Chemical Society.

To prevent premature drug release, strategies were undertaken to directly integrate the prodrug in the carrier architectures. Commonly, the Pt(IV) prodrug is

fabricated as self-assembling polymer-drug conjugate which provides enhanced chemical stability, further post surface functionalization and permits co-delivery of other therapeutics agents for combinational therapy. For example, Farokhzad et al. derivatized a poly(lactic acid) polymer with cisplatin(IV) prodrug pendants as the carrier material to co-deliver hydrophobic docetaxel for combination drug therapy (Figure 17A).¹⁵⁵ Microfluidic technology was harnessed to prepare the nanoparticles, which was post-modified with A10 aptamer for active targeting prostate-specific membrane antigens on prostate cancer cells. The polymer-prodrug conjugate approach minimized batch-to-batch variability of nanoparticles during synthesis with temporally controllable drug release rates for superior in vitro toxicity over single drug nanoparticles. Huang et al. demonstrated a dual drug built-in polymeric theranostic system with exact drugs composition and fixed drugs ratio.¹⁵⁷ DNA-damaging agent, demethylcantharidin was conjugated onto cisplatin in an optimal 2:1 fixed ratio, and subsequently polymerized with ethylenediamine to form a polymeric-prodrug material that can self-assemble into nanoparticles. In another work, Johnson and co-workers utilized the “brush-first” ring-opening metathesis polymerization method to prepare triple drugs-conjugated polymeric nanoparticles (Figure 17B).¹⁵⁶ The unique convergent polymer synthesis allowed tunable and precise drugs loading ratio with orthogonally triggered drug release. Compared to the one- and two-drug-loaded systems, their three-drug-loaded nanoparticles led to at least two-folds reduction in IC₅₀ for treatment with OVCAR3 human ovarian cancer cells.

Interestingly, Zheng’s group reported a coordination-driven self-assembly strategy for Pt(IV) drug delivery.¹⁵⁸ Specifically, a Pt(IV)-conjugated dipyriddy building block self-assemble with organoplatinum linkers to give supramolecular hexagon, each of which could deliver three equivalence of cisplatin(IV) prodrug. Cytotoxicity assays

of the supramolecular DDS tested on various cancer cell lines demonstrated a more superior therapeutic efficacy over cisplatin. Mechanistic studies indicated that the complex had preferential entry into cancer cells and were reduced intracellularly to release free cisplatin which damaged DNA and caused apoptosis.

However, more investigations detailing the therapeutic performance of these above-mentioned system in-vivo is required.

1.8 Research gaps and objectives

As discussed in the previous sections, constructing Pt(IV) prodrug has been a new direction in the development of platinum based anticancer drug. Although this approach has greatly overcome the limitations of typical Pt(II) drugs, these molecular Pt(IV) prodrugs could not fully escape the fate of premature degradation or rapid clearance. Consequently, inadequate tumour accumulation of such molecular complexes fails to fully exploit its latent potential. On the other hand, delivering platinum drugs encapsulated in traditional DDS suffers from low drug loading efficiency, undesirable premature leakage during systemic circulation and potential toxicity from the carrier material.

Inspired by the natural self-assembly processes and the advent of nanomedicine, the author wondered if it is possible to construct novel Pt(IV) prodrug-based delivery systems outside of the typical Pt(IV)-polymer conjugate micelles category. Particularly, the author chose to focus on oxaliplatin(IV) prodrug as it is the newest platinum derivative in standard platinum-based chemotherapy and not much research concerning the development of its prodrug applications was conducted. Primarily used in colorectal cancer therapy, improving treatment efficacy of oxaliplatin via prodrug-based nano-delivery systems could be a hopeful approach to expand its applications in different

types of cancer. Taking this forward, the main objective of this dissertation is to develop different oxaliplatin(IV) prodrug self-assembled nanocarrier for cancer therapy.

In Chapter 2, the author first explores the idea of constructing oxaliplatin(I) prodrug-based self-delivery nanocarrier based on host-guest complex and self-assembly. As a proof-of-concept, a redox-responsive supramolecular nanocarrier was constructed between a colorectal cancer-targeting ligand conjugated on cyclodextrin and a cholesterol-modified oxaliplatin(IV) prodrug. The resulting amphiphilic supramolecular complex self-assembled into nano-sized carrier, which preferentially accumulated in polyamine transporter over-expressing HCT116 colorectal cancer cells and disassembled under highly reducing intracellular environment to maximize anticancer efficacy.

In Chapter 3, to provide a facile strategy for delivering combination therapeutics for synergistic cancer therapy, a nanocarrier self-assembled with oxaliplatin(IV) prodrug and porphyrin photosensitizer components via host-guest chemistry is proposed. Motivated by the excellent spatial control of binding ratio with host-guest complexes, porphyrin and oxaliplatin were separately modified with cyclodextrin and adamantane respectively and the resulting amphiphilic host-guest complex can easily self-assemble into nanomaterials that act as both carriers and cargoes. Our approach allows precise control over the drug and photosensitizer loading ratios; eliminates the toxicity concerns of the drug carrier; and at the same time, achieves therapeutic synergy from chemotherapy and photodynamic therapy.

In Chapter 4, the potential of concurrent delivery of anticancer proteins and chemotherapy drugs as a self-delivery nanocomplex is studied. Delivering cytotoxic proteins is an emerging field in cancer treatment. Currently, the delivery of such proteins is highly compromised by multiple successive physiological barriers which reduce the

therapeutic efficacy. Hence, there is a need to design a robust and sustainable nanocarrier which not only provide good protection of the protein but also improve cellular accumulation. Here, an oxaliplatin(IV) polymeric prodrug was used to fabricate a redox-responsive nanocarrier to deliver bioreversibly caged RNase A protein to realize cancer-targeted treatment. The results obtained indicates that this system facilitates preferential cancer cells accumulation of the protein and was more potent compared to the free therapeutics in anticancer proliferation.

In Chapter 5, we summarize how the proposed oxaliplatin(IV) prodrug self-assembly systems discussed in this dissertation could provide a prospective future for precision theranostics in nanomedicine.

Chapter 2

An oxaliplatin(IV) prodrug-based supramolecular self-delivery nanocarrier for targeted colorectal cancer treatment

2.1 Introduction

Oxaliplatin is a commonly used chemotherapeutics for treating colorectal cancer. Despite their success, these drugs often suffer from some drawbacks such as poor tumor selectivity and accumulation as well as dosage-dependent drug resistance, thus limiting their applications to some extent.¹⁵⁹⁻¹⁶⁰ To address these problems, considerable research has been made to develop sophisticated drug carriers for targeted delivery to tumors.^{139, 161-162} However, low drug loading capacity, potential systemic toxicity, complex fabrication, and related metabolism concern of these multifunctional nanocarriers have hampered their clinical translations toward practical uses.¹⁶³

Carrier-free Pt(IV) prodrug delivery systems are a novel kind of therapeutics.¹⁶⁴⁻¹⁶⁶ Such prodrugs are easily prepared by introducing two axial ligands to a Pt(II) drug to not only tune the redox potential of the platinum core to Pt(IV) but also change the hydrophobicity/hydrophilicity balance.^{101, 106, 113} The amphiphilic molecular prodrugs could in turn form nanostructures that act as both carriers and cargoes.^{76, 167} Compared to traditional carrier-based drug delivery systems, such carrier-free self-delivery nanosystems demonstrate much higher drug loading capacity and eliminate the toxicity concerns of the drug carriers, whilst having the nanoscale advantage of enhanced permeability and retention (EPR) effect to achieve passive accumulation at the tumor site.¹⁶⁸⁻¹⁷³ To become pharmacologically active, Pt(IV) prodrugs need to undergo intracellular reduction discriminatingly in cancer cells containing high concentrations of endogenous reducing agents.^{115, 174} This feature could reduce side effects to normal cells, as the activation of the active drug is selective in cancer cells.

In this chapter, we report the fabrication of self-assembled nanoparticles through the host-guest complexation between spermine-functionalized β -cyclodextrin (CD-spermine) and cholesterol-conjugated oxaliplatin(IV) (oxliPt(IV)-chol) for colorectal cancer-targeted and redox-responsive drug delivery (Scheme 1).

Scheme 1. Schematic illustration of (A) self-assembly of CD-spermine and oxliPt(IV)-chol for the formation of CD-spermine: oxliPt(IV)-chol nanoparticles and (B) redox-responsive drug release in cancer cells.

Each component in the nanocarrier has irreplaceable function, i.e., oxaliplatin(IV) as a prodrug, spermine as a targeting ligand, and cholesterol and β -

cyclodextrin for the formation of inclusion complex with inherent amphiphilicity for the self-assembly.¹⁷⁵ As natural biomolecules involved in cellular biochemical pathways such as signaling and protein biosynthesis, polyamine spermine exhibits favorable biocompatibility and minimal side effects.¹⁷⁶⁻¹⁷⁷ Colorectal cancer cells were found to overexpress polyamine transportation receptors in order to scavenge polyamines from exogenous sources for sustaining rapid cell division.¹⁷⁸⁻¹⁷⁹ Hence, spermine could function as an active targeting ligand for enhanced accumulation of the nanocarrier in colorectal cancer cells.

2.2 Experimental Section

2.2.1 Materials and reagents

All solvents and chemicals were purchased from Sigma-Aldrich. Dulbecco's Modified Eagle's medium (DMEM) and fetal bovine serum (FBS) were purchased from PAA Laboratories. 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide was bought from Alfa Aesar. HCT116 cells were obtained from American Type Culture Collection (ATCC).

2.2.2 Characterization

¹H NMR spectra were probed on 300 MHz Bruker spectrometer using CDCl₃, deuterated DMSO or D₂O as the solvent. ThermoFinnigan LCQ quadrupole ion trap mass spectrometer was used to record electronic spray ionization (ESI) mass spectra. High-resolution mass spectrometry (HR-MS) was performed on a Waters Q-tof Premier MS spectrometer. Fourier transform infrared (FTIR) spectra were recorded as KBr-pellet on a SHIMADZU IR Prestige-21 spectrophotometer. High performance liquid chromatography (HPLC) analysis was performed with Shimadzu LC2010 using the Waters XSelectTM HSS C18 (5 μm, 4.6 × 250 mm) column with UV detection at 240 nm. Acetonitrile/water binary system containing 0.5% trifluoroacetic acid was used as the mobile phase with a flow rate at 1 mL min⁻¹. The analysis was performed with gradient elution method starting with 95% water to 5% water and back to 95% water at 30 °C over a period of 40 min. Transmission electron microscope (TEM) images were collected on a JEM-1400 (JEOL) operated at 100 kV. Zeta potential and dynamic light scattering (DLS) measurements were measured using Zetasizer, Malvern Instrument Ltd. UV-Vis spectra were obtained on a Shimadzu UV-3600 UV-Vis-NIR spectrophotometer (1 mm quartz cell used). Inductively coupled plasma mass spectrometry (ICP-MS DRCE Elan, Perkin Elmer) was used for quantitative

determination of platinum. Confocal laser scanning microscopy (CLSM) images were acquired by a Carl ZEISS LSM 800. Flow cytometry was recorded on a BD LSR Rortessa X20 (3 lasers).

2.2.3 Synthesis of chol-NH₂

Chol-NH₂ was synthesized according to a modified procedure.¹⁸⁰

Scheme 2. Synthesis of chol-NH₂.

Triethylamine (0.259 mL, 1.86 mmol) and N-Boc-ethylenediamine (149 mg, 0.93 mmol) dissolved in DCM (10 mL) were stirred at 0 °C. Following which, cholesteryl chloroformate (500 mg, 1.11 mmol) in DCM (10 mL) was added dropwise into the reaction solution over 30 min. The reaction was left to stir for 18 hr. DCM was then added, and the solution was washed with dilute HCl and brine. The organic layer was collected, dried over sodium sulfate and filtered. The Boc-protected product was purified with column chromatography, with the product eluting at 20% ethyl acetate/hexane. Deprotection was performed by dissolving the obtained product (200 mg, 0.35 mmol) in TFA (0.27 mL, 3.50 mmol) and DCM (6 mL) and stirred at 25°C for 6 hr. The solvent was removed, affording a colorless oil used without further purification. ¹H-NMR (300 MHz) δ (ppm) = 7.87 (bs, 3H), 5.37 (s, 1H), 5.30 (s, 1H), 4.44 (bs, 1H), 3.48– 3.23 (m, 4H), 2.30–2.19 (m, 2H), 2.02–1.81 (m, 6H), 1.56–0.85 (m, 37H). LC-MS m/z [M + H]⁺ calcd. For [C₃₀H₅₂N₂O₂]⁺ 472.76; found 473.18.

2.2.4 Synthesis of oxliPt(IV)-chol

The compound oxliPt(IV)-COOH was synthesized as reported in literature.¹³⁴

Scheme 3. Synthesis of oxliPt(IV)-chol.

For the synthesis of oxliPt(IV)-chol, a two-necked round bottle flask containing oxliPt(IV)-COOH (57 mg, 90.3 μmol) and 1,1'-carbonyldiimidazole (44 mg, 0.271 mmol) was flushed with nitrogen to maintain an inert environment. Anhydrous DMF (2 mL) was added and the solution was stirred at 60 $^{\circ}\text{C}$ for 10 min. After which, the solution was cooled to room temperature followed by the addition of chol-NH₂ (205 mg, 0.44 mmol) and triethylamine (88 mg, 0.87 mmol). After stirring overnight, solvent was removed. The obtained residue was dissolved in acetone and then precipitated with diethyl ether to give the final product. ¹H-NMR (300 MHz) δ (ppm) = 8.33 (m, 2H), 8.12 (m, 2H), 7.87 (m, 2H), 7.67 (m, 1H), 7.03 (t, 2H), 5.33 (m, 2H), 4.33–4.30 (m, 4H), 3.05–2.67 (m, 12H), 2.32–1.78 (m, 24H), 1.54–0.89 (m, 73H). LC-MS m/z [$\text{M} + 3\text{H}$]³⁺ calcd. For [C₇₆H₁₂₄N₆O₁₄Pt]⁺ 514.29; found 513.33.

2.2.5 Synthesis of CD-spermine

CD-Tos was prepared according to procedures reported in literature.¹⁸¹

Scheme 4. Synthesis of CD-spermine.

Spermine (1.53 g, 7.76 mmol) in anhydrous DMF (3 mL) was added to a solution of CD-Tos (1.0 g, 0.78 mmol) in anhydrous DMF (5 mL). The reaction was stirred under inert condition at 80 °C overnight. The resulting solution was then cooled to room temperature and the solvent was evaporated. The crude solid obtained was dissolved in deionized (DI) H₂O. The product was obtained by precipitation with acetone and collected by filtration. ¹H-NMR (300 MHz) δ (ppm) = 5.10– 5.08 (m, 7H), 3.97–3.30 (m, 47H), 2.76–2.61 (m, 18H), 2.23 (m, 12H), 1.73–1.71 (m, 10H). HRMS m/z [M + H]⁺ calcd. For [C₅₂H₉₄N₄O₃₄]⁺ 1319.57; found 1319.58.

2.2.6 Synthesis of CD-NH₂

The compound CD-NH₂ was prepared according to procedures reported in literature.¹⁸²

Scheme 5. Synthesis of CD-NH₂.

Ethylenediamine (0.311 mL, 23.3 mmol) was added to a solution of CD-Tos (300 mg, 0.23 mmol) in anhydrous DMF (3 mL). The reaction was stirred under inert conditions at 80 °C overnight. The resulting solution was then cooled to room temperature and the solvent was evaporated. The crude solid obtained was dissolved in DI H₂O. The product was obtained by precipitation with acetone and collected by filtration.

2.2.7 Reduction of oxliPt(IV)-chol prodrug and its chelation with 5'-GMP

To study the reduction products of the prodrug, oxliPt(IV)-chol (1 mM) was dissolved in sodium ascorbate aqueous solution (2.5 mM) and left to stir overnight. The resulting mixture was appropriately diluted with methanol and analyzed by HPLC and ESI-MS.

For the chelation study of Pt species with 5'-GMP, a solution of oxliPt(IV)-chol, 5'-GMP, and sodium ascorbate at final concentrations of 1 mM, 5 mM and 5 mM, respectively, was prepared and incubated at 37 °C for 12 hr. The reaction mixture was then diluted with methanol and analyzed by HPLC and LC-MS. For comparison, a solution of oxliPt(II) and 5'-GMP at final concentrations of 1 mM and 5 mM, respectively, was prepared and incubated at 37 °C for 12 hr. The solution was then diluted to suitable concentration for HPLC and ESI-MS measurements.

2.2.8 Critical aggregation concentration (CAC) determination of pre-formed nanoparticles

Stock solutions of oxliPt(IV)-chol and CD-spermine were prepared in DMSO and distilled water, respectively. oxliPt(IV)-chol was titrated into a solution of CD-spermine (5 μ M) in 1 cm cuvette to achieve different ratios. Each mixture solution was stabilized for 5 min before recording the UV-vis curve. The transmittance at 240 nm was plotted as a function of the concentration of oxliPt(IV)-chol.

2.2.9 Preparation of prodrug nanocarrier

The prodrug nanocarrier was prepared by a nanoprecipitation method. Briefly, stock solution of oxliPt(IV)-chol in DMSO was added dropwise to CD-spermine or CD-NH₂ dissolved in DI water. The mixture was sonicated at room temperature for 30 min, and CD-spermine: oxliPt(IV)-chol or CD-NH₂: oxliPt(IV)-chol nanoparticles were

formed spontaneously. The prepared nanoparticles were then dialyzed (MWCO = 2000) against deionized water to remove DMSO.

To prepare FITC-labelled nanoparticles for imaging purposes, the above procedure was carried out with the addition of FITC-labelled CD-spermine. FITC-labelled CD-spermine was obtained by reacting FITC isomer I with CD-spermine in DMSO for 2 hr, dialyzed against DI H₂O for 2-3 days and finally lyophilized. Stock solutions of oxliPt(IV)-chol in DMSO, CD-spermine (or CD-NH₂) and FITC-labelled CD-spermine in H₂O were mixed in the molar ratio 1: 1.4: 0.6 and then sonicated at room temperature for 30 min. The prepared nanoparticles were then dialyzed (MWCO = 2000) against DI water. The amount of FITC dye adsorbed on the nanoparticles was indirectly calculated using FITC calibration curve based on the amount of free FITC in dialysate solution.

2.2.10 Drug release study

To study the drug release behavior in physiological conditions and cellular environments, CD-spermine: oxliPt(IV)-chol (1 mM) was dissolved in phosphate buffer solution (PBS, pH 7.4, 2 mL). A dialysis bag (MWCO = 2000 Da) used to contain the solution was immersed into PBS (78 mL) comprising either 0 mM or 5 mM sodium ascorbate. The release was conducted at 37 °C. The dialysate (1 mL) was withdrawn at predefined time and equivalent volume of incubation medium was added to ensure the total volume of solution was maintained. The amount of Pt in the collected samples was measured by ICP-MS. The percentage of drug released is expressed as the amount of accumulative drug in the incubation medium to the total drug in the original nanoparticles.

2.2.11 Cell culture

HCT116 cells were cultured in DMEM medium supplemented with 10% (v/v) FBS, 0.03% L-glutamine and 1% penicillin-streptomycin under 5% CO₂ atmosphere at 37 °C. DMEM was replaced every two days.

2.2.12 In vitro cytotoxicity assay

HCT116 cells were seeded into 96-well plate with an initial cell seeding density of 1×10^4 cells/well. The cells were then incubated with various drug formulations of oxliPt(II) and CD-spermine : oxliPt(IV)-chol at final equivalent Pt concentrations ranging from 1 to 100 μ M. After incubation at 37 °C for 24 and 48 hr, the incubation medium was removed. Fresh culture medium containing MTT (5 mg/mL, 100 μ L) was added into each well, and the cells were incubated at 37 °C for another 4 hr. DMSO (100 μ L) was added to dissolve the formazan crystals. The optical density (OD) was read at 490 nm using a microplate reader (infinite M200, TECAN). The control was taken as the culture medium in the absence of samples. The normalized cell viability in % was calculated according to cell viability = $\frac{OD_{\text{sample}} - OD_{\text{blank}}}{OD_{\text{control}} - OD_{\text{blank}}} \times 100\%$. The results were obtained from 5 replicates for each sample and expressed as means.

2.2.13 Determination of cellular platinum contents

HCT116 cells were seeded in 24-well plates at a density of 10^5 cells per well. Fresh culture medium containing oxliPt(II) or CD-spermine : oxliPt(IV)-chol was incubated with the cells at 37 °C for 2 or 12 hr. The platinum concentration of each sample was regulated at 25 μ M. After the stipulated incubation period, the cells were washed three times with PBS, harvested by trypsination and collected by ultra-centrifugation. Thereafter, 1 mL cell lysis buffer (Invitrogen cell extraction buffer) was used to lyse the cell pellets. The cell lysis solution (100 μ L) of each sample was used in

the measurement of Pt content by ICP-MS. The cell lysis solution (25 μ L) was used for the determination of the protein content by bicinchoninic acid (BCA) protein assay kit (Pierce Thermo Scientific). The amount of Pt content was expressed as nanograms of Pt per milligram of total proteins.

2.2.14 Cellular uptake study

HCT116 cells were seeded on glass cover slip and allowed to grow for 24 hr. The cells were then incubated with cell culture medium containing 10 μ M FITC-labelled CD-spermine: oxliPt(IV)-chol or CD-NH₂: oxliPt(IV)-chol for 30 min and 4 hr. After which, the cells were washed with PBS three times, and Hoechst 33342 was used to stain the nucleus before fixation. After the treatment, cells were imaged by CLSM (ZEISS LSM 800). Fluorescence was examined under excitation at 405 nm for Hoechst 33342 and 488 nm for FITC.

For the flow cytometric study, HCT116 cells were seeded in a 6-well tissue culture plate (2 mL medium) and cultured for 24 hr. Samples were added in the culture medium and incubated for 4 and 18 hr. After the treatment, samples-containing medium was removed, and the cells were harvested. The cells collected were washed with PBS twice, centrifuged, finally resuspended in PBS (1 mL) and subjected to flow cytometry analysis.

Experiments for the spermine competition assay were carried out as described above, except that cells were pretreated with spermine for 4 hr before incubation with the nanoparticles.

2.3 Results and Discussion

2.3.1 Synthesis and characterization

The oxliPt(IV)-chol prodrug as the guest molecule was prepared by reacting ethylenediamine-modified cholesterol with oxliPt(IV)-COOH (Schemes 2 and 3). The chemical structures of the intermediates and final product were fully characterized by ^1H NMR, HR-MS, FTIR and elemental analysis (Figure 1, Table 1).

Figure 1. FTIR spectra for intermediate and final products in the synthesis of oxliPt(IV)-chol.

Table 1. CHN elemental analysis for intermediate and final products in the synthesis of oxliPt(IV)-chol.

Anal. Calcd. for	Calcd [%]			Found [%]		
	C	H	N	C	H	N
oxliPt(II)	24.18	3.53	7.05	20.59	3.85	6.09
oxliPt(IV)-OH	22.27	3.71	6.5	18.83	3.72	5.32
oxliPt(IV)-COOH	30.43	3.8	4.44	23.13	3.70	4.04
oxliPt(IV)-chol	59.26	8.06	5.46	48.64	7.57	4.91

CD-spermine as the host molecule was attained by reacting excess spermine with tosylated β -cyclodextrin (Scheme 4). Its structure was well characterized with ^1H NMR and mass spectrometry. A decrease in zeta potential from -20.1 mV of β -cyclodextrin to

-6.8 mV of CD-spermine also indicated the successful synthesis of CD-spermine. Such changes can be attributed to the reduction of the surface anionic hydroxyl groups on β -cyclodextrin after conjugation to spermine which carries highly positive amino functionalities.

Before preparing the self-assembled nanoparticles, we investigated whether the molecular prodrug could undergo the reduction to give active oxaliplatin(II) species.

Scheme 6. Schematic illustration of possible reduction products of oxliPt(IV)-chol.

High-performance liquid chromatography (HPLC) and liquid chromatography–mass spectrometry (LC-MS) were used to study the reduction reaction products of oxliPt(IV)-chol incubated with sodium ascorbate. The reduction of the prodrug was expected to lead to the dissociation of the axial cholesterol ligand and produce active oxaliplatin (Scheme 6).

Figure 2. HPLC analysis of oxliPt(II), and oxliPt(IV)-chol with and without the treatment of sodium ascorbate. The retention time at 9.5 min corresponds to oxliPt(II), while the retention time at 17 min corresponds to oxliPt(IV)-chol. The absorbance channel was set at 240 nm.

HPLC monitoring (Figure 2) revealed that the reaction outcome was indeed consistent with our expectation. Peak corresponding to oxliPt(IV)-chol (retention time: 17 min) disappeared after the reduction with sodium ascorbate, whilst there was an emergence of a new peak corresponding to oxliPt(II) (retention time: 9.5 min). In addition, the LC-MS spectrum (not shown) of the reaction solution confirmed the presence of active oxaliplatin drug and cholesterol ligand after the reduction reaction. These results indicate that our designed molecule could indeed function as a redox-responsive prodrug. It is pharmacologically inactive and requires reductive elimination to achieve the active form.

The cytotoxicity of oxaliplatin arises from the formation of an adduct with DNA nucleobases, specifically binding to the N7 guanine.¹⁸³ Hence, the ability of the released oxaliplatin drug binding with DNA was investigated. 5'-GMP was used to stimulate the interaction mechanism of the drugs with DNA (Scheme 7), where the adduct products were studied by HPLC and ESI-MS (Figure 3).

Scheme 7. Schematic illustration of possible Pt conjugates upon reduction of oxliPt(IV)-chol in the presence of 5'-GMP and sodium ascorbate.

Figure 3. (A) HPLC chromatograms of 5'-GMP, oxliPt(II) and prodrug oxliPt(IV)-chol with and without the incubation with sodium ascorbate. (B) Tabulated mass of the chromatogram peaks observed at 8.4 and 10.2 min measured with ESI-MS.

Taking the result of direct reaction of oxaliplatin(II) with 5'-GMP as the reference, HPLC peaks at the retention time of 8.4 min and 10.2 min were found, corresponding to the adduct of oxaliplatin(II) with one and two 5'-GMP, respectively. The same peaks appeared for the reaction between the reduced oxaliPt(IV)-chol prodrug and 5'-GMP, indicating that the released platinum species from the prodrug was most likely oxaliplatin(II) and could possibly exhibit similar behavior in vitro.

2.3.2 Preparation and characterization of nanocarrier

Figure 4. (A) Job's plot indicating a 2:1 binding ratio between CD-spermine and oxliPt(IV)-chol, (B) ^1H NMR comparison of CD-spermine and CD-spermine: oxliPt(IV)-chol complex in $\text{D}_2\text{O}/\text{DMSO}-d_6$ (5/5, v/v), (C) Plot of concentration-dependent optical transmittance changes of oxliPt(IV)-chol at 240 nm in the nanoparticle formation, and (D) TEM image of CD-spermine: oxliPt(IV)-chol nanoparticles.

To prepare CD-spermine: oxliPt(IV)-chol nanoparticles, we first quantitatively investigated the binding stoichiometry between CD-spermine and oxliPt(IV)-chol. Job's plot was obtained to determine the host-guest inclusion ratio of 2:1 (Figure 4A). ^1H NMR spectra of CD-spermine revealed changes in the chemical shifts of characteristic protons before and after binding with 0.5 equivalent oxliPt(IV)-chol, attributing to the complexation-induced shielding effect (Figure 4B). This result demonstrated that the

cholesterol moiety of the prodrug was included into the hydrophobic cavity of the cyclodextrin host to form a supramolecular complex. Based on the obtained binding ratio, CD-spermine: oxliPt(IV)-chol nanoparticles were prepared by mixing oxliPt(IV)-chol in DMSO with CD-spermine aqueous solution. The critical aggregation concentration (CAC) of the complex was calculated to be 64.9 μM by monitoring the change in the optical transmittance at 240 nm upon increasing the concentration of oxliPt(IV)-chol (Figure 4C). Transmission electron microscopy (TEM) images revealed a well-dispersed spherical morphology of our nanoparticles (Figure 4D). The nanoparticles exhibited a narrow size distribution with an average hydrodynamic diameter of 130 nm as measured by dynamic light scattering. The oxaliplatin encapsulation efficiency was 22.7 wt % as measured by inductively coupled plasma - mass spectrometry (ICP-MS).

2.3.3 Drug release study

To study the redox-triggered drug release property of the CD-spermine: oxliPt(IV)-chol nanoparticles, cumulative drug release of Pt from the nanoparticles was studied using ICP-MS (Figure 5).

Figure 5. Cumulative platinum drug release profile from CD-spermine: oxliPt(IV)-chol nanoparticles in PBS solution (pH 7.4) and in 5 mM sodium ascorbate solution. Vc: sodium ascorbate.

5 mM sodium ascorbate (Vc) solution was used to simulate the reducing environment inside cancer cells.¹⁸⁴⁻¹⁸⁵ In the absence of sodium ascorbate, there was only around 25% Pt detected after 24 hr, suggesting that the nanoparticles were relatively stable in physiological conditions. In comparison, in 5 mM sodium ascorbate, the released amount of Pt drastically increased to 63%, illustrating that the CD-spermine: oxliPt(IV)-chol nanoparticles could be an effective drug delivery system. The encapsulated Pt(IV) prodrug would only be released in the presence of enough amount of reducing agents in the intercellular matrix of cancer cells. This feature ensured minimal drug leakage before the cellular uptake for reduced side effects.

2.3.4 In-vitro anticancer activity

Figure 6. (A) CLSM images of HCT116 cells incubated with FITC-labelled CD-spermine: oxliPt(IV)-chol nanoparticles. Cell nuclei were stained by H33342. NP: nanoparticles. Scale bar: 20 μm. (B) Cellular accumulation of CD-spermine: oxliPt(IV)-chol in HCT116 cells after 2 and 18 hr incubation, measured by flow cytometry.

The cellular uptake of CD-spermine: oxliPt(IV)-chol nanoparticles was evaluated using fluorescein isothiocyanate (FITC)-labelled nanoparticles. Figure 6A

shows the confocal laser scanning microscopy (CLSM) images of colorectal cancer HCT116 cells after the incubation with the nanoparticles at 2 and 18 hr. After 2 hr of incubation, a little green fluorescence was observed inside the cells, implying that the nanoparticles hardly entered the cells. On the other hand, enhanced cellular FITC fluorescence after 18 hr incubation was evident that the CD-spermine: oxliPt(IV)-chol nanoparticles could be internalized upon time. Flow cytometry also confirmed that the uptake of the nanoparticles was time-dependent (Figure 6B).

Figure 7. Cell viability of HCT116 cells after incubation with different concentrations of free CD-spermine for 24 and 48 hr.

Subsequently, the anticancer efficacy of the nanoparticles was evaluated by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. The cytotoxicity of CD-spermine alone on HCT116 cells was first evaluated and results showed that it exhibited high biocompatibility and hence would have minimal effect on the cytotoxicity results of the nanocarrier system (Figure 7).

Figure 8. In vitro cytotoxicity of oxliPt(II) and CD-spermine: oxliPt(IV)-chol nanoparticles incubated with HCT116 cells for (A) 24 hr and (B) 48 hr.

Then, oxliPt(II) and CD-spermine: oxliPt(IV)-chol nanoparticles with various Pt concentrations from 1 to 100 μM were incubated with HCT116 cells for 24 and 48 hr. As shown in Figure 8, CD-spermine: oxliPt(IV)-chol nanoparticles exhibited both concentration- and time-dependent cytotoxicity. The nanoparticles presented higher cytotoxicity to HCT116 cells as compared to free oxliPt(II) possibly due to enhanced cellular accumulation of the nanoparticles. The internalization of the drug-containing nanoparticles via transporter-mediated process is more efficient than simple molecular diffusion of the free platinum drug within the same incubation time. Thereby, a better cell killing effect was observed for the nanoparticles.

Figure 9. Cellular Pt accumulation in HCT116 cells after 2 and 12 hr incubation with CD-spermine : oxliPt(IV)-chol, measured by ICP-MS.

ICP-MS results of the cellular Pt uptake showed that the accumulation of CD-spermine: oxliPt(IV)-chol nanoparticles was about 30-folds greater than that of oxliPt(II) at 12 hr incubation (Figure 9), verifying that the enhanced endocytosis of the nanoparticles was due to its nanosize and targeting effect of spermine. Under longer incubation time, however, the therapeutics effect of CD-spermine: oxliPt(IV)-chol nanoparticles ($IC_{50} = 41.2 \mu M$ at 48 hr) was not significantly better than the free oxaliplatin ($IC_{50} = 37.7 \mu M$ at 48 hr). This observation might be the result of time-dependent drug release characteristics of the nanoparticles. Despite enhanced cellular uptake, the nanoparticles had to undergo the reduction of the Pt(IV) species to its active Pt(II) form and subsequent dissociation and diffusion from the nanoparticles before inhibiting the cell growth. Moreover, it was plausible that not all of oxliPt(IV)-chol prodrug was reduced under this condition, offsetting the effect of increased nanoparticle accumulation and mitigating corresponding cytotoxicity in vitro.

Figure 10. CLSM images of HCT116 cells incubated with 10 μM CD-spermine: oxliPt(IV)-chol or CD-NH₂: oxliPt(IV)-chol nanoparticles for 4 hr. Nuclei were stained with H33342. NP: nanoparticles. Scale bar: 20 μm .

To confirm the targeting ability of the spermine ligand, FITC labelled CD-NH₂: oxliPt(IV)-chol nanoparticles as a control group were also prepared for comparison. It was anticipated that the cellular uptake of these nanoparticles would be lower as the

ethylenediamine ligand of CD-NH₂ cannot be recognized by polyamine transporter, thereby reducing the rate of cellular internalization. As shown in Figure 10, the fluorescence intensity of FITC was stronger in the CLSM images of the CD-spermine: oxliPt(IV)-chol treated cells than those treated with the control nanoparticles, indicating that the spermine modification could promote cellular uptake probably through the transporter-mediated pathway.

Figure 11. (A) In vitro cytotoxicity of CD-spermine: oxliPt(IV)-chol and CD-NH₂: oxliPt(IV)-chol nanoparticles incubated with HCT116 colorectal cancer cells for 48 hr. Competitive intracellular uptake of CD-spermine: oxliPt(IV)-chol nanoparticles by spermine: (B) CLSM images and (C) flow cytometry profiles of HCT116 cells without and with pre-treatment of spermine prior to the incubation with FITC-labelled CD-spermine: oxliPt(IV)-chol nanoparticles. Cell nuclei were stained by H33342. NP: nanoparticles. Scale bar: 50 μm.

Furthermore, the HCT116 cell viability after incubating with CD-spermine: oxliPt(IV)-chol nanoparticles for 48 hr were shown to be lower than that of CD-NH₂: oxliPt(IV)-chol nanoparticles (Figure 11A). The higher cytotoxicity of the CD-spermine: oxliPt(IV)-chol nanoparticles could be due to their increased cellular uptake via polyamine transporter-mediated endocytosis. In addition, competitive inhibition of the polyamine transporter was carried out by pretreating cells with spermine prior to the incubation with CD-spermine: oxliPt(IV)-chol nanoparticles. As observed in Figure 11B and 11C, fluorescence intensity of cells incubated with FITC labelled CD-spermine: oxliPt(IV)-chol nanoparticles was significantly higher than that of the

spermine-pretreated group, once again demonstrating that the uptake of the nanoparticles was closely related to the polyamine transporter.

2.4 Conclusion

In summary, redox-responsive supramolecular oxaliplatin(IV) prodrug nanoparticles were successfully constructed by the host-guest interaction. OxliPt(IV)-chol and CD-spermine not only serve as the building units, but also the cargo and active targeting component, respectively. Thus, the supramolecular nanoparticles prepared using this unique strategy integrated the advantages of active targeting, prodrug, and carrier-free, while simplifying the assembly complexity. The prodrug could be released under reductive cellular microenvironment, allowing for controlled drug release. Importantly, in vitro experiments have confirmed that the CD-spermine: oxliPt(IV)-chol nanoparticles offered comparable cytotoxicity as free platinum drugs and provided the specificity to cancer cells based on their redox-responsiveness and targeting capability. This work presented an alternative strategy to the development of smart supramolecular nanocarriers, showing a great potential in the field of controlled drug release and delivery toward the cancer treatment.

Chapter 3

Self-assembled Oxaliplatin(IV) Prodrug-Porphyrin Conjugate for Combinational Photodynamic Therapy and Chemotherapy

3.1 Introduction

Despite being the mainstay in cancer therapy, chemotherapy still suffers from drawbacks such as poor aqueous solubility of some chemotherapeutic drugs and poor tumor specificity leading to systemic cytotoxicity, greatly impairing the treatment efficiency.¹⁸⁶⁻¹⁸⁷ Extensive research have been devoted to developing drug delivery systems ranging from polymers, liposomes to inorganic nanomaterials to achieve tumor-targeted drug accumulation and controlled drug release.^{14, 40-41, 188} To reduce systemic toxicity and enable stimuli responsive activation of therapeutic drugs, prodrugs are harnessed in place of conventional drugs. In particular, the fabrication of prodrug-based self-delivery systems allows high drug loading efficiency, possesses intrinsic biodegradability, and has low/no carrier-induced immunogenicity.^{76, 189-190} Furthermore, the inclusion of prodrugs in the delivery systems, which remain inactive until activated by specific stimuli in the tumor microenvironment, ensures controlled tumor-selective drug release.^{87, 191-194}

Chemotherapy alone is often insufficient to ensure optimal success for cancer treatment due to the drug resistance that arises from tumor heterogeneity and diverse pathogenesis pathways.¹⁹⁵⁻¹⁹⁶ To address such dilemma, combination therapy has been established to be a promising strategy, and many studies have reported the combination of two or more therapeutic treatments to increase therapeutic efficiency.¹⁹⁷⁻¹⁹⁹ A common strategy is to combine photodynamic therapy (PDT) with chemotherapy.²⁰⁰⁻²⁰² PDT involves the excitation of photosensitizers (PSs) with light to produce cytotoxic reactive oxygen species (ROS) to induce significant oxidative damage of biological substrates, thereby leading to ultimate cell death via apoptosis or necrosis.²⁰³⁻²⁰⁵ Largely

different yet complementary mechanisms of action from chemotherapeutic drugs and PDT agents could maximize therapeutic efficacy with lesser systemic toxicity through administration of lower drug dosage. In addition, PDT depends on external stimuli for the activation of PSs, allowing spatial and temporal control of the treatment. Collectively, combining PDT and chemotherapy has become an attractive combinational therapy regime for cancer.

In spite of numerous efforts made to construct therapeutic platforms that co-deliver PSs and prodrugs,²⁰⁶⁻²¹⁰ little research has been conducted on investigating an effective way to realize the optimal loading ratio of PSs and prodrugs for satisfactory cancer treatment. Some nanosystems reported are based on the covalent conjugation of drugs to PSs.^{88, 165, 211} Although high loading amount is achieved, it only allowed a fixed loading ratio between drugs and PSs, which might not be optimal for the treatment. Physical encapsulation of the payloads in organic nanocarriers such as polymeric nanoparticles or liposomes often suffers from unpredictable drug loading amount and premature leakage.²¹²⁻²¹³ In addition, the mass ratio of loaded drugs to nanocarriers is also low. To circumvent these issues, inorganic materials such as titanium dioxide and zinc oxide with intrinsic photosensitizing capabilities have been developed as platforms to deliver chemodrugs.²¹⁴⁻²¹⁷ While high drug encapsulation efficiency and superior therapeutic efficacy of these systems seem advantageous, recent research uncovered that the use of such inorganic materials might accelerate cancer metastasis.¹¹ On the other hand, the π - π stacking and rigid structures of typical organic PSs may lead to aggregation in aqueous media, resulting in reduced ROS production efficiency. Fortunately, modifications of the hydrophobic PSs with spacer molecules or hydrophilic polymers could prevent them from self-quenching.²¹⁸⁻²¹⁹ Taking these challenges into

account, it is thus urgent to fabricate an organic nanocarrier capable of co-loading chemotherapeutic drugs and PSs with a predictable and controllable ratio.

Scheme 1. Chemical structures and schematic illustration of PEG-Por-CD and oxliPt(IV)-ada.

Here, we design a self-assembled supramolecular nanocarrier (PEG-Por-CD: oxliPt(IV)-ada) that can simultaneously deliver PS and oxaliplatin(IV) prodrug in a controlled fashion for combinational therapy. Utilizing the host-guest complexation strategy, the nanocarrier was assembled between adamantane-modified oxaliplatin prodrug (oxliPt(IV)-ada) and β -cyclodextrin-conjugated porphyrin PS (PEG-Por-CD) (Scheme 1). The well-established host-guest binding between adamantane and β -cyclodextrin allows convenient control of the ratio between the two therapeutic components.^{168, 220-221} By converting to a prodrug, oxliPt(IV)-ada remains inactive until it is reduced specifically in tumor cells, thus decreasing side effects to normal cells.¹¹⁴⁻¹¹⁵ Polyethylene glycol (PEG) was conjugated to the porphyrin core, which not only enhances the water solubility and maintains the porphyrin photoactivity by eliminating

self-quenching in aqueous medium but also contributes to the amphiphilicity of the host-guest complex to facilitate the self-assembly into nanoparticles (Scheme 2A).²²²⁻²²⁵

Scheme 2. (A) Schematic illustration for the self-assembly formation of PEG-Por-CD: oxliPt(IV)-ada nanoparticles, and (B) Its application for combinational photodynamic and chemotherapy of cancer.

Upon the internalization into cancer cells, intracellular bioreductants could reduce the prodrug to release active oxaliplatin for chemotherapy,²²⁶⁻²²⁸ while light irradiation on the porphyrin PS generates ROS for PDT (Scheme 2B). The preparation of PEG-Por-CD: oxliPt(IV)-ada with an optimized drug loading ratio was studied, and its anticancer efficacy was investigated both in-vitro and in-vivo. The PEG-Por-CD: oxliPt(IV)-ada nanocarrier displays a number of outstanding features as the next

generation therapeutics, including: (1) precise control over the drug and PS loading ratios, (2) fabrication as self-delivery nanoparticles to eliminate the toxicity concern of additional carrier, (3) high efficiency of $^1\text{O}_2$ generation in aqueous medium due to the introduction of PEG and β -cyclodextrin to prevent the self-aggregation of original hydrophobic PSs, and (4) site-specific cytotoxicity based on redox-responsiveness for programmed degradation of the nanocarrier and activation of the prodrug payload. Therefore, the obtained PEG-Por-CD: oxliPt(IV)-ada nanoparticles with an optimized loading ratio show to be a promising therapeutic system in-vitro and in-vivo, which benefitted from the superiority of unique nanostructure and combinational therapy.

3.2 Experimental Section

3.2.1 Materials and reagents

All solvents and inorganic salts were purchased from Sigma-Aldrich. 1,1'-Carbonyldiimidazole (CDI), hydrogen peroxide (H_2O_2), poly(ethylene glycol) methyl ether (mPEG, MW = 2000), sodium L-ascorbate, succinic anhydride, 5,10,15,20-tetrakis(4-hydroxyphenyl)-21H,23H-porphine (THPP), p-toluenesulfonyl chloride (p-TsCl), triethylamine (TEA), and zinc acetate were purchased from Sigma-Aldrich. Propargyl bromide was purchased from Alfa Aesar. Oxaliplatin was purchased from Aaron Pharmatech Ltd. (Shanghai, China). 1-Adamantanecarbonyl chloride, N-Boc-ethylenediamine, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide and disodium dihydrogen ethylenediaminetetracetate dihydrate (EDTA) were bought from TCI. DMEM high glucose, fetal bovine serum (FBS), penicillin-streptomycin solution and RPMI 1640 w/ L-glutamine medium were purchased from Alantis Bioscience. 4T1 and HT116 cells were obtained from ATCC. Hoechst 33342 staining dye was bought from Abcam. Annexin V-FITC Apoptosis Detection Kit, Calcein AM, Image-iT™ LIVE Green Reactive Oxygen Species Detection Kit, propidium iodide (PI) and Singlet Oxygen Sensor Green (SOSG) were purchased from Invitrogen, ThermoFisher Scientific.

3.2.2 Characterization

^1H NMR spectra were probed on 300 MHz or 400 MHz Bruker spectrometer using CDCl_3 or deuterated DMSO as solvent. ThermoFinnigan LCQ quadrupole ion trap mass spectrometer was used to record electronic spray ionization (ESI) mass spectra. Transmission electron microscope (TEM) images were collected on a JEM-1400 (JEOL) operated at 100 kV. Zeta potential and dynamic light scattering (DLS) measurements were measured using Zetasizer, Malvern Instrument Ltd. High performance liquid

chromatography (HPLC) analysis was performed with Agilent Technologies 1200 Series using the zorbax RX-C8 (5 μ m, 4.6 $\times$ 150 mm) column with UV detection at 230 nm. Gel permeation chromatography (GPC) analysis with DMF as an eluent was performed on a Shimadzu i-Series Plus liquid chromatograph LC-2030c Plus (Kyoto, Japan) equipped with two Shodex KF-804 mixed gel columns (300 $\times$ 8.0 mm; bead size = 7 mm). The flow rate was 0.5 mL/min (DMF) (40 $^{\circ}$ C). The DMF eluent contained LiBr (10 mM). Sample detection and quantification were conducted using a refractive index detector (RID-20A) for DMF-GPC. The column system was calibrated with standard linear polyethylene glycol (PEGs) in the DMF systems. UV-vis measurements were obtained on a Shimadzu UV-3600 UV-Vis-NIR spectrophotometer (1-mm quartz cell used). Fluorescent measurements were obtained using Shimadzu RF-5301 spectrofluorophotometer. Inductively coupled plasma mass spectrometry (ICP-MS DRCE Elan, Perkin Elmer) was used for quantitative determination of platinum. Confocal laser scanning microscopy (CLSM) images were acquired by a ZEISS LSM 800 confocal microscope. Flow cytometry was recorded on a BD LSR Rortessa X20 (3 lasers). Hematoxylin and eosin (H&E) staining images were obtained with a Leica DM5500B upright wide-field fluorescence and BF microscope.

Schematic synthetic procedures of oxliPt(IV)-ada and PEG-Por-CD are shown in Schemes 3 and 4.

3.2.3 Synthesis of oxliPt(IV)-ada

Scheme 3. Synthesis route of oxliPt(IV)-ada.

oxliPt(IV)-ada was obtained by coupling oxliPt(IV)-COOH with ada-NH₂. ada-NH₂ was prepared by reacting 1-adamantanecarbonyl chloride with N-Boc-ethylenediamine. Briefly, N-Boc-ethylenediamine (336 mg, 2.10 mmol) in DCM (10 mL) was added TEA (0.58 mL, 4.19 mmol) and stirred at 0 °C. 1-adamantanecarbonyl chloride in DCM (10 mL) was added dropwise to the reaction mixture and the reaction was stirred overnight. Subsequently, DCM was added to dilute the reaction mixture, followed by washing with dilute HCl and brine. The organic layer was dried with sodium sulfate, filtered and evaporated to dryness. The crude product was purified by column (hexane: EA, 100:45) to give ada-NH-Boc as a white product. ¹H-NMR (300 MHz; CDCl₃, δ): 6.34 (s, 1H), 4.87 (s, 1H), 3.31 (m, 4H), 2.03 (m, 3H), 1.84 (m, 6H), 1.68 (m, 6H), 1.45 (s, 9H); ESI-MS *m/z* [M + H]⁺ calcd for [C₁₈H₃₀N₂O₃]⁺ 322.45; found 322.13. ada-NH-Boc was then deprotected with TFA in DCM for 3 hr before reaction with oxliPt(IV)-COOH.

oxliPt(IV)-COOH was synthesized using a reported procedure.¹³⁴ oxliPt(IV)-COOH (20 mg, 31.7 μ mol) and CDI (15.4 mg, 95.1 μ mol) were dissolved in anhydrous DMF (2 mL) and stirred under nitrogen for 10 mins at 60 °C. After which, the solution was cooled to room temperature followed by the addition of ada-NH₂ (34 mg, 152.9 μ mol) and triethylamine (31 mg, 0.306 mmol). After stirring overnight, solvents were removed. The obtained residue was first dissolved in acetone and then precipitated thrice with diethyl ether to give oxliPt(IV)-ada. ¹H-NMR (300 MHz; d-DMSO, δ): 8.35 (m, 2H), 8.13 (m, 2H), 7.87 (m, 2H), 7.36 (m, 2H), 3.06 (m, 8H), 2.64 (m, 2H), 2.27 (m, 4H), 2.08 (m, 4H), 1.95 (m, 6H), 1.73 (m, 12H), 1.66 (m, 12H), 1.55- 1.04 (m, 10H); ESI-MS m/z [M + H]⁺ calcd for [C₄₂H₆₄N₆O₁₂Pt]⁺ 1040.09; found 1040.18.

3.2.4 Synthesis of PEG-Por-CD

Scheme 4. Synthesis route of PEG-Por-CD.

THPP was first modified with a terminal alkyne to give THPP- $\equiv$. Briefly, THPP (500 mg, 0.74 mmol) and K₂CO₃ (305 mg, 2.21 mmol) was stirred in anhydrous DMF (10 mL) under reflux for 1 hr. Propargyl bromide (105 mg, 0.88 mmol) in DMF was then dropwise added and the reaction was stirred overnight. The crude reaction was

quenched with water, followed by extraction with DCM. The organic layers collected was washed with brine, dried with anhydrous sodium sulfate and filtered. The crude product was purified with column (DCM: MeOH, 100: 2) to give purple-blue solid as THPP- $\equiv$. $^1\text{H-NMR}$ (400 MHz; CDCl_3 , δ): 8.86 (s, 8H), 8.14 (d, 2H, 8.8 Hz), 8.07 (d, 6H, 8.4 Hz), 7.37 (d, 2H, 8.8 Hz), 7.23 (d, 6H, 8.4 Hz), 4.99 (d, 2H, 2.4 Hz), 2.69 (t, 1H, 2.4 Hz); ESI-MS m/z $[\text{M} + \text{H}]^+$ calcd for $[\text{C}_{47}\text{H}_{32}\text{N}_4\text{O}_4]^+$ 716.8; found 716.4.

The obtained THPP- $\equiv$ was next reacted with tosylated mPEG₂₀₀₀. Tosylated mPEG₂₀₀₀ was prepared according to a reported procedure.²²⁶ THPP- $\equiv$ (300 mg, 0.42 mmol) was mixed with K_2CO_3 (525 mg, 3.8 mmol) in anhydrous DMF (6 mL) for 1 hr under reflux. Then, tosylated mPEG₂₀₀₀ (2.9 g, 1.34 mmol) in anhydrous DMF (3 mL) was slowly added to the above-activated THPP molecules. After reacting for 24 hr under reflux, the solvent was removed and resulting residue was dissolved in deionized water and dialyzed (MWCO: 3000 Da) against deionized water for three days before lyophilization to give PEG-Por- $\equiv$ as a purple powder. $^1\text{H-NMR}$ (300 MHz; $d\text{-DMSO}$, δ): 8.86 (s, 8H), 8.16- 8.09 (m, 8H), 7.45- 7.38 (m, 8H), 5.10 (m, 2H), 4.40- 3.23 (m, 560H).

Finally, a click reaction was performed to obtain PEG-Por-CD. PEG-Por- $\equiv$ was chelated with Zn^{2+} to prevent copper-induced PDT quenching in the subsequent click reaction. Briefly, PEG-Por- $\equiv$ (355 mg, 52.9 μmol) was heated under reflux with excess zinc acetate (100 mg, 0.529 mmol) in chloroform: MeOH (1:1, 14 mL) for 3 hr. The solvents were then removed under reduced pressure to give PEG-ZnPor- $\equiv$ without further purification.

CD- N_3 was prepared according to procedures reported in literature.²²⁹ PEG-ZnPor- $\equiv$ (400 mg, 59 μmol), CD- N_3 (137 mg, 0.118 mmol) and $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (88 mg, 0.354 mmol) were dissolved in THF: H_2O (1:1, 40 mL) under N_2 protection. Sodium

ascorbate (234 mg, 1.18 mmol) was added and the reaction proceeded at 50 °C for 2 days. The reaction mixture was filtered, and the collected filtrate was rotavaped. The obtained residue was dissolved in deionised water and dialyzed against EDTA-containing water for three days, then against deionised water for three days. The final product PEG-Por-CD was obtained by lyophilization. ¹H-NMR (300 MHz; d-DMSO, δ): 8.86 (s, 8H), 8.16- 8.09 (m, 8H), 7.45- 7.38 (m, 8H), 5.10 (m, 2H), 5.80- 5.60 (m, 7H), 4.84- 4.40 (m, 32H), 3.94- 3.23 (m, 575H); ESI-MS *m/z* calcd for C₃₆₂H₆₄₇N₇O₁₇₄ (M+11H)¹¹⁺ 717.45; found 717.41.

3.2.5 Reduction of oxliPt(IV)-ada prodrug

To study the reduction products of the prodrug, oxliPt(IV)-ada (1 mM) was dissolved in sodium ascorbate aqueous solution (2.5 mM) and left to stir at 37 °C. Aliquots were taken at 12 and 24 hr, diluted with methanol and analyzed by high-performance liquid chromatography (HPLC). The run was conducted using acetonitrile/water binary system containing 0.05% trifluoroacetic acid as the mobile phase with a flow rate at 2 mL min⁻¹. The analysis was performed with gradient elution method starting with 95% water to 30% water over a period of 10 min.

3.2.6 Preparation of nanocarriers

The nanocarrier was prepared by a nanoprecipitation method. Briefly, stock solutions of oxliPt(IV)-ada and PEG-Por-CD were prepared in dimethyl sulfoxide (DMSO) and deionized water, respectively. Appropriate volumes were taken from the stock solutions and mixed in glass vials, which were sonicated at room temperature for 10 min and dialyzed (molecular weight cut-off (MWCO) = 2000) against deionized water to remove DMSO.

3.2.7 Critical aggregation concentration determination of nanoparticles

Stock solutions of oxliPt(IV)-ada and PEG-Por-CD were prepared in DMSO and distilled water, respectively. Solutions with varying concentrations of oxliPt(IV)-ada: PEG-Por-CD at a fixed ratio of 1:2 were prepared. Each mixture solution was stabilized for 5 min before the UV-vis curve was recorded. The transmittance at 430 nm was plotted as a function of the concentration of PEG-Por-CD.

3.2.8 Optimization of drug loading ratio

Nanoparticles containing oxliPt(IV)-ada and PEG-Por-CD were prepared as the abovementioned procedures, with final molar ratios of the drug : PS ranging from 0 to 32 : 1. Cell viability of the prepared nanoparticles, PEG-Por-CD plus light irradiation, and oxliPt(IV)-ada was tested with the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Results obtained were analyzed with the CalcuSyn software (Biosoft, Cambridge, UK) to calculate the combination index (CI) value.²³⁰ CI values of less than 1 indicate drug synergism, value equal to 1 indicates additive activity, and values more than 1 indicate antagonism.

3.2.9 Detection of singlet oxygen

Method for singlet oxygen detection was carried out according to the manufacturer's instructions. Stock singlet oxygen sensor green (SOSG) solution (12.5 μ L) was added to the nanoparticle solution (2 mL) containing porphyrin unit (6.25 μ M). Next, the sample was irradiated with a 630 nm laser at a power density of 50 mW cm⁻². The same concentration of free PEG-Por-CD in water under laser irradiation and water were used as the control samples. The fluorescence intensity of SOSG at 525 nm was recorded with an excitation wavelength of 494 nm.

3.2.10 Drug release study

To study the drug release behavior in physiological conditions, a dialysis bag (MWCO: 1000 Da) containing the nanoparticles (containing 20 μ M Pt in 1 mL phosphate buffer solution (PBS), pH 7.4) was immersed in PBS (49 mL). The release was conducted at 37 °C. Fixed volume of the incubation medium (1 mL) was withdrawn at predefined time and equivalent volume of fresh PBS was added to ensure the total volume of the solution was maintained. To mimic cellular reducing conditions, similar procedure was followed in the presence of 5 and 10 mM sodium ascorbate aqueous solution. The amounts of Pt in the collected samples were measured by inductively coupled plasma mass spectrometry (ICP-MS). The percentage of released drug is expressed as the amount of accumulative drug in the incubation medium to the total drug in the original nanoparticles.

3.2.11 Cell culture

4T1 cells were cultured in RPMI-1640 medium supplemented with 10% (v/v) FBS, 0.03% L-glutamine and 1% penicillin-streptomycin within a humidified atmosphere containing 5% CO₂ at 37 °C. The cell culture medium was replaced every two days.

3.2.12 Cellular uptake study

The time-dependent internalization and intracellular distribution of PEI-Por-CD: oxlit(IV)-ada nanoparticles were visualized by confocal laser scanning microscopy (CLSM). 4T1 cells were seeded in 6-well plate (1×10^5 cells/well) on glass cover slip and cultured for 24 hr. The cell culture medium was then replaced with fresh ones and cells were incubated with the nanoparticles with porphyrin concentration of 6.25 μ M for different time durations. Following which, cells were washed three times with PBS, and nucleus was stained with H33342 for 5 min and fixed with 4% paraformaldehyde

before CLSM observation. Fluorescence was examined under excitation at 405 nm for Hoechst 33342 and 488 nm for porphyrin.

3.2.13 Cellular ROS generation

Cellular ROS generation was detected using Image-iTTM LIVE Green Reactive Oxygen Species Detection Kit. 4T1 cells (1×10^5 cells/well) were seeded on disposable petri dish and cultured for 24 hr. The culture medium was then removed and replaced with fresh culture medium containing nanoparticles with porphyrin concentration of 6.25 μ M. After 4 hr incubation, cells were illuminated with 630 nm laser (150 mW/cm²) for 20 min. Cells incubated with fresh culture medium with and without laser treatment or incubated with nanoparticles without laser illumination were used as control groups. After treatments, samples were incubated with 2',7'-dichlorofluorescein diacetate (DCF-DA) according to the standard protocol provided by the suppliers. Fluorescence images of the cells were acquired under a ZEISS LSM 800.

3.2.14 In vitro cytotoxicity assay

The cytotoxicity of PEG-Por-CD, PEG-Por-CD: oxliPt(IV)-ada nanoparticles in the dark and under light irradiation against 4T1 cells was evaluated with the MTT assay. Briefly, 4T1 cells were seeded into 96-well plate (1×10^4 cells/well) and cultured overnight. The cell culture medium was replaced with 100 μ L fresh medium containing samples with final equivalent Pt and porphyrin concentrations ranging from 0 to 100 μ g/mL and 0 to 12.5 μ M respectively. After 4 hr incubation, cells undergoing PDT treatment were illuminated with 630 nm laser (50 mW/cm²) for 30 mins. Following incubation at 37 °C for another 44 hr, the incubation medium was removed. Fresh culture medium containing MTT (100 μ L, 5 mg/ml) were added into each well and the cells were incubated at 37 °C for another 4 hr. DMSO (100 μ L) was added to dissolve the formazan crystals. The optical density (OD) was read at 490 nm using a microplate

reader (infinite M200, TECAN). The control was taken as the culture medium in the absence of samples. The normalized cell viability in % was calculated as cell viability $= \frac{OD_{\text{sample}} - OD_{\text{blank}}}{OD_{\text{control}} - OD_{\text{blank}}} \times 100\%$. The results were obtained from 5 replicates for each sample and expressed as means.

3.2.15 Live/dead assay

To visualize the cytotoxicity effect using CLSM, 4T1 cells were seeded into 6 well plate (2×10^5 cells/well) and cultured overnight. The culture medium was then replaced with fresh medium containing nanoparticles with final porphyrin concentration of 6.25 μM and incubated in the dark for 4 hr. Following which, cells undergoing PDT treatment were illuminated with 630 nm laser (50 mW/cm^2) for 20 min. After another 44 hr of incubation in dark, the cells were washed with PBS and stained with calcein-AM (3 μM) and PI (8 μM) at 37 °C for 20 min before visualizing under the microscope.

3.2.16 Apoptosis study

4T1 cells were seeded into 6-well plate (1×10^6 cells/well) and cultured at 37 °C overnight. After replacing the culture medium, cells were incubated with different samples at final porphyrin concentration of 6.25 μM . After 4 hr incubation, cells undergoing PDT treatment were illuminated with 630 nm laser (50 mW/cm^2) for 30 min and further incubated for another 44 hr. To quantitatively measure the percentage of apoptotic cells, cells after treatments were harvested by trypsin, washed thrice with PBS, resuspended in 100 μL binding buffer and stained with Annexin V-FITC/PI at room temperature in the dark for 15 min. Finally, the cells were analyzed using a flow cytometer (excitation wavelength: 488 nm).

3.2.17 Animal models

Female BALB/c mice (9 weeks old) were purchased from InVivos Pte Ltd and used under approved protocols of the Institutional Animal Care and Use Committee (IACUC) at Nanyang Technological University, Singapore. A total of 1×10^6 cells suspended in PBS (40 μ L) were subcutaneously injected into the right flank of each mouse. The mice were used in experiments when the tumor volumes reached 60- 70 mm^3 .

3.2.18 In vivo combination therapy

When the mean tumor size reached 60- 70 mm^3 , the mice were divided into 5 treatment groups (n= 5 per group) at random: (a) control; (b) free drugs mixture (oxaliplatin meso-Tetra(4-hydroxyphenyl)porphine (THPP)) with light irradiation, (c) PEG-Por-CD nanoparticles with light irradiation, (d) PEG-Por-CD: oxliPt(IV)-ada nanoparticles in the dark, and (e) PEG-Por-CD: oxliPt(IV)-ada nanoparticles with light irradiation. The treatment was conducted on day 1 and 4. Light irradiation was conducted locally with a white light source (50 mW cm^{-2} , 30 mins) after the injection. The tumor volume was measured daily for 14 days and calculated using the following formula: $\text{volume} = 0.5 \times L \times W^2$, where L is the tumor length and W is the tumor width. The relative tumor volume was calculated where $\text{relative tumor volume} = (\text{volume of tumor on day } n / \text{volume of tumor on day } 0)$. On day 14 post-treatment, the mice were euthanized by CO_2 inhalation. Tumors were excised and fixed with 4% formaldehyde, followed by embedment in a paraffin block, sectioned, stained with hematoxylin and eosin (H&E) and mounted on glass slides before imaging. For systemic toxicity study, the mice were either injected with PBS or PEG-Por-CD: oxliPt(IV)-ada nanoparticles. On day 14 post-injection, the major organs were excised, processed and stained for H&E imaging.

3.2.19 Statistical Analysis

All data were expressed as mean $\pm$ standard error of the mean. The statistical comparison between two sets of data was determined by one-tailed student's t-test. Values with $p < 0.05$ were considered to be statistically significant.

3.3 Results and Discussion

3.3.1 Synthesis and characterization

The chemical structures of the therapeutic components are depicted in Scheme 1. OxliPt(IV)-ada prodrug functioning as the guest molecule was obtained by reacting ethylenediamine-modified adamantane with oxliPt(IV)-COOH (Scheme 3). Successful synthesis of the prodrug was confirmed with electrospray ionization mass spectrometry (ESI-MS) and ^1H NMR.

Figure 1. HPLC monitoring for the reduction of oxliPt(IV)-ada to give active oxaliptatin(II).

To investigate the redox-sensitive property of the prodrug, reverse-phase HPLC was used to evaluate the reduction reaction products of oxliPt(IV)-ada incubated with sodium ascorbate. Sodium ascorbate as a reducing agent was used to mimic the reducing environment inside cancer cells.¹⁸⁴⁻¹⁸⁵ The reduction of oxliPt(IV)-ada prodrug was expected to produce the cytotoxic oxaliptatin. As shown in Figure 1, free oxaliptatin exhibited a retention time peak at 1.1 min, whereas the peak representing oxliPt(IV)-ada prodrug appeared at 7.6 min in the HPLC chromatogram. After the incubation of the prodrug with sodium ascorbate over time, the intensity of the peak at 7.6 min decreased, along with the emergence of a peak at 1.1 min. This observation verified that the as-

synthesized oxliPt(IV)-ada could function as a prodrug, becoming pharmaceutical active after the reduction.

Figure 2. (A) FTIR spectra of different intermediate compounds to confirm the successful synthesis of PEG-Por-CD. (B) Gel permeation chromatography traces of PEG₂₀₀₀ and PEG-Por-CD. (C) Absorbance spectrum of PEG-Por-CD. Inset shows the absorbance peaks at 564 nm and 623 nm. (D) Generation of singlet oxygen by measuring the fluorescence intensity changes of SOSG in water.

THPP was modified with PEG and cyclodextrin to give PEG-Por-CD as the host molecule (Scheme 4). ^1H NMR and Fourier-transform infrared spectroscopy (FTIR) were used to monitor the step-wise synthesis of PEG-Por-CD. According to the FTIR analysis (Figure 2A), the appearance of a new signal at 3320 cm^{-1} corresponding to alkyne C-H confirms the formation of alkyne-functionalized THPP. Subsequently, the emergence of a peak at 2885 cm^{-1} assigned to the C-H stretch of the PEG unit indicates successful conjugation of PEG onto the porphyrin core. After conducting a click reaction with azide-modified cyclodextrin, the disappearance of the azide peak at 2110 cm^{-1} confirms the formation of PEG-Por-CD. Gel permeation chromatography (GPC) traces of PEG-Por-CD and initial PEG reactant are shown in Figure 2B. After the reaction of THPP with PEG and cyclodextrin, the GPC traces of the obtained PEG-Por-CD shifted to the higher molecular weight region, supporting the fact that PEG-Por-CD was successfully synthesized. In addition, UV-Vis absorbance spectrum of PEG-Por-CD displays typical absorption peaks of the porphyrin unit, including the Soret band at 430 nm, and the Q bands at 564 and 623 nm (Figure 2C).²³¹ As compared to unmodified THPP, there are slight red shifts of these absorption bands for PEG-Por-CD. On the other hand, the shape and intensity ratio of these absorbance peaks remain constant after the modifications, suggesting that the optical properties and integrity of the porphyrin core are largely retained.

To investigate photodynamic therapeutic ability of PEG-Por-CD, its singlet-oxygen generation was monitored by the fluorescence change of SOSG, a singlet-oxygen responsive dye (Figure 2D). Continuously increasing fluorescence of SOSG at 525 nm when PEG-Por-CD is irradiated with 630 nm light at different times indicates the generation of singlet oxygen. Hence, both the optical properties and singlet oxygen

generation capability are highly retained even after the modifications to the pristine porphyrin.

3.3.2 Preparation and characterization of nanocarrier

It was expected that the functionalization of hydrophilic PEG would render the porphyrin core with good aqueous solubility and amphiphilic character when complexed with oxliPt(IV)-ada, facilitating the formation of nanoparticles via the self-assembly in aqueous media.

Figure 3. (A) ¹H NMR comparison of PEG-Por-CD and PEG-Por-CD: oxliPt(IV)-ada in D₂O/d-DMSO mixture (5/5, v/v). (B) Enlarged region of the spectra from 3.30 to 4.00 ppm. Changes in the chemical shifts of the cyclodextrin in the nanoparticles are denoted by the arrows.

To demonstrate this point, ¹H NMR spectrum of the PEG-Por-CD: oxliPt(IV)-ada complex was recorded and compared to PEG-Por-CD (Figure 3). Obvious chemical shifts for the key protons of the cyclodextrin ring in the complex were observed,

indicating the formation of an inclusion complex between the adamantane unit of the prodrug and the cyclodextrin ring of PEG-Por-CD.

Figure 4. (A) Absorbance spectra of the nanocarrier obtained at different concentrations. (B) Critical aggregation concentration of the nanocarrier determined by monitoring the dependence of the optical transmittance at 430 nm.

The critical aggregation concentration of the nanocarrier was determined by monitoring the change in the optical transmittance at different concentrations of the complex (Figure 4). The optical transmittance at 430 nm decreased gradually upon increasing the concentration, and a great decrease was observed at the concentration of 0.35 μM on account of the formation of larger particles. Knowing that the complex could indeed form nanoparticles, we then investigated the optimal loading ratio of the PS and drug for the preparation of PEG-Por-CD: oxliPt(IV)-ada nanoparticles.

Since the aim of the nanocarrier design was to achieve a synergistic effect from chemotherapy and PDT, we evaluated the optimal loading ratio of the PS and prodrug in PEG-Por-CD: oxliPt(IV)-ada nanoparticles. The determining factors for the optimal ratio were low dark phototoxicity and a favorable combination index for synergistic tumor cell inhibition.²³⁰ Hence, the cytotoxicity of oxliPt(IV)-ada and PEG-Por-CD in the dark was first studied using the standard MTT assay.

Figure 5. MTT viability assay of 4T1 cells 48 hr post-incubation with (A) PEG-Por-CD in dark, (B) oxliPt(IV)-ada and (C) oxliPt(II).

As shown in Figure 5A, 4T1 cells exhibited high cell viability when incubated with up to 12.5 μM of PEG-Por-CD in the dark. The oxliPt(IV)-ada prodrug ($IC_{50} = 141$ μM, Figure 5B) showed lower cytotoxicity than unmodified oxaliplatin ($IC_{50} = 85.3$ μM, Figure 5C) since its reactivity was minimized. Based on the concentrations used in the above cell viability assay, nanoparticles consisting of varying ratios of oxliPt(IV)-ada and PEG-Por-CD (0.5-32 : 1) were prepared. 4T1 cells were incubated with the nanoparticles containing different molar ratios and irradiated with 630 nm light (50 mW cm^{-2} , 20 mins), followed by evaluating the cytotoxicity of these samples with the MTT assay.

Figure 6. Graph of combination index for the treatment of 4T1 cells with different molar ratios of PEG-Por-CD to oxliPt(IV)-ada.

As shown in Figure 6, synergistic therapeutic analysis (combination index < 1) under the tested ratios indicates that the prodrug/PS ratio of 8 : 1 exhibits the strongest

effect. Based on this result, we employed this optimal ratio for the fabrication of PEG-Por-CD: oxliPt(IV)-ada nanoparticles in the following cancer treatment studies.

Figure 7. (A) DLS data and corresponding TEM image (inset) of PEG-Por-CD: oxliPt(IV)-ada in water. (B) UV-vis and fluorescence spectra PEG-Por-CD: oxliPt(IV)-ada nanoparticles in water. The concentration of PEG-Por-CD was $16 \mu\text{g mL}^{-1}$. (C) DLS size of PEG-Por-CD: oxliPt(IV)-ada nanoparticles in different media over time.

The transmission electron microscopy (TEM) image in Figure 7A reveals that the obtained PEG-Por-CD: oxliPt(IV)-ada nanoparticles have a uniform spherical morphology with a diameter of about 164 nm. UV-vis absorption and fluorescence spectra of PEG-Por-CD: oxliPt(IV)-ada in aqueous solution show prominent peaks of the porphyrin unit, confirming that the porphyrin moiety is incorporated in the nanocarrier (Figure 7B). Similar hydrodynamic size in water and PBS was observed, with slightly smaller size in RPMI medium (Figure 7C). Importantly, the size in the different solutions remains relatively constant over three days, indicating good

physiological stability of the nanocarrier. The oxaliplatin loading capacity in the nanocarrier was 19.6 wt % as measured by ICP-MS.

3.3.3 ROS generation and drug release study

Figure 8. Fluorescence intensity changes of SOSG for control, PEG-Por-CD and PEG-Por-CD: oxliPt(IV)-ada over time. NP represents PEG-Por-CD: oxliPt(IV)-ada nanoparticles..

We then assessed the PDT effect of PEG-Por-CD: oxliPt(IV)-ada nanoparticles by measuring the ROS generation. The continuous increase in the SOSG fluorescence when incubated with the nanoparticles under 630 nm light irradiation demonstrated its capability to produce singlet oxygen for PDT (Figure 8). The singlet-oxygen production from the nanocarrier was less efficient than free PEG-Por-CD at the same porphyrin concentration likely due to the self-quenching of the generated singlet oxygen in compact nanoparticle formulation. Nonetheless, compared to the control group with no notable generation of ROS, the PDT efficacy of PEG-Por-CD: oxliPt(IV)-ada is highly sufficient.

Figure 9. (A) Redox-responsive Pt drug release from PEG-Por-CD: oxliPt(IV)-ada nanoparticles when incubated with different concentrations of sodium ascorbate (Vc). (B) TEM of PEG-Por-CD: oxliPt(IV)-ada nanoparticles after incubation with sodium ascorbate for 12 hr.

Since an intracellular reducing environment was expected to facilitate the drug release, we studied the oxaliplatin release from PEG-Por-CD: oxliPt(IV)-ada nanoparticles in different concentrations of sodium ascorbate at 37 °C using ICP-MS (Figure 9A). As anticipated, cumulative Pt drug release increases upon increasing the concentration of the reducing agent. In the absence of sodium ascorbate, only about 20% Pt was detected after 12 hr. In contrast, when the concentration of sodium ascorbate was increased to 10 mM, approximately 70% Pt release was observed within the same time period. The original spherical morphology of the nanoparticles was not observed after the reduction as seen from the TEM image (Figure 9B), since the reduction of the prodrug led to concomitant dissociation of the nanocarrier. The redox-triggered drug release is highly desirable for drug delivery since it can reduce premature drug release during blood circulation and, consequently, decrease side effects to normal cells.

3.3.4 Cellular accumulation and in vitro ROS generation

The intrinsic fluorescence of PEG-Por-CD was utilized to investigate the intracellular uptake of the nanoparticles in 4T1 cells. As shown from CLSM images

(Figure 10A), obvious red fluorescence from the porphyrin moiety was observed in the cytoplasm, indicating efficient cellular uptake of the nanocarrier. Particularly, the fluorescence intensity increases upon the incubation time, suggesting that the cellular accumulation of PEG-Por-CD: oxliPt(IV)-ada nanoparticles occurs in a time-dependent manner.

Figure 10. (A) Time-dependent intracellular accumulation of PEG-Por-CD: oxliPt(IV)-ada nanoparticles. Nucleus was stained with Hoechst 33342. NP: nanoparticles. Scale bar: 20 μm . (B) Detection for intracellular generation of reactive oxygen species under different treatments using DCFH-DA. Nucleus was stained with Hoechst 33342. NP: nanoparticles. Scale bar: 50 μm .

The capability of PEG-Por-CD: oxliPt(IV)-ada nanoparticles to generate ROS in-vitro was evaluated using ROS sensitive DCFH-DA dye (Figure 10B). The generation of intracellular ROS results in green fluorescence of the dye. In the absence and presence of light irradiation, almost no fluorescence was observed when the cells were not incubated with any samples. In contrast, cells treated with PEG-Por-CD: oxliPt(IV)-ada nanoparticles under light irradiation displayed significantly stronger

green fluorescence, indicating the ability of nanoparticles to generate cytotoxic ROS for PDT.

3.3.5 In vitro combination therapy

Encouraged by efficient cellular accumulation and ROS generation property, the cytotoxicity of PEG-Por-CD: oxliPt(IV)-ada nanoparticles was then investigated in 4T1 cells, and its capability to inhibit 4T1 cancer cell proliferation by combinational PDT and chemotherapy was conducted by the MTT assay. The PDT and chemotherapy efficacies were assessed by treating 4T1 cells with PEG-Por-CD plus light irradiation (PDT control group) and PEG-Por-CD: oxliPt(IV)-ada in the dark (chemotherapy control group), respectively. For the PDT group, 630 nm light (50 mW cm^{-2} , 20 mins) was used to irradiate the cells after 4 hr incubation with the samples.

Figure 11. In-vitro cell viability of 4T1 cells following 48 hr treatment with PEG-Por-CD or PEG-Por-CD: oxliPt(IV)-ada nanoparticles in the absence and presence of light. Data points represent mean $\pm$ SD ($n = 5$).

Figure 11 shows that the cell viability decreases upon increasing PS and drug concentrations. Particularly, cells treated with PEG-Por-CD: oxliPt(IV)-ada nanoparticles under light irradiation present the lowest viability at equivalent

concentrations, indicating that the combined PDT and chemotherapy provided by PEG-Por-CD: oxliPt(IV)-ada nanoparticles is more effective than the monotherapy, that is, PDT with PEG-Por-CD (light) or chemotherapy with PEG-Por-CD: oxliPt(IV)-ada (dark). The IC_{50} values of PEG-Por-CD and oxliPt(IV)-ada against 4T1 cells were determined to be 15.6 μ M and 79.8 μ g/mL respectively, while IC_{50} of PEG-Por-CD: oxliPt(IV)-ada nanoparticles was significantly lowered to 4.7 μ M (IC_{50} in terms of Pt = 37.5 μ g/mL). The highly improved anticancer effect of PEG-Por-CD: oxliPt(IV)-ada nanoparticles could be attributed to the combined cell killing mechanisms, where the released oxaliplatin induces DNA damage and PDT causes oxidative apoptosis.²³²

Figure 12. (A) Live/dead cell assay for cells treated with PBS, free photosensitizer, and PEG-Por-CD: oxliPt(IV)-ada in the absence and presence of light irradiation. Green: live cells, red: dead cells. Scale bar: 100 μ m. (B) Bar chart representing the percentage of viable, early apoptotic, late apoptotic and necrotic cells after the treatment with the indicated samples for 48 hr, measured by flow cytometry. NP: nanoparticles.

To visually detect the cytotoxicity, 4T1 cells treated with different formulations were co-stained with calcein-AM (green) and PI (red) for the live/dead assay (Figure 12A). As control groups, cells without any treatment in the dark or under light irradiation showed strong green fluorescence, thereby demonstrating no marked loss of cell viability. The ratio of dead/live cells increased when cells were treated with PEG-Por-

CD under light irradiation or PEG-Por-CD: oxliPt(IV)-ada in the dark, indicating certain cytotoxicity of the monotherapy. Notably, increased red fluorescence and negligible green fluorescence were observed when cells were treated with PEG-Por-CD: oxliPt(IV)-ada nanoparticles under light irradiation, demonstrating the enhanced therapeutic efficiency as compared with corresponding monotherapy, consistent with the result of MTT assay.

To gain more insight on the cytotoxicity, the apoptosis assay of 4T1 cells treated with the different formulations were analyzed by flow cytometry. The incubation of 4T1 cells with PEG-Por-CD under laser irradiation or PEG-Por-CD: oxliPt(IV)-ada in the dark shows low cell apoptosis of only 12.4 % and 44 %, respectively (Figure 12B). On the contrary, cells treated with PEG-Por-CD: oxliPt(IV)-ada nanoparticles under light irradiation display the highest apoptosis with the rate of 92.6 %. The enhanced cytotoxicity of PEG-Por-CD: oxliPt(IV)-ada nanoparticles further confirmed the synergistic effect of combinational chemotherapy and PDT.

3.3.6 In vivo anticancer treatment efficacy

Finally, the combined antitumor efficacy of PEG-Por-CD: oxliPt(IV)-ada nanoparticles in 4T1 tumor-bearing mice was assessed. When the sizes of the tumor grew to be around 70 mm³, mice were randomly divided into five treatment groups (n=5) including: (1) PBS control, (2) free drugs mixture comprising oxaliplatin and THPP, (3) PEG-Por-CD plus light (PDT), (4) PEG-Por-CD: oxliPt(IV)-ada nanoparticles in the dark (chemotherapy), and (5) PEG-Por-CD: oxliPt(IV)-ada nanoparticles plus light (PDT+ chemotherapy), where the corresponding porphyrin and oxaliplatin dose were kept at 2.5 mg kg⁻¹ and 5 mg kg⁻¹ respectively. These samples were administrated intratumorally on day 1 and 4 during the treatment period. For mice receiving PDT, the tumors were irradiated with a 630 nm LED light (50 mW cm⁻², 30 min). The tumor sizes

and body weights of the mice were recorded every day for 2 weeks after receiving treatments.

Figure 13. (A) Relative tumor volume growth curves of 4T1 tumor-bearing BALB/c mice treated with indicated samples over 14 days. (B) Weight of tumors excised from the mice at 14 days after initial treatment with different samples. Data are expressed as means $\pm$ SD; p value

was determined by a student's t-test: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.005$. (C) Representative images of 4T1 xenograft tumors of the mice after different treatments. (D) Body weight variation of the mice during the indicated treatment. (E) H&E images of tumor excised from mice 14 days post-treatment with different samples. Scale bar: 100 μm . The treatment groups were: (i) PBS control, (ii) free drugs mixture comprising oxaliplatin and THPP, (iii) PEG-Por-CD NP plus light (PDT), (iv) PEG-Por-CD: oxliPt(IV)-ada NP (chemotherapy), and (v) PEG-Por-CD: oxliPt(IV)-ada NP plus light (PDT+ chemotherapy). (F) H&E micrographs of representative major organs at 14 days post-treatment with PBS and PEG-Por-CD: oxliPt(IV)-ada nanoparticles. Scale bar: 100 μm . NP: nanoparticles.

As expected, mice without any treatment (group 1) exhibited the greatest tumor growth. As shown in Figure 13A, the free drug mixture (group 2) could only partially inhibit the tumor growth because of its poor tumor retention. In addition, PDT (group 3) or chemotherapy (group 4) treatment slightly suppressed the tumor growth. Remarkably, the treatment with PEG-Por-CD: oxliPt(IV)-ada nanoparticles plus light irradiation offered the greatest tumor growth inhibition.

In addition, the average tumor weight (Figure 13B) and representative photographs (Figure 13C) of the tumor excised from the different treatment groups verified that the treatment with PEG-Por-CD: oxliPt(IV)-ada nanoparticles under light irradiation provided the best antitumor efficacy. Tumors obtained from each treatment groups were evaluated histologically by hematoxylin and eosin (H&E) staining. Notably, the H&E stained tumor tissues from mice treated with PEG-Por-CD: oxliPt(IV)-ada nanoparticles under light irradiation led to more severe tumor cell apoptosis, indicating a synergistic effect of combination PDT and chemotherapy (Figure 13E). Body weights of the mice in all groups monitored during the treatment period exhibited similar changes, suggesting no major systemic toxicity arising from these formulations (Figure 13D). In addition, the tissue histologies of major organs (i.e. heart, liver, spleen, lung,

kidney, and small intestine) in each treatment group were investigated by H&E staining. As shown in Figure 13F, H&E stained images of the major organs from the mice treated by PEG-Por-CD: oxliPt(IV)-ada nanoparticles plus light irradiation showed no noticeable tissue damage or pathological variation, demonstrating its good biocompatibility.

3.4 Conclusion

In summary, we have utilized a simple strategy based on host-guest chemistry to achieve synergistic PDT and chemotherapy for cancer treatment. Results obtained have clearly demonstrated that the combination therapy was more effective than monotherapy alone. By integrating the therapeutic molecules with supramolecular self-assembly properties, the present oxaliplatin(IV) prodrug self-delivery nanosystem allowed precise control over the drug and PS loading ratio while eliminating the toxicity concern of the drug carrier to achieve programmable drug release on the tumor sites. Such a design strategy may pave a new avenue for the fabrication of prodrug-based delivery systems toward efficient combination therapy for cancer.

Chapter 4

Redox-responsive polymeric nanocomplex for delivery of cytotoxic protein & chemotherapeutics

4.1 Introduction

Protein-based therapeutics such as enzymes, antibodies, and recombinant DNA have played an important role in biomedicine on account of their pharmacological potency and highly selective bioactivities²³³⁻²³⁴ Several protein-based drugs have also demonstrated anticancer properties, which include cell growth inhibition, apoptosis activation, and anti-angiogenesis, and thus have been used for cancer treatment.²³⁵⁻²³⁶ Even though such protein-based therapeutics are attractive as the treatment approach for cancer, the challenge of delivering them efficiently into cancer cells to reach their intracellular targets still hindered their clinical translations.²³⁷ Issues such as their low stability and vulnerability to protease degradation and denaturation often compromise the amount of proteins that can accumulate selectively in the tumor.^{233, 238-239} Furthermore, the hydrophilicity, charges and sizes of native protein drugs make them unfavorable for active uptake into cancer cells.²⁴⁰⁻²⁴² Over the last decade, the most thoroughly explored protein delivery strategy is the modification of target proteins with protein transduction domains or membrane transport signals.²⁴³⁻²⁴⁹ However, the delivery efficiency varies with the protein type, and the method often lacks the specificity for tumor tissues.

To minimize unnecessary toxicity to off-target normal tissues, it is useful to mask the activity of proteins before systemic administration. Upon the accumulation at the tumor site, the function of proteins should be spatiotemporally activated by endogenous stimuli. To realize on-demand activation of the protein function in cells, different triggers for uncaging the modified proteins have been developed. Intracellular stimuli such as endosomal acidic pH,²⁵⁰⁻²⁵² as well as overproduction of reactive oxygen

species (ROS)^{229, 253} and bio-reductants²⁵⁴⁻²⁵⁵ were used for the removal of blockage moieties in order to uncage proteins such that the protein activities could be recovered.

In light of poor accumulation of the protein therapeutics, nanosized delivery systems such as inorganic nanocarriers,²⁵⁶⁻²⁵⁷ polymeric nanoparticles,²⁵⁸⁻²⁶⁰ nanogels,²⁶¹⁻²⁶² and liposomes^{64, 263-264} have been employed for better spatial and temporal delivery of proteins. While some of the nanocarriers can prevent protein proteolysis during the blood circulation and improve the tumor accumulation, loss in protein activity and complicated material preparation limit their further applications.²⁶⁵ Therefore, a facile and convenient strategy to fabricate novel nanosystems for efficient cellular protein delivery is highly desirable.

Here, we demonstrate a simple but efficacious strategy to deliver cytotoxic protein and drug using cationic polymeric nanocarrier for combinational cancer therapy. RNase A, a clinically used therapeutic protein for treating refractory cancer by cleaving cellular RNA and inhibiting cell proliferation,²⁶⁶ was used as the model protein. To realize masking of the protein activity, the activity of RNase A was blocked by the conjugation with phenylboronic acid to its lysine residue that accounts for its protein activity, forming the deactivated RNase A-nitrophenylboronic conjugate (RNBC).²⁶⁷ Phenylboronic acid could then be cleaved in the presence of hydrogen peroxide (H₂O₂) to regain the activity of RNase A.^{229, 268}

Scheme 1. Schematic illustration of redox-responsive protein delivery by PEI-oxliPt(IV)@RNBC/GOD nanocomplex toward synergistic cancer therapy. (A) Formation of PEI-oxliPt(IV)@RNBC/GOD through electrostatic interactions between PEI-oxliPt(IV), GOD and RNBC, and subsequent cargo release by redox stimuli. (B) Processes involved in synergistic cancer therapy after cellular internalization. a: Cellular accumulation of PEI-oxliPt(IV)@RNBC/GOD; b: Redox-responsive degradation of the nanocomplex, releasing RNBC and oxaliplatin; c: RNA degradation by the active RNase A that is uncaged by enzymatically produced H_2O_2 from GOD and glucose; d: Inhibition of DNA synthesis by oxaliplatin.

To achieve efficient restoration of the protein biological function, glucose oxidase (GOD) was co-delivered in the nanocarrier. GOD is an enzymatic biocatalyst that oxidizes endogenous glucose into gluconic acid and H_2O_2 .²⁶⁹⁻²⁷⁰ Highly elevated glucose level inside cancer cells enables GOD to continuously generate H_2O_2 , which subsequently restores the RNase A activity.²⁷¹⁻²⁷² Intracellular delivery of both GOD

and RNase A proteins was facilitated by forming the nanocomplex (PEI-oxliPt(IV)@RNBC/GOD) with cationic oxaliplatin prodrug cross-linked polyethylenimine (PEI-oxliPt(IV)). Polyethylenimine (PEI) was chosen as the main polymeric carrier on account of its positive charge that could form electrostatic interaction with negatively charged proteins and enhance cellular internalization with negatively charged cell membrane (Scheme 1A). Compared to high molecular PEI (i.e. 10- 25 kDa) whose large charge and non-degradability make them cytotoxic, low molecular weight PEI (i.e. 0.8- 2 kDa) exhibits innocuous toxic profile.²⁷³⁻²⁷⁴ Therefore we strategically crosslinked low molecular weight PEI with the reduction-sensitive oxaliplatin(IV) prodrug to give a cationic polymer suitable to complex with negatively charged proteins. Furthermore, this process ensures reductive degradability of the Pt prodrug-integrated PEI to minimize its cytotoxicity. In addition, oxaliplatin which treats colorectal cancer and metastatic breast cancer,²⁷⁵ has orthogonally different cell killing mechanisms from RNase A. Hence, the simultaneous delivery of drugs and proteins would cooperatively overcome cancer.^{150, 276-277}

The resulted PEI-oxliPt(IV) was kinetically inert and non-toxic which complexed with GOD and RNBC to give PEI-oxliPt(IV)@RNBC/GOD. Upon accumulation in cancer cells, the nanocomplex could be reduced to give activated RNase A and its cytotoxic Pt(II) counterpart. The inhibition of cellular protein synthesis by RNase A and the suppression of DNA replication by active oxaliplatin drug lead to synergistic cancer treatment (Scheme 1B).²⁷⁸ The proposed design of the nanocomplex facilitates 1) facile preparation of PEI-oxliPt(IV)@RNBC/GOD nanocomplex by electrostatic interactions, 2) concurrent delivery of drug and protein with high efficiency, and 3) responsiveness to overexpressed cellular bio-reductants (e.g. glutathione and ascorbic acid) for the release of RNase A and active platinum drug.¹⁰⁶

4.2 Experimental Section

4.2.1 Materials and reagents

All solvents were purchased from Sigma-Aldrich. Alizarin Red S, bovine pancreatic ribonuclease A (RNase A), 1,1'-carbonyldiimidazole (CDI), 3,3'-dithiodipropionic acid, fluorescein isothiocyanate isomer 1 (FITC), L-glutathione reduced, D-(+)-glucose, glucose oxidase from aspergillus niger (GOD), heparin, 4-methoxycarbonylphenylboronic acid pinacol ester, 4-nitrophenyl chloroformate, polyethyleneimine (branched PEI, MW 800), succinic anhydride, and triethylamine (TEA) were purchased from Sigma-Aldrich. Ethidium bromide was brought from Biorad. Oxaliplatin was purchased from Aaron Pharmatech Ltd. (Shanghai, China). Cy-3 labeled RNA with the sequence of 5'-UAGAGAGAGAAAGUUUCGACUUUCUCUCUCUA-3' was kindly provided by Professor Gang Chen group. RPMI 1640 w/ L-glutamine medium, penicillin-streptomycin solution and fetal bovine serum (FBS) were purchased from Alantis Bioscience. 4T1 cells were obtained from ATCC. Coomassie Brilliant Blue G-250 and 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide was bought from TCI. Hoechst 33342 staining dye and Caspase 3 assay kit (colorimetric) were bought from Abcam. RNase A activity was assayed using RNaseAlert® Kit (Integrated DNA Technologies, Inc., IA). LysoTracker Deep Red, Annexin V-FITC Apoptosis Detection Kit, Click-iT® RNA Alexa Fluor® 594 Imaging Kit, Pierce™ BCA Protein Assay Kit, Pierce™ Quantitative Peroxide Assay Kit, Quanti-IT Ribogreen RNA Assay kit, PureLink® RNA Mini Kit, Spectra™ multicolor broad range protein ladder, pre-cast PAGE gels and Ribonuclease inhibitor (RiboLock, E. coli, 2500 units) were purchased from ThermoFisher Scientific.

4.2.2 Characterization

¹H NMR spectra were probed on 300 MHz Bruker spectrometer using deuterated DMSO or D₂O as solvent. Transmission electron microscope (TEM) images were collected on a JEM-1400 (JEOL) operated at 100 kV. Zeta potential and dynamic light scattering (DLS) measurements were measured with Zetasizer, Malvern Instrument Ltd. UV-vis and fluorescent spectra were obtained with a microplate reader (infinite M200, TECAN). Inductively coupled plasma mass spectrometry (ICP-MS DRCE Elan, Perkin Elmer) was used for quantitative determination of platinum. Circular dichroism (CD) spectra were measured with a Jasco J-810 CD spectrophotometer. Sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed using OmniPAGE mini vertical system (Cleaver Scientific). Briefly, 20 µL protein sample (1 mg/mL) was heated at 90 °C for 5 mins, mixed with 5 µL loading buffer and loaded onto the pre-cast PAGE gel. Electrophoresis was conducted at room temperature for 2 h at constant voltage (100 V) in 1× running buffer (containing 250 mM glycine, 3.5 mM SDS and 25 mM Tris). The obtained gel was stained with Coomassie Brilliant Blue G-250 for 1 h and destained with destaining buffer (containing 10% acetic acid and 5% ethanol) for 2 h. Matrix-assisted laser desorption/ionization- time-of-flight (MALDI-TOF) spectra were obtained with a JMS-S3000 (JEOL) mass spectrometer. Native polyacrylamide gel electrophoresis (Native-PAGE) experiment was conducted with incubation buffer containing 200 mM NaCl, 0.5 mM EDTA, 20 mM PBS (pH 7.5). The loading volume for each sample was 20 µL (containing 20% glycerol). 1× Tris–borate–EDTA (TBE) buffer, pH 8.3 was used as the running buffer. The gel was run at 4°C at 250 V for 5 hr and imaged by the Typhoon Trio Variable Mode Imager. Confocal laser scanning microscopy (CLSM) images were acquired by a ZEISS LSM 800 confocal microscope. Flow cytometry was recorded on a BD LSR Rortessa X20 (3 lasers).

4.2.3 Synthesis of RNBC and FITC-labelled RNBC

NBC-modified RNase A was prepared according to procedures reported in literature.²²⁹

Scheme 2. Schematic illustration for the synthesis route of RNBC.

4-nitrophenyl 4(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl carbonate (NBC) was synthesized as described in the literature.²⁷⁹ Briefly, 4-(Hydroxymethyl)phenylboronic acid pinacol ester (0.5 g, 2.14 mmol) dissolved in anhydrous THF (20 mL) was added TEA (0.6 mL) and 4-nitrophenyl chloroformate (0.47 g, 2.35 mmol). The reaction mixture was left to stir at room temperature for 1 hr. After which, the solvent was removed by rotary evaporation, and the residue was dissolved in ethyl acetate and washed with water, 1 M HCl and NaHCO₃. The organic phase was combined and dried over anhydrous MgSO₄. The solution was filtered, concentrated by rotary evaporation and purified by silica column (hexane:ethyl acetate, 5:1) to yield NBC as white powder. ¹H-NMR (500 MHz; CDCl₃) δ (ppm) = 8.27 (d, 2H, 9 Hz), 7.85 (d, 2H, 8 Hz), 7.44 (d, 2H, 8 Hz), 7.38 (m, 2H, 9 Hz), 5.31 (s, 2H), 1.35 (s, 12H). ESI-MS m/z [M + H]⁺ calcd. For [C₂₀H₂₂BNO₇]⁺ 399.21; found 399.25.

NBC was then modified onto RNase A via reacting with the primary amine groups of the lysine residues. Briefly, RNase A (3 mg) and NBC (9.6 mg) separately dissolved in 0.1 M NaHCO₃ buffer solution (1.5 mL, pH 8.5) and DMSO were mixed

and stirred at room temperature for 10 hr. The obtained mixture was purified using ultrafiltration (MWCO = 10 kDa) and lyophilized to give RNBC as white powder.

To allow the observation and quantification of RNBC and RNase A *in vitro*, the proteins were labeled with FITC dye. Protein (2 mg) was dissolved in 0.1 M NaHCO₃ solution (1 mL, pH 9.5), followed by the addition of FITC solution (4 mg/mL in DMSO, 25 μ L). The solution was stirred in the dark at room temperature overnight. The FITC-labelled protein was collected by ultrafiltration (MWCO = 10 kDa), washed with deionized water and obtained by lyophilization.

4.2.4 Characterization of RNBC

Zeta potential, CD measurements and SDS-PAGE were carried out for native RNase A and RNBC. MALDI-TOF was used to calculate the number of NBC caging groups on RNBC.

Fluorometric Alizarin Red S (ARS) assay was used to verify the presence of boronic acid moiety in RNBC. Protein samples (0.2 mg/mL), including RNase A and RNBC with or without H₂O₂ treatment for 3 hr were incubated with 0.02% w/v ARS solution. After 15 min incubation, ARS fluorescence emission was recorded on a microplate reader (infinite M200, TECAN). Emission was monitored at 600 nm with the excitation at 490 nm. In addition, the mass of RNBC after the treatment with H₂O₂ was measured by MALDI-TOF and compared to RNBC without H₂O₂ treatment or RNase A.

4.2.5 Uncaging of RNBC

The uncaging of RNBC by GOD and glucose was studied by RNA degradation. RNase A activity assay kit, RNA Native-PAGE electrophoresis and ethidium bromide assay were employed to monitor the RNA degradation.

For RNase A activity assay, RNBC or RNase A (1 μ L, 1 mg/mL) was mixed with GOD (1 μ L, 10 μ g/mL), and then incubated with or without glucose (1 μ L, 1 mg/mL) for 1 hr (total volume of solution: 80 μ L). The samples were then mixed with RNaseAlert®Substrate (10 μ L) and assay buffer (10 μ L) in 96-well plate, in which the fluorescence intensity at 520 nm (excitation 490 nm) was monitored within 25 min of incubation.

For RNA Native-PAGE electrophoresis, the protein samples were prepared as mentioned previously. Following 30 min incubation with Cy 3-labelled RNA (0.2 μ M), RNase inhibitor (1 μ L, 40 U/mL) was added to inhibit enzymatic activity of the proteins. The samples were then loaded on a 12.5% Native-PAGE gel and ran in the dark at 240 V for 5 hr. The gel bands were imaged directly by Typhoon Trio Variable Mode Imager.

For ethidium bromide assay, EtBr (10 μ L, 10 μ g/mL) was added into PBS buffer (70 μ L, pH 7.4) as blank control. Then, RNA (10 μ L, 10 μ M) was added, and the mixture was left to equilibrate for 5 min. Protein samples (10 μ L, prepared as mentioned previously) were added and mixed for 30 min before reading the EtBr fluorescence (λ_{ex} = 510 nm, λ_{em} = 590 nm) with a microplate reader (infinite M200, TECAN).

4.2.6 Synthesis of PEI-oxliPt(IV)

Scheme 3. Schematic illustration for the synthesis route of PEI-oxliPt(IV).

The compound oxliPt(IV)-COOH was prepared according to procedures reported in literature.¹³⁴ oxliPt(IV)-COOH (10 mg, 15.8 μ mol) and CDI (5.14 mg, 31.7

μmol) dissolved in anhydrous DMF (1 mL) was stirred at 60 °C under nitrogen for 30 min. PEI₈₀₀ (9.5 mg, 15.8 μmol) dissolved in anhydrous DMF and TEA were then added to the reaction. The mixture was left to stir at 45 °C for 24 hr. Following which, the reaction solvent was removed using rotary evaporation. Resulting crude product was dissolved in deionized water, precipitated with acetone and collected by centrifugation (9000 rpm, 10 min). Final product was obtained by drying in vacuum oven. Yield: 10 mg. The % Pt loading was determined by ¹H NMR and ICP-OES.

4.2.7 Synthesis of PEI-SS

Scheme 4. Schematic illustration for the synthesis route of PEI-SS.

3,3'-Dithiodipropionic acid (7 mg, 33.3 μmol) and CDI (10.8 mg, 66.7 μmol) dissolved in anhydrous DMF (1 mL) was stirred at 60 °C under nitrogen for 30 min. PEI₈₀₀ (20 mg, 33.3 μmol) dissolved in anhydrous DMF and TEA were then added to the reaction. The reaction mixture was left to stir at 45 °C for 24 hr. Then, the reaction solvent was removed using rotary evaporation. Resulting crude product was dissolved in deionized water, precipitated with acetone and collected by centrifugation (9000 rpm, 10 mins). Final product was obtained by drying in vacuum oven. Yield: 5.7 mg. Structure of PEI-SS was determined by ¹H NMR.

4.2.8 Optimization and preparation of prodrug nanocarrier

PEI-oxliPt(IV), GOD, and RNBC were separately dissolved in deionized water at concentrations of 100 mg/mL, 800 mU/mL and 1 mg/mL, respectively. GOD and

RNBC were mixed and vortexed at the GOD/RNBC weight ratio of 1/250 for 10 s. PEI-oxliPt(IV) was then added to the mixture at different PEI-oxliPt(IV):RNBC weight ratios of 5:1 - 40:1. The mixtures were further incubated at room temperature for 1 hr, followed by the dialysis to allow the formation of PEI-oxliPt(IV)@RNBC/GOD. FITC-labeled nanoparticles were prepared by loading FITC-RNBC using the same method as described above. Protein encapsulation efficiency and Pt loading efficiency were measured with UV-Vis absorbance and ICP-OES, respectively. The % protein encapsulation efficiency was calculated as $\frac{(\text{final protein encapsulated})}{(\text{initial protein added})} \times 100$. The % Pt loading efficiency was calculated as $\frac{(\text{final Pt content of nanocarrier})}{(\text{initial Pt of PEI-oxliPt(IV) added})} \times 100$. The particle size and zeta potential of obtained nanoparticles were measured with a zetasizer.

4.2.9 Protein activity assay

Pierce™ quantitative peroxide assay kit was used according to the manufacturer's instruction to determine the enzymatic activity of the encapsulated GOD in various nanocarriers. Samples with GOD concentration of 10 mU/mL were prepared and incubated with different concentrations of glucose (0, 0.1, 0.5 and 1 mg/mL) for 1 hr. Each sample (10 µL) was then mixed with the working reagent solution (100 µL) in a 96-well plate at room temperature for 15 min. The absorbance of the assay components was measured at 595 nm with a microplate reader.

To study the enzymatic activity recovery of RNBC in PEI-oxliPt(IV)@RNBC/GOD, ethidium bromide assay was carried out. PEI-oxliPt(IV)@RNBC and PEI-oxliPt(IV)@GOD were prepared as controls. The nanocarriers were first incubated with 1 mg/mL glucose for 1 hr, then with EtBr-intercalated RNA (10 µM) for 30 min before reading the fluorescence ($\lambda_{\text{ex}} = 510 \text{ nm}$, $\lambda_{\text{em}} = 590 \text{ nm}$) with a microplate reader (infinite M200, TECAN).

RNaseAlert® kit was used to study the kinetic enzymatic activity recovery of RNBC in PEI-oxliPt(IV)@RNBC/GOD. The nanocarrier was incubated with or without glucose (0.1 or 1 mg/mL) for 1 hr (total volume of solution: 80 µL) before the addition of RNaseAlert® Substrate (10 µL) and assay buffer (10 µL) placed in a 96-well plate. The fluorescence intensity at 520 nm (excitation 490 nm) was recorded within 25 min of incubation.

The stability of the enzymes encapsulated in PEI-oxliPt(IV)@RNBC/GOD in serum and lysosomal environment was studied up to 24 hr. FBS and PBS (pH 5) was used to stimulate serum and lysosomal environment respectively. Enzymatic activities of the nanoparticle samples after incubation in the stated test solutions were tested with the Pierce™ quantitative peroxide assay kit and RNaseAlert® kit. Results were expressed as means $\pm$ S.D. in terms of percentage, obtained from 3 replicates for each sample.

4.2.10 Redox-responsive degradation and drug release study

The degradation of protein-loaded nanocarrier was evaluated by studying the size change of the nanoparticles, as well as the release behavior of RNBC and Pt drug from the nanoparticles when incubated in highly reducing environment.

For the size change study, nanocarrier samples containing RNBC (20 µg) and GOD (30 mU) were dispersed in phosphate buffer solution (PBS, pH 7.4, 2 mL) comprising either 0 mM or 10 mM sodium ascorbate. The hydrodynamic size of the samples was recorded at predetermined time intervals.

To investigate the protein release profile, freshly prepared nanocarrier samples (200 µL) containing fluorescein isothiocyanate (FITC) labelled RNBC (FITC-RNBC, 0.2 mg/mL) were contained in the dialysis bag (molecular weight cut-off (MWCO) = 50 kDa) and immersed in PBS (pH 7.4, 2 mL) comprising either 0 mM or 10 mM sodium

ascorbate at 37 °C in the dark. At predefined time intervals, the dialysate (100 µL) was collected with an equivalent volume of incubation medium added to maintain the total solution volume. The accumulative RNBC release concentration was quantified by spectrofluorometer ($\lambda_{\text{ex}} = 490 \text{ nm}$, $\lambda_{\text{em}} = 525 \text{ nm}$).

For the oxaliplatin release study, freshly prepared nanocarrier samples (200 µL) were placed inside a dialysis bag (MWCO = 1000 Da) and incubated in PBS (pH 7.4, 5 mL) comprising either 0 mM or 10 mM sodium ascorbate at 37 °C. At predefined time intervals, the dialysate (100 µL) was collected and replaced with an equivalent volume of incubation medium. Pt content in the collected samples was determined with inductively coupled plasma mass spectrometry (ICP-MS). The percentage drug release was calculated as
$$\frac{(\text{accumulative drug amount in incubation medium})}{(\text{initial drug amount in nanoparticles})} \times 100.$$

Additionally, to confirm that the Pt drug was indeed cleaved from the polymer, ^1H NMR spectra were recorded for PEI-oxliPt(IV) incubated with sodium ascorbate at different periods of time. Briefly, PEI-oxliPt(IV) (8 mg/mL) was dissolved in D_2O (pD = 7.4) and mixed with 10 mM ascorbic acid solution. The reduction products were measured by ^1H NMR spectroscopy at different time points over 24 hr.

4.2.11 Cell culture

Murine 4T1 breast cancer cells were cultured in RPMI-1640 medium supplemented with 10% (v/v) FBS, 0.03% L-glutamine and 1% penicillin-streptomycin under 5% CO_2 atmosphere at 37 °C. The cell culture medium was replaced every two days.

4.2.12 Cellular uptake study

The time-dependent cellular accumulation of PEI-oxliPt(IV)@RNBC/GOD was visualized with confocal laser scanning microscopy (CLSM). 4T1 cells seeded in 8-well

chamber slides (1×10^5 cells/well) were incubated with fresh cell culture medium containing the nanocarrier (10 μ g/mL FITC-RNBC) for different time durations. Before the CLSM observation, cells were first washed with 20 U/mL heparin in PBS thrice. The endosomes/lysosomes were stained with LysoTracker Deep Red (50nM) for 1 hr and nuclei were stained with Hoechst 33342 for 5 min and cells were fixed with 4% paraformaldehyde. Fluorescence was observed for Hoechst 33342 and FITC at 405 nm and 488 nm, respectively. Fluorescence was imaged for LysoTracker Deep Red at 640 nm.

The uptake level was quantified by flow cytometry. After the sample incubation, the culture medium was removed, and the cells were washed with 20 U/mL heparin in PBS thrice before harvested. Cells were collected via centrifugation (3000 rpm, 5 min) and washed with PBS twice, which were then resuspended in PBS (1 mL) and subjected to flow cytometry analysis (excitation wavelength: 488 nm).

The cellular uptake efficiency between PEI-oxliPt(IV)@RNBC/GOD and native RNase A was investigated using CLSM and flow cytometry. Following the incubation with FITC-labelled PEI-oxliPt(IV)@RNBC/GOD or RNase A at protein concentration of 10 μ g/mL for 4 hr, the cells were prepared according to above mentioned procedures.

4.2.13 Determination of cellular platinum contents

4T1 cells seeded in 24-well plates (1×10^5 cells/ well) were incubated with different samples (5 μ g/mL RNBC, 50 μ g/mL Pt drug) at 37 °C for 4 hr. After the stipulated incubation period, cells were washed three times with PBS, harvested by trypsination and collected by centrifugation. Thereafter, 1 mL cell lysis buffer (Invitrogen cell extraction buffer) was used to lyse the cell pellets. The cell lysis solution (100 μ L) of each sample was used in the measurement of Pt content by ICP-MS.

4.2.14 In-vitro cytotoxicity assay

Cytotoxicity of RNase A, PEI-oxliPt(IV), PEI-SS@RNBC/GOD (PEI-SS is a disulfide cross-linked PEI), and PEI-oxliPt(IV)@RNBC/GOD in 4T1 cells was investigated using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. 4T1 cells were first seeded into 96-well plate (1×10^4 cells/well) and incubated with sample-containing medium (100 μ L) at 37 °C for 48 hr. Next, the sample-containing culture medium was replaced with 5 mg/mL MTT solution (100 μ L) and incubated for 4 hr. Finally, the formazan crystal was dissolved in dimethyl sulfoxide (100 μ L). The optical density (OD) was measured at 490 nm using a microplate reader (infinite M200, TECAN). Cells cultured in the absence of samples were taken as the control. The normalized cell viability in % was calculated according to $\frac{OD_{\text{sample}}}{OD_{\text{control}}} \times 100\%$. Results were expressed as means $\pm$ S.D. obtained from 5 replicates for each sample. The combination index (CI) value was calculated using the CalcuSyn software (Biosoft, Cambridge, UK).²³⁰

4.2.15 In-vitro RNA degradation study

Click-iT® RNA Alexa Fluor® 594 imaging kit, prepared according to the manufacturer's instructions, was used to visualize the change in cellular RNA levels after the treatment with PEI-oxliPt(IV)@RNBC/GOD. Briefly, cells seeded in 8-well chamber slides (1×10^4 cells/well) were incubated with PEI-oxliPt(IV)@RNBC/GOD or native RNase A at the protein concentration of 20 μ g/mL for 4 hr. One hour before the fixation, 2x EU working solution was added and the cells were incubated further at 37 °C. The incubation medium was removed, and cells were washed with 20 U/mL heparin in PBS and fixed in 4% paraformaldehyde. Fixed cells were washed with PBS and permeabilized using 0.5% Triton X-100 at room temperature for 10 min, followed by the incubation with Click-iT reaction cocktail in dark at room temperature for 30

min. After the removal of the reaction cocktail, cells were washed once with Click-iT reaction rinse buffer, then thrice with PBS and stained with H33342 for nucleus visualization with CLSM (excitation wavelength 405 nm for Hoechst 33342 and 561 nm for Alexa Fluor™ 594).

The quantification of the cellular RNA level was measured with the Quant-iT™ RiboGreen® RNA assay kit. Cells were seeded in 12-well plate and incubated with different samples as described previously. Cellular RNA was extracted with the PureLink® RNA mini kit and the RNA purified was quantified with Quant-iT™ RiboGreen® RNA kit. All procedures were performed according to the manufacturers' instructions. Results were expressed as mean $\pm$ S.D. obtained from 3 replicates of each sample.

The intracellular cellular of active RNase A was investigated by measuring the RNase A activity in cell lysate using the RNaseAlert® kit. Briefly, 4T1 cells were seeded in 12-well plate and incubated with PEI-oxliPt(IV)@RNBC/GOD and the control samples for 8 hr. The samples-containing medium was removed, and cells were washed with PBS twice. Cells were harvested via centrifugation and dispersed in ice-cold cell lysis buffer (80 μ L) for 30 mins. After centrifugation at 12000 $\times$ g for 1 min, the supernatant was collected for measurement with the assay kit. Results were expressed as mean $\pm$ S.D. obtained from 3 replicates of each sample.

4.2.16 Apoptosis study

4T1 cells seeded into 6-well plate (1×10^6 cells/well) were incubated with PEI-oxliPt(IV), PEI-SS@RNBC/GOD or PEI-oxliPt(IV)@RNBC/GOD at the RNBC concentration of 5 μ g/mL for 24 hr. Cells were then collected with trypsin and washed three times with PBS. After resuspended in the binding buffer (100 μ L), cells were

stained with Annexin V-FITC/propidium iodide dyes at room temperature in the dark for 15 min. Finally, the number of apoptotic cells was analyzed using a flow cytometer.

4.2.17 Caspase-3 assay

Caspase-3 activity in the 4T1 cell lysate was assessed with a colorimetric assay kit (Abcam) prepared according to the manufacturer's instructions. 4T1 cells were seeded into 12-well plate (1×10^6 cells/well) and cultured for 24 hr. After replacing the culture medium, cells were incubated with PEI-oxliPt(IV), PEI-SS@RNBC/GOD or PEI-oxliPt(IV)@RNBC/GOD at the RNBC concentration of 5 $\mu\text{g/mL}$ for 4 hr. Cells were harvested by trypsin, collected by centrifugation and lysed with the lysis buffer on ice for 10 min. Samples were then centrifuged at 10,000 g for 5 min and the supernatant was collected. Final protein concentration was determined by the BCA kit, in which an equivalent amount of protein (100 μg) from each sample was incubated with the caspase reaction buffer and substrate at 37°C for 2 hr. The absorbance was read at 490 nm using a microplate reader and the results were obtained from 3 replicates.

4.3 Results and Discussion

4.3.1 Synthesis and characterization

For the fabrication of a cationic oxaliplatin(IV) prodrug-based polymer to complex and encapsulate the RNBC, PEI 800 Da was cross-linked by a oxaliplatin(IV) prodrug (Scheme 3) to obtain PEI-oxliPt(IV) with a zeta potential of +12.5 mV and a molecular weight of 2220 Da as measured by MALDI-TOF. Characteristics resonance peaks between 1.1 to 1.5 ppm ascribed to the 1,2-diaminocyclohexane ligand of oxaliplatin could be found in the ^1H NMR spectrum of PEI-oxliPt(IV) (Figure 1). In addition, the emergence of new peaks between 3.0 to 3.4 ppm which were attributed to the methylene protons of (-N-CH₂CH₂NH-CO-) in the side chain of PEI confirmed the successful crosslinking reaction. The Pt content in the polymer was found to be 25 wt% as measured by ICP-MS.

Figure 1. ^1H NMR spectrum of PEI-oxliPt(IV), PEI and oxliPt(IV)-COOH.

RNBC was synthesized by caging the lysine residue on RNase A with H₂O₂-sensitive aryl boronic acid via a covalent carbamate linker (Scheme 2) and characterized by zeta potential, circular dichroism (CD), sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), Alizarin Red S (ARS) assay and matrix-assisted laser desorption ionization time-of-flight mass spectrometry (MALDI-TOF MS). Zeta potential changed from + 4 mV to – 25 mV after the reaction, owing to the consumption of free positively charged lysine after the conjugation, thereby demonstrating successful surface modification on the protein with the caged group. The modification with aryl boronic acid reduces the isoelectric point of RNase A, facilitating electrostatic interactions with positively charged PEI-oxliPt(IV) to form the nanocomplex.

The CD spectrum of RNBC was similar to that of native RNase A (Figure 2A), suggesting that the conjugation process did not modify the secondary structure of the protein. This was further supported by the SDS-PAGE results (Figure 2B), where the band corresponding to RNBC appeared at slightly higher molecular weight than RNase A. The presence of this gel band signifies that the NBC modification was not detrimental to the protein structure.

To confirm the presence of surface aryl boronic acid group on RNBC, the ARS assay based on the binding between the catechol moieties in ARS and aryl boronic acid residues to generate fluorescent complexes was conducted (Figure 2C).²⁸⁰

Figure 2. Characterization of RNBC (A) Circular dichroism spectra of free RNase A and RNBC. (B) SDS-PAGE of free RNase A and RNBC. The protein ladder lane contains a mixture of proteins (10 to 260 kDa) for use as molecular weight standards. Concentrations of RNase A and RNBC were 1 mg/mL. (C) Schematic illustration for the fluorescence detection of boronic acid by Alizarin Red S (ARS). (D) Fluorescence of ARS (0.0025% w/v) in the presence of H₂O₂, deionized (DI) H₂O, native RNase A, and RNBC with or without the treatment with H₂O₂. (E) MALDI-TOF mass spectra of native RNase A, and RNBC before and after the H₂O₂ treatment.

Figure 2D shows that the fluorescence intensity of ARS when added with RNBC was notably higher than that of RNase A, which proved that boronic acid was present in RNBC. Comparatively, the fluorescence enhancement of ARS was not observed when RNBC was pretreated with H_2O_2 , indicating the absence of aryl boronic group on the uncaged RNBC.

MALDI-TOF analysis (Figure 2E) showed an increase in molecular weight from 13235 Da (RNase A) to 14800 Da (RNBC), indicating the conjugation of eight NBC molecules per protein molecule. Upon treatment with H_2O_2 , the MALDI-TOF mass peaks decreased back to molecular weight that corresponded to native RNase A, indicating the cleavage of NBC and restoration to the native protein. Combining these results, we have validated that the modified RNBC was successfully synthesized and could be uncaged by H_2O_2 treatment.

4.3.2 Uncaging of RNBC by GOD and glucose

Before the preparation of the nanocomplex, we investigated the efficiency of uncaging RNBC by the enzymatic production of H_2O_2 from free GOD and glucose.

Figure 3. Concentration- dependent study of H_2O_2 generation from free GOD and glucose.

Figure 3 shows the increase in absorbance of H_2O_2 indicator dye when free GOD was incubated with increasing concentrations of glucose, demonstrating that the generation of H_2O_2 by GOD is dependent on glucose concentration.

Figure 4. Assessing RNA-degrading activity of uncaged RNBC (A) Native-PAGE analysis of 32-mer RNA treated with different samples. Lane 1: control; Lane 2: free RNase A; Lane 3: RNBC; Lane 4: RNBC pretreated with GOD and glucose; Lane 5: GOD incubated with glucose; Lane 6: RNBC pretreated with H_2O_2 ; Lane 7: H_2O_2 . (B) Uncaging RNBC with GOD and glucose oxidase. Enzymatic activity of uncaged RNBC measured by an ethidium bromide assay. Data points represent mean $\pm$ SD ($n = 3$). (C) RNase A activity assay of native RNase A, and RNBC with and without GOD/glucose treatment.

The RNA-degrading capability of RNBC was first evaluated using the Native-polyacrylamide gel electrophoresis (Native-PAGE). As shown in Figure 4A, the RNA band disappeared when RNA was treated with free RNase A, indicating the capability of free RNase A to degrade RNA. Conversely, the band was still conspicuous after the incubation with RNBC, demonstrating that the aryl boronic acid unit could cage the activity of the protein to inhibit its RNA degradation capacity. After the incubation with H_2O_2 , however, the band was no longer visualized as RNA was cleaved by H_2O_2 -uncaged RNBC. Importantly, the preincubation of RNBC with 1 mg/mL glucose and GOD also showed the cleavage of free RNA, confirming that GOD was able to produce

H₂O₂ to uncage RNBC. Additionally, two-fold decrease in the fluorescence intensity of ethidium bromide when RNA was incubated with GOD and glucose pretreated RNBC proved the cascade recovery of the protein activity by GOD and glucose integrated system (Figure 4B).²⁸¹

The enzyme activity of uncaged RNBC was then measured by a commercially available fluorometry-based RNase A kit. The kit consists of an oligonucleotide substrate with an attached fluorophore and a quencher. Its incubation with RNase A-containing sample would result in the cleavage of oligonucleotide and the recovery of fluorescence. The greater the fluorescence intensity, the higher the RNase A activity. As shown in Figure 4C, the enzymatic activity of RNBC was drastically diminished (~ 80 %) as compared to native RNase A, which was recovered after the treatment with GOD and glucose over time. Thus, the suppression of the enzymatic activity and its site-specific restoration were successfully realized through the caging and uncaging strategy.

4.3.3 Preparation and characterization of prodrug nanocomplex

After verifying that RNBC could be uncaged to form RNase A, the nanocomplex synthesis was optimized. The negatively charged RNBC and GOD facilitate the formation of nanocomplex with PEI-oxliPt(IV) via electrostatic interactions to yield PEI-oxliPt(IV)@RNBC/GOD. The main role of GOD in the nanocomplex was to produce sufficient H₂O₂ to recover the activity of RNBC and not to potentiate the oxidation or starvation therapy. Hence, it was important to control the GOD concentration and minimize its ROS-related cytotoxicity

Figure 5. Cell viability of 4T1 cells after treatment with free GOD for 24 and 48 hr. Data points represent mean $\pm$ SD (n = 5).

The concentration of GOD below 3 mU/mL was deemed to have negligible cytotoxicity, and this concentration (3 mU/mL GOD) was employed in the fabrication of the nanocomplex (Figure 5).

Table 1. Properties of PEI-oxliPt(IV)@RNBC/GOD nanocomplex comprising different ratios of PEI-oxliPt(IV) prodrug to RNBC protein.

PEI-oxliPt(IV) : RNBC	5:1	10:1	20:1	40:1
Protein encapsulation efficiency	30 %	67 %	82 %	94 %
Pt loading efficiency	84.0 %	86.5 %	86.8 %	86.9 %
Size / nm (PDI)	290 (0.26)	219 (0.23)	217 (0.24)	203 (0.11)
Zeta / mV	3.3 $\pm$ 1.0	8.8 $\pm$ 1.9	10.9 $\pm$ 0.4	12.5 $\pm$ 0.3

Nanocomplex samples with different weight ratio of PEI-oxliPt(IV) to RNBC were prepared and their properties were assessed (Table 1). Considering higher protein encapsulation efficiency, Pt loading content and suitable hydrodynamic size, the

nanocarrier comprising weight ratio of 40 : 1 PEI-oxliPt(IV): RNBC was chosen for further studies.

Figure 6. Characterization of PEI-oxliPt(IV)@RNBC/GOD nanocomplex. (A) Fluorescence spectra of RhB-GOD, FITC-RNBC and PEI-oxliPt(IV)@RNBC/GOD with the same protein content. GOD and RNBC were pre-labelled with RhB and FITC, respectively. The spectra were recorded with the excitation at 450 nm. (B) TEM image of the PEI-oxliPt(IV)@RNBC/GOD nanoparticles.

Successful encapsulation of proteins in the nanocomplex was verified using FITC-RNBC and rhodamine-B labelled GOD (RhB-GOD) based on Förster resonance energy transfer (FRET) phenomenon.²⁸² Figure 6A presents the fluorescence spectra of individual proteins as well as the PEI-oxliPt(IV)@RNBC/GOD nanocomplex when excited at 490 nm. The excitation maxima of FITC and RhB occur at 495 and 540 nm, respectively. Thus, only FITC-RNBC exhibits fluorescence, while negligible fluorescence is observed from RhB-GOD. On the contrary, the nanocomplex spatially makes two proteins in close proximity, resulting in intense emission from both FITC and RhB at 525 and 585 nm via effective FRET from FITC-RNBC to RhB-GOD.

Figure 7. DLS of PEI-oxliPt(IV)@RNBC/GOD nanoparticles in water.

Transmission electron microscopy (TEM, Figure 6B) image shows a spherical morphology of PEI-oxliPt(IV)@RNBC/GOD with an average hydrodynamic size of 203 nm measured from dynamic laser scattering (Figure 7).

The enzymatic activities of the encapsulated proteins in PEI-oxliPt(IV)@RNBC/GOD were then studied. The production efficiency of H_2O_2 by PEI-oxliPt(IV)@RNBC/GOD was carried out by varying the concentrations of glucose (0 – 1 mg/mL). As illustrated in Figure 8A, free GOD showed a glucose-concentration dependent production of H_2O_2 . Comparatively, PEI-oxliPt(IV)@RNBC/GOD demonstrated lower catalytic production of H_2O_2 than free GOD, but significantly higher than sole PEI-oxliPt(IV) or PEI-oxliPt(IV)@RNBC (Figure 8B). Up to 65 % activity of GOD in PEI-oxliPt(IV)@RNBC/GOD was still retained, indicating that the polymeric carrier only partially shielded the catalytic site of GOD after the encapsulation, and high enzymatic production of H_2O_2 could still be achieved.

Figure 8. (A) Glucose concentration- dependent generation of H₂O₂ from native GOD and PEI-oxliPt(IV)@RNBC/GOD measured with a Quantitative Peroxide Assay Kit. (B) H₂O₂ production from native GOD, PEI-oxliPt(IV), PEI-oxliPt(IV)@GOD, PEI-oxliPt(IV)@RNBC and PEI-oxliPt(IV)@RNBC/GOD nanoparticles incubated with 0.5 mg/mL glucose, measured

with a peroxide assay kit. (C) Examination of RNA degradation efficiency by PEI-oxliPt(IV)@RNBC/GOD when incubated with different glucose concentrations, measured by RNase A activity assay kit. (D) Enzymatic activity of uncaged RNBC in PEI-oxliPt(IV)@RNBC, PEI-oxliPt(IV)@GOD and PEI-oxliPt(IV)@RNBC/GOD, measured by an ethidium bromide assay. Data points represent mean $\pm$ SD (n = 3). (E) Activity of GOD and RNase A in PEI-oxliPt(IV)@RNBC/GOD nanocomplex after 24 hr incubation in FBS and simulated lysosome environment (pH 5).

The feasibility of the tandem uncaging of RNBC in PEI-oxliPt(IV)@RNBC/GOD was subsequently investigated by conducting the kinetic profile of RNA degradation when incubated with the nanocomplex. The RNA having fluorescein and dark quencher on its two ends would not fluoresce under intact state. However, the fluorescence could be regained when it is degraded. As illustrated in Figure 8C, the fluorescence was greatly enhanced when PEI-oxliPt(IV)@RNBC/GOD was incubated with higher glucose concentrations, demonstrating the recovery of RNBC enzymatic capability and its reliance on the glucose level to degrade RNA. Comparing with control carriers that only encapsulate single protein species, PEI-oxliPt(IV)@RNBC/GOD pretreated with glucose showed the greatest fluorescence decrease in the ethidium bromide assay (Figure 8D), validating the vital role of GOD in the nanocomplex to reactivate RNBC.

A major function of a nanocarrier for protein delivery is to provide protection of the protein's activity before reaching the target site. Hence, the stability of the encapsulated proteins inside PEI-oxliPt(IV)@RNBC/GOD was investigated in stimulated serum and lysosomal environment. The nanoparticles were incubated up to 24 hr in fetal bovine serum (FBS) and PBS (pH 5), to mimic stimulated serum and lysosomal environment respectively. As shown in Figure 8E, the capabilities of (1) GOD to produce H_2O_2 and (2) RNase A to degrade RNA were largely well-maintained after

prolonged incubation in physiological conditions, indicating the dependable ability of PEI-oxliPt(IV)@RNBC/GOD to deliver functional proteins intracellularly.

4.3.4 Redox-responsive degradation and drug release study

Figure 9. (A) Time-dependent reduction of PEI-oxliPt(IV) by sodium ascorbate monitored with ^1H NMR spectra. Nanoparticle size changes of PEI-oxliPt(IV)@RNBC/GOD with time in (B) PBS and (C) 10 mM sodium ascorbate at 37 °C monitored by DLS. (D) FITC-tagged RNBC protein release profile from PEI-oxliPt(IV)@RNBC/GOD in the absence and presence of sodium ascorbate, measured by fluorescence spectroscopy. (E) Pt drug release profile from PEI-oxliPt(IV)@RNBC/GOD in the absence and presence of sodium ascorbate (Vc), measured with ICP-MS.

To verify the redox-responsiveness of the nanocomplex originating from the reducible oxaliplatin prodrug, the reduction of PEI-oxliPt(IV) was monitored over time with ^1H NMR. As shown in the ^1H NMR spectra of PEI-oxliPt(IV) incubated with 10 mM sodium ascorbate (Figure 9A), the chemical shift of sodium ascorbate at 4.23 ppm disappeared over time, implying the depletion of the reducing agent in the mixture. Furthermore, the concomitant time-dependent decrease of peak signals in 2.3 to 3.0 ppm region attributed to the $-\text{CH}_2$ of PEI backbone confirmed the reduction of PEI-oxliPt(IV).

To demonstrate intracellular responsive disassembly of PEI-oxliPt(IV)@RNBC/GOD, the size change of the nanoparticles was monitored in the presence and absence of sodium ascorbate, an overexpressed cellular reducing agent.¹⁸⁵ In PBS only, PEI-oxliPt(IV)@RNBC/GOD showed relatively unchanged DLS size over time (Figure 9B), indicating the stability of the nanocomplex. In contrast, the nanocomplex presented a progressive size increment within 12 hr, and then decreased to less than 5 nm when incubated with 10 mM sodium ascorbate for 24 hr (Figure 9C). This observation could be explained by the reduction of the polymer carrier with an initial volume expansion followed by the release of the encapsulated proteins upon the disruption of the polymer coating.

We then studied how the redox-induced dissociation of the nanocomplex would influence the release of the encapsulated proteins and drug. As shown in Figure 9D, the release of RNBC in the absence of sodium ascorbate was negligible, and only 6.2 % protein release was attained within 56 hr. In comparison, a four-fold increase in the accumulative release of RNBC was obtained in the presence of 10 mM sodium ascorbate within the same incubation time. Similarly, an accelerated release of Pt drug species was observed when PEI-oxliPt(IV)@RNBC/GOD was incubated under highly reducing

condition (Figure 9E). Such redox-responsive RNBC and Pt drug release upon the degradation of PEI-oxliPt(IV)@RNBC/GOD under the reducing condition is useful for controlled release of the therapeutics upon internalization into cancer cells. Hence, the introduction of oxliPt(IV) as a crosslinker in the polymeric carrier is favorable for the encapsulation of protein therapeutics and could facilitate site-specific release of active protein and oxaliplatin(II) drug.

4.3.5 Cellular uptake study and Pt accumulation

Figure 10. Intracellular accumulation of PEI-oxliPt(IV)@RNBC/GOD nanoparticles. (A) Time-dependent cellular uptake of FITC-labelled PEI-oxliPt(IV)@RNBC/GOD. Nucleus was stained with H33342. NP: nanoparticles. Scale bar: 20 μ m. (B) Quantification of time-dependent intracellular accumulation by flow cytometry. (C) CLSM images of 4T1 cells after incubation with PEI-oxliPt(IV)@RNBC/GOD nanoparticles. Nucleus was stained with H33342; FITC-

labelled nanoparticles give green fluorescence; Endosomes and lysosomes stained by LysoTracker Deep Red. NP: nanoparticles. Scale bar: 20 μ m. White arrows show co-localization of nanoparticles and endolysosomes. Red arrows show escape of nanoparticles from the endolysosomes.

Cellular accumulation of PEI-oxliPt(IV)@RNBC/GOD in 4TI murine breast cancer cells was evaluated by CLSM imaging and flow cytometry using FITC-labelled proteins. The cellular uptake of PEI-oxliPt(IV)@RNBC/GOD was shown to be time-dependent (Figure 10A). Upon increasing time, the fluorescence observed from the nanocomplex increased, demonstrating the capability of the nanocomplex to accumulate in cells (Figure 10B). Furthermore, PEI-oxliPt(IV)@RNBC/GOD also showed the ability to escape the endosomes to get into the cytosol after endocytosed into the cells. As shown in Figure 10C, the co-localization of endolysosomes stained with lysotracker Deep Red and the FITC-labelled nanoparticles indicated the subcellular localization of the nanocomplex in the endosomes and lysosomes after 4 hr incubation. At 8 hr post-incubation, the nanocomplex escaped from the endolysosomes to the cytosol, as observed by the separation of green fluorescence (FITC-labelled nanoparticles) from the red fluorescence (lysotracker-stained endolysosomes), likely due to the PEI carrier-mediated “proton sponge” effect.²⁸³

Figure 11. Intracellular accumulation of PEI-oxliPt(IV)@RNBC/GOD nanoparticles. (A) Enhanced cellular uptake of FITC-labelled PEI-oxliPt(IV)@RNBC/GOD as compared to carrier-free RNase A. Nucleus was stained with H33342. NP: nanoparticles. Scale bar: 20 μ m. (B) Quantification of intracellular accumulation of FITC-labelled nanoparticles and RNase A after 4 hr incubation with 4T1 cells measured by flow cytometry. (C) Quantification of intracellular Pt accumulation measured by ICP-MS.

The efficiency of cellular uptake of free RNase A and PEI-oxliPt(IV)@RNBC/GOD was also evaluated. As shown in Figure 11A, negligible green fluorescence was observed from cells treated with free RNase A. In the case of PEI-oxliPt(IV)@RNBC/GOD, the fluorescence was greatly enhanced, indicating that the protein was notably internalized and accumulated in 4T1 cells. The preferential uptake of PEI-oxliPt(IV)@RNBC/GOD nanoparticles as compared to free RNase A was also confirmed by flow cytometry analysis (Figure 11B). RNase A treated cells showed comparable mean fluorescence intensity to that of the control, while cells incubated with PEI-oxliPt(IV)@RNBC/GOD exhibited four-fold fluorescence intensity enhancement, suggesting that the protein delivered by the nanocomplex had more efficient cellular internalization. Similar results were obtained from cells treated with PEI-

oxliPt(IV)@RNBC/GOD using ICP-MS analysis, whereby the presence of cellular Pt accumulation from the nanocarrier validated the internalization into cells (Figure 11C).

4.3.6 In-vitro cytotoxicity assay

Encouraged by the promising in-vitro accumulation results, cytotoxic effect of PEI-oxliPt(IV)@RNBC/GOD in 4T1 cells was fully investigated by cell viability assay, cellular RNA degradation, and apoptosis-related protein caspase-3 activity assay. The capability of PEI-oxliPt(IV)@RNBC/GOD to inhibit cancer cell proliferation by combined chemotherapy and protein therapeutics was assessed by the standard MTT assay. To evaluate the effectiveness of combinational therapy by PEI-oxliPt(IV)@RNBC/GOD over monotherapy, PEI-oxliPt(IV) and PEI-SS@RNBC/GOD were prepared as control groups that deliver only the drug and protein cargoes, respectively. PEI-SS is a control polymer carrier without therapeutic efficacy, but degradable via the disulfide linkage (Scheme 4).

Figure 12. ^1H NMR spectrum of PEI-SS.

The successful synthesis of PEI-SS polymer was confirmed by ^1H NMR (Figure 12) before employed in the fabrication of the proteins-encapsulated nanocarriers.

Figure 13. In-vitro cell viability of 4T1 cells after 48 hr treatment with PEI-SS control polymer and single or dual protein loaded nanocarriers.

The incubation of 4T1 cells with PEI-SS@RNBC and PEI-SS@RNBC/GOD resulted in 80% and 40% cell viability respectively, with no significant cytotoxicity from the PEI-SS carrier itself (Figure 13). This result reveals that the inclusion of GOD in the nanocomplex is important, as the reliance on intracellular H_2O_2 is insufficient for effective uncaging of RNBC.

Figure 14. Cellular studies of treatment efficacy of PEI-oxliPt(IV)@RNBC/GOD. (A) Cell viability of 4T1 cells after 48 hr incubation with concentration-dependent PEI-oxliPt(IV), PEI-SS@RNBC/GOD and PEI-oxliPt(IV)@RNBC/GOD. Data points represent mean $\pm$ SD (n = 5). (B) Combination index (CI) analysis. CI versus Fraction affected (Fa) plots were generated

using the method of Chou and Talalay to determine the extent of synergism for PEI-oxliPt(IV)@RNBC/GOD in 4T1 cells. Dotted line indicates a reference point of CI value of 1. (C) Cell viability of 4T1 cells after 48 hr incubation with native RNase A.

As illustrated in Figure 14A, the cell viability of 4T1 cells treated with PEI-oxliPt(IV)@RNBC/GOD decreased in a concentration-dependent manner. As compared to PEI-oxliPt(IV) and PEI-SS@RNBC/GOD, PEI-oxliPt(IV)@RNBC/GOD displayed stronger cytotoxicity at equivalent therapeutics concentrations. The half maximal inhibitory concentration (IC_{50}) of PEI-oxliPt(IV)@RNBC/GOD against 4T1 cells was determined to be 0.23 $\mu\text{g/mL}$ with an improvement to that of PEI-oxliPt(IV) ($IC_{50} = 0.42 \mu\text{g/mL}$) and PEI-SS@RNBC/GOD ($IC_{50} = 0.36 \mu\text{g/mL}$). In addition, PEI-oxliPt(IV)@RNBC/GOD exhibited combination index (CI) values below 1 at tested concentrations, indicating the synergism between the prodrug and the protein therapeutics (Figure 14B). The improved anticancer proliferation effect could be attributed to synergistic mechanisms of cell killing by oxaliplatin and RNase A. Notably, PEI-oxliPt(IV)@RNBC/GOD treated cells attained 80 % cell killing efficiency at a low protein concentration (5 $\mu\text{g/mL}$). On the other hand, the cell viability was consistently high even when treated with free RNase A at high concentrations (Figure 14C). This comparison indicates the vital role of this delivery carrier for intracellular delivery of protein, as free protein could not traffic into cells easily.

4.3.7 In-vitro RNA degradation study

Figure 15. In-vitro RNA degradation efficiency. (A) Confocal images of cellular RNA content in 4T1 cells after the incubation with indicated samples for 4 hr. RNA (red) was detected with the Click-iT® RNA Alexa Fluor® 594 Imaging Kit, and nucleus (blue) was stained with Hoechst 33342. NP represents PEI-oxliPt(IV)@RNBC/GOD nanoparticles. Scale bar: 50 μ m. (B) Quantification of RNA content in 4T1 cells after the incubation with free RNase A or PEI-oxliPt(IV)@RNBC/GOD for 4 hr. Data points represent mean $\pm$ SD (n = 3).

Then, the RNA degradation in cells was evaluated using a cellular RNA imaging assay, which allows the detection of intracellular mRNA spatially. Compared with untreated cells, the fluorescence intensity of the RNA imaging probe in PEI-oxliPt(IV)@RNBC/GOD treated cells was greatly reduced, indicating that a large proportion of mRNA was degraded (Figure 15A). On the other hand, the fluorescence intensity in free RNase A treated cells was comparable to the control group. Quantitatively, the treatment with RNase A and PEI-oxliPt(IV)@RNBC/GOD showed one-fold and seven-fold reduction of RNA levels in 4T1 cells, respectively (Figure 15B).

Figure 16. Examination of RNA degradation efficiency by RNase A contained in cell lysate extracted from 4T1 cells incubated with the different samples, measured by RNase A activity assay kit.

The intracellular release of RNase A was indirectly confirmed by examining the RNase A activity of the cell lysate extracted from 4T1 cells. As shown in Figure 16, the fluorescence intensity recovery due to cleavage of RNA probe was distinctly higher from cell lysate extracted from PEI-oxliPt(IV)@RNBC/GOD treated 4T1 cells than the control samples, indicating the successful release of active RNase A from the nanocomplex after endocytosis. High cellular RNA cleavage after the incubation with the nanocomplex clearly indicated that the therapeutic protein was successfully delivered into the cells, with good retention of its activities when released from the carrier.

4.3.8 Apoptosis study and Caspase-3 assay

Figure 17. (A) Bar chart showing the percentage of viable, apoptotic and necrotic cells after the incubation with indicated samples for 48 hr, measured by flow cytometry. (B) Caspase 3 levels in 4T1 cells after 4 hr exposure to indicated samples. Data points represent mean $\pm$ SD (n = 3).

Apoptosis assay revealed that PEI-oxliPt(IV)@RNBC/GOD achieved much higher apoptosis rate than that of PEI-SS@RNBC/GOD and PEI-oxliPt(IV). While PEI-SS@RNBC/GOD and PEI-oxliPt(IV) were able to induce 60.4 % and 77.1 % apoptotic and necrotic cells respectively, PEI-oxliPt(IV)@RNBC/GOD displayed highly enhanced apoptosis-inducing ability as evidenced by its apoptosis rate of 94 % (Figure 17A). The enhanced cytotoxicity of the nanocomplex could be attributed to synergistic effect of chemotherapy and cytotoxic protein treatment, which was consistent with the MTT results.

To confirm the apoptotic effect of PEI-oxliPt(IV)@RNBC/GOD, the activity of caspase-3 enzyme was analyzed (Figure 17B). The examination of caspase-3 activity, a marker for apoptosis, revealed that PEI-oxliPt(IV)@RNBC/GOD could induce caspase-3 activity drastically as compared to untreated control group, promoting high activity of

apoptosis. Consistent with above data, the delivery of PEI-oxliPt(IV)@RNBC/GOD into the cancer cells induced obvious cell death.

4.4 Conclusion

In summary, we have developed a nanocomplex (PEI-oxliPt(IV)@RNBC/GOD) that can adaptively modulate its own property by responding to endogenous stimuli in order to achieve controlled drug release and activation of protein activity. Incorporated with a degradable oxaliplatin prodrug polymer carrier, reversibly caged RNase A protein could be mediated for the RNA cleavage after being uncaged with self-supplied H₂O₂ catalyzed by the encapsulated GOD. Contrary to conventional protein delivery systems, PEI-oxliPt(IV)@RNBC/GOD is able to ensure the integrity and activity retention of proteins during the delivery and then enable facile restoration of the protein activity after the cleavage of H₂O₂ responsive phenylboronic acid unit. Furthermore, highly reducing cellular environment could convert the oxaliplatin prodrug to active oxaliplatin drug. The degradation of the nanocomplex in vitro leads to simultaneous release of both the platinum drug and therapeutic RNase A. Thus, the smartly designed nanocomplex could enhance cellular accumulation, promote redox-responsive release of the therapeutics, initiate caspase-3 activity, and result in effective anticancer effect. The present work would provide a new strategy for the delivery of combinational anticancer therapeutics.

Chapter 5

Conclusion and Future Outlook

In view that newly designed platinum complexes functioning on DNA binding are unlikely to have actual therapeutic relevance as they could not provide obvious clinical advantages over existing ones; it is more advantageous to improve on the anticancer efficacy of current platinum-based drugs. This is judiciously achieved by selective accumulation and site-specific cytotoxicity activation of the platinum-based drugs. Accordingly, constructing nanosized prodrug aggregate delivery systems appears to be a strategic approach by combining the advantages of both prodrugs and nanotechnology. With supramolecular chemistry and nature of self-assembly, nanoscale multicomponent assemblies could be designed and developed with increased system complexities together with enhanced integrated functions of the molecular components. In this dissertation, the author proposed and evaluated different approaches to preparing self-assembled delivery systems to better deliver the platinum-based drug and other therapeutics.

In Chapter 2, the preparation of novel oxaliplatin(IV) prodrug and formulation into self-delivery nanocarrier on the basis of self-assembly chemistry was explored. Generally, the synthesis and purification of amphiphilic prodrugs is tedious and complex. Instead, the author proposed the use of host-guest chemistry to develop supramolecular amphiphiles nanocarriers for targeted cancer therapy. By choosing an appropriate cancer cells targeting ligand and a hydrophobic moiety, namely spermine and cholesterol in this study to fabricate the host molecule and prodrug guest respectively, the inherently amphiphilic supramolecular complex aggregated to form nanoparticles. The spermine targeting ligand facilitated effective accumulation of the nanoparticles in cancer cells with overexpressed polyamine transporters accompanied by controlled drug release, thereby successfully improved treatment efficacy.

In addition, to overcome the challenge of realizing optimal loading ratio of the therapeutics for combinational therapy, a design strategy in Chapter 3 to construct a carrier-free therapeutic system with control over the drug loading amount by leveraging the specificity of host-guest chemistry was presented. The oxaliplatin drug and a porphyrin photosensitizer were modified to form a host-guest complex and subsequently self-assembled to give the resulting drug carrier, which encompassed chemotherapy and photodynamic therapy. Our system achieved synergistic therapy in-vitro and in-vivo with precise ratiometric control over drug loading, paving the way for potential clinical translation of nano-size carrier in combination therapy.

Lastly in Chapter 4, an oxaliplatin(IV) polymeric prodrug-based nanocarrier to facilitate effective delivery of protein therapeutics was developed. The treatment capabilities of therapeutic proteins have been highly sought after but the efficacy has been seriously impeded by the low tumour accumulation. Typically, proteins are genetically altered to be super positively charged or encapsulated in cationic polymers for effective cellular translocation. Adapting this concept, a cationic oxaliplatin(IV) polymeric prodrug and modified therapeutic proteins was prepared, which can interact to form a hierarchical self-assembled nanocomplex. Results obtained demonstrated superior cellular accumulation of the therapeutic proteins when encapsulated in our proposed system and exhibited better treatment efficacy as utilized concurrently with the controlled-released oxaliplatin drug. This strategy offers an alternative outlook for combinational chemo-protein therapy.

In conclusion, this dissertation unravelled new approaches to designing platinum prodrug-based delivery systems for cancer therapy. The facile integration of different axial ligands to give novel oxaliplatin prodrugs offers supramolecular self-assembly opportunity or self-complexation with therapeutic agents to form nano-drug delivery

vehicles. The customizable properties and varied therapies combinations that can be designed with our approach will be promising for future drug delivery research. Still, various issues regarding developing a prodrug-based delivery system are left unanswered. Major challenges include scaling up the synthesis and production of the chemicals and nanoformulations; long-term chemical stability; and understanding the relevant pharmacokinetics in vivo. In addition, current prodrug-based assemblies often undergo disassembly upon introduction into blood plasma. More research is required to improve the structural integrity of self-assembled systems following systemic administration. Ultimately, it is of great hope to push the development of prodrug-based delivery system past the infancy stage and be applied clinically in the future.

List of publications

1. **Lim, W. Q.**; Phua, S. Z. F.; Zhao, Y. *ACS Appl. Mater. Interfaces* **2019**, *11* (35), 31638-31648.
2. **Lim, W. Q.**; Yang, G.; Phua, S. Z. F.; Chen, H.; Zhao, Y. *ACS Appl. Mater. Interfaces* **2019**, *11* (18), 16391-16401.
3. Nakayama, M.; **Lim, W. Q.**; Kajiyama, S.; Kumamoto, A.; Ikuhara, Y.; Kato, T.; Zhao, Y. *ACS Appl. Mater. Interfaces* **2019**, *11* (19), 17759-17765.
4. **Lim, W. Q.**; Phua, S. Z. F.; Chen, H.; Zhao, Y. *Chem. Commun.* **2018**, *54* (90), 12762-12765.
5. **Lim, W. Q.**; Phua, S. Z. F.; Xu, H. V.; Sreejith, S.; Zhao, Y. *Nanoscale* **2016**, *8* (25), 12510-12519.
6. Wang, D.; Wu, H.; **Lim, W. Q.**; Phua, S. Z. F.; Xu, P.; Chen, Q.; Guo, Z.; Zhao, Y. *Adv. Mater.* **2019**, *31* (27), 1901893.
7. Yang, G.; Phua, S. Z. F.; **Lim, W. Q.**; Zhang, R.; Feng, L.; Liu, G.; Wu, H.; Bindra, A. K.; Jana, D.; Liu, Z.; Zhao, Y. *Adv. Mater.* **2019**, *31* (25), 1901513.
8. Phua, S. Z. F.; Xue, C.; **Lim, W. Q.**; Yang, G.; Chen, H.; Zhang, Y.; Wijaya, C. F.; Luo, Z.; Zhao, Y. *Chem. Mater.* **2019**, *31* (9), 3349-3358.
9. Phua, S. Z. F.; Yang, G.; **Lim, W. Q.**; Verma, A.; Chen, H.; Thanabalu, T.; Zhao, Y. *ACS Nano* **2019**, *13* (4), 4742-4751.
10. Su, Y.; Yu, J.; Li, Y.; Phua, S. F. Z.; Liu, G.; **Lim, W. Q.**; Yang, X.; Ganguly, R.; Dang, C.; Yang, C.; Zhao, Y. *Commun. Chem.* **2018**, *1* (1), 12.
11. Tham, H. P.; Xu, K.; **Lim, W. Q.**; Chen, H.; Zheng, M.; Thng, T. G. S.; Venkatraman, S. S.; Xu, C.; Zhao, Y. *ACS Nano* **2018**, *12* (12), 11936-11948.
12. Zhao, L.; Yuan, W.; Li, J.; Yang, L.; Su, Y.; Peng, J.; Chen, R.; Tham, H. P.; Chen, H.; **Lim, W. Q.**; Xiang, H.; Xing, P.; Li, F.; Zhao, Y. *Adv. Funct. Mater.* **2018**, *28* (50), 1806162.
13. Su, Y.; Phua, S. Z. F.; Li, Y.; Zhou, X.; Jana, D.; Liu, G.; **Lim, W. Q.**; Ong, W. K.; Yang, C.; Zhao, Y. *Sci. Adv.* **2018**, *4* (5), eaas9732.
14. Zhang, Y.; Yang, D.; Chen, H.; **Lim, W. Q.**; Phua, F. S. Z.; An, G.; Yang, P.; Zhao, Y. *Biomaterials* **2018**, *163*, 14-24.
15. Zhou, J.; Li, M.; **Lim, W. Q.**; Luo, Z.; Phua, S. Z. F.; Huo, R.; Li, L.; Li, K.; Dai, L.; Liu, J.; Cai, K.; Zhao, Y. *Theranostics* **2018**, *8* (2), 518-532.

16. Zheng, C.; Wang, Y.; Phua, S. Z. F.; **Lim, W. Q.**; Zhao, Y. *ACS Biomater. Sci. Eng.* **2017**, *3* (10), 2223-2229.
17. Xiang, H.-J.; Tham, H. P.; Nguyen, M. D.; Fiona Phua, S. Z.; **Lim, W. Q.**; Liu, J.-G.; Zhao, Y. *Chem. Commun.* **2017**, *53* (37), 5220-5223.
18. Bai, L.; Phua, S. Z. F.; **Lim, W. Q.**; Jana, A.; Luo, Z.; Tham, H. P.; Zhao, L.; Gao, Q.; Zhao, Y. *Chem. Commun.* **2016**, *52* (22), 4128-4131.

References

1. Thun, M. J.; DeLancey, J. O.; Center, M. M.; Jemal, A.; Ward, E. M. *Carcinogenesis* **2010**, *31* (1), 100-110.
2. DeVita, V. T. J.; Rosenberg, S. A. *N. Engl. J. Med.* **2012**, *366* (23), 2207-2214.
3. Mehlen, P.; Puisieux, A. *Nat. Rev. Cancer* **2006**, *6* (6), 449-458.
4. Khalil, D. N.; Smith, E. L.; Brentjens, R. J.; Wolchok, J. D. *Nat. Clin. Pract. Oncol.* **2016**, *13* (5), 273-290.
5. Vanneman, M.; Dranoff, G. *Nat. Rev. Cancer* **2012**, *12* (4), 237-251.
6. Leader, B.; Baca, Q. J.; Golan, D. E. *Nat. Rev. Drug Discovery* **2008**, *7* (1), 21-39.
7. Pieters, R.; Hunger, S. P.; Boos, J.; Rizzari, C.; Silverman, L.; Baruchel, A.; Goekbuget, N.; Schrappe, M.; Pui, C.-H. *Cancer* **2011**, *117* (2), 238-249.
8. Cheng, L.; Wang, C.; Feng, L.; Yang, K.; Liu, Z. *Chem. Rev.* **2014**, *114* (21), 10869-10939.
9. Dolmans, D. E. J. G. J.; Fukumura, D.; Jain, R. K. *Nat. Rev. Cancer* **2003**, *3* (5), 380-387.
10. Liu, Y.; Bhattarai, P.; Dai, Z.; Chen, X. *Chem. Soc. Rev.* **2019**, *48* (7), 2053-2108.
11. Peng, F.; Setyawati, M. I.; Tee, J. K.; Ding, X.; Wang, J.; Nga, M. E.; Ho, H. K.; Leong, D. T. *Nat. Nanotechnol.* **2019**, *14* (3), 279-286.
12. Easwaran, H.; Tsai, H.-C.; Baylin, Stephen B. *Mol. Cell* **2014**, *54* (5), 716-727.
13. Nam, J.; Son, S.; Park, K. S.; Zou, W.; Shea, L. D.; Moon, J. J. *Nat. Rev. Mater.* **2019**, *4* (6), 398-414.
14. Sun, T.; Zhang, Y. S.; Pang, B.; Hyun, D. C.; Yang, M.; Xia, Y. *Angew. Chem., Int. Ed.* **2014**, *53* (46), 12320-12364.
15. Shi, J.; Votruba, A. R.; Farokhzad, O. C.; Langer, R. *Nano Lett.* **2010**, *10* (9), 3223-3230.
16. Abdalla, A. M. E.; Xiao, L.; Ullah, M. W.; Yu, M.; Ouyang, C.; Yang, G. *Theranostics* **2018**, *8* (2), 533-548.
17. Barreto, J. A.; O'Malley, W.; Kubeil, M.; Graham, B.; Stephan, H.; Spiccia, L. *Adv. Mater.* **2011**, *23* (12), H18-40.
18. Kobayashi, H.; Watanabe, R.; Choyke, P. L. *Theranostics* **2013**, *4* (1), 81-89.

19. Zhang, L.; Chan, J. M.; Gu, F. X.; Rhee, J.-W.; Wang, A. Z.; Radovic-Moreno, A. F.; Alexis, F.; Langer, R.; Farokhzad, O. C. *ACS Nano* **2008**, *2* (8), 1696-1702.
20. Yu, T.; Liu, X.; Bolcato-Bellemin, A.-L.; Wang, Y.; Liu, C.; Erbacher, P.; Qu, F.; Rocchi, P.; Behr, J.-P.; Peng, L. *Angew. Chem., Int. Ed.* **2012**, *51* (34), 8478-8484.
21. Kataoka, K.; Harada, A.; Nagasaki, Y. *Adv. Drug Delivery Rev.* **2001**, *47* (1), 113-131.
22. Li, M.; Teh, C.; Ang, C. Y.; Tan, S. Y.; Luo, Z.; Qu, Q.; Zhang, Y.; Korzh, V.; Zhao, Y. *Adv. Funct. Mater.* **2015**, *25* (35), 5602-5610.
23. Liu, D.; Auguste, D. T. *J. Controlled Release* **2015**, *219*, 632-643.
24. Jiang, Q.; Zhao, S.; Liu, J.; Song, L.; Wang, Z.-G.; Ding, B. *Adv. Drug Delivery Rev.* **2019**.
25. Elzoghby, A. O.; Samy, W. M.; Elgindy, N. A. *J. Controlled Release* **2012**, *161* (1), 38-49.
26. Bhaskar, S.; Lim, S. *NPG Asia Mater.* **2017**, *9*, e371.
27. Yang, B.; Chen, Y.; Shi, J. *Adv. Mater.* **2019**, *31* (2), 1802896.
28. Dai, L.; Zhang, Q.; Shen, X.; Sun, Q.; Mu, C.; Gu, H.; Cai, K. *J. Mater. Chem. B* **2016**, *4* (26), 4594-4604.
29. Ni, D.; Zhang, J.; Wang, J.; Hu, P.; Jin, Y.; Tang, Z.; Yao, Z.; Bu, W.; Shi, J. *ACS Nano* **2017**, *11* (4), 4256-4264.
30. Min, H.; Wang, J.; Qi, Y.; Zhang, Y.; Han, X.; Xu, Y.; Xu, J.; Li, Y.; Chen, L.; Cheng, K.; Liu, G.; Yang, N.; Li, Y.; Nie, G. *Adv. Mater.* **2019**, *31* (15), 1808200.
31. Park, H.; Kim, J.; Jung, S.; Kim, W. J. *Adv. Funct. Mater.* **2018**, *28* (5), 1705416.
32. Nicolas, J.; Mura, S.; Brambilla, D.; Mackiewicz, N.; Couvreur, P. *Chem. Soc. Rev.* **2013**, *42* (3), 1147-235.
33. Danhier, F.; Feron, O.; Preat, V. *J. Controlled Release* **2010**, *148* (2), 135-146.
34. Chen, F.; Hong, H.; Zhang, Y.; Valdovinos, H. F.; Shi, S.; Kwon, G. S.; Theuer, C. P.; Barnhart, T. E.; Cai, W. *ACS Nano* **2013**, *7* (10), 9027-9039.
35. Aravind, A.; Jeyamohan, P.; Nair, R.; Veerananarayanan, S.; Nagaoka, Y.; Yoshida, Y.; Maekawa, T.; Kumar, D. S. *Biotechnol. Bioeng.* **2012**, *109* (11), 2920-2931.
36. Shao, K.; Zhang, Y.; Ding, N.; Huang, S.; Wu, J.; Li, J.; Yang, C.; Leng, Q.; Ye, L.; Lou, J.; Zhu, L.; Jiang, C. *Adv. Healthcare Mater.* **2015**, *4* (2), 291-300.

37. Majumdar, S.; Siahaan, T. J. *Med. Res. Rev.* **2012**, *32* (3), 637-658.
38. Wang, X.; Li, S.; Shi, Y.; Chuan, X.; Li, J.; Zhong, T.; Zhang, H.; Dai, W.; He, B.; Zhang, Q. *J. Controlled Release* **2014**, *193*, 139-153.
39. Yu, M. K.; Park, J.; Jon, S. *Theranostics* **2012**, *2* (1), 3-44.
40. Wong, P. T.; Choi, S. K. *Chem. Rev.* **2015**, *115* (9), 3388-3432.
41. Pacardo, D. B.; Ligler, F. S.; Gu, Z. *Nanoscale* **2015**, *7* (8), 3381-3391.
42. Mura, S.; Nicolas, J.; Couvreur, P. *Nat. Mater.* **2013**, *12*, 991-1003.
43. Fleige, E.; Quadir, M. A.; Haag, R. *Adv. Drug Delivery Rev.* **2012**, *64* (9), 866-884.
44. Warburg, O. *Science* **1956**, *123* (3191), 309-314.
45. Murthy, N.; Campbell, J.; Fausto, N.; Hoffman, A. S.; Stayton, P. S. *J. Controlled Release* **2003**, *89* (3), 365-374.
46. Liu, J.; Huang, Y.; Kumar, A.; Tan, A.; Jin, S.; Mozhi, A.; Liang, X.-J. *Biotechnol. Adv.* **2014**, *32* (4), 693-710.
47. Allocati, N.; Masulli, M.; Di Ilio, C.; Federici, L. *Oncogenesis* **2018**, *7* (1), 8.
48. Trachootham, D.; Alexandre, J.; Huang, P. *Nat. Rev. Drug Discovery* **2009**, *8*, 579.
49. Gorrini, C.; Harris, I. S.; Mak, T. W. *Nat. Rev. Drug Discovery* **2013**, *12*, 931.
50. Saravanakumar, G.; Kim, J.; Kim, W. J. *Adv. Sci.* **2017**, *4* (1), 1600124.
51. Karimi, M.; Sahandi Zangabad, P.; Baghaee-Ravari, S.; Ghazadeh, M.; Mirshekari, H.; Hamblin, M. R. *J. Am. Chem. Soc.* **2017**, *139* (13), 4584-4610.
52. Zhang, Y.; Yu, J.; Bomba, H. N.; Zhu, Y.; Gu, Z. *Chem. Rev.* **2016**, *116* (19), 12536-12563.
53. Fan, W.; Yung, B.; Huang, P.; Chen, X. *Chem. Rev.* **2017**, *117* (22), 13566-13638.
54. Yang, D.; Yang, G.; Yang, P.; Lv, R.; Gai, S.; Li, C.; He, F.; Lin, J. *Adv. Funct. Mater.* **2017**, *27* (18), 1700371.
55. Webber, M. J.; Appel, E. A.; Meijer, E. W.; Langer, R. *Nat. Mater.* **2015**, *15*, 13-26.
56. Zhu, H.; Shangguan, L.; Shi, B.; Yu, G.; Huang, F. *Mater. Chem. Front.* **2018**, *2* (12), 2152-2174.
57. Xing, P.; Zhao, Y. *Adv. Mater.* **2016**, *28* (34), 7304-7339.
58. Qi, G.-B.; Gao, Y.-J.; Wang, L.; Wang, H. *Adv. Mater.* **2018**, *30* (22), 1703444.
59. Li, Y.; Wang, Y.; Huang, G.; Gao, J. *Chem. Rev.* **2018**, *118* (11), 5359-5391.

60. Cook, T. R.; Zheng, Y.-R.; Stang, P. J. *Chem. Rev.* **2013**, *113* (1), 734-777.
61. Luo, Q.; Hou, C.; Bai, Y.; Wang, R.; Liu, J. *Chem. Rev.* **2016**, *116* (22), 13571-13632.
62. Yu, G.; Jie, K.; Huang, F. *Chem. Rev.* **2015**, *115* (15), 7240-7303.
63. Thota, B. N. S.; Urner, L. H.; Haag, R. *Chem. Rev.* **2016**, *116* (4), 2079-2102.
64. Wang, M.; Alberti, K.; Sun, S.; Arellano, C. L.; Xu, Q. *Angew. Chem., Int. Ed.* **2014**, *53* (11), 2893-2898.
65. Davis, M. E.; Brewster, M. E. *Nat. Rev. Drug Discovery* **2004**, *3* (12), 1023-1035.
66. Cao, L.; Hettiarachchi, G.; Briken, V.; Isaacs, L. *Angew. Chem., Int. Ed.* **2013**, *52* (46), 12033-12037.
67. Zhou, M.; Zhang, X.; Yang, Y.; Liu, Z.; Tian, B.; Jie, J.; Zhang, X. *Biomaterials* **2013**, *34* (35), 8960-8967.
68. Dan, Z.; Cao, H.; He, X.; Zhang, Z.; Zou, L.; Zeng, L.; Xu, Y.; Yin, Q.; Xu, M.; Zhong, D.; Yu, H.; Shen, Q.; Zhang, P.; Li, Y. *Theranostics* **2016**, *6* (3), 435-445.
69. Zhang, J.; Liang, Y.-C.; Lin, X.; Zhu, X.; Yan, L.; Li, S.; Yang, X.; Zhu, G.; Rogach, A. L.; Yu, P. K. N.; Shi, P.; Tu, L.-C.; Chang, C.-C.; Zhang, X.; Chen, X.; Zhang, W.; Lee, C.-S. *ACS Nano* **2015**, *9* (10), 9741-9756.
70. Liu, X.; Zhou, J.; Yu, T.; Chen, C.; Cheng, Q.; Sengupta, K.; Huang, Y.; Li, H.; Liu, C.; Wang, Y.; Posocco, P.; Wang, M.; Cui, Q.; Giorgio, S.; Fermeglia, M.; Qu, F.; Priol, S.; Shi, Y.; Liang, Z.; Rocchi, P.; Rossi, J. J.; Peng, L. *Angew. Chem. Int. Ed.* **2014**, *53* (44), 11822-11827.
71. Rautio, J.; Kumpulainen, H.; Heimbach, T.; Oliyai, R.; Oh, D.; Järvinen, T.; Savolainen, J. *Nat. Rev. Drug Discovery* **2008**, *7*, 255.
72. Kuang, Y.; Balakrishnan, K.; Gandhi, V.; Peng, X. *J. Am. Chem. Soc.* **2011**, *133* (48), 19278-19281.
73. Thapa, P.; Li, M.; Bio, M.; Rajaputra, P.; Nkepang, G.; Sun, Y.; Woo, S.; You, Y. *J. Med. Chem.* **2016**, *59* (7), 3204-3214.
74. Legigan, T.; Clarhaut, J.; Tranoy-Opalinski, I.; Monvoisin, A.; Renoux, B.; Thomas, M.; Le Pape, A.; Lerondel, S.; Papot, S. *Angew. Chem. Int. Ed.* **2012**, *51* (46), 11606-11610.
75. Liu, P.; Xu, J.; Yan, D.; Zhang, P.; Zeng, F.; Li, B.; Wu, S. *Chem. Commun.* **2015**, *51* (46), 9567-9570.

76. Cheetham, A. G.; Chakraborty, R. W.; Ma, W.; Cui, H. *Chem. Soc. Rev.* **2017**, *46* (21), 6638-6663.
77. Li, T.; Smet, M.; Dehaen, W.; Xu, H. *ACS Appl. Mater. Interfaces* **2016**, *8* (6), 3609-3614.
78. Xue, P.; Liu, D.; Wang, J.; Zhang, N.; Zhou, J.; Li, L.; Guo, W.; Sun, M.; Han, X.; Wang, Y. *Bioconjugate Chem.* **2016**, *27* (5), 1360-1372.
79. Liu, J.; Pang, Y.; Zhu, Z.; Wang, D.; Li, C.; Huang, W.; Zhu, X.; Yan, D. *Biomacromolecules* **2013**, *14* (5), 1627-1636.
80. Liu, J.; Pang, Y.; Chen, J.; Huang, P.; Huang, W.; Zhu, X.; Yan, D. *Biomaterials* **2012**, *33* (31), 7765-7774.
81. Cao, W.; Li, Y.; Yi, Y.; Ji, S.; Zeng, L.; Sun, Z.; Xu, H. *Chem. Sci.* **2012**, *3* (12), 3403-3408.
82. Chung, J. E.; Tan, S.; Gao, S. J.; Yongvongsoontorn, N.; Kim, S. H.; Lee, J. H.; Choi, H. S.; Yano, H.; Zhuo, L.; Kurisawa, M.; Ying, J. Y. *Nat. Nanotechnol.* **2014**, *9* (11), 907-912.
83. Shen, Y.; Jin, E.; Zhang, B.; Murphy, C. J.; Sui, M.; Zhao, J.; Wang, J.; Tang, J.; Fan, M.; Van Kirk, E.; Murdoch, W. J. *J. Am. Chem. Soc.* **2010**, *132* (12), 4259-4265.
84. Chen, W.-H.; Lei, Q.; Yang, C.-X.; Jia, H.-Z.; Luo, G.-F.; Wang, X.-Y.; Liu, G.; Cheng, S.-X.; Zhang, X.-Z. *Small* **2015**, *11* (39), 5230-5242.
85. Cheetham, A. G.; Zhang, P.; Lin, Y.-a.; Lock, L. L.; Cui, H. *J. Am. Chem. Soc.* **2013**, *135* (8), 2907-2910.
86. Peng, M.; Qin, S.; Jia, H.; Zheng, D.; Rong, L.; Zhang, X. *Nano Res.* **2016**, *9* (3), 663-673.
87. Huang, P.; Wang, D.; Su, Y.; Huang, W.; Zhou, Y.; Cui, D.; Zhu, X.; Yan, D. *J. Am. Chem. Soc.* **2014**, *136* (33), 11748-11756.
88. Xue, X.; Huang, Y.; Bo, R.; Jia, B.; Wu, H.; Yuan, Y.; Wang, Z.; Ma, Z.; Jing, D.; Xu, X.; Yu, W.; Lin, T.-y.; Li, Y. *Nat. Commun.* **2018**, *9* (1), 3653.
89. Wang, Y.; Huang, P.; Hu, M.; Huang, W.; Zhu, X.; Yan, D. *Bioconjugate Chem.* **2016**, *27* (11), 2722-2733.
90. Ji, C.; Gao, Q.; Dong, X.; Yin, W.; Gu, Z.; Gan, Z.; Zhao, Y.; Yin, M. *Angew. Chem., Int. Ed.* **2018**, *57* (35), 11384-11388.
91. Hall, M. D.; Mellor, H. R.; Callaghan, R.; Hambley, T. W. *J. Med. Chem.* **2007**, *50* (15), 3403-3411.

92. Alderden, R. A.; Hall, M. D.; Hambley, T. W. *J. Chem. Educ.* **2006**, 83 (5), 728.
93. Einhorn, L. H. *J. Clin. Oncol.* **1990**, 8 (11), 1777-1781.
94. Eastman, A., The Mechanism of Action of Cisplatin: From Adducts to Apoptosis. In *Cisplatin*, Verlag Helvetica Chimica Acta: 2006; pp 111-134.
95. Jamieson, E. R.; Lippard, S. J. *Chem. Rev.* **1999**, 99 (9), 2467-2498.
96. Legendre, F.; Chottard, J.-C., Kinetics and Selectivity of DNA-Platination. In *Cisplatin*, Verlag Helvetica Chimica Acta: 2006; pp 223-245.
97. Kelland, L. *Nat. Rev. Cancer* **2007**, 7 (8), 573-584.
98. McWhinney, S. R.; Goldberg, R. M.; McLeod, H. L. *Mol. Cancer Ther.* **2009**, 8 (1), 10-16.
99. Hydes, P. C.; Russell, M. J. H. *Cancer Metastasis Rev.* **1988**, 7 (1), 67-89.
100. Wang, X.; Guo, Z. *Dalton Trans.* **2008**, (12), 1521-1532.
101. Hall, M. D.; Hambley, T. W. *Coord. Chem. Rev.* **2002**, 232 (1-2), 49-67.
102. Ho, Y.-P.; Au-Yeung, S. C. F.; To, K. K. W. *Med. Res. Rev.* **2003**, 23 (5), 633-655.
103. Lovejoy, K. S.; Todd, R. C.; Zhang, S.; McCormick, M. S.; D'Aquino, J. A.; Reardon, J. T.; Sancar, A.; Giacomini, K. M.; Lippard, S. J. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, 105 (26), 8902-8907.
104. van der Vijgh, W. J. F. *Clin. Pharmacokinet.* **1991**, 21 (4), 242-261.
105. Rixe, O.; Ortuzar, W.; Alvarez, M.; Parker, R.; Reed, E.; Paull, K.; Fojo, T. *Biochem. Pharmacol.* **1996**, 52 (12), 1855-1865.
106. Johnstone, T. C.; Suntharalingam, K.; Lippard, S. J. *Chem. Rev.* **2016**, 116 (5), 3436-3486.
107. Tseng, C.-L.; Su, W.-Y.; Yen, K.-C.; Yang, K.-C.; Lin, F.-H. *Biomaterials* **2009**, 30 (20), 3476-3485.
108. Avgoustakis, K.; Beletsi, A.; Panagi, Z.; Klepetsanis, P.; Karydas, A. G.; Ithakissios, D. S. *J. Controlled Release* **2002**, 79 (1), 123-135.
109. Guo, S.; Lin, C. M.; Xu, Z.; Miao, L.; Wang, Y.; Huang, L. *ACS Nano* **2014**, 8 (5), 4996-5009.
110. Khiati, S.; Luvino, D.; Oumzil, K.; Chauffert, B.; Camplo, M.; Barthélémy, P. *ACS Nano* **2011**, 5 (11), 8649-8655.
111. Chen, H.; He, W.; Guo, Z. *Chem Commun (Camb)* **2014**, 50 (68), 9714-9717.
112. Miao, L.; Guo, S.; Zhang, J.; Kim, W. Y.; Huang, L. *Adv. Funct. Mater.* **2014**, 24 (42), 6601-6611.

113. Wilson, J. J.; Lippard, S. J. *Chem. Rev.* **2014**, *114* (8), 4470-4495.
114. Lemma, K.; Sargeson, A. M.; Elding, L. I. *J. Chem. Soc., Dalton Trans.* **2000**, (7), 1167-1172.
115. Graf, N.; Lippard, S. J. *Adv. Drug Delivery Rev.* **2012**, *64* (11), 993-1004.
116. Bhargava, A.; Vaishampayan, U. N. *Expert Opin. Invest. Drugs* **2009**, *18* (11), 1787-1797.
117. Huang, X.; Huang, R.; Gou, S.; Wang, Z.; Liao, Z.; Wang, H. *Bioconjugate Chem.* **2016**, *27* (9), 2132-2148.
118. Pathak, R. K.; Marrache, S.; Choi, J. H.; Berding, T. B.; Dhar, S. *Angew. Chem., Int. Ed.* **2014**, *53* (7), 1963-1967.
119. Ang, W. H.; Khalaila, I.; Allardyce, C. S.; Juillerat-Jeanneret, L.; Dyson, P. J. *J. Am. Chem. Soc.* **2005**, *127* (5), 1382-1383.
120. Dai, Y.; Xiao, H.; Liu, J.; Yuan, Q.; Ma, P.; Yang, D.; Li, C.; Cheng, Z.; Hou, Z.; Yang, P.; Lin, J. *J. Am. Chem. Soc.* **2013**, *135* (50), 18920-18929.
121. Min, Y.; Li, J.; Liu, F.; Yeow, E. K.; Xing, B. *Angew. Chem., Int. Ed.* **2014**, *53* (4), 1012-1016.
122. Mackay, F. S.; Woods, J. A.; Heringova, P.; Kasparkova, J.; Pizarro, A. M.; Moggach, S. A.; Parsons, S.; Brabec, V.; Sadler, P. J. *Proc Natl Acad Sci U S A* **2007**, *104* (52), 20743-20748.
123. Al-Baker, S.; Dabrowiak, J. C. *Inorg. Chem.* **1987**, *26* (4), 613-617.
124. Giandomenico, C. M.; Abrams, M. J.; Murrer, B. A.; Vollano, J. F.; Rheinheimer, M. I.; Wyer, S. B.; Bossard, G. E.; Higgins, J. D. *Inorg. Chem.* **1995**, *34* (5), 1015-1021.
125. Galanski, M.; Keppler, B. K. *Inorg. Chem.* **1996**, *35* (6), 1709-1711.
126. Zhang, J. Z.; Bonnitche, P.; Wexselblatt, E.; Klein, A. V.; Najajreh, Y.; Gibson, D.; Hambley, T. W. *Chem. Eur. J.* **2013**, *19* (5), 1672-1676.
127. Ang, W. H.; Pilet, S.; Scopelliti, R.; Bussy, F.; Juillerat-Jeanneret, L.; Dyson, P. J. *J. Med. Chem.* **2005**, *48* (25), 8060-8069.
128. Perez, J. M.; Camazón, M.; Alvarez-Valdes, A.; Quiroga, A. G.; Kelland, L. R.; Alonso, C.; Navarro-Ranninger, M. C. *Chem.-Biol. Interact.* **1999**, *117* (2), 99-115.
129. Reithofer, M.; Galanski, M.; Roller, A.; Keppler, B. K. *Eur. J. Inorg. Chem.* **2006**, *2006* (13), 2612-2617.
130. Barnes, K. R.; Kutikov, A.; Lippard, S. J. *Chem. Biol.* **11** (4), 557-564.

131. Gaviglio, L.; Gross, A.; Metzler-Nolte, N.; Ravera, M. *Metallomics* **2012**, *4* (3), 260-266.
132. Abramkin, S.; Valiahdi, S. M.; Jakupec, M. A.; Galanski, M.; Metzler-Nolte, N.; Keppler, B. K. *Dalton Trans.* **2012**, *41* (10), 3001-3005.
133. Mukhopadhyay, S.; Barnés, C. M.; Haskell, A.; Short, S. M.; Barnes, K. R.; Lippard, S. J. *Bioconjugate Chem.* **2008**, *19* (1), 39-49.
134. Reithofer, M. R.; Valiahdi, S. M.; Jakupec, M. A.; Arion, V. B.; Egger, A.; Galanski, M.; Keppler, B. K. *J. Med. Chem.* **2007**, *50* (26), 6692-6699.
135. Varbanov, H. P.; Göschl, S.; Heffeter, P.; Theiner, S.; Roller, A.; Jensen, F.; Jakupec, M. A.; Berger, W.; Galanski, M.; Keppler, B. K. *J. Med. Chem.* **2014**, *57* (15), 6751-6764.
136. Theiner, S.; Varbanov, H. P.; Galanski, M.; Egger, A. E.; Berger, W.; Heffeter, P.; Keppler, B. K. *J. Biol. Inorg. Chem.* **2015**, *20* (1), 89-99.
137. Pichler, V.; Mayr, J.; Heffeter, P.; Domotor, O.; Enyedy, E. A.; Hermann, G.; Groza, D.; Kollensperger, G.; Galanksi, M.; Berger, W.; Keppler, B. K.; Kowol, C. R. *Chem Commun (Camb)* **2013**, *49* (22), 2249-51.
138. Chin, C. F.; Tian, Q.; Setyawati, M. I.; Fang, W.; Tan, E. S. Q.; Leong, D. T.; Ang, W. H. *J. Med. Chem.* **2012**, *55* (17), 7571-7582.
139. Zheng, Y. R.; Suntharalingam, K.; Johnstone, T. C.; Lippard, S. J. *Chem. Sci.* **2015**, *6* (2), 1189-1193.
140. Li, S.; Li, C.; Jin, S.; Liu, J.; Xue, X.; Eltahan, A. S.; Sun, J.; Tan, J.; Dong, J.; Liang, X.-J. *Biomaterials* **2017**, *144*, 119-129.
141. Johnstone, T. C.; Kulak, N.; Pridgen, E. M.; Farokhzad, O. C.; Langer, R.; Lippard, S. J. *ACS Nano* **2013**, *7* (7), 5675-5683.
142. Liu, D.; Poon, C.; Lu, K.; He, C.; Lin, W. *Nat. Commun.* **2014**, *5*, 4182.
143. Ling, X.; Tu, J.; Wang, J.; Shajii, A.; Kong, N.; Feng, C.; Zhang, Y.; Yu, M.; Xie, T.; Bharwani, Z.; Aljaeid, B. M.; Shi, B.; Tao, W.; Farokhzad, O. C. *ACS Nano* **2019**, *13* (1), 357-370.
144. Liu, H.; Li, Y.; Lyu, Z.; Wan, Y.; Li, X.; Chen, H.; Chen, H.; Li, X. *J. Mater. Chem. B* **2014**, *2* (47), 8303-8309.
145. Pathak, R. K.; Dhar, S. *J. Am. Chem. Soc.* **2015**, *137* (26), 8324-8327.
146. Wan, X.; Min, Y.; Bludau, H.; Keith, A.; Sheiko, S. S.; Jordan, R.; Wang, A. Z.; Sokolsky-Papkov, M.; Kabanov, A. V. *ACS Nano* **2018**, *12* (3), 2426-2439.

147. Xu, X.; Xie, K.; Zhang, X.-Q.; Pridgen, E. M.; Park, G. Y.; Cui, D. S.; Shi, J.; Wu, J.; Kantoff, P. W.; Lippard, S. J.; Langer, R.; Walker, G. C.; Farokhzad, O. C. *Proc. Natl. Acad. Sci.* **2013**, *110* (46), 18638-18643.
148. He, C.; Liu, D.; Lin, W. *Biomaterials* **2015**, *36*, 124-133.
149. Zhang, W.; Shen, J.; Su, H.; Mu, G.; Sun, J.-H.; Tan, C.-P.; Liang, X.-J.; Ji, L.-N.; Mao, Z.-W. *ACS Appl. Mater. Interfaces* **2016**, *8* (21), 13332-13340.
150. He, C.; Lu, K.; Liu, D.; Lin, W. *J. Am. Chem. Soc.* **2014**, *136* (14), 5181-5184.
151. Wang, Z.; Ma, R.; Yan, L.; Chen, X.; Zhu, G. *Chem. Commun.* **2015**, *51* (58), 11587-11590.
152. Dai, Y.; Cheng, S.; Wang, Z.; Zhang, R.; Yang, Z.; Wang, J.; Yung, B. C.; Wang, Z.; Jacobson, O.; Xu, C.; Ni, Q.; Yu, G.; Zhou, Z.; Chen, X. *ACS Nano* **2018**, *12* (1), 455-463.
153. Lin, Y.-X.; Wang, Y.; An, H.-W.; Qi, B.; Wang, J.; Wang, L.; Shi, J.; Mei, L.; Wang, H. *Nano Lett.* **2019**, *19* (5), 2968-2978.
154. Yang, Z.; Dai, Y.; Yin, C.; Fan, Q.; Zhang, W.; Song, J.; Yu, G.; Tang, W.; Fan, W.; Yung, B. C.; Li, J.; Li, X.; Li, X.; Tang, Y.; Huang, W.; Song, J.; Chen, X. *Adv. Mater.* **2018**, *30* (23), 1707509.
155. Kolishetti, N.; Dhar, S.; Valencia, P. M.; Lin, L. Q.; Karnik, R.; Lippard, S. J.; Langer, R.; Farokhzad, O. C. *Proc. Natl. Acad. Sci.* **2010**, *107* (42), 17939-17944.
156. Liao, L.; Liu, J.; Dreaden, E. C.; Morton, S. W.; Shopsowitz, K. E.; Hammond, P. T.; Johnson, J. A. *J. Am. Chem. Soc.* **2014**, *136* (16), 5896-5899.
157. Cong, Y.; Xiao, H.; Xiong, H.; Wang, Z.; Ding, J.; Li, C.; Chen, X.; Liang, X.-J.; Zhou, D.; Huang, Y. *Adv. Mater.* **2018**, *30* (11), 1706220.
158. Yue, Z.; Wang, H.; Li, Y.; Qin, Y.; Xu, L.; Bowers, D. J.; Gangoda, M.; Li, X.; Yang, H. B.; Zheng, Y. R. *Chem Commun (Camb)* **2018**, *54* (7), 731-734.
159. Kweekel, D. M.; Gelderblom, H.; Guchelaar, H. J. *Cancer Treat. Rev.* **2005**, *31* (2), 90-105.
160. Alcindor, T.; Beauger, N. *Curr. Oncol.* **2011**, *18* (1), 18-25.
161. Della Rocca, J.; Huxford, R. C.; Comstock-Duggan, E.; Lin, W. *Angew Chem Int Ed* **2011**, *50* (44), 10330-10334.
162. Xu, J.; Kuang, Y.; Lv, R.; Yang, P.; Li, C.; Bi, H.; Liu, B.; Yang, D.; Dai, Y.; Gai, S.; He, F.; Xing, B.; Lin, J. *Biomaterials* **2017**, *130*, 42-55.
163. Ma, P. a.; Xiao, H.; Li, C.; Dai, Y.; Cheng, Z.; Hou, Z.; Lin, J. *Mater Today Chem.* **2015**, *18* (10), 554-564.

164. Guo, Z.; Zou, Y.; He, H.; Rao, J.; Ji, S.; Cui, X.; Ke, H.; Deng, Y.; Yang, H.; Chen, C.; Zhao, Y.; Chen, H. *Adv. Mater.* **2016**, *28* (46), 10155-10164.
165. Yuan, Y.; Zhang, C.-J.; Liu, B. *Chem. Commun.* **2015**, *51* (41), 8626-8629.
166. He, S.; Li, C.; Zhang, Q.; Ding, J.; Liang, X.-J.; Chen, X.; Xiao, H.; Chen, X.; Zhou, D.; Huang, Y. *ACS Nano* **2018**, *12* (7), 7272-7281.
167. Wang, C.; Wang, Z.; Zhang, X. *Acc. Chem. Res* **2012**, *45* (4), 608-618.
168. Zhou, J.; Yu, G.; Huang, F. *Chem. Soc. Rev.* **2017**, *46* (22), 7021-7053.
169. Yu, G.; Zhao, X.; Zhou, J.; Mao, Z.; Huang, X.; Wang, Z.; Hua, B.; Liu, Y.; Zhang, F.; He, Z.; Jacobson, O.; Gao, C.; Wang, W.; Yu, C.; Zhu, X.; Huang, F.; Chen, X. *J. Am. Chem. Soc.* **2018**, *140* (25), 8005-8019.
170. Yu, G.; Yang, Z.; Fu, X.; Yung, B. C.; Yang, J.; Mao, Z.; Shao, L.; Hua, B.; Liu, Y.; Zhang, F.; Fan, Q.; Wang, S.; Jacobson, O.; Jin, A.; Gao, C.; Tang, X.; Huang, F.; Chen, X. *Nat. Commun.* **2018**, *9* (1), 766.
171. Zhou, J.; Zhang, Y.; Yu, G.; Crawley, M. R.; Fulong, C. R. P.; Friedman, A. E.; Sengupta, S.; Sun, J.; Li, Q.; Huang, F.; Cook, T. R. *J. Am. Chem. Soc.* **2018**, *140* (24), 7730-7736.
172. Danhier, F.; Feron, O.; Préat, V. *J. Controlled Release* **2010**, *148* (2), 135-146.
173. Greish, K., Enhanced Permeability and Retention (EPR) Effect for Anticancer Nanomedicine Drug Targeting. In *Cancer Nanotechnology: Methods and Protocols*, Grobmyer, S. R.; Moudgil, B. M., Eds. Humana Press: Totowa, NJ, 2010; pp 25-37.
174. Lemma, K.; Sargeson, A. M.; Elding, L. I. *J. Chem. Soc., Dalton Trans.* **2000**, (7), 1167-1172.
175. Williams Iii, R. O.; Mahaguna, V.; Sriwongjanya, M. *Eur. J. Pharm. Biopharm* **1998**, *46* (3), 355-360.
176. Pegg, A. E. *J. Biol. Chem.* **2016**, *291* (29), 14904-14912.
177. Murray-Stewart, T. R.; Woster, P. M.; Casero, R. A. *Biochem J.* **2016**, *473* (19), 2937-2953.
178. Casero, R. A., Jr.; Marton, L. J. *Nat. Rev. Drug Discovery* **2007**, *6* (5), 373-390.
179. Palmer, A. J.; Wallace, H. M. *Amino Acids* **2010**, *38* (2), 415-422.
180. Sengupta, P.; Basu, S.; Soni, S.; Pandey, A.; Roy, B.; Oh, M. S.; Chin, K. T.; Paraskar, A. S.; Sarangi, S.; Connor, Y.; Sabbisetti, V. S.; Koppam, J.; Kulkarni, A.; Muto, K.; Amarasiriwardena, C.; Jayawardene, I.; Lupoli, N.;

- Dinulescu, D. M.; Bonventre, J. V.; Mashelkar, R. A.; Sengupta, S. *Proc. Natl. Acad. Sci.* **2012**, *109* (28), 11294-11299.
181. Matsui, Y.; Yokoi, T.; Mochida, K. *Chem. Lett.* **1976**, *5* (10), 1037-1040.
 182. L. May, B.; D. Kean, S.; J. Easton, C.; F. Lincoln, S. *J. Chem. Soc., Perkin Trans. I* **1997**, (21), 3157-3160.
 183. Woynarowski, J. M.; Chapman, W. G.; Napier, C.; Herzig, M. C. S.; Juniewicz, P. *Mol. Pharmacol.* **1998**, *54* (5), 770-777.
 184. Beevi, S. S. S.; Rasheed, A. M. H.; Geetha, A. *Jpn. J. Clin. Oncol.* **2004**, *34* (7), 379-385.
 185. Fiaschi, A. I.; Cozzolino, A.; Ruggiero, G.; A Giorgi, G. *Eur. Rev. Med. Pharmacol. Sci.* **2005**, *9* (6), 361-367.
 186. Savjani, K. T.; Gajjar, A. K.; Savjani, J. K. *ISRN Pharm.* **2012**, *2012*, 195727-195727.
 187. Narvekar, M.; Xue, H. Y.; Eoh, J. Y.; Wong, H. L. *AAPS PharmSciTech* **2014**, *15* (4), 822-833.
 188. Liang, X.-J.; Chen, C.; Zhao, Y.; Wang, P. C. *Methods Mol. Biol.* **2010**, *596*, 467-488.
 189. Luo, C.; Sun, J.; Sun, B.; He, Z. *Trends Pharmacol. Sci.* **2014**, *35* (11), 556-566.
 190. Mishra, A. P.; Chandra, S.; Tiwari, R.; Srivastava, A.; Tiwari, G. *Open Med Chem J* **2018**, *12*, 111-123.
 191. Li, R.; Xie, Y. *J. Controlled Release* **2017**, *251*, 49-67.
 192. Qin, S.-Y.; Zhang, A.-Q.; Cheng, S.-X.; Rong, L.; Zhang, X.-Z. *Biomaterials* **2017**, *112*, 234-247.
 193. Ge, Y.; Ma, Y.; Li, L. *Colloids Surf., B* **2016**, *146*, 482-489.
 194. Taresco, V.; Alexander, C.; Singh, N.; Pearce, A. K. *Adv. Ther.* **2018**, *1* (4), 1800030.
 195. Housman, G.; Byler, S.; Heerboth, S.; Lapinska, K.; Longacre, M.; Snyder, N.; Sarkar, S. *Cancers* **2014**, *6* (3), 1769-1792.
 196. Alfarouk, K. O.; Stock, C.-M.; Taylor, S.; Walsh, M.; Muddathir, A. K.; Verduzco, D.; Bashir, A. H. H.; Mohammed, O. Y.; Elhassan, G. O.; Harguindey, S.; Reshkin, S. J.; Ibrahim, M. E.; Rauch, C. *Cancer Cell Int.* **2015**, *15*, 71.
 197. Miles, D.; Von Minckwitz, G.; Seidman, A. D. *The Oncologist* **2002**, *7* (suppl 6), 13-19.
 198. He, C.; Lu, J.; Lin, W. *J. Controlled Release* **2015**, *219*, 224-236.

199. Mignani, S.; Bryszewska, M.; Klajnert-Maculewicz, B.; Zablocka, M.; Majoral, J.-P. *Biomacromolecules* **2015**, *16* (1), 1-27.
200. Luo, D.; Carter, K. A.; Miranda, D.; Lovell, J. F. *Adv. Sci.* **2017**, *4* (1), 1600106.
201. Kimura, M.; Miyajima, K.; Kojika, M.; Kono, T.; Kato, H. *Int. J. Mol. Sci.* **2015**, *16* (10), 25466-25475.
202. Davids, L. M.; Kleemann, B. *Cancer Treat. Rev.* **2011**, *37* (6), 465-475.
203. Macdonald, I. J.; Dougherty, T. J. *J. Porphyrins Phthalocyanines* **2001**, *5* (2), 105-129.
204. Bacellar, I. O. L.; Tsubone, T. M.; Pavani, C.; Baptista, M. S. *Int. J. Mol. Sci.* **2015**, *16* (9), 20523-20559.
205. Agostinis, P.; Berg, K.; Cengel, K. A.; Foster, T. H.; Girotti, A. W.; Gollnick, S. O.; Hahn, S. M.; Hamblin, M. R.; Juzeniene, A.; Kessel, D.; Korbelik, M.; Moan, J.; Mroz, P.; Nowis, D.; Piette, J.; Wilson, B. C.; Golab, J. *CA: Cancer J. Clin.* **2011**, *61* (4), 250-281.
206. Khdair, A.; Handa, H.; Mao, G.; Panyam, J. *Eur. J. Pharm. Biopharm.* **2009**, *71* (2), 214-222.
207. Pei, Q.; Hu, X.; Zheng, X.; Liu, S.; Li, Y.; Jing, X.; Xie, Z. *ACS Nano* **2018**, *12* (2), 1630-1641.
208. Zhou, F.; Feng, B.; Wang, T.; Wang, D.; Cui, Z.; Wang, S.; Ding, C.; Zhang, Z.; Liu, J.; Yu, H.; Li, Y. *Adv. Funct. Mater.* **2017**, *27* (46), 1703674.
209. Zhao, N.; Wu, B.; Hu, X.; Xing, D. *Biomaterials* **2017**, *141*, 40-49.
210. Xu, X.; Saw, P. E.; Tao, W.; Li, Y.; Ji, X.; Bhasin, S.; Liu, Y.; Ayyash, D.; Rasmussen, J.; Huo, M.; Shi, J.; Farokhzad, O. C. *Adv. Mater.* **2017**, *29* (33), 1700141.
211. Naik, A.; Rubbiani, R.; Gasser, G.; Spingler, B. *Angew. Chem., Int. Ed.* **2014**, *53* (27), 6938-6941.
212. Zhao, Y.; Chen, F.; Pan, Y.; Li, Z.; Xue, X.; Okeke, C. I.; Wang, Y.; Li, C.; Peng, L.; Wang, P. C.; Ma, X.; Liang, X.-J. *ACS Appl. Mater. Interfaces* **2015**, *7* (34), 19295-19305.
213. Conte, C.; Ungaro, F.; Maglio, G.; Tirino, P.; Siracusano, G.; Sciortino, M. T.; Leone, N.; Palma, G.; Barbieri, A.; Arra, C.; Mazzaglia, A.; Quaglia, F. *J. Controlled Release* **2013**, *167* (1), 40-52.

214. You, D. G.; Deepagan, V. G.; Um, W.; Jeon, S.; Son, S.; Chang, H.; Yoon, H. I.; Cho, Y. W.; Swierczewska, M.; Lee, S.; Pomper, M. G.; Kwon, I. C.; Kim, K.; Park, J. H. *Sci. Rep.* **2016**, *6*, 23200.
215. Hou, Z.; Zhang, Y.; Deng, K.; Chen, Y.; Li, X.; Deng, X.; Cheng, Z.; Lian, H.; Li, C.; Lin, J. *ACS Nano* **2015**, *9* (3), 2584-2599.
216. Li, Q.; Wang, X.; Lu, X.; Tian, H.; Jiang, H.; Lv, G.; Guo, D.; Wu, C.; Chen, B. *Biomaterials* **2009**, *30* (27), 4708-4715.
217. Zhao, Q.-g.; Wang, J.; Zhang, Y.-p.; Zhang, J.; Tang, A.-n.; Kong, D.-m. *J. Mater. Chem. B* **2018**, *6* (47), 7898-7907.
218. Jin, J.; Zhu, Y.; Zhang, Z.; Zhang, W. *Angew. Chem.* **2018**, *130* (50), 16592-16596.
219. Wu, J.; Tian, J.; Rui, L.; Zhang, W. *Chem. Commun.* **2018**, *54* (55), 7629-7632.
220. Granadero, D.; Bordello, J.; Pérez-Alvite, M. J.; Novo, M.; Al-Soufi, W. *Int. J. Mol. Sci.* **2010**, *11* (1), 173.
221. Carrazana, J.; Jover, A.; Meijide, F.; Soto, V. H.; Vázquez Tato, J. *J. Phys. Chem. B* **2005**, *109* (19), 9719-9726.
222. Yu, B.; Wei, H.; He, Q.; Ferreira, C. A.; Kuttyreff, C. J.; Ni, D.; Rosenkrans, Z. T.; Cheng, L.; Yu, F.; Engle, J. W.; Lan, X.; Cai, W. *Angew. Chem., Int. Ed.* **2018**, *57* (1), 218-222.
223. Chung, C. Y.-S.; Fung, S.-K.; Tong, K.-C.; Wan, P.-K.; Lok, C.-N.; Huang, Y.; Chen, T.; Che, C.-M. *Chem. Sci.* **2017**, *8* (3), 1942-1953.
224. Cheng, L.; Jiang, D.; Kamkaew, A.; Valdovinos, H. F.; Im, H.-J.; Feng, L.; England, C. G.; Goel, S.; Barnhart, T. E.; Liu, Z.; Cai, W. *Adv. Funct. Mater.* **2017**, *27* (34), 1702928.
225. Rui, L.; Xue, Y.; Wang, Y.; Gao, Y.; Zhang, W. *Chem. Commun.* **2017**, *53* (21), 3126-3129.
226. Pompella, A.; Visvikis, A.; Paolicchi, A.; Tata, V. D.; Casini, A. F. *Biochem. Pharmacol.* **2003**, *66* (8), 1499-1503.
227. Saito, G.; Swanson, J. A.; Lee, K.-D. *Adv. Drug Delivery Rev.* **2003**, *55* (2), 199-215.
228. Huo, M.; Yuan, J.; Tao, L.; Wei, Y. *Polym. Chem.* **2014**, *5* (5), 1519-1528.
229. Wang, M.; Sun, S.; Neufeld, C. I.; Perez-Ramirez, B.; Xu, Q. *Angew. Chem., Int. Ed.* **2014**, *53* (49), 13444-13448.
230. Chou, T.-C. *Cancer Res.* **2010**, *70* (2), 440-446.

231. Giovannetti, R., *The Use of Spectrophotometry UV-Vis for the Study of Porphyrins*. INTECH Open Access Publisher: 2012.
232. Lin, S.; Lei, K.; Du, W.; Yang, L.; Shi, H.; Gao, Y.; Yin, P.; Liang, X.; Liu, J. *Int. J. Biochem. Cell Biol.* **2016**, *71*, 24-34.
233. Leader, B.; Baca, Q. J.; Golan, D. E. *Nat. Rev. Drug Discovery* **2008**, *7*, 21.
234. Lagassé, H. A. D.; Alexaki, A.; Simhadri, V. L.; Katagiri, N. H.; Jankowski, W.; Sauna, Z. E.; Kimchi-Sarfaty, C. *F1000Research* **2017**, *6*, 113.
235. Dimitrov, D. S., Therapeutic Proteins. In *Therapeutic Proteins: Methods and Protocols*, Voynov, V.; Caravella, J. A., Eds. Humana Press: Totowa, NJ, 2012; pp 1-26.
236. Serna, N.; Sánchez-García, L.; Unzueta, U.; Díaz, R.; Vázquez, E.; Manges, R.; Villaverde, A. *Trends Biotechnol.* **2018**, *36* (3), 318-335.
237. Harada, H.; Kizaka-Kondoh, S.; Hiraoka, M. *Breast Cancer* **2006**, *13* (1), 16-26.
238. Fu, A.; Tang, R.; Hardie, J.; Farkas, M. E.; Rotello, V. M. *Bioconjugate Chem.* **2014**, *25* (9), 1602-1608.
239. Torchilin, V. P.; Lukyanov, A. N. *Drug Discovery Today* **2003**, *8* (6), 259-266.
240. Lu, Y.; Yang, J.; Segal, E. *AAPS J.* **2006**, *8* (3), E466-E478.
241. Vaishya, R.; Khurana, V.; Patel, S.; Mitra, A. K. *Expert Opin. Drug Delivery* **2015**, *12* (3), 415-440.
242. Leich, F.; Stöhr, N.; Rietz, A.; Ulbrich-Hofmann, R.; Arnold, U. *J. Biol. Chem.* **2007**, *282* (38), 27640-27646.
243. D'Astolfo, Diego S.; Pagliero, Romina J.; Pras, A.; Karthaus, Wouter R.; Clevers, H.; Prasad, V.; Lebbink, Robert J.; Rehmann, H.; Geijsen, N. *Cell* **2015**, *161* (3), 674-690.
244. Mix, K. A.; Lomax, J. E.; Raines, R. T. *J. Am. Chem. Soc.* **2017**, *139* (41), 14396-14398.
245. Brodin, J. D.; Sprangers, A. J.; McMillan, J. R.; Mirkin, C. A. *J. Am. Chem. Soc.* **2015**, *137* (47), 14838-14841.
246. Futaki, S.; Nakase, I. *Acc. Chem. Res* **2017**, *50* (10), 2449-2456.
247. Li, M.; Tao, Y.; Shu, Y.; LaRochelle, J. R.; Steinauer, A.; Thompson, D.; Schepartz, A.; Chen, Z.-Y.; Liu, D. R. *J. Am. Chem. Soc.* **2015**, *137* (44), 14084-14093.
248. Cronican, J. J.; Thompson, D. B.; Beier, K. T.; McNaughton, B. R.; Cepko, C. L.; Liu, D. R. *ACS Chem. Biol.* **2010**, *5* (8), 747-752.

249. McNaughton, B. R.; Cronican, J. J.; Thompson, D. B.; Liu, D. R. *Proc. Natl. Acad. Sci.* **2009**, *106* (15), 6111-6116.
250. Hu, S.; Chen, X.; Lei, C.; Tang, R.; Kang, W.; Deng, H.; Huang, Y.; Nie, Z.; Yao, S. *Chem. Commun.* **2018**, *54* (56), 7806-7809.
251. Liu, X.; Zhang, P.; He, D.; Rödl, W.; Preiß, T.; Rädler, J. O.; Wagner, E.; Lächelt, U. *Biomacromolecules* **2016**, *17* (1), 173-182.
252. Liu, X.; Zhang, P.; Rödl, W.; Maier, K.; Lächelt, U.; Wagner, E. *Mol. Pharmaceutics* **2017**, *14* (5), 1439-1449.
253. He, H.; Chen, Y.; Li, Y.; Song, Z.; Zhong, Y.; Zhu, R.; Cheng, J.; Yin, L. *Adv. Funct. Mater.* **2018**, *28* (14), 1706710.
254. Dutta, K.; Hu, D.; Zhao, B.; Ribbe, A. E.; Zhuang, J.; Thayumanavan, S. *J. Am. Chem. Soc.* **2017**, *139* (16), 5676-5679.
255. Wang, M.; Zuris, J. A.; Meng, F.; Rees, H.; Sun, S.; Deng, P.; Han, Y.; Gao, X.; Pouli, D.; Wu, Q.; Georgakoudi, I.; Liu, D. R.; Xu, Q. *Proc. Natl. Acad. Sci.* **2016**, *113* (11), 2868-2873.
256. Cheng, G.; Li, W.; Ha, L.; Han, X.; Hao, S.; Wan, Y.; Wang, Z.; Dong, F.; Zou, X.; Mao, Y.; Zheng, S.-Y. *J. Am. Chem. Soc.* **2018**, *140* (23), 7282-7291.
257. Shao, D.; Li, M.; Wang, Z.; Zheng, X.; Lao, Y.-H.; Chang, Z.; Zhang, F.; Lu, M.; Yue, J.; Hu, H.; Yan, H.; Chen, L.; Dong, W.-f.; Leong, K. W. *Adv. Mater.* **2018**, *30* (29), 1801198.
258. Qiu, M.; Zhang, Z.; Wei, Y.; Sun, H.; Meng, F.; Deng, C.; Zhong, Z. *Chem. Mater.* **2018**, *30* (19), 6831-6838.
259. Li, B.; Yuan, Z.; Zhang, P.; Sinclair, A.; Jain, P.; Wu, K.; Tsao, C.; Xie, J.; Hung, H.-C.; Lin, X.; Bai, T.; Jiang, S. *Adv. Mater.* **2018**, *30* (14), 1705728.
260. Lv, J.; He, B.; Yu, J.; Wang, Y.; Wang, C.; Zhang, S.; Wang, H.; Hu, J.; Zhang, Q.; Cheng, Y. *Biomaterials* **2018**, *182*, 167-175.
261. Zhu, Q.; Chen, X.; Xu, X.; Zhang, Y.; Zhang, C.; Mo, R. *Adv. Funct. Mater.* **2018**, *28* (17), 1707371.
262. Liu, M.; Shen, S.; Wen, D.; Li, M.; Li, T.; Chen, X.; Gu, Z.; Mo, R. *Nano Lett.* **2018**, *18* (4), 2294-2303.
263. Wang, X.; Li, Y.; Li, Q.; Neufeld, C. I.; Pouli, D.; Sun, S.; Yang, L.; Deng, P.; Wang, M.; Georgakoudi, I.; Tang, S.; Xu, Q. *J. Controlled Release* **2017**, *263*, 39-45.

264. Jiang, T.; Mo, R.; Bellotti, A.; Zhou, J.; Gu, Z. *Adv. Funct. Mater.* **2014**, *24* (16), 2295-2304.
265. Gu, Z.; Biswas, A.; Zhao, M.; Tang, Y. *Chem. Soc. Rev.* **2011**, *40* (7), 3638-3655.
266. Leland, P. A.; Raines, R. T. *Chem. Biol.* **2001**, *8* (5), 405-413.
267. Raines, R. T. *Chem. Rev.* **1998**, *98* (3), 1045-1066.
268. Vancoillie, G.; Hoogenboom, R. *Sensors (Basel, Switzerland)* **2016**, *16* (10), 1736.
269. Bankar, S. B.; Bule, M. V.; Singhal, R. S.; Ananthanarayan, L. *Biotechnol. Adv.* **2009**, *27* (4), 489-501.
270. Wong, C. M.; Wong, K. H.; Chen, X. D. *Appl. Microbiol. Biotechnol.* **2008**, *78* (6), 927-938.
271. Hamanaka, R. B.; Chandel, N. S. *J. Exp. Med.* **2012**, *209* (2), 211-215.
272. Warburg, O.; Wind, F.; Negelein, E. *J. Gen. Physiol* **1927**, *8* (6), 519-530.
273. Ahn, C.-H.; Chae, S. Y.; Bae, Y. H.; Kim, S. W. *J. Controlled Release* **2002**, *80* (1), 273-282.
274. Godbey, W. T.; Wu, K. K.; Mikos, A. G. *J. Biomed. Mater. Res.* **1999**, *45* (3), 268-275.
275. Garufi, C.; Nisticò, C.; Vaccaro, A.; D'Ottavio, A.; Zappalà, A. R.; Aschelter, A. M.; Terzoli, E. *Ann. Oncol.* **2001**, *12* (2), 179-182.
276. Vondálová Blanářová, O.; Jelínková, I.; Szöör, Á.; Skender, B.; Souček, K.; Horváth, V.; Vaculová, A.; Anděra, L.; Sova, P.; Szöllösi, J.; Hofmanová, J.; Vereb, G.; Kozubík, A. *Carcinogenesis* **2010**, *32* (1), 42-51.
277. Serova, M.; Calvo, F.; Lokiec, F.; Koepfel, F.; Poindessous, V.; Larsen, A. K.; Laar, E. S. V.; Waters, S. J.; Cvitkovic, E.; Raymond, E. *Cancer Chemother. Pharmacol.* **2006**, *57* (4), 491-499.
278. Zhang, R. X.; Wong, H. L.; Xue, H. Y.; Eoh, J. Y.; Wu, X. Y. *J. Controlled Release* **2016**, *240*, 489-503.
279. Major Jourden, J. L.; Cohen, S. M. *Angew. Chem., Int. Ed.* **2010**, *49* (38), 6795-6797.
280. Tomsho, J. W.; Benkovic, S. J. *J. Org. Chem.* **2012**, *77* (5), 2098-2106.
281. Tripathy, D. R.; Dinda, A. K.; Dasgupta, S. *Anal. Biochem.* **2013**, *437* (2), 126-129.

282. Liu, Y.; Du, J.; Yan, M.; Lau, M. Y.; Hu, J.; Han, H.; Yang, O. O.; Liang, S.; Wei, W.; Wang, H.; Li, J.; Zhu, X.; Shi, L.; Chen, W.; Ji, C.; Lu, Y. *Nat. Nanotechnol.* **2013**, 8, 187.
283. Benjaminsen, R. V.; Matthebjerg, M. A.; Henriksen, J. R.; Moghimi, S. M.; Andresen, T. L. *Mol. Ther.* **2013**, 21 (1), 149-157.