The integrin α L leg region controls the Mg/EGTA mediated activation of LFA-1

Guan, Siyu; Cheng, Ming; Law, S.K. Alex

2015

Guan, S., Cheng, M., & Law, S. K. A. (2015). The integrin α L leg region controls the Mg/EGTA mediated activation of LFA-1. Biochemical and biophysical research communications, 458(2), 251-255.

<https://hdl.handle.net/10356/106917>

<https://doi.org/10.1016/j.bbrc.2015.01.094>

© 2015 Elsevier Inc. This is the author created version of a work that has been peer reviewed and accepted for publication by Biochemical and Biophysical Research Communications, Elsevier Inc. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [<http://dx.doi.org/10.1016/j.bbrc.2015.01.094>].

Downloaded on 26 Apr 2023 18:58:32 SGT

The integrin α L leg region controls the Mg/EGTA mediated activation of LFA-1

Siyu Guan, Ming Cheng¹, and S. K. Alex Law*

School of Biological Sciences, Nanyang Technological University, Singapore

¹ Current address: Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, New York, USA

* Corresponding author at: School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore 637551. Fax: +65 6515 1632

Email address: alaw@ntu.edu.sg

Abstract

We have shown that Mg/EGTA (5 mM Mg^{2+} and 1.5 mM EGTA) could effectively promote the adhesion of integrin $\alpha\text{L}\beta_2$ to its ligand ICAM-1 but could not promote that of the $\alpha\text{M}\beta_2$ to denatured BSA. In order to determine the structural differences between αL and αM that specifically contribute to Mg/EGTA sensitivity, a series of $\alpha\text{L}/\alpha\text{M}$ chimeras were constructed. Our results showed that $\alpha\text{L}\beta_2$ with αM calf-1 domain completely lost the response to Mg/EGTA activation. In the reverse experiment, $\alpha\text{M}\beta_2$ would require the presence of both the αL calf-1 and calf-2 domain to initiate the Mg/EGTA sensitivity.

Keywords

Divalent cation
Integrin activation
Domain swapping
Cell Adhesion

Abbreviations

EGTA, Ethylene glycol tetra-acetic acid; LFA-1, Leukocyte function associated antigen-1 ($\alpha\text{L}\beta_2$); Mac-1, Macrophage antigen-1 ($\alpha\text{M}\beta_2$); ICAM, Intercellular adhesion molecule; BSA, Bovine serum albumin; WT, Wild-type.

Highlights

- Addressed the difference of Mg/EGTA sensitivity in the activation between $\alpha\text{L}\beta_2$ and $\alpha\text{M}\beta_2$, by domain swapping.
- The leg region, rather than the headpiece causes the Mg/EGTA sensitivity.
- αL calf-1 domain is responsible for Mg/EGTA regulated activation.
- The presence of both calf-1 and calf-2 domains of αL restored the Mg/EGTA sensitivity of $\alpha\text{M}\beta_2$.

1. Introduction

Integrins are a family of membrane glycoproteins that serve as bi-directional signaling molecules between the interior of the cell and its extracellular environment. Integrins are heterodimers, each member is consisted of one α subunit and one β subunit, which associate with each other non-covalently [1]. In human, there are 24 integrin heterodimers formed by selective combinations from 18 different α subunits and 8 β subunits. Of specific interest to us are the members of the β_2 integrin family. Four α subunits, namely α_L , α_M , α_X , and α_D , combine with the common β_2 subunit to form a closed set of $\alpha_L\beta_2$, $\alpha_M\beta_2$, $\alpha_X\beta_2$, and $\alpha_D\beta_2$ integrins. They are expressed exclusively on leukocytes. Specifically, $\alpha_L\beta_2$, $\alpha_M\beta_2$, and $\alpha_X\beta_2$ are expressed on peripheral blood leukocytes and the regulation of their adhesion properties has been studied extensively [2,3,4,5]. Generally, these integrin molecules are at a resting state, i.e. incapable to mediate cell adhesion to their ligands.

Integrin activation is regulated bi-directionally. The inside-out signal can be from another receptor on the cell, such as the CD14 on monocytes and CD40 on B cells [4]. The signal is transmitted to the cytoplasmic tail of the β_2 subunit by intracellular proteins such as talin and kindlin [6]. Upon such interaction, the ectodomain of the integrin changes from a folded compact conformation to an extended and open conformation. Functionally, the open integrin can adhere to ligand with high affinity. Activation can also be induced from outside via ligand binding. The signal can be manifested to cellular activities such as cell spreading and cell migration. The α subunits of the β_2 integrin family belong to the subset that has an inserted I-domain, which we shall refer to as the α_I domain. For these integrins, the α_I domain is the major ligand binding site [7].

In the laboratory, the transition of integrin activity states can be studied by the manipulation of extracellular conditions: either by antibodies that promote the conformational change associated with activation, or by the alteration of the concentrations of divalent cations. Ca^{2+} , Mg^{2+} and Mn^{2+} are capable in regulating the adhesion between leukocyte integrins and ligands, among which Ca^{2+} generally inhibits the ligand-receptor binding, whereas Mg^{2+} and Mn^{2+} generally promote the adhesion [8,9,10]. Divalent cation binding sites have been identified in the headpiece of the integrin heterodimer, i.e. in the β -propeller of the α subunits, the βI domain of the β subunits, and also on the αI domain for those α subunits with this domain. As of today, the structure of the ectodomain of three integrin heterodimers had been solved by X-ray crystallography; they are the $\alpha\text{V}\beta_3$ [11], $\alpha\text{IIb}\beta_3$ [12] and $\alpha\text{X}\beta_2$ [13] integrins. For αIIb and αV , a Ca^{2+} binding site has also been found in the genu region between the thigh and the calf-1 domain. However, this Ca^{2+} binding site was not identified in the αX subunit.

It was discovered in 1992 that the high concentration of Mg^{2+} coupled with the depletion of Ca^{2+} by the inclusion of EGTA in the medium would activate LFA-1 from a resting to a ligand binding state [9]. In contrast, it appears that both Ca^{2+} and Mg^{2+} are required to activate Mac-1 to adhere to iC3b [8]. In this study, we focus to discern the difference of Mg/EGTA mediated activation between LFA-1 and Mac-1 and to identify the domain responsible for this difference.

2. Material and Methods

2.1. Antibodies and reagents

Hybridomas of mAbs were obtained from different sources: MHM24 (anti- α L blocking mAb) and MHM23 (β_2 heterodimer specific mAb) [14] were provided by Professor AJ McMichael (John Radcliff Hospital, Oxford, UK). KIM185 (anti- β_2 activating mAb) [15] was obtained from Dr. MK Robinson (UCB, CellTech, UK). LPM19c (anti- α M blocking mAb) [16] was purchased from ATCC. All mAbs were purified from hybridoma supernatant using Hi-Trap protein G or A columns (Amersham Pharmacia Biotech). Preparation of recombinant human ICAM-1-Fc was described previously [17]. Bovine serum albumin (fraction V, BSA) was bought from Sigma-Aldrich.

2.2 *plasmids*

The pcDNA3 expression plasmids containing α L, α M and β_2 cDNA were described previously [18]. Each domain swapping plasmid was constructed by a two-step PCR. In the 1st step, a pair of 64-70-mer primers containing donor sequences at 5' tails that were ~30 bases in length and receptor sequences at 3' tails that were ~35 bases in length was designed. A mega primer containing the domain of interest from the donor in the middle, and flanked by the receptor sequences were synthesized by PCR using Advantage® -GC cDNA Polymerase Mix kit (ClonTech), followed the manufacturer's instruction. The mega primer was obtained by gel purification. In the 2nd step, the purified mega primer was used to clone the chimera using Pfu Turbo polymerase (Stratagene). Setting of thermo cycling parameters was followed the manufacturer's instruction with the exception of using a gradually decreased annealing temperature which dropped 1°C every 30 seconds from 76°C to 45°C. Subsequent procedures after PCR were followed by the instruction from the manufacturer. All constructs were verified by sequencing (Axil Scientific, Singapore).

2.3. *Cell culture and transfection*

Human embryonic kidney 293T (HEK293T) cells were purchased from ATCC, cells were cultured in DMEM (Hyclone) supplemented with 10% (v/v) heat-inactivated FBS (Hyclone), 100 IU/ml penicillin and 100 µg/ml streptomycin (Hyclone) at 37°C in 5% CO₂ humidified incubator. HEK293T cells were transfected with expression plasmids using polyethylenimine (PEI) according to a previously described method [19].

2.4. *Flow cytometry*

Surface expression detection was described previously [17] using a β₂ heterodimer specific mAb MHM23 followed by FITC-conjugated sheep anti-mouse IgG [20]. Background histograms were obtained using the LPM19c mAb (αM specific) for LFA-1, and MHM24 mAb (αL specific) for Mac-1. Stained cells were collected on a FACS Calibur (BD Biosciences). Data were analyzed using Flowjo (Tree Star).

2.5. *Cell adhesion assay*

For ICAM-1 adhesion, 100 µl of 5 µg/ml goat anti-human IgG (Fc specific) (Sigma-Aldrich) in bicarbonate buffer (50 mM, pH 9.2) was introduced into each well in 96-well microtiter plates (Polysorb, Nunc Immno-Plate) and left overnight at 4°C. Non-specific sites were blocked with 130 µl of 0.5% (w/v) BSA in PBS for each well. 50 µl of ICAM-1-Fc in PBS with 0.1% BSA (w/v) at 1 µg/ml was added into each well and left at RT for 2 hours. For BSA adhesion assay, 100 µl of 100 µg/ml BSA in bicarbonate buffer was coated on each well and the plate was left overnight at 4°C. Non-specific sites were blocked with 130 µl of 0.2% (w/v) polyvinylpyrrolidone (PVP) in PBS for each well. In both cases the wells were washed twice with wash buffer (RPMI1640 with 10 mM HEPES and 5% v/v FBS) before adding cells. Transfected cells were harvested, washed and incubated with 1 µg/ml BCECF (Invitrogen) in wash buffer at 37°C for 20 min. Labeled cells were transferred to each wells

with or without activation reagents (Mg/EGTA: 5 mM MgCl₂ and 1.5 mM EGTA; Mn²⁺: 0.5 mM MnCl₂; mAb KIM185: 10 µg/ml) and incubated at 37°C for 30 min. To establish specificity, appropriate antibodies were added to block adhesion. Unbound cells were removed by washing twice with wash buffer. Fluorescence signal, which correlates with the number of cells adhering to the ligand-coated well, was measured using a FL600 fluorescence plate reader (Bio-Tek instruments, Winooski, VT).

3. Results

3.1 The effect of Mg/EGTA on α L β_2 and α M β_2 mediated adhesion

Manipulation of divalent cation concentrations, generally Ca²⁺, Mg²⁺ and Mn²⁺, in the cell medium had been found to affect their integrin mediated adhesion to ligands. Mn²⁺, at sub mM level, has been shown to promote α L β_2 and α M β_2 mediated adhesion to their ligands. However, the requirements for Mg²⁺ and Ca²⁺ are different for the two integrins. It was shown that both Mg²⁺ and Ca²⁺ are required for α M β_2 mediated adhesion [8,21] whereas Ca²⁺ is inhibitory to α L β_2 mediated adhesion [9]. This observation is confirmed in an experiment shown in Figure 1. α L β_2 and α M β_2 transfectants were assayed for their adhesion to their respective ligands ICAM-1 and denatured BSA. Adhesion can be promoted for both transfectants with the β_2 -specific activating mAb KIM185, α L β_2 transfectants, but not the α M β_2 can be activated by Mg/EGTA. Since the two integrins share the common β_2 subunit, it is logical to argue that this difference is due to certain structural features that are different in α L and α M subunits.

3.2 α L calf-1 domain is required for Mg/EGTA mediated activation

A series of α L/ α M chimera were constructed such that the domains of the leg region of the α M subunit were into the α L background. These chimeras, together with the β_2 subunit, were co-transfected into HEK293T cells. Expression of the chimeric integrins was monitored by flow cytometry using the mAb MHM23, and adhesion to ICAM-1 was performed. The results are shown in Figure 2. Chimeras with the calf-1 domain of α M would render the integrin α M β_2 -like, in that Mg/EGTA cannot stimulate the adhesion of the transfectants to ICAM-1.

Significantly, the integrin with only the α M calf-1 domain on an α L β_2 background is not responsive to Mg/EGTA stimulation, suggesting that Mg/EGTA response element is located in the calf-1 domain of α L subunit.

3.3 The restoration of Mg/EGTA mediated activation on α M β_2

The converse series of chimera were constructed where domains of the leg region of α L were introduced into the α M background. Mg/EGTA stimulation can be demonstrated if the region C-terminal and inclusive of the calf-1, calf-2 and cytoplasmic segment is from α L. Replacement with just the calf-1 or calf-2 region with those of α L result in integrins that are constitutively active in binding to the α M β_2 ligand denatured BSA. In these cases, Mg/EGTA stimulation cannot be determined. Nevertheless, these results do not contradict those obtained with the series with α L background.

4. Discussion

The requirement of divalent cations for integrin-mediated adhesion has long been investigated [2,10,22,23]. However, the distinct divalent cation requirements of α L β_2 and α M β_2 had not been addressed. Some studies had chose to add to the cell medium with millimolar level of EGTA and MgCl₂ as a standard condition for α L β_2 -mediated adhesion

[24,25,26,27,28], while millimolar level of MnCl_2 was used to promote $\alpha\text{M}\beta_2$ mediated adhesion [27,29,30]. In this article we have established, Mg/EGTA is not able to activate $\alpha\text{M}\beta_2$ transfectants to adhere to denatured BSA.

We argued that the functional difference must be conferred by the αL and αM subunit. We therefore constructed a series of chimeric $\alpha\text{L}/\alpha\text{M}$ subunit and pinpoint the domain that is responsible for this functional difference. In the series using αL as the background, we found that replacement of the calf-1 domain with that of αM would abolish the activation by Mg/EGTA . The reverse series was less clear-cut. Using an αM background, we can only demonstrate that the Mg/EGTA sensitivity can be observed if both the calf-1 and calf-2 of αL were introduced. Introduction of either calf-1 or calf-2 did not yield any informative results as the integrins with these two $\alpha\text{L}/\alpha\text{M}$ chimeras are constitutively active. Thus the effect of Mg/EGTA cannot be determined. This is not a totally unexpected result. Keeping an integrin in a resting state would require the coordinated interactions of its domains. These interactions would be disrupted if segments from another integrin were introduced to replace the corresponding segments. It was demonstrated that if we replaced the leg region of the β_2 subunit with that of β_1 , the resultant $\alpha\text{L}\beta_{2/1}$ chimera would be constitutively active [31]. Indeed, the $\alpha\text{X}\beta_2$ integrin was shown to be constitutively active integrins if the αX subunit was from human and β_2 subunit from chicken [32].

Similar observation was found with the $\alpha\text{V}\beta_3$ and $\alpha\text{IIb}\beta_3$ integrins, in which 1 mM Mn^{2+} was able to activate $\alpha\text{V}\beta_3$, but not $\alpha\text{IIb}\beta_3$, to bind fibrinogen. Using the similar domain swapping strategy, the domain responsible for Mn^{2+} sensitivity was located to the calf-2 domain of αV [33]. The ectodomain structures of both $\alpha\text{V}\beta_3$ and $\alpha\text{IIb}\beta_3$ had been solved, but no divalent cation (and such binding site) was found in the calf-2 domain [11,12]. In both the

αV and αIIb subunits, a Ca^{2+} ion was found to be chelated with the residues in the genu and the calf-1 domain [11,12]. However, such Ca^{2+} ion binding site was not found in the $\alpha X\beta_2$ integrin [13], although the potential residues can be identified when aligned with those in the αV and αIIb subunits. On detailed analyses of the structure, it was concluded that the backbone structure of αX is sufficiently different such that the geometry is not appropriate to facilitate Ca^{2+} binding [13].

αL is different from αX , which is more closely related to αM . Thus, it is not possible to predict if such a divalent binding site is present in αL . We have attempted to express the various combinations of the domains in leg region of αL and αM (from thigh to calf-2) but with no success. Until such information becomes available, it will remain a puzzle as to why the two integrins $\alpha L\beta_2$ and $\alpha M\beta_2$ would respond differently to the exposure to Mg/EGTA.

It was widely believed that the divalent cation binding sites at integrin αI domain and βI domain are the sites that account for the divalent cation mediated integrin adhesion. However, mutation on these sites, namely, MIDAS of αI , MIDAS, ADMIDAS and SyMBS of βI usually lead to completely abolished ligand binding or even surface expression [34,35,36,37]. Our results, together with those published by Kamata et al [34] on the $\alpha IIb\beta_3$ and $\alpha V\beta_3$ integrins, suggest that there may be a remote action of the α subunit leg domain that controls the divalent cation binding sites in the headpiece. Alternatively, there may be other yet to be identified divalent cation binding site in the leg region.

Acknowledgements

This work was supported by Tier 1 grant from Nanyang Technological University Academic Research Fund (AcRF) RG 93/07.

Figure Legends

- Fig. 1.** Adhesion of integrin $\alpha\text{L}\beta_2$ (A) and $\alpha\text{M}\beta_2$ (B) transfectants binding to their respective ligands ICAM-1 and denatured BSA, in the presence of Mg/EGTA. 10 $\mu\text{g/ml}$ activating mAb KIM185 was included as a positive control.
- Fig. 2.** Upper panel: Adhesion of wild-type $\alpha\text{L}\beta_2$ and chimeric $\alpha\text{LM}\beta_2$ integrins to ICAM-1 with no activating agents (white), in the presence of Mg/EGTA (grey), and the activating mAb KIM185 (black). For each condition, the mAb MHM24, at 10 $\mu\text{g/ml}$, was used to block the adhesion and shown on the right. Middle panel: Expression profile of the wild-type and chimeric integrins using the mAb MHM23 (β_2 heterodimer specific). Background histograms were obtained using the mAb LPM19c (αM specific). Lower panel: domain organisation of the α subunits of the chimeric integrins – αL domains in white, and αM domains in black. A letter “I” is assigned to represent for the αI domain, “ $\beta\text{-P}$ ” represents for the β -propeller domain, “T” represents for the thigh domain, “1” represents for the calf-1 domain and “2” represents for the calf-2 domain.
- Fig. 3.** Upper panel: Adhesion of wild-type $\alpha\text{M}\beta_2$ and chimeric $\alpha\text{ML}\beta_2$ integrins to denatured BSA with no activating agents (white), in the presence of Mg/EGTA (grey), and the activating mAb KIM185 (black). For each condition, the mAb LPM19c, at 10 $\mu\text{g/ml}$, was used to block the adhesion and shown on the right. Middle panel: Expression profile of the wild-type and chimeric integrins using the mAb MHM23 (β_2 heterodimer specific). Background histograms were obtained using the mAb MHM24 (αL specific). Lower panel: domain organisation of the α subunits of the chimeric integrins.

Reference

- [1] R.O. Hynes, Integrins: Bidirectional, allosteric signaling machines, *Cell* 110 (2002) 673-687.
- [2] B.H. Luo, C.V. Carman, T.A. Springer, Structural basis of integrin regulation and signaling, *Annual Review of Immunology* 25 (2007) 619-647.
- [3] N. Hogg, I. Patzak, F. Willenbrock, The insider's guide to leukocyte integrin signalling and function, *Nature Reviews Immunology* 11 (2011) 416-426.
- [4] C.L. Abram, C.A. Lowell, The ins and outs of leukocyte integrin signaling, *Annual review of immunology* 27 (2009) 339-362.
- [5] T.A. Springer, M.L. Dustin, Integrin inside-out signaling and the immunological synapse, *Current Opinion in Cell Biology* 24 (2012) 107-115.
- [6] S.J. Shattil, C. Kim, M.H. Ginsberg, The final steps of integrin activation: the end game, *Nature reviews. Molecular cell biology* 11 (2010) 288-300.
- [7] M.J. Humphries, E.J. Symonds, A.P. Mould, Mapping functional residues onto integrin crystal structures, *Current opinion in structural biology* 13 (2003) 236-243.
- [8] S.D. Wright, S.C. Silverstein, Tumor-Promoting Phorbol Esters Stimulate C3b and C3b' Receptor-Mediated Phagocytosis in Cultured Human-Monocytes, *Journal of Experimental Medicine* 156 (1982) 1149-1164.
- [9] I. Dransfield, C. Cabanas, A. Craig, N. Hogg, Divalent-Cation Regulation of the Function of the Leukocyte Integrin Lfa-1, *Journal of Cell Biology* 116 (1992) 219-226.
- [10] E.F. Plow, T.K. Haas, L. Zhang, J. Loftus, J.W. Smith, Ligand binding to integrins, *Journal of Biological Chemistry* 275 (2000) 21785-21788.
- [11] J.P. Xiong, T. Stehle, B. Diefenbach, R.G. Zhang, R. Dunker, D.L. Scott, A. Joachimiak, S.L. Goodman, M.A. Arnaout, Crystal structure of the extracellular segment of integrin alpha V beta 3, *Science* 294 (2001) 339-345.
- [12] J.H. Zhu, B.H. Luo, T. Xiao, C.Z. Zhang, N. Nishida, T.A. Springer, Structure of a Complete Integrin Ectodomain in a Physiologic Resting State and Activation and Deactivation by Applied Forces, *Molecular Cell* 32 (2008) 849-861.
- [13] C. Xie, J.H. Zhu, X. Chen, L.Z. Mi, N. Nishida, T.A. Springer, Structure of an integrin with an alpha I domain, complement receptor type 4, *Embo Journal* 29 (2010) 666-679.
- [14] J.E.K. Hildreth, F.M. Gotch, P.D.K. Hildreth, A.J. Mcmichael, A Human Lymphocyte-Associated Antigen Involved in Cell-Mediated Lympholysis, *European Journal of Immunology* 13 (1983) 202-208.
- [15] M.K. Robinson, D. Andrew, H. Rosen, D. Brown, S. Ortlepp, P. Stephens, E.C. Butcher, Antibody against the Leu-Cam Beta-Chain (Cd18) Promotes Both Lfa-1-Dependent and Cr3-Dependent Adhesion Events, *Journal of Immunology* 148 (1992) 1080-1085.
- [16] P. Uciechowski, R.E. Schmidt, NK and non-lineage antigens cluster report: CD11. , In *Leukocyte Typing IV: White Cell Differentiation Antigens*. W. Knapp, editor . Oxford University Press, Oxford . (1989) 543-551.
- [17] S.M. Tan, R.H. Hyland, A. Al-Shamkhani, W.A. Douglass, J.M. Shaw, S.K.A. Law, Effect of integrin beta(2) subunit truncations on LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18) assembly, surface expression, and function, *Journal of Immunology* 165 (2000) 2574-2581.

- [18] J.M. Shaw, A. Al-Shamkhani, L.A. Boxer, C.D. Buckley, A.W. Dodds, N. Klein, S.M. Nolan, I. Roberts, D. Roos, S.L. Scarth, D.L. Simmons, S.M. Tan, S.K.A. Law, Characterization of four CD18 mutants in leucocyte adhesion deficient (LAD) patients with differential capacities to support expression and function of the CD11/CD18 integrins LFA-1, Mac-1 and p150,95, *Clinical and Experimental Immunology* 126 (2001) 311-318.
- [19] Y. Durocher, S. Perret, A. Kamen, High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells, *Nucleic Acids Research* 30 (2002).
- [20] J.E.K. Hildreth, J.T. August, The Human-Lymphocyte Function-Associated (Hlfa) Antigen and a Related Macrophage Differentiation Antigen (Hmac-1) - Functional-Effects of Subunit-Specific Monoclonal-Antibodies, *Journal of Immunology* 134 (1985) 3272-3280.
- [21] D.C. Altieri, Occupancy of Cd11b/Cd18 (Mac-1) Divalent Ion Binding Site(S) Induces Leukocyte Adhesion, *Journal of Immunology* 147 (1991) 1891-1898.
- [22] J.P. Xiong, T. Stehle, S.L. Goodman, M.A. Arnaout, Integrins, cations and ligands: making the connection, *Journal of Thrombosis and Haemostasis* 1 (2003) 1642-1654.
- [23] M.E. Binnerts, Y. van Kooyk, How LFA-1 binds to different ligands, *Immunology Today* 20 (1999) 240-245.
- [24] S.M. Tan, M.K. Robinson, K. Drbal, Y. van Kooyk, J.M. Shaw, S.K.A. Law, The N-terminal region and the mid-region complex of the integrin beta(2) subunit, *Journal of Biological Chemistry* 276 (2001) 36370-36376.
- [25] A. Smith, M. Bracke, B. Leitinger, J.C. Porter, N. Hogg, LFA-1-induced T cell migration on ICAM-1 involves regulation of MLCK-mediated attachment and ROCK-dependent detachment, *Journal of cell science* 116 (2003) 3123-3133.
- [26] E. Tng, S.M. Tan, S. Ranganathan, M. Cheng, S.K.A. Law, The integrin alpha(L)beta(2) hybrid domain serves as a link for the propagation of activation signal from its stalk regions to the I-like domain, *Journal of Biological Chemistry* 279 (2004) 54334-54339.
- [27] M. Cheng, S.Y. Foo, M.L. Shi, R.H. Tang, L.S. Kong, S.K.A. Law, S.M. Tan, Mutation of a conserved asparagine in the I-like domain promotes constitutively active integrins alpha(L)ss(2) and alpha(IIb)ss(3), *Journal of Biological Chemistry* 282 (2007) 18225-18232.
- [28] A. Vararattanavech, M.L. Tang, H.Y. Li, C.H. Wong, S.K.A. Law, J. Torres, S.M. Tan, Permissive transmembrane helix heterodimerization is required for the expression of a functional integrin, *Biochemical Journal* 410 (2008) 495-502.
- [29] D. Ehrichtiou, Dual function for a unique site within the beta I-2 domain of integrin alpha(M)beta(2), *Circulation* 110 (2004) 312-312.
- [30] Z.H. Xue, C.Q. Zhao, G.L. Chua, S.W. Tan, X.Y. Tang, S.C. Wong, S.M. Tan, Integrin alphaMbeta2 clustering triggers phosphorylation and activation of protein kinase C delta that regulates transcription factor Foxp1 expression in monocytes, *Journal of immunology* 184 (2010) 3697-3709.
- [31] W.A. Douglass, R.H. Hyland, C.D. Buckley, A. Al-Shamkhani, J.M. Shaw, S.L. Scarth, D.L. Simmons, S.K. Law, The role of the cysteine-rich region of the beta2 integrin subunit in the leukocyte function-associated antigen-1 (LFA-1, alphaLbeta2, CD11a/CD18) heterodimer formation and ligand binding, *FEBS letters* 440 (1998) 414-418.
- [32] C.A. Bilsland, M.S. Diamond, T.A. Springer, The leukocyte integrin p150,95 (CD11c/CD18) as a receptor for iC3b. Activation by a heterologous beta subunit

- and localization of a ligand recognition site to the I domain, *Journal of immunology* 152 (1994) 4582-4589.
- [33] T. Kamata, M. Handa, Y. Sato, Y. Ikeda, S. Aiso, Membrane-proximal {alpha}/{beta} stalk interactions differentially regulate integrin activation, *The Journal of biological chemistry* 280 (2005) 24775-24783.
- [34] N. Hogg, M.P. Stewart, S.L. Scarth, R. Newton, J.M. Shaw, S.K.A. Law, N. Klein, A novel leukocyte adhesion deficiency caused by expressed but nonfunctional beta 2 integrins Mac-1 and LFA-1, *Journal of Clinical Investigation* 103 (1999) 97-106.
- [35] E.C. Mathew, J.M. Shaw, F.A. Bonilla, S.K.A. Law, D.A. Wright, A novel point mutation in CD18 causing the expression of dysfunctional CD11/CD18 leucocyte integrins in a patient with leucocyte adhesion deficiency (LAD), *Clinical and Experimental Immunology* 121 (2000) 133-138.
- [36] J. Chen, W. Yang, M. Kim, C.V. Carman, T.A. Springer, Regulation of outside-in signaling and affinity by the beta(2) I domain of integrin alpha(L)beta(2), *Proceedings of the National Academy of Sciences of the United States of America* 103 (2006) 13062-13067.
- [37] M.L. Bajt, T. Goodman, S.L. Mcguire, Beta(2) (Cd18) Mutations Abolish Ligand Recognition by I-Domain Integrins Lfa-1 (Alpha(L)Beta(2), Cd11a/Cd18) and Mac-1 (Alpha(M)Beta(2), Cd11b/Cd18), *Journal of Biological Chemistry* 270 (1995) 94-98.

Fig.1

Fig. 2

Fig. 3