Quantitative proteomic study of *Aspergillus fumigatus* secretome revealed deamidation of secretory enzymes

Adav, Sunil S.; Ravindran, Anita; Sze, Siu Kwan

2015

Adav, S. S., Ravindran, A., & Sze, S. K. (2015). Quantitative proteomic study of *Aspergillus fumigatus* secretome revealed deamidation of secretory enzymes. *Journal of proteomics*, 119, 154-168.

<https://hdl.handle.net/10356/107411>

<https://doi.org/10.1016/j.jprot.2015.02.007>

© 2015 Elsevier B.V. This is the author created version of a work that has been peer reviewed and accepted for publication by *Journal of Proteomics*, Elsevier B.V. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [Article DOI: <http://dx.doi.org/10.1016/j.jprot.2015.02.007>].

Downloaded on 26 Apr 2023 20:04:59 SGT

1 **Quantitative Proteomic Study of *Aspergillus Fumigatus* Secretome**
2 **Revealed Deamidation of Secretory Enzymes**

3 **Sunil S. Adav#, Anita Ravindran#, Siu Kwan Sze***

4 School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,
5 Singapore 637551

10 **Key words:** *Aspergillus Fumigatus*, bioenergy, biorefinery, cellulases, deamidation

12 #Authors contributed equally

14 *** Corresponding author**

15 Siu Kwan SZE, PhD
16 School of Biological Sciences
17 Division of Chemical Biology & BioTechnology
18 Nanyang Technological University,
19 60 Nanyang drive, Singapore 637551
20 Tel: +65: 6514-1006
21 Fax: +65: 6791-3856
22 Email: sksze@ntu.edu.sg

Abstract

Aspergillus sp. plays an essential role in lignocellulosic biomass recycling and is also exploited as cell factories for the production of industrial enzymes. This study profiled the secretome of *Aspergillus fumigatus* when grown with cellulose, xylan and starch by high throughput quantitative proteomics using isobaric tags for relative and absolute quantification (iTRAQ). Post translational modifications (PTMs) of proteins play a critical role in protein functions. However, our understanding of the PTMs in secretory proteins is limited. Here, we present the identification of PTMs such as deamidation of secreted proteins of *A. fumigatus*. This study quantified diverse groups of extracellular secreted enzymes and their functional classification revealed cellulases and glycoside hydrolases (32.9%), amylases (0.9%), hemicellulases (16.2%), lignin degrading enzymes (8.1%), peptidases and proteases (11.7%), chitinases, lipases and phosphatases (7.6%), and proteins with unknown function (22.5%). The comparison of quantitative iTRAQ results revealed that cellulose and xylan stimulates expression of specific cellulases and hemicellulases, and their abundance level as a function of substrate. In-depth data analysis revealed deamidation as a major PTM of key cellulose hydrolyzing enzymes like endoglucanases, cellobiohydrolases and glucosidases. Hemicellulose degrading endo-1,4-beta-xylanase, monosidases, xylosidases; lignin degrading laccase, isoamyl alcohol oxidase and oxidoreductases were also found to be deamidated.

Introduction

Lignocellulosic biomass comprising mainly of cellulose, hemicellulose, and lignin is the largest sink of global carbon. With the growing demand and shortage of fuel, exhausting fossil fuel reservoirs, rising CO₂ emissions and natural environmental deterioration; replacement of fossil fuels by renewable energy sources is an urgent global agenda. Biomasses such as agricultural wastes, crop residues, forest wastes etc. constitute potential sources of feedstock and could be targeted as a potential raw material for biofuels and new biomaterials portfolios including feed, specialty chemicals, and other value-added products [1]. These biomasses are natural, abundant, and renewable resources of carbon. Their full exploitation remains critical to the development of a sustainable global economy. Carbohydrate moieties of biomass such as cellulose, hemicellulose and pectin can be hydrolyzed into monomeric sugars and subsequently fermented into bioethanol, biogases, organic acids, chemicals, or biofuel, while the aromatic compounds, vanillin and gallic acids, can also be produced from lignin [2]. Xylitol and furfural can be derived from xylose [3]. In the biofuel production sector, the USA has developed a relatively mature technology for the production of corn ethanol and they produce more than 52.6 billion liters of ethanol per year [4, 5].

The carbohydrate fractions of the biomass can be converted into fermentable monomeric sugars through enzymatic (hemicellulase/cellulase) reactions. It has been exploited to produce ethanol, xylitol, and 2, 3-butanediol via microbial fermentation processes [6]. The conventional conversion of starch to glucose requires a two-step process, namely, liquefaction and saccharification, but this process is energy-intensive [5]. On the contrary, raw starch degrading enzymes (RSDE) can directly degrade raw starch granules below the gelatinization temperature of starch and thus significantly reduce

energy requirements, simplifying the process in the industry [7]. As reviewed by Sun et al, [5] several strains of *Aspergillus* sp. has potential to produce RSDE. To explore the cellulose and lignocellulosic biomass hydrolysis potential and to investigate the abundance of lignocellulolytic enzymes, the proteomic analysis of secretome by biomass degrading microbes such as *Thermobifida fusca* [8], *Trichoderma reesei* [9-11], *Phanerochaete chrysosporium* [12, 13], *Botrytis cinerea* [14, 15], *Postia placenta* [16], *Aspergillus niger* [17] *Fusarium graminearum* [18] and many more have been documented. However, the post translational modifications (PTMs) of these proteins that could result in a huge impact on their functions have not been studied so far. Although, proteomic analysis of secretomes from biomass degrading microbes has been performed, but very limited literature exists on PTMs of lignocellulolytic enzymes.

The fungal species like *T. reesei* is extensively studied and also used in the industry as a source of cellulases and hemicellulases. However, with a total of 200 glycosyl hydrolases (GH) encoding genes, *T. reesei* has fewer GHs than ascomycetes fungi of the eurotiomycetes lineage such as *A. nidulans* (247GHs), *A. fumigatus* (263 GHs). *A. oryzae* (185 GHs) has fewer GHs than *T. reesei* [19]. Similarly, with 16 hemicellulase genes, *T. reesei* has the smallest set of hemicellulases in comparison to other fungi like *A. nidulans* (35 genes), *A. fumigatus* (36 genes), *A. oryzae* (40 genes). *Aspergillus* sp. therefore represents a family of fungi with more diverse range of lignocellulosic hydrolyzing enzymes for industrial application. iTRAQ based proteomic analysis of the secretome of thermophilic *A. fumigatus* Z5 in the presence of different carbon sources (glucose, Avicel and rice straw) revealed the carbon source dependent production of cellulases and hemicellulases [20]. *Aspergillus niger* NS-2 grown in

1 presence of wheat bran following solid state fermentation techniques produced good
2 yields of lignocellulolytic enzymes [21]. Thus, studies of *Aspergillus* sp. are gaining
3 more attention due to its ability to secrete lignocellulolytic enzymes that are useful in
4 lignocellulosic biorefinery. However, recent research has mostly focused on the
5 evaluation of lignocellulolytic enzyme production or purification of the main components
6 [22, 23], and limited literature exists on quantitative profiling of the secretome.
7 Thermostable enzymes from thermophilic fungi has its own industrial advantages since
8 thermostable enzymes use shorter reaction times for the complete saccharification of
9 plant polysaccharides as compared to hydrolytic enzymes of mesophilic fungi [24-26].
10 Thermal pretreatment of biomass loosens biomass structure and enhance penetration of
11 hydrolytic enzymes [27]. The comprehensive quantitative profiling of lignocellulolytic
12 enzymes of thermophilic fungi on different carbon sources could be a practical approach
13 for understanding unique enzyme systems. Coupling of thermal biomass treatment with
14 thermotolerant enzymes could enhance biomass hydrolysis significantly. It is also noted
15 that most of glycosyl hydrolases undergo modification like glycosylation, deamidation,
16 phosphorylation etc. and these PTMs play important role in enzyme function, stability
17 and interaction with lignocellulose [28]. PTMs like glycosylation and sites of
18 modification in endoglucanase I, II and exoglucanase I, II from *T. reesei* and some
19 biomass degrading enzymes of *Phanerochaete chrysosporium* etc. have been documented
20 [29-32]. Most of these studies indicated that glycosylation plays an important role in
21 glycosyl hydrolase function, biomass conversion and is influenced by culture conditions.
22 Thus, information on PTMs of lignocellulolytic enzymes will make it possible to
23 interpret their biological role, their expression regulation, interaction with substrates and

1 their role in regulating metabolism with new insights. Therefore, the main aim of this
2 study was to explore the quantitative production of lignocellulolytic enzymes by *A.*
3 *fumigatus* LF9 in presence of different carbon sources by adopting isobaric tag for
4 relative and absolute quantitation (iTRAQ)-based quantitative proteomics using liquid
5 chromatography tandem mass spectrometry (LC–MS/MS). Furthermore, this study
6 focused on the characterization of deamidation of quantified enzymes.

8 MATERIALS AND METHODS

9 Microorganism Isolation, Cultivation and Secretome Extraction

10 The fungal strains were isolated from biofertilizer compost from Singapore. The
11 procedure adopted for the isolation and identification of strain has been detailed in our
12 earlier publication that investigated abundances of lignocellulolytic enzymes at different
13 temperatures using saw dust as a substrate [33]. The culture of *A. fumigatus* LF9 was
14 prepared by using potato dextrose broth composed of 20 g L⁻¹ D-glucose and 4 g L⁻¹
15 potato extract (Sigma, St. Louis, MO, USA) at 40° C. Cell biomass was collected by
16 centrifugation at 4° C(Beckman Coulter, Brea, CA, USA), washed with sterilized Milli-Q
17 water and inoculated in flask containing medium with the following composition:
18 ammonium sulfate 3.1 g L⁻¹ , sodium chloride 1.5 g L⁻¹ , dipotassium phosphate 1.2 g
19 L⁻¹, monopotassium phosphate 0.9 g L⁻¹, magnesium sulfate 0.3 g L⁻¹ and 5.0 g L⁻¹
20 glucose. The isolate was grown at 50° C, and the biomass was collected by centrifugation
21 and inoculated into test flasks containing medium as stated above but with 5 g L⁻¹
22 glucose, cellulose, xylan and starch was used as a carbon source. The test flasks were
23 incubated at 50° C at 100 rpm and growth was monitored by analyzing total protein

content to determine the harvesting period in mid exponential phase (4 days). For proteomic analysis of the secretome from glucose, cellulose, xylan and starch condition, three flasks for each carbon source was used. The secretome was collected by centrifugation at 7500 g at 4 °C (Beckman Coulter, Brea, CA, USA) for 7 min, mycelium was removed and supernatant (secretome) was filtered through 0.25 µm filters. The supernatant (10–15 mL from ~200 mL) was retained and stored at –80°C for the enzyme assay. The remaining secretome (~180 mL) was concentrated by lyophilization. Proteins were precipitated by ice-cold acetone and protein content was determined by the Bradford method [34]. Biosafety guidelines were strictly followed while working with this fungal isolate.

Protein Separation, Protein Digestion, and Peptide Extraction

The protein digestion, peptide extraction, and LC-MS/MS analysis were performed as described previously [8, 10, 35]. Briefly, proteins (200µg) from each test sample were separated on 10% SDS-PAGE at 100 V, and protein bands were visualized by staining with Coomassie Brilliant blue G-250. Each sample lane was sliced and separately washed with 75% acetonitrile (ACN) containing Triethylammonium bicarbonate buffer (TEAB, 25mM), and then destained using TEAB (25 mM) alone and TEAB with 50% ACN. Following destaining, gel pieces were reduced with Tris 2-carboxyethyl phosphine hydrochloride (5 mM) and alkylated with methyl methanethiosulfonate (10 mM). The gel pieces were then washed with TEAB (25 mM) alone and then dehydrated using 100% ACN. Protein digestion was performed by subjecting dehydrated gel pieces to sequencing grade modified trypsin (Promega,

Madison, WI) at 37 °C for 12 h. The peptides were extracted using 50% ACN, 5% acetic acid. Tryptic peptides derived from each lane were combined separately and concentrated using a concentrator (Eppendorf AG, Hamburg, Germany) for further iTRAQ labeling.

iTRAQ Labeling, LC-MS/MS Analysis and data analysis

The iTRAQ labeling of peptides from test samples from different carbon sources were performed using a 4-plex iTRAQ reagent multiplex kit (Applied Biosystems, Foster City, CA) according to the manufacturer's protocol. The peptides from each test condition were labeled with respective isobaric tags, incubated for 2 h, quenched with water, combined together and vacuum-centrifuged to dryness. The main advantage of iTRAQ technique is the multiplex comparison of four to eight different conditions in single mass spectrometry-based experiment. The iTRAQ labeling for different conditions were assigned as follows- 113: *A. fumigatus* LF9 glucose (control); 114: *A. fumigatus* LF9 cellulose; 115: *A. fumigatus* LF9 xylan; 116: *A. fumigatus* LF9 starch. The iTRAQ ratios of 114/113 (cellulose/glucose), 115/113 (xylan/glucose), and 116/113 (starch/glucose) are presented in this study. The iTRAQ labeled peptides were reconstituted in buffer A (10 mM ammonium acetate, 85% acetonitrile, 0.1% formic acid) and fractionated using ERLIC column (200 × 4.6 mm, 5µm particle size, 300 Å pore size) by a HPLC system (Shimadzu, Japan) at flow rate of 1.0 mL min⁻¹ using our previously optimized protocol [36]. The HPLC chromatograms were recorded at 280 nm, and fractions were collected using an automated fraction collector and vacuum dried. The vacuum dried peptides were reconstituted in 3% ACN with 0.1% formic acid for LC-MS/MS analysis. For proteomic analysis of secretome, the experimental design included

1 three flaks for each carbon source. Set 1 sample was analyzed in duplicate (technical
2 replicates). The samples of set 2 and set 3 were prepared as stated above, analyzed
3 individually and mean data with standard deviation were reported.

4 The LC-MS/MS analysis of labeled peptides was performed using a QStar Elite
5 mass spectrometer (Applied Biosystems/MDS Sciex) coupled with online microflow
6 HPLC system (Shimadzu). The labeled peptides were separated on a home-packed
7 nanobored C18 column with a picofrit nanospray tip (New Objectives, Wubrun, MA)
8 coupled to the LC-MS/MS system at a constant flow rate of 300 nl min⁻¹. Mobile phase
9 A (0.1% formic acid in H₂O) and mobile phase B (0.1% formic acid in 80% ACN) were
10 used to establish the 90 min gradient comprising 15 min of 12–23% B, 55 min 23-40% B,
11 5 min 40-65% B, 3 min 65-90% B, maintained at 90% B for 2 min and finally re-
12 equilibration at 12% B for 10 min. The mass spectrometer was set in positive ion mode
13 using Analyst QS 2.0 software (Applied Biosystems) and data was acquired with a
14 selected mass range of 300 –1600m/z. The peptides with +2 and above were selected for
15 MS/MS. The three most abundant peptides above a five-count threshold were selected for
16 MS/MS, and dynamically excluded for 30 s with ±30 mDa mass tolerance. Smart
17 information-dependent acquisition (IDA) was activated with automatic collision energy
18 and automatic MS/MS accumulation. The fragment intensity multiplier was set to 20 and
19 maximum accumulation time was 2 s. The peak areas of the iTRAQ reporter ions reflect
20 the relative abundance of the proteins in the samples.

21 The data acquisition was performed with Analyst QS 2.0 software (Applied
22 Biosystems/MDS SCIEX). The peak list generation, protein identification, and peptide
23 quantification were performed using ProteinPilotTM software 3.0 (revision number

1 114732; Applied Biosystems, Foster City, CA). The Paragon algorithm in the
2 ProteinPilotTM software was used for the peptide identification. The data was further
3 processed by Pro Group algorithm where isoform-specific quantification was adopted to
4 trace the differences between expressions of various isoforms. The defined parameters
5 were as follows: (i) Sample Type, iTRAQ 4-plex (Peptide Labeled); (ii) Cysteine
6 alkylation, MMTS; (iii) Digestion, Trypsin; (iv) Instrument, QSTAR Elite ESI; (v)
7 Special factors, None; (vi) Species, None; (vii) Specify Processing, Quantitate; (viii) ID
8 Focus, biological modifications, amino acid substitutions; (ix) Database, concatenated
9 JGI downloaded from (<http://www.aspgd.org/>, [http://genome.jgi.doe.gov/Aspfu1/](http://genome.jgi.doe.gov/Aspfu1/Aspfu1.home.html)
10 [Aspfu1.home.html](http://genome.jgi.doe.gov/Aspfu1/Aspfu1.home.html)); (x) Search effort, thorough. The default precursors and fragment
11 mass tolerances for QStar Elite MS instrument were adopted by the software.
12 ProteinPilotTM software automatically determined various PTMs based on the HUPO-PSI
13 Modification Nomenclature for Mass Spectrometry standard (<http://www.unimod.org>).
14 The peptides for iTRAQ quantitation were automatically selected by Pro Group
15 algorithm (at least one peptide with 99% confidence) for calculations of the reporter peak
16 area, p-value etc. The resulting data set was auto bias-corrected to get rid of any
17 variations imparted due to the unequal mixing during combining different labeled
18 samples. A strict cut-off of unused ProteinScore ≥ 2 , which corresponds to a confidence
19 limit of 99%, and at least two peptides with 95% confidence were considered for protein
20 identifications. False discovery rate (FDR) was calculated using in-house developed
21 program as $FDR = 2 \times M_d / (M_d + M_t)$, where M_d represents the number of decoy
22 matches, and M_t is the number of target matches. Proteins identified with $\leq 1\%$ FDR was
23 further analyzed in this study.

Zymogram Development and Enzyme Assays

The zymographic analysis of test samples was performed using 10% SDS-PAGE gel that contained 1% carboxymethyl cellulose as described earlier [13]. In brief, SDS-PAGE gel was loaded with 10 µg of test protein and electrophoresis was performed at 90 V for 120 min. The gels were washed overnight with 50 mM phosphate buffer (pH 7.2±0.2) containing 25% isopropanol, incubated at 50 °C for 1 h and stained with 0.1% (w/v) Congo red for 15 min. The gel was washed with 1 M NaCl until bands became visible. The enzyme activities were determined on the supernatant samples using a protocol reported by Ghose [37]. The endo-1,4-β-glucanase, exo-1,4-β-glucanase, and β-glucosidase were assayed using carboxymethyl cellulose (CMC), Avicel, and salicin (β-salicyl alcohol glucoside) as substrates respectively. Xylan (from birchwood (Sigma, CAS: 9014-63-5) were used as substrate for xylanase activity. Reducing sugars were determined by using 3,5-dinitrosalicylic acid (DNSA)[38] One international unit (IU) of enzyme activity was defined as the amount of enzyme releasing 1 µmol of reducing sugar per minute.

Results

This work focused on the secretome of *A. fumigatus* LF9 during cellulose, xylan and starch degradation following iTRAQ labeling. With a 99% confidence level (cut off of unused protein score ≥ 2.0) and false discovery rate of ≤ 1.0 , 429±26 proteins were identified. These proteins were sorted using N-terminal Sec-dependant secretion signal using SignalP 3.0 (<http://www.cbs.dtu.dk/services/SignalP/>) [39]. This study identified

1 some intracellular proteins, suggesting cell lysis or secretion through unknown
2 mechanism. Based on the iTRAQ ratio distribution frequency and percentage variation of
3 reporter ion intensities, the significance threshold for proteins with iTRAQ ratios ≤ 0.7
4 was set for down- and ≥ 1.4 for up-regulation. To gain biological insights into regulated
5 proteins, the iTRAQ quantified proteins were functionally classified. The functional
6 classification revealed 32.9, 0.9, 16.2, 8.1, 11.7, 7.6 and 22.5% cellulases and glycoside
7 hydrolases, amylases, hemicellulases, lignin degrading enzymes, peptidases and proteases,
8 chitinases, lipases and phosphatases, and proteins with unknown function respectively
9 (Fig 1).

11 **Abundance of cellulases and glycoside hydrolases**

12 This iTRAQ based study quantified 73 proteins belonging to cellulase, glycoside
13 hydrolase and amylases. Their comparative iTRAQ ratios were listed in Table1. The
14 analysis of these proteins showed up-regulation of 27 (36.9%), 41 (56.1%) and 30 (41.1%)
15 of 73 proteins when cellulose, xylan and starch were used as substrates respectively.
16 When cellulose was used as a major substrate, enzymes like GH12, GH5 GH45
17 endoglucanase, (Afu7g06150, Afu5g01830, Afu7g06740), GH7 cellobiohydrolase D
18 (Afu6g07070), GH3 beta-D-glucoside glucohydrolase (Afu7g06140), GH7 1,4-beta-D-
19 glucan-cellobiohydrolyase (Afu6g11610), GH3 beta-glucosidase (Afu5g07080,
20 Afu1g17410), GH30 beta-1,6-glucanase (Afu8g07120), GH15glucan1,4-alpha-
21 glucosidase (Afu2g00690) were significantly up-regulated with iTRAQ ratios ranging
22 between 7.05 and 1.81. Thus, analysis of the secretome of *A. fumigatus* LF9 revealed a
23 significant up-regulation of cellulolytic enzymes. Some enzymes like GH16 endo-1,3(4)-

1 beta-glucanase (Afu1g05290), GH16 cell wall glucanase (Afu6g08510) etc. were down-
2 regulated when the strain was cultivated in cellulosic medium.

3 When cellulose was used as a main carbon source, iTRAQ ratios of GH12, GH5,
4 and GH45 endoglucanase were 7.05, 6.57 and 3.96 respectively. Enzymes GH7
5 cellobiohydrolase D (Afu6g07070) and 1-4-beta-D-glucan-cellobiohydrolase
6 (Afu6g11610) were quantified with iTRAQ ratios of 3.97 and 3.29. GH3 beta-
7 glucosidases (Afu7g06140, Afu5g07080 and Afu1g17410) were quantified with iTRAQ
8 ratios between 3.34 and 1.81. Taken together, endoglucanases, exocellobiohydrolases and
9 beta-glucosidases were significantly up-regulated when cellulose was used as a substrate,
10 suggesting the cellulose degradation potential of this isolated strain. Cluster analysis of
11 the proteins having cellulolytic activities showed five different clusters (Fig 2). Proteins
12 clustered under cluster C3 and C5 were up-regulated when cellulose was used as a
13 substrate.

14 When xylan was used as the main carbon source, enzymes like GH61, GH5
15 endoglucanase (Afu4g07850, Afu2g09520, Afu6g11600), GH7 endoglucanase
16 (Afu6g01800), GH61 endo-1,4-beta-glucanase (Afu3g03870), GH5 glucanase,
17 (Afu7g05610), GH71 alpha-1,3-glucanase (Afu2g03980), GH62, GH31 GH43, GH10,
18 GH2, GH76 glycosyl hydrolase (Afu2g00920, Afu1g03140, Afu3g01660, Afu3g15210,
19 Afu2g14520, Afu3g00340), GH3, GH1, GH3, GH35 beta glucosidase, (Afu6g08700,
20 Afu5g07080, Afu1g14710, Afu1g05770, Afu1g17410, Afu1g16700), GH7 1,4-beta-D-
21 glucan-cellobiohydrolase (Afu6g11610), cellobiose dehydrogenase (Afu2g17620),
22 GH72 1,3-beta-glucanosyltransferase (Afu2g05340) were significantly up-regulated, with
23 iTRAQ ratios in the range of 6.86 to 2.02. Interestingly, these enzymes except GH7 1,4-

1 beta-D-glucan-cellobiohydrolase (Afu6g11610) were not up-regulated when cellulose
2 was used as a substrate. This suggests that xylan can stimulate the expression of cellulases
3 and revealed the abundance level of enzymes as a function of substrate used. The cluster
4 analysis of cellulolytic proteins as presented in Fig 2 suggested that the proteins clustered
5 under cluster C2 were significantly up-regulated when xylan was used as a substrate.
6 Enzymes like GH31, GH1 glucosidase, GH81 endo-1,3-beta-glucanase, GH76 glycosyl
7 hydrolase were up-regulated when starch was used as a substrate.

8

9 **Abundance of Hemicellulases**

10 A particularly noteworthy observation is that 26 of the 36 iTRAQ quantified (or
11 72.2%) hemicellulases were significantly up-regulated when xylan was used as substrate.
12 Fascinatingly, enzymes GH10 endo-1,4-beta-xylanase (Afu4g09480, Afu6g13610),
13 GH11 endo-1,4-beta-xylanase (Afu3g00320, Afu3g00470), GH43 xylosidase
14 (Afu8g04710), acetyl xylan esterase (Afu3g00420), GH43 xylosidase/arabinosidase,
15 (Afu2g04480), alpha-L-arabinofuranosidase (Afu6g14620, Afu2g15160), alpha-
16 galactosidase (Afu4g03580) GH3 beta-xylosidase (Afu3g02090), cell wall
17 galactomannoprotein (Afu4g03240) GH62 arabinofuranosidase (Afu2g12770) were the
18 most significantly up-regulated with iTRAQ ratios between 7.06 and 3.36, when xylan
19 was used as a substrate (Table S1, supplementary). Again, with iTRAQ ratios ranging
20 between 2.7 to 1.9, enzymes including carboxylesterase, (Afu3g09070), GH92 alpha-1,2-
21 mannosidase (Afu6g13760), GH35 beta-galactosidase, (Afu1g14170) GH43 beta-
22 xylosidase (Afu2g00930), GH2 beta-mannosidase (Afu6g08840), alpha-galactosidase
23 (Afu6g02560), GH68 arabinosidase (Afu2g00650) were up-regulated in xylan containing
24 culture condition (Fig 3 and 4). Thus, based on the abundance level and regulation of

1 hemicellulases, *A. fumigatus* LF9 has the potential to hydrolyze hemicellulose completely.
2 The hierarchical clustering of iTRAQ quantified hemicellulolytic proteins demonstrated
3 four different clusters. The proteins of cluster H3 were significantly expressed in the
4 xylan containing culture media (Fig 5).

5 Hemicellulases such as GH28 extracellular polygalacturonase, (Afu1g17220),
6 alpha-L-arabinofuranosidase (Afu6g14620), GH11 endo-1,4-beta-xylanase (Afu3g00320),
7 GH10 extracellular endo-1,4-beta-xylanase, (Afu4g09480), acetyl xylan esterase
8 (Afu3g00420), GH11 endo-1,4-beta-xylanase, (Afu3g00470) were up-regulated when
9 cellulose was used as a substrate. The iTRAQ ratios of these enzymes were in the range
10 of 8.53 to 1.51. Enzymes such as GH2 beta-mannosidase (Afu6g08840), GH47 alpha-
11 mannosidase (Afu1g14560), GH92 alpha-1,2-mannosidase (Afu5g10520, Afu6g13760),
12 GH43 xylosidase/arabinosidase (Afu2g04480), alpha-galactosidase (Afu6g02560),
13 esterase (Afu5g09860), GH68 arabinosidase (Afu2g00650) were up-regulated when
14 starch was used as a substrate. Thus, cellulose and starch can provoke expression of
15 specific hemicellulases. Proteins clustered under cluster H1 were up-regulated when
16 starch was used as a main carbon source.

17

18 **Abundance of lignin degrading and other enzymes**

19 Enzymes involved in lignin degradation such as laccase, isoamyl alcohol oxidase,
20 alcohol dehydrogenase, bifunctional catalase-peroxidase and enzymes belonging to
21 oxidoreductase family were identified in this study (Table S2, supplementary). Amylases
22 that hydrolyze starch and pectate lyases that carried out eliminative cleavage of pectate
23 yielding oligosaccharides with 4-deoxy- α -D-mann-4-enuronosyl groups at their non-

1 reducing ends were also iTRAQ quantified and regulated. The iTRAQ quantified
2 amylases (Afu3g00900, Afu2g00710) were up-regulated only in starch containing medium and
3 not in other substrate conditions, and their up-regulation suggested that amylases of *A. fumigatus*
4 could have potential to degrade raw starch (Table S3, supplementary). The cellulolytic proteins
5 of cluster C1 were up regulated in starch containing culture condition. Enzymes like GH5
6 endoglucanase (Afu2g09520) and diene lactone hydrolase (Afu6g12740) were
7 significantly up-regulated when starch was main carbon source. This indicates that starch
8 can stimulate expression of some specific cellulolytic enzymes.

9 Peptidases and protease like carboxypeptidase S1, aspartyl aminopeptidase,
10 dipeptidyl peptidase, alkaline serine protease and aspartyl aminopeptidase were up-
11 regulated when cellulose and xylan were used as a substrate. While enzymes like
12 aminopeptidase, metallopeptidase, inorganic diphosphatase, aminopeptidase B, dipeptidyl
13 peptidase III, dipeptidyl peptidase, aminopeptidase P, aspartyl aminopeptidase etc. were
14 up-regulated when starch was used as a substrate. The enzymes like tripeptidyl peptidase
15 A, aspartic endopeptidase, carboxypeptidase S1, tripeptidyl peptidase A, serine
16 carboxypeptidase were down-regulated when starch were used as a substrate.

18 **Zymograph and Enzyme Activities**

19 As seen from zymogram (Fig 6B), some intense bands of proteins with
20 cellulolytic activity were noted when SDS-PAGE containing CMC was used for cellulase
21 activity. The prominent bands such as B3 were excised, tryptic digested and further
22 analyzed by LC-MS/MS. The LC-MS/MS analysis confirmed abundance of
23 endoglucanase (Afu7g06150), cell wall glucanase (Afu1g16190) endo-1,4-beta-xylanase
24 (Afu3g00320) and several hypothetical proteins (Table S4, Supplementary). Enzyme

activities under different substrates suggested high endoglucanase and exoglucanase activity in cellulosic conditions, followed by xylan and then starch, while xylanase activity was higher in xylan culture condition (Fig 6A).

Post translational modifications

This study presents the deamidation of lignocellulolytic enzymes of thermophilic fungal strain *A. fumigatus* LF9. Based on the functional groups, 15% of identified cellulases and glycosyl hydrolases, 6% hemicellulases, 22% lignin degrading enzymes, 3% peptidases and proteases, and 2% of proteins with unknown function was deamidated. Based on the number of available deamidation sites in native protein and identified deamidation sites, the degree of deamidation varies from protein to protein. The list of modified enzymes, peptide sequences and sites of the modifications were listed in Table 2. The MS and MS/MS spectra, peptide sequence and modification sites are provided as supplementary information (MS and MS/MS spectra, and fragment ions assignment). The enzymes like endoglucanase (Afu7g06150|), glucanase (Afu7g05610|), exo-beta-1,3-glucanase (Afu6g13270|), endo-1,3-beta-glucanase (Afu1g04260|), glycosyl hydrolase (Afu2g00920|), cellobiohydrolase D (Afu6g07070|), endo-1,4-beta-xylanase (Afu3g00320|), alpha-mannosidase (Afu1g14560|), glucan 1,4-alpha-glucosidase (Afu2g00690|), beta-xylosidase (Afu3g02090|), isoamyl alcohol oxidase (Afu6g03620|), oxidoreductase (Afu2g14480|) etc. were detected with deamidation (Table 2). To elucidate the detailed biochemical functions of these PTMs and their impact on the fungal physiology and synthesis of these lignocellulolytic enzymes, a more detailed study with utilizing a high-throughput approach integrating proteomic data, metabolic

network analysis, protein structure analysis, and targeted genome engineering is required.

Discussion

Rapid and efficient enzymatic conversion of lignocellulosic biomass into fermentable sugars is a major challenge for the sustainable production of bio-chemicals and biofuels. The thermal pretreatment of biomass, followed by enzymatic treatment using thermotolerant enzymes could speed up the biomass hydrolysis rate and improve process efficiency. Hence, several researchers attempted to improve the thermotolerance of cellulases and hemicellulases by adopting techniques such as introduction of disulfide bridges, ionic pairs, tightly bound water molecules, protein packing and oligomerization [27]. Despite the improvement in the thermoactivity of these engineered cellulases and hemicellulases, their instability remains a major issue. These issues can be resolved by searching for naturally occurring thermostable plant biomass degrading fungi. The main advantages of thermotolerant enzymes from thermostable fungi for bioprocesses are their prolonged stability at room temperature, a reduced risk of process contamination, and an increased tolerance to organic solvents [40]. Our previous study [33] isolated *A. fumigatus* LF9 from compost and profiled its secretome at different temperatures between 20 and 70 °C using saw dust as a substrate, and adopting solid state fermentation technique. The enzyme activities and protein abundances as determined by exponentially modified protein abundance index (emPAI) indicated the maximum activities at the range of temperature 40–60°C, demonstrating the thermophilic nature of this strain.

1 Characterization of the thermostability of the secretome suggested the thermotolerant
2 nature of biomass degrading enzymes [33].

3 As of now, little work has been done on the proteomic analysis of the secretome
4 of *A. fumigatus* and their applications in recycling of lignocellulosic biomass. *A.*
5 *fumigatus* is pathogenic species but its potential biomass hydrolyzing enzymes could be
6 useful if these enzymes are heterologously expressed in a GRAS (generally recognized as
7 safe) organism and then used. Several strains of *Aspergillus* species are extremely useful
8 in biotechnological industry and biomass recycling. Hence, the genomes of *A. clavatus*
9 [41], *A. flavus* [42], *A.fumigatus* [43], *A. nidulans* [44], *A. niger* [45], *A. oryzae* [46], *A.*
10 *terreus* etc. have been sequenced. Some *Aspergillus* species have been exploited for
11 extracellular glucohydrolases secretion, which includes enzymes such as cellulases,
12 hemicellulases, pectinases, and β -glucosidase [33, 47-51]. In this study, we profiled
13 global view of the secretome of *A. fumigatus* LF9 in cellulose, xylan and starch
14 containing culture media using iTRAQ-based quantitative proteomics. The efficient
15 hydrolysis of lignocellulosic biomass requires several types of glycohydrolases, including
16 different cellulases and hemicellulases. iTRAQ quantitative proteomics quantified 65
17 cellulolytic proteins, comparable to 65 proteins expressed in *T. reesei* but were higher
18 than 52 in *P. chrysosporium* when cultivated in cellulosic media [12, 35]. The
19 comparisons of cellulolytic enzymes encoded in the genome of *T. reesei* and *A. fumigatus*
20 support a higher cellulolytic activity in *A. fumigatus* with 19 cellulases genes as
21 compared to 10 in *T. ressei* [19]. Although the *P. chrysosporium* genome encodes 27
22 genes for cellulolytic enzymes [19] but, gene expression depends on carbon source and
23 several other factors [9, 13]. The significant up-regulation of endoglucanases (GH12,

GH5 and GH45), cellobiohydrolase (GH7), beta-glucosidase (GH3) suggested potential candidates for lignocellulosic bioenergy. Two-dimensional gel and MALDI-TOF-MS analysis of the secretome of *A. oryzae* with wheat bran as a major carbon source demonstrated the abundance of 43 and 37 cellulolytic proteins in solid states and submerged cultures [52]. Further classification of these iTRAQ quantified 65 cellulolytic proteins revealed twenty endo-1-3(4)-beta-glucanase, six exo-glucanase, sixteen different glycosyl hydrolases, three alpha-glucosidases, six beta glucosidases, one beta-1-6-glucanase, one cellobiose dehydrogenase, and seven glucanotransferases.

Numerous processes during substrate utilization are governed not only by the relative abundance of proteins but also by their regulation, localization, protein-protein interactions and post translational modifications (PTMs). Glycosylation of secreted enzymes is one of the most frequent and abundant enzyme modifications and is well studied in *Aspergillus* sp., *T. reesei* and *Phanerochaete chrysosporium* [29, 32, 53]. According to Robinson [54, 55], protein deamidation serve as molecular clocks and plays a major role in protein turnover, organism development and aging. However, no literature exists on the presence and extent of fungal protein deamidation.

The impact of protein deamidation on their function and biomass hydrolysis potential requires more research that can integrate proteomic data, metabolic network analysis, protein structure analysis, and targeted genome engineering. It is essential to know which enzymes are deamidated and this study reports the deamidated enzymes. The three-dimensional structure of cellobiohydrolase I resolved by Divne et al [56] reveals a 40 angstrom long active site tunnel that plays a major role in its interaction with solid cellulose. Using this three dimensional structure, we found that deamidation sites

are located away from the enzyme active sites. Recently, it has been documented that low molecular weight monosaccharides or disaccharides produced by hydrolysis of cellulose and hemicellulose triggers enzyme production [57, 58]. However, in some cases, these compounds need to be further modified by glycosylation prior to functioning as physiological inducers [58]. Thus, PTMs play important role in biomass hydrolysing enzyme production and more focused research needs to be done. Also, the impact of deamidation and its role in secreted enzymes has not been documented.

Xylan belongs to a group of hemicelluloses and are abundant in hardwood and agricultural residues [59]. Xylan is almost as ubiquitous as cellulose in plant cell wall, so its fermentation is essential for the economic conversion of lignocellulose to ethanol [60, 61]. Interestingly, 60% (39 of 65) of iTRAQ quantified cellulolytic enzymes were up-regulated when xylan was used as a substrate. The enzymes belonging to GH families including GH7, GH62, GH43, GH5, GH31, GH71, GH1, GH61, GH5 etc. were up regulated and their iTRAQ ratios ranged between 6.12 ± 0.74 and 2.68 ± 1.76 , when xylan was used as a substrate. Surprisingly, these cellulolytic enzymes were not up regulated when cellulose was used as a substrate. This indicates that cellulose and xylan induces the expression of some specific cellulolytic enzymes. The enhanced production of cellulases in xylan culture condition could be explained by the currently documented role of transcriptional activator (XlnR) in coordinating and controlling expression of both xylanolytic and cellulolytic enzymes [62, 63]. Current literature on cellulase and hemicellulases [57, 58] indicated that monosaccharides or disaccharides produced by hydrolysis of cellulose and hemicellulose triggers enzyme production. The saccharides derived from cellulosic biomass act as an inducer for cellulase and hemicellulases

1 production, e.g. D-xylose in *A. niger* [64], cellobiose in *A. oryzae* [65] and sophorose in
2 *A. terreus* and *T. Reesei* [66]. When xylan was used as a substrate, 70.58% of iTRAQ
3 quantified hemicellulases were up regulated. This significant up-regulation of
4 hemicellulases could also be due to the thermotolerant nature of the enzymes secreted by
5 this thermostable fungal strain because *Aspergilli* endoxylanases demonstrates the
6 maximal activities at a range of 42°C–60°C, and a pH range of 4.0–7.0 [33, 67].
7 Xylanolytic enzymes have great biotechnological potential and are exploited for various
8 industrial processes, including feed, food, pulp, and others [68-70]. Interestingly,
9 enzymes like endo-1,4-beta-xylanase, beta-xylosidase, glucuronidase,
10 arabinofuranosidase and acetyl xylan esterases that act on different heretopolymers were
11 expressed and significantly up-regulated when xylan was used as a substrate, indicating
12 their potential use in lignocellulosic biorefineries. The most significantly up-regulated
13 enzymes likes GH10 endo-1,4-beta-xylanase, GH11 endo-1,4-beta-xylanase, GH43
14 xylosidase, acetyl xylan esterase, GH43 xylosidase/arabinosidase, alpha-L-
15 arabinofuranosidase, alpha-galactosidase, GH62 arabinofuranosidase could be potential
16 enzymes for biomass degradation and hence more detailed and focused research is
17 required on these enzymes. Beta-mannosidase and alpha-1, 2-mannosidase, which plays
18 major roles in glucomannan degradation were also up-regulated when xylan was used as
19 a substrate. Thus, taken together, xylanolytic enzymes from *A. fumigatus* LF9 could be
20 exploited for several industrial processes including biomass hydrolysis.

21 Although GH10 and GH11 xylanases differ in produced xylo-oligosaccharides
22 [67, 71], both were quantified with iTRAQ ratios of 7.58 and 6.9, showing a significantly
23 higher abundance of these enzymes produced by *A. fumigatus* LF9. With 36

1 hemicellulase genes, *A. fumigatus* has largest set of hemicellulases , as compared to *T.*
2 *reesei* (16 genes), *Neurospora crassa* (19 genes) and *P. chrysosporium* (19 genes).
3 Interestingly, when starch was used as a substrate, hemicellulolytic enzymes belonging to
4 GH families GH2, GH92, GH43, GH47, GH43 etc. were up-regulated. Enzymes such as
5 GH11 endo-1,4-beta-xylanase (Afu3g00320|), GH11 endo-1,4-beta-xylanase
6 (Afu3g00470|), GH3 beta-xylosidase (Afu3g02090|), alpha-L-arabinofuranosidase
7 (Afu6g14620|) and alpha-galactosidase (Afu4g03580|) that were quantified with iTRAQ
8 ratios of 6.90 ± 0.51 , 4.49 ± 0.98 , 5.03 ± 0.25 , 6.18 ± 0.54 and 6.17 ± 0.72 respectively, were
9 identified to be deamidated. The role of deamidation in secretory enzymes and their
10 impact on biomass hydrolysis requires further detailed research.

11 In addition to cellulose and hemicellulose degrading enzymes, this study also
12 identified and quantified lignin degrading enzymes like laccase, isoamyl alcohol oxidase,
13 aryl-alcohol dehydrogenase, glutathione reductase, Cu-Zn superoxide dismutase and
14 proteins of oxidoreductase family. Although laccase was iTRAQ quantified, but it was
15 found to be down-regulated, indicating that the substrates like cellulose, xylan or starch
16 could not induce its production. Further analysis of PTMs revealed the deamidation of
17 laccase (Afu1g15670|), isoamyl alcohol oxidase (Afu6g03620|) and oxidoreductase
18 (Afu2g14480|). Peptidases such as aspartyl aminopeptidase (Afu5g03990) and secreted
19 dipeptidyl peptidase (Afu2g09030) were up-regulated in all studied substrate conditions.
20 In xylan containing substrate conditions, 66.6% of identified peptidases and proteases
21 were up-regulated while aspartic endopeptidase (Afu5g13300), serine carboxypeptidase
22 (Afu6g00310) and tripeptidyl-peptidase (Afu4g03490) were down-regulated. Of these
23 iTRAQ quantified peptidases, aminopeptidase (Afu4g09030|) and secreted dipeptidyl

1 peptidase (Afu2g09030) were identified and found to be deamidated. In starch
2 containing culture conditions, peptidases such as aminopeptidase (Afu4g09030),
3 metallopeptidase (Afu7g05930), dipeptidyl peptidase III (Afu4g06140), aspartyl
4 aminopeptidase (Afu5g03990) etc. were up-regulated. Of the 50 iTRAQ quantified
5 proteins with unknown functions, 11, 26 and 23 proteins were up-regulated when
6 cellulose, xylan and starch respectively were used as substrates, suggesting these
7 enzymes contribute in biomass degradation. This study showed a significant up-
8 regulation of cellulases, hemicellulases and lignin degrading enzymes of *A. fumigatus*
9 LF9 in cellulose and xylan culture conditions. A previous study[33] of this strain (*A.*
10 *fumigatus* LF9) in presence of saw dust as a substrate identified significant up-
11 regulation of lignocellulolytic enzymes. Thus, *A. fumigatus* LF9 has the potential to
12 degrade lignocellulosic biomass and can be exploited in lignocellulosic biorefinery.

14 **Conclusions**

15 Thermostable enzymes from thermophilic microbes have potential applications in
16 the biotechnological industries and biomass hydrolysis. Proteomic characterization of the
17 secretome of thermophilic *A. fumigatus* LF9 in cellulose, xylan and starch culture
18 conditions suggested the up-regulation of a large number of cellulases, hemicellulases,
19 and lignin degrading proteins, demonstrating biomass degradation potential of this strain.
20 The comparison of quantitative iTRAQ ratios suggests that cellulose and xylan causes
21 expression and up-regulation of specific cellulases and hemicellulases. Post translational
22 modifications analysis of the secretome of *A. fumigatus* revealed deamidation of key
23 cellulases, hemicellulases and lignin degrading enzymes. The knowledge of deamidated

enzymes along with the specific sites of modifications could be crucial information for further functional studies of these enzymes in *A. fumigatus*.

Conflict of interest

None for this study.

ACKNOWLEDGMENTS

This work is in part supported by grants from the Singapore Ministry of Education (Tier 1: RGT15/13) and NTU-NHG Ageing Research Grant (ARG/14017) and NTU iFood (Grant #: S006).

References

- [1] Pérez J, Muñoz-Dorado J, De La Rubia T, Martínez J. Biodegradation and biological treatments of cellulose, hemicellulose and lignin: An overview. *Int Microbiol* 2002;5:53-63.
- [2] Walton NJ, Mayer MJ, Narbad A. Vanillin. *Phytochemistry* 2003;63:505-15.
- [3] Porro D, Bianchi MM, Brambilla L, Menghini R, Bolzani D, Carrera V, et al. Replacement of a metabolic pathway for large-scale production of lactic acid from engineered yeasts. *Appl Environ Microbiol* 1999;65:4211-15.
- [4] Renewable fuels association: Accelerating industry innovation. Washington: 2012. Ethanol industry outlook; 2012.
- [5] Sun H, Zhao P, Ge X, Xia Y, Hao Z, Liu J, et al. Recent advances in microbial raw starch degrading enzymes. *Appl Biochem Biotechnol* 2010;160:988-1003.
- [6] Chandel AK, Singh OV, Rao LV. Biotechnological applications of hemicellulosic derived sugars: State-of-the-art. In: Singh ov, harvey sp (eds) sustainable biotechnology: Renewable resources and new perspectives. Springer, netherland, pp 63–81. 2010.
- [7] Robertson GH, Wong DW, Lee CC, Wagschal K, Smith MR, Orts WJ. Native or raw starch digestion: A key step in energy efficient biorefining of grain. *J Agric Food Chem* 2006;54:353-65.
- [8] Adav SS, Ng CS, Arulmani M, Sze SK. Quantitative itraq secretome analysis of cellulolytic *Thermobifida fusca*. *J Proteome Res* 2010;9:3016-24.
- [9] Adav SS, Ravindran A, Chao LT, Tan L, Singh S, Sze SK. Proteomic analysis of pH and strains dependent protein secretion of *Trichoderma reesei*. *J Proteome Res* 2011;10:4579-96.
- [10] Adav SS, Chao LT, Sze SK. Protein abundance in multiplexed samples (PAMUS) for quantitation of *Trichoderma reesei* secretome. *J Proteomics* 2013;83:180-96.

- [11] Gómez-Mendoza DP, Junqueira M, Do Vale LHF, Domont GB, Ferreira Filho EX, Sousa MVD, et al. Secretomic survey of trichoderma harzianum grown on plant biomass substrates. J Proteome Res 2014;13:1810-22.
- [12] Manavalan A, Adav SS, Sze SK. iTRAQ-based quantitative secretome analysis of phanerochaete chrysosporium. J Proteomics 2011;75:642-54.
- [13] Adav SS, Ravindran A, Sze SK. Quantitative proteomic analysis of lignocellulolytic enzymes by *Phanerochaete chrysosporium* on different lignocellulosic biomass. J Proteomics 2012;75:1493-504.
- [14] Shah P, Atwood JA, Orlando R, Mubarek HE, Podila GK, Davis MR. Comparative proteomic analysis of botrytis cinerea secretome. J Proteome Res 2009;8:1123-30.
- [15] González-Fernández R, Aloria K, Valero-Galván J, Redondo I, Arizmendi JM, Jorrín-Novo JV. Proteomic analysis of mycelium and secretome of different *Botrytis cinerea* wild-type strains. J Proteomics 2014;97:195-221.
- [16] Martinez D, Challacombe J, Morgenstern I, Hibbett D, Schmoll M, Kubicek CP, et al. Genome, transcriptome, and secretome analysis of wood decay fungus *Postia placenta* supports unique mechanisms of lignocellulose conversion. Proc Natl Acad Sci U S A 2009;106:1954-59.
- [17] Ferreira de Oliveira JMP, van Passel MWJ, Schaap PJ, de Graaff LH. Proteomic analysis of the secretory response of *Aspergillus niger* to D-maltose and D-xylose. PLoS ONE 2011;6.
- [18] Ji XL, Yan M, Yang ZD, Li AF, Kong LR. Shotgun analysis of the secretome of *Fusarium graminearum*. Indian J Microbiol 2013;53:400-09.
- [19] Martinez D, Berka RM, Henrissat B, Saloheimo M, Arvas M, Baker SE, et al. Genome sequencing and analysis of the biomass-degrading fungus trichoderma reesei (syn. Hypocrea jecorina). Nat Biotechnol 2008;26:553-60.
- [20] Liu D, Li J, Zhao S, Zhang R, Wang M, Miao Y, et al. Secretome diversity and quantitative analysis of cellulolytic *Aspergillus fumigatus* z5 in the presence of different carbon sources. Biotechnol for Biofuels 2013;6.
- [21] Bansal N, Tewari R, Soni R, Soni SK. Production of cellulases from *Aspergillus niger* ns-2 in solid state fermentation on agricultural and kitchen waste residues. Waste Manage (Oxford) 2012;32:1341-46.
- [22] Narra M, Dixit G, Divecha J, Madamwar D, Shah AR. Production of cellulases by solid state fermentation with *Aspergillus terreus* and enzymatic hydrolysis of mild alkali-treated rice straw. Bioresour Technol 2012;121:355-61.
- [23] Cunha FM, Esperança MN, Zangirolami TC, Badino AC, Farinas CS. Sequential solid-state and submerged cultivation of aspergillus niger on sugarcane bagasse for the production of cellulase. Bioresour Technol 2012;112:270-74.
- [24] Baldrian P, Valášková V. Degradation of cellulose by Basidiomycetous fungi. FEMS Microbiol Rev 2008;32:501-21.
- [25] Louise Glass N, Schmoll M, Cate JHD, Coradetti S. Plant cell wall deconstruction by Ascomycete fungi. Annual Review of Microbiology 2013. p. 477-98.
- [26] Van Den Brink J, De Vries RP. Fungal enzyme sets for plant polysaccharide degradation. Appl Microbiol Biotechnol 2011;91:1477-92.
- [27] Paës G, O'Donohue MJ. Engineering increased thermostability in the thermostable GH-11 xylanase from *Thermobacillus xylanilyticus*. J Biotechnol 2006;125:338-50.

1 [28] Hamada JS. Deamidation of food proteins to improve functionality. Crit Rev Food
2 Sci Nutr 1994;34:283-92.

3 [29] Hui JPM, White TC, Thibault P. Identification of glycan structure and glycosylation
4 sites in cellobiohydrolase ii and endoglucanases I and II from *Trichoderma reesei*.
5 Glycobiology 2002;12:837-49.

6 [30] Stals I, Sandra K, Geysens S, Contreras R, Van Beeumen J, Claeysens M. Factors
7 influencing glycosylation of *Trichoderma reesei* cellulases. I: Postsecretorial changes of
8 the o- and n-glycosylation pattern of cel7a. Glycobiology 2004;14:713-24.

9 [31] Sandra K, Van Beeumen J, Stals I, Sandra P, Claeysens M, Devreese B.
10 Characterization of cellobiohydrolase i n-glycans and differentiation of their
11 phosphorylated isomers by capillary electrophoresis - Q-trap mass spectrometry. Anal
12 Chem 2004;76:5878-86.

13 [32] Adav SS, Ravindran A, Sze SK. Study of *Phanerochaete chrysosporium* secretome
14 revealed protein glycosylation as a substrate-dependent post-translational modification. J
15 Proteome Res 2014;13:4272-80.

16 [33] Adav SS, Ravindran A, Sze SK. Proteomic analysis of temperature dependent
17 extracellular proteins from *Aspergillus fumigatus* grown under solid-state culture
18 condition. J Proteome Res 2013;12:2715-31.

19 [34] Hammond JB, Kruger NJ. The bradford method for protein quantitation. Methods
20 Mol Biol 1988;3:25-32.

21 [35] Adav SS, Chao LT, Sze SK. Quantitative secretomic analysis of *Trichoderma reesei*
22 strains reveals enzymatic composition for lignocellulosic biomass degradation. Mol Cell
23 Proteomics 2012;11.

24 [36] Hao P, Guo T, Li X, Adav SS, Yang J, Wei M, et al. Novel application of
25 electrostatic repulsion-hydrophilic interaction chromatography (ERLIC) in shotgun
26 proteomics: Comprehensive profiling of rat kidney proteome. J Proteome Res
27 2010;9:3520-6.

28 [37] Ghose TK. Measurement of cellulase activities. Pure Appl Chem 1987;59:257-68.

29 [38] Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugar.
30 Anal Chem 1959;31:426-28.

31 [39] Bendtsen JD, Nielsen H, Von Heijne G, Brunak S. Improved prediction of signal
32 peptides: Signalp 3.0. J Mol Biol 2004;340:783-95.

33 [40] Vieille C, Zeikus GJ. Hyperthermophilic enzymes: Sources, uses, and molecular
34 mechanisms for thermostability. Microbiol Mol Biol Rev 2001;65:1-43.

35 [41] Wortman JR, Fedorova N, Crabtree J, Joardar V, Maiti R, Haas BJ, et al. Whole
36 genome comparison of the a. Fumigatus family. Med Mycol 2006;44:3-7.

37 [42] Payne GA, Nierman WC, Wortman JR, Pritchard BL, Brown D, Dean RA, et al.
38 Whole genome comparison of *Aspergillus flavus* and *A. Oryzae*. Med Mycol 2006;44:9-
39 11.

40 [43] Nierman WC, Pain A, Anderson MJ, Wortman JR, Kim HS, Arroyo J, et al.
41 Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus*
42 *fumigatus*. Nature 2005;438:1151-56.

43 [44] Galagan JE, Calvo SE, Cuomo C, Ma LJ, Wortman JR, Batzoglou S, et al.
44 Sequencing of *aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A.*
45 *oryzae*. Nature 2005;438:1105-15.

- [45] Pel HJ, De Winde JH, Archer DB, Dyer PS, Hofmann G, Schaap PJ, et al. Genome sequencing and analysis of the versatile cell factory *Aspergillus niger* cbs 513.88. *Nat Biotechnol* 2007;25:221-31.
- [46] Machida M, Asai K, Sano M, Tanaka T, Kumagai T, Terai G, et al. Genome sequencing and analysis of *Aspergillus oryzae*. *Nature* 2005;438:1157-61.
- [47] De Nicolás-Santiago S, Regalado-González C, García-Almendárez B, Fernández FJ, Téllez-Jurado A, Huerta-Ochoa S. Physiological, morphological, and mannanase production studies on *Aspergillus niger* uam-gs1 mutants. *Electron J Biotechnol* 2006;9:50-60.
- [48] Ward OP, Qin WM, Dhanjoon J, Ye J, Singh A. Physiology and biotechnology of *Aspergillus*. *Advances in Applied Microbiology* 2005. p. 1-75.
- [49] Adav SS, Li AA, Manavalan A, Punt P, Sze SK. Quantitative itraq secretome analysis of *Aspergillus niger* reveals novel hydrolytic enzymes. *J Proteome Res* 2010;9:3932-40.
- [50] Adav SS, Sze SK. Effective biocatalysts for lignocellulosic biofuel by proteomics study of fungal secretome. In *Bioenergy Systems, Biological Sources and Environmental Impact* 2013. p. 59-76.
- [51] Adav SS, Sze SK. Fungal secretome for biorefinery: Recent advances in proteomic technology. *Mass Spectrometry Letters* 2013;4:1-9.
- [52] Oda K, Kakizono D, Yamada O, Iefuji H, Akita O, Iwashita K. Proteomic analysis of extracellular proteins from *Aspergillus oryzae* grown under submerged and solid-state culture conditions. *Appl Environ Microbiol* 2006;72:3448-57.
- [53] Wang L, Aryal UK, Dai Z, Mason AC, Monroe ME, Tian ZX, et al. Mapping n-linked glycosylation sites in the secretome and whole cells of *Aspergillus niger* using hydrazide chemistry and mass spectrometry. *J Proteome Res* 2012;11:143-56.
- [54] Robinson NE, Robinson AB. Deamidation of human proteins. *Proc Natl Acad Sci U S A* 2001;98:12409-13.
- [55] Robinson NE, Robinson AB. Amide molecular clocks in drosophila proteins: Potential regulators of aging and other processes. *Mech Ageing Dev* 2004;125:259-67.
- [56] Divne C, Ståhlberg J, Reinikainen T, Ruohonen L, Pettersson G, Knowles JKC, et al. The three-dimensional crystal structure of the catalytic core of cellobiohydrolase I from *Trichoderma reesei*. *Science* 1994;265:524-28.
- [57] Tani S, Kawaguchi T, Kobayashi T. Complex regulation of hydrolytic enzyme genes for cellulosic biomass degradation in filamentous fungi. *Appl Microbiol Biotechnol* 2014;98:4829-37.
- [58] Kubicek CP, Messner R, Gruber F, Mach RL, Kubicek-Pranz EM. The trichoderma cellulase regulatory puzzle: From the interior life of a secretory fungus. *Enzyme Microb Technol* 1993;15:90-99.
- [59] Jeffries TW. Engineering yeasts for xylose metabolism. *Curr Opin Biotechnol* 2006;17:320-26.
- [60] Saha BC. A-L-arabinofuranosidases: Biochemistry, molecular biology and application in biotechnology. *Biotechnol Adv* 2000;18:403-23.
- [61] Saha BC, Dien BS, Bothast RJ. Fuel ethanol production from corn fiber current status and technical prospects. *Appl Biochem Biotechnol- Part A Enzyme Engineering and Biotechnology* 1998;70-72:115-25.

- [62] Klaubauf S, Narang HM, Post H, Zhou M, Brunner K, Mach-Aigner AR, et al. Similar is not the same: Differences in the function of the (hemi-)cellulolytic regulator xlnr (xlr1/xyr1) in filamentous fungi. *Fungal Genet Biol* 2014;72:73-81.
- [63] Aro N, Pakula T, Penttilä M. Transcriptional regulation of plant cell wall degradation by filamentous fungi. *FEMS Microbiol Rev* 2005;29:719-39.
- [64] De Souza WR, De Gouveia PF, Savoldi M, Malavazi I, De Souza Bernardes LA, Goldman MHS, et al. Transcriptome analysis of *Aspergillus niger* grown on sugarcane bagasse. *Biotechnology for Biofuels* 2011;4.
- [65] Marui J, Kitamoto N, Kato M, Kobayashi T, Tsukagoshi N. Transcriptional activator, aoxlnr, mediates cellulose-inductive expression of the xylanolytic and cellulolytic genes in *aspergillus oryzae*. *FEBS Lett* 2002;528:279-82.
- [66] Hrmova M, Petrakova E, Biely P. Induction of cellulose- and xylan-degrading enzyme systems in *Aspergillus terreus* by homo- and heterodisaccharides composed of glucose and xylose. *J Gen Microbiol* 1991;137:541-47.
- [67] Teixeira RSS, Siqueira FG, De Souza MV, Filho EXF, Da Silva Bon EP. Purification and characterization studies of a thermostable β -xylanase from *Aspergillus awamori*. *J Ind Microbiol Biotechnol* 2010;37:1041-51.
- [68] Akpınar O, Ak O, Kavas A, Bakır U, Yılmaz L. Enzymatic production of xylooligosaccharides from cotton stalks. *J Agric Food Chem* 2007;55:5544-51.
- [69] Akpınar O, Erdoğan K, Bostancı S. Enzymatic production of xylooligosaccharide from selected agricultural wastes. *Food Bioprod Process* 2009;87:145-51.
- [70] Twomey LN, Pluske JR, Rowe JB, Choct M, Brown W, McConnell MF, et al. The effects of increasing levels of soluble non-starch polysaccharides and inclusion of feed enzymes in dog diets on faecal quality and digestibility. *Anim Feed Sci Technol* 2003;108:71-82.
- [71] Takahashi Y, Kawabata H, Murakami S. Analysis of functional xylanases in xylan degradation by *Aspergillus niger* e-1 and characterization of the gh family 10 xylanase xynvii. *SpringerPlus* 2013;2:1-11.

1
2 Figure 1. Functional classification of iTRAQ quantified proteins. The secretome of *A.*
3 *fumigatus* LF9 in glucose, cellulose, xylan and starch containing culture media were
4 characterized for protein composition.

Fig 2. Hierarchical clustering of cellulolytic protein by *A. fumigatus* LF9, exhibiting their expressions when grown on cellulose, xylan and starch as the major carbon source. Up-regulated protein expression values are displayed in red, while the down-regulation values are in blue, and the intermediate values are in shades of red and blue.

Figure3. Comparative abundance of different hemicellulose degrading enzymes of *A. fumigatus* LF9 in different carbon sources (Afu5g10520: alpha-1,2-mannosidase, Afu1g14560: alpha-mannosidase, Afu1g10790: alpha-1,2-mannosidase, Afu6g08840: beta-mannosidase, Afu6g13760: alpha-1,2-mannosidase, Afu3g00320: endo-1,4-beta-xylanase, Afu6g13610: endo-1,4-beta-xylanase, Afu4g09480: extracellular endo-1,4-beta-xylanase, Afu3g00470: endo-1,4-beta-xylanase, Afu2g04480: xylosidase, Afu8g04710: xylosidase, Afu3g02090: beta-xylosidase, Afu2g00930: beta-xylosidase, Afu6g14620: alpha-L-arabinofuranosidase, Afu1g09900: alpha-L-arabinofuranosidase, Afu2g15160: alpha-L-arabinofuranosidase A, Afu2g12770: arabinofuranosidase)

Figure 4. Comparative abundance of different hemicellulose degrading enzymes of *A. fumigatus* LF9 in different carbon sources (Afu4g03580: alpha-galactosidase, Afu1g14170: beta-galactosidase, Afu6g02560: alpha-galactosidase, Afu5g02130: alpha-galactosidase, Afu1g06910: arabinogalactan endo-1,4-beta-galactosidase, Afu1g16700: beta galactosidase, Afu5g09860: esterase, Afu3g00420: acetyl xylan esterase, Afu3g09070: carboxylesterase, Afu1g03170: esterase, Afu3g14510: rhamnogalacturonan acetylerase, Afu1g17220: extracellular polygalacturonase, Afu8g01970: extracellular endo-polygalacturonase, Afu7g06410: exo-polygalacturonase, Afu4g03240: cell wall galactomannoprotein Mp1, Afu6g00770: extracellular arabinanase, Afu2g15430: L-xylulose reductase, Afu1g04820: D-xylulose reductase (Xyl1), Afu2g00650: arabinosidase, Afu5g00910: mannose binding protein, Afu2g14750: endo-arabinase)

Fig 5. Hierarchical clustering of hemicellulolytic proteins by *A. fumigatus* LF9 exhibiting their expressions when grown on cellulose, xylan and starch as the major carbon source. Up-regulated protein expression values are displayed in red, while the down-regulation values are in blue, and the intermediate values are in shades of red and blue.

Fig 6. Activities of enzymes in the secretome of *A. fumigatus* LF9 when grown in glucose, cellulose, xylan and starch culture media. A) Activities were determined by spectrometric methods; B) Zymogram showing cellulolytic activity in the secretome of *A. fumigatus*. SDS zymogram was performed using carboxymethyl cellulose as a substrate, stained with Congo red and destained by 1 M NaCl. The cellulase activity can be seen as a white band in zymogram analysis. The bands were further analyzed by LC-MS/MS. (M: Protein Marker, X: xylan, C: cellulose, S: starch and G: glucose)

Table 1. Comparative quantitative expression of cellulases, glycosyl hydrolases and amylases by *Aspergillus fumigatus* LF9 in different carbon sources

Unused Prot-Score	Total	% Cov	Accession No.	Name	Family	Peptide (95%)	Cellulose	Xylan	Starch
68.95±14.02	68.95±14.02	72.51±4.60	Afu2g00690	glucan 1,4-alpha-glucosidase, putative	GH15	123±45	1.84±0.03	1.95±0.15	1.39±0.11
65.34±8.92	7.15±2.28	33.21±18.28	Afu6g12740	dienelactone hydrolase family protein	NA	4±2	0.73±0.02	1.53±1.54	12.26±5.65
45.17±6.20	45.17±6.20	61.21±5.99	Afu3g00270	cell wall glucanase, putative	NA	77±25	0.52±0.03	0.95±0.11	0.98±0.07
43.24±6.92	43.24±6.92	54.88±4.44	Afu2g05340	1,3-beta-glucanosyltransferase, putative	GH72	48±14	0.82±0.06	2.02±0.58	2.14±0.64
42.74±6.79	42.74±6.79	64.97±4.24	Afu2g03980	alpha-1,3-glucanase, putative	GH71	59±18	0.97±0.05	4.18±0.41	1.97±0.22
39.27±7.95	39.27±7.95	49.06±6.31	Afu4g10150	alpha-glucosidase	GH31	33±10	1.34±0.22	1.06±0.23	3.14±0.70
37.28±6.38	37.28±6.38	40.86±3.81	Afu6g13270	exo-beta-1,3-glucanase, putative	PL3	49±15	1.21±0.03	1.37±0.15	0.83±0.09
35.43±3.57	35.43±3.57	81.84±2.93	Afu7g06150	endoglucanase, putative	GH12	80±26	7.05±1.00	0.53±0.05	0.50±0.04
34.88±7.94	34.88±7.94	63.98±7.70	Afu6g06770	enolase	NA	37±13	1.06±0.08	0.59±0.06	2.21±0.28
33.20±9.34	33.20±9.34	60.84±5.67	Afu2g01170	1,3-beta-glucanosyltransferase Gel1	GH72	34±11	0.59±0.01	0.68±0.07	1.31±0.12
32.96±6.95	32.96±6.95	56.80±8.47	Afu7g06140	beta-D-glucoside glucohydrolase	GH3	27±7	3.34±0.17	0.31±0.05	0.61±0.05
31.56±6.89	31.56±6.89	55.80±9.55	Afu7g05610	glucanase, putative	GH5	20±6	0.89±0.04	4.89±0.38	0.73±0.08
30.92±4.85	30.92±4.85	37.96±2.86	Afu1g04260	endo-1,3-beta-glucanase Eng11	GH81	35±13	1.31±0.11	1.91±0.32	2.58±0.23
24.03±3.79	24.03±3.79	55.44±2.04	Afu1g16190	cell wall glucanase Crf1	GH16	27±9	1.07±0.04	1.81±0.20	2.39±0.30
22.43±2.35	22.43±2.35	61.64±4.95	Afu1g11460	1,3-beta-glucanosyltransferase Bgt1	NA	23±6	0.56±0.02	0.86±0.10	1.44±0.10
22.20±6.11	22.20±6.11	57.38±9.21	Afu8g07120	beta-1,6-glucanase, putative	GH30	27±8	1.85±0.33	0.85±0.20	1.49±0.11
21.83±5.72	21.83±5.72	37.83±5.08	Afu6g11610	1,4-beta-D-glucan-cellobiohydrolase, putative	GH7	24±10	3.29±0.06	3.60±0.55	0.54±0.05
21.32±3.08	21.32±3.08	48.66±8.98	Afu3g00900	alpha-amylase AmyA	NA	17±4	0.76±0.04	1.10±0.16	1.32±0.21
21.04±3.91	21.04±3.91	62.89±13.36	Afu5g09230	transaldolase	NA	17±4	0.96±0.03	1.70±0.21	4.02±0.64
20.32±4.13	20.32±4.13	37.59±6.08	Afu5g14380	alpha-glucuronidase	GH67	11±3	1.31±0.08	5.16±0.56	0.75±0.06
20.26±4.16	20.26±4.16	41.49±4.26	Afu2g17620	cellobiose dehydrogenase	CDH	16±5	1.03±0.08	3.34±0.32	0.84±0.08
19.10±2.19	19.10±2.19	35.35±2.20	Afu3g02280	alpha,alpha-trehalose glucohydrolase, putative	GH65	13±3	1.50±0.04	1.76±0.18	1.13±0.10
18.87±4.26	18.88±4.26	31.86±3.83	Afu2g00710	alpha-amylase, putative	NA	15±5	1.06±0.08	0.57±0.05	1.25±0.23

16.88±6.66	16.88±6.66	31.39±5.29	Afu1g14450	exo-beta-1,3-glucanase Exg0	PL3	10±3	0.77±0.13	0.98±0.10	1.24±0.15
16.29±3.14	16.29±3.14	45.99±7.16	Afu8g06880	pectin methylesterase	NA	24±7	0.31±0.02	0.27±0.04	0.16±0.03
15.13±2.52	15.13±2.52	50.95±7.37	Afu2g00920	glycosyl hydrolase, putative	GH62	14±4	0.86±0.17	5.85±0.71	0.89±0.11
15.01±3.19	15.01±3.19	44.78±5.06	Afu5g01830	extracellular endoglucanase, putative	GH5	10±3	6.57±0.25	0.64±0.07	0.49±0.04
13.75±2.26	14.12±2.27	42.65±7.79	Afu2g12850	1,3-beta-glucanosyltransferase Gel3	GH72	10±1	0.72±0.07	0.94±0.15	0.88±0.11
12.76±2.08	12.76±2.08	25.34±3.13	Afu2g03120	cell wall glucanase	NA	21±7	0.66±0.05	1.14±0.20	3.18±0.95
12.68±5.49	12.68±5.49	36.09±11.81	Afu3g00310	extracellular phytase, putative	NA	9±4	0.52±0.02	0.24±0.05	0.79±0.10
12.62±3.06	12.62±3.06	39.69±9.30	Afu2g00820	extracellular GDSL-like lipase/acylhydrolase,	NA	7±2	1.20±0.06	3.94±1.17	0.71±0.11
12.58±3.13	12.58±3.13	37.20±10.89	Afu5g07080	beta-glucosidase	GH3	8±2	1.92±0.34	2.22±0.61	0.81±0.13
12.47±3.77	12.47±3.77	33.70±3.85	Afu4g03350	exo-beta-1,3-glucanase, putative	PL3	8±3	0.52±0.04	0.40±0.04	0.61±0.04
12.29±1.13	13.29±2.08	28.84±3.97	Afu7g01240	phytase, putative	NA	16±4	0.97±0.04	0.60±0.09	2.37±0.51
11.97±3.09	11.97±3.09	57.31±8.74	Afu2g02100	dihydrolipoamide dehydrogenase	NA	5±2	1.00±0.08	0.78±0.06	3.01±0.76
11.64±1.41	11.64±1.41	31.80±2.38	Afu3g00340	glycosyl hydrolase, putative	GH76	9±1	0.74±0.04	2.07±0.28	2.27±0.25
11.44±2.89	11.44±2.89	38.28±3.55	Afu6g11600	endoglucanase, putative	GH5	6±2	1.31±0.09	3.11±0.51	0.82±0.14
10.05±1.34	10.05±1.34	50.78±5.43	Afu2g00760	pectate lyase A	NA	6±1	0.88±0.03	0.99±0.14	1.40±0.18
9.01±1.74	9.01±1.74	29.97±6.23	Afu1g03140	glycosyl hydrolase, family 31	GH31	5±1	0.89±0.10	4.61±1.18	0.86±0.11
8.96±2.75	8.96±2.75	38.07±2.89	Afu3g01660	putative hydrolase	GH43	9±3	0.89±0.10	5.33±0.80	1.01±0.21
8.66±2.65	8.66±2.65	29.37±14.92	Afu6g11390	1,3-beta-glucanosyltransferase Gel2	GH72	7±2	0.84±0.08	1.68±0.20	2.08±0.22
8.31±3.51	8.32±3.52	32.44±6.73	Afu6g08510	cell wall glucanase, putative	GH16	5±2	0.58±0.11	0.33±0.08	0.48±0.02
8.06±1.22	8.06±1.22	34.06±3.10	Afu1g05290	endo-1,3(4)-beta-glucanase	GH16	7±2	0.33±0.01	0.19±0.02	0.34±0.03
8.05±3.80	8.05±3.80	33.87±11.43	Afu5g08780	cell wall glucanase, putative	NA	5±2	0.94±0.13	1.42±0.22	1.32±0.25
8.02±1.43	8.02±1.43	38.30±8.20	Afu3g15210	Glycosyl hydrolase family 10	GH10	5±1	1.26±0.06	3.53±2.45	0.66±0.11
8.02±2.47	8.02±2.47	22.72±4.31	Afu3g03080	endo-1,3(4)-beta-glucanase	GH16	5±2	1.15±0.12	1.68±0.20	1.57±0.21
7.79±2.23	7.79±2.23	31.50±8.42	Afu3g01910	cellobiohydrolase, putative	GH6	5±2	1.01±0.36	0.76±0.43	0.43±0.20
7.71±2.04	5.39±2.03	22.91±11.56	Afu1g16700	beta galactosidase, putative	GH35	2±1	2.92±0.11	2.48±1.58	1.30±0.99
7.70±1.53	7.70±1.53	19.00±4.00	Afu8g02130	1,3-beta-glucanosyltransferase, putative	GH72	5±1	0.65±0.04	1.70±0.33	0.94±0.10
7.58±2.85	5.98±2.21	22.37±10.56	Afu4g07850	endoglucanase, putative	GH61	3±2	1.07±0.13	2.69±0.18	0.45±0.01
7.52±1.55	7.52±1.55	40.48±7.94	Afu6g08700	beta glucosidase, putative	GH3	4±1	1.30±0.13	4.10±0.47	1.10±0.12

7.42±3.13	7.42±3.13	24.67±5.11	Afu6g01800	endoglucanase, putative	GH7	5±2	1.05±0.06	6.12±0.74	0.78±0.16
7.30±2.20	7.30±2.20	31.50±7.51	Afu1g16420	GPI anchored protein, putative	NA	4±1	1.53±0.01	0.73±0.11	1.41±0.14
7.05±1.05	7.05±1.05	33.05±4.93	Afu1g07130	dienelactone hydrolase family protein	NA	5±2	0.78±0.03	0.57±0.15	1.13±0.12
6.43±1.42	6.43±1.42	29.06±3.11	Afu1g17410	beta-glucosidase, putative	GH3	3±1	1.81±0.19	2.88±0.47	0.97±0.08
6.35±1.60	6.35±1.60	26.61±8.92	Afu2g14520	hydrolase, putative	GH2	3±1	1.04±0.17	2.76±0.53	1.13±0.17
6.02±0.02	6.02±0.02	32.76±3.92	Afu5g14510	beta-lactamase	NA	3±0	0.62±0.05	1.43±0.24	0.50±0.04
6.00±0.84	7.51±1.41	24.01±3.58	Afu6g07070	cellobiohydrolase D	GH7	4±1	3.97±1.15	0.83±0.21	0.42±0.03
5.52±1.69	5.52±1.69	29.20±8.72	Afu5g08270	HAD superfamily hydrolase, putative	NA	3±1	1.58±0.68	1.68±0.39	1.66±0.41
4.27±0.86	4.27±0.86	38.39±4.65	Afu2g09350	endo-beta-1,6-glucanase, putative	GH5	2±0	0.82±0.06	0.68±0.11	1.08±0.14
4.27±1.10	4.27±1.10	27.09±4.95	Afu1g05770	beta-glucosidase, putative	GH3	2±1	1.36±0.21	2.99±1.23	0.61±0.15
4.09±1.54	4.09±1.54	24.72±5.62	Afu1g16250	alpha-glucosidase B	GH31	2±1	0.94±0.07	0.52±0.07	1.34±0.22
3.79±2.37	3.79±2.37	22.28±8.03	Afu4g00390	glycosyl hydrolase, putative	GH2	2±1	0.99±0.08	1.85±0.23	1.34±0.22
3.55±1.23	3.55±1.23	27.77±4.61	Afu2g00220	hydrolase	NA	2±1	1.32±0.22	1.05±0.15	0.71±0.15
2.80±1.45	3.60±1.40	11.03±4.45	Afu1g14710	beta-glucosidase 1	GH1	2±0	0.86±0.00	3.97±2.86	3.29±2.01
2.72±1.77	3.19±0.75	7.83±5.83	Afu3g03870	endo-1,4-beta-glucanase, putative	GH61	2±0	1.21±0.00	6.85±0.00	0.74±0.00
2.69±0.76	2.84±0.23	5.73±4.56	Afu2g00510	cellulose-binding GDSL lipase/acylhydrolase,	NA	2±0	0.91±0.08	1.68±1.11	0.45±0.00
2.67±1.57	3.34±0.95	13.29±3.13	Afu1g11400	alpha/beta hydrolase, putative	GH4	2±0	1.33±0.00	0.46±0.00	0.47±0.00
2.67±0.97	2.00±0.00	12.64±7.54	Afu5g00670	glycosyl hydrolase family 35, putative	GH35	2±0	1.10±0.21	1.53±0.11	0.66±0.13
2.6±1.27	2.69±0.95	4.89±2.75	Afu2g09520	endoglucanase, putative	GH5	2±0	0.77±0.00	2.68±1.76	16.64±6.12
2.49±1.23	3.38±1.63	12.23±4.72	Afu3g07520	exo-beta-1,3-glucanase, putative	PL3	2±1	0.89±0.00	1.12±0.00	0.52±0.00
2.27±0.16	2.09±0.00	11.46±3.25	Afu1g01730	glycosyl hydrolase, putative	GH76	3±0		0.92±0.00	1.54±0.00
2.00±0.00	2.28±1.00	12.17±4.16	Afu2g14630	cell wall glycosyl hydrolase family 88	NA	3±0	0.37±0.00	0.17±0.00	0.23±0.00
2.00±0.00	2.00±0.00	17.19±0.96	Afu7g06740	endoglucanase, putative	GH45	2±0	3.96±0.36	0.88±0.09	1.00±0.21
2.00±0.00	2.51±0.50	16.76±4.76	Afu3g03950	endoglucanase, putative	GH61	2±0	1.06±0.00	1.29±0.00	1.49±0.00

NA: Data not available

PL: Pectate lyase

Table 2. Deamidated enzymes and their modification with peptide sequence showing modification site (MS spectrum showing the precursor ion mass, and MS/MS spectrum showing peptide fragmentation and site of modification are provided in supplementary information)

Accession No.	Name	Peptide sequence and modification site
Afu7g06150	Endoglucanase, putative (235 aa)	LVSQISSIPTTVQWSYDNTN*TR
Afu7g06150	Endoglucanase, putative (235 aa)	N*HGYPASSQYLINMQFGTEPFTGGPATLR
Afu7g06150	Endoglucanase, putative (235 aa)	VSQWTASVN
Afu7g05610	Glucanase, putative (471 aa)	YSLLEQ*LR
Afu1g05290	Endo-1,3(4)-beta-glucanase, putative (357 aa)	NTAQDTGLIFAN*GAVYMGVDHTNVAGSSGR
Afu6g07070	Cellobiohydrolase D (453 aa)	GSVVIDAN*WR
Afu6g13270	Exo-beta-1,3-glucanase, putative (805 aa)	ATPGFTGLGLIDGDQ*YQGDGNQGWISTNVFFR
Afu1g04260	Endo-1,3-beta-glucanase Eng11 (975 aa)	GN*LMLGILR
Afu8g07120	Beta-1,6-glucanase, putative (489 aa)	FMPPGAIVLNGSGSYTYSGGGGIQSVASLNPDGTR
Afu8g07120	Beta-1,6-glucanase, putative (489 aa)	WSGNAPSQ*SVTTWVLPSA
Afu3g00270	Cell wall glucanase, putative (447 aa)	ANAGIGTN*PDEIVSYIN*EVR
Afu2g03980	Alpha-1,3-glucanase, putative (497 aa)	WQEILN*LGPR
Afu2g00690	Glucan 1,4-alpha-glucosidase, putative (632 aa)	ATALIDFGNWLIDNGYSSYAVN*NIWPIVR
Afu2g00690	Glucan 1,4-alpha-glucosidase, putative (632 aa)	DLTWSYAAFLTAN*MR
Afu2g00690	Glucan 1,4-alpha-glucosidase, putative (632 aa)	SVYAIN*SGIPQGAASVAGR
Afu2g00920	Glycosyl hydrolase, putative (397 aa)	SQTTLNFPNGMGN*TVIALSDSNPNNLFEASNVR
Afu3g00320	Endo-1,4-beta-xylanase (XlnA), putative (229 aa)	GTVN*TDGGTYNIYTAVR
Afu3g00320	Endo-1,4-beta-xylanase (XlnA), putative (229 aa)	LGMNLGTHNYQIVATEGYQSSGSASITVY
Afu3g00320	Endo-1,4-beta-xylanase (XlnA), putative (229 aa)	TFTQ*YWSVR
Afu3g00470	Endo-1,4-beta-xylanase, putative (314 aa)	GTVTSDGSTYDIYEHQQ*VNQPSIVGTATFN*QYWSIR
Afu3g02090	Beta-xylosidase (772 aa)	HFAAYDLEDWN*GVVR
Afu1g10790	Alpha-1,2-mannosidase (840 aa)	SIN*GYPLPGGAFVR
Afu1g14560	Alpha-mannosidase (494 aa)	N*ATIVSQILDHIK
Afu1g14560	Alpha-mannosidase (494 aa)	N*GWGASAVDALSTAIVMR
Afu1g14560	Alpha-mannosidase (494 aa)	AGFYVQ*SGAYILRPEVIESFYAYR
Afu6g14620	Alpha-L-arabinofuranosidase (507 aa)	GANAAASGSLSTFYN*GAR
Afu4g03580	Alpha-galactosidase (892 aa)	IGN*DIIPAWR
Afu1g15670	Laccase (597 aa)	MILT*GQFPAPPLYVR
Afu6g03620	Isoamyl alcohol oxidase (572 aa)	HGLAADQVLEWEVIDGQ*GNFLVANR
Afu6g03620	Isoamyl alcohol oxidase (572 aa)	SVVQTN*NAELTAAYR

Afu2g14480	Oxidoreductase, FAD-binding, putative (474 aa)	GGANN*FGIVTNFIFR
Afu4g09030	Aminopeptidase (954 aa)	LTFSGILN*DNMAGFYR
Afu4g09030	Aminopeptidase (954 aa)	QGIDEN*TMLTER
Afu2g09030	Secreted dipeptidyl peptidase (722 aa)	LPVAEGLSLFN*VLQER
Afu4g11800	Alkaline serine protease Alp1 (404 aa)	ASFSNYGSVVDIFAPGQDILSAWIGSTTATN*TISGTSMATPH
Afu6g03150	Hypothetical protein (431 aa)	DQTQYLQ*MWGR
Afu6g0315	Hypothetical protein (431 aa)	DQTQ*YLQMWGR
Afu6g03150	Hypothetical protein (431 aa)	DQ*TQYLQMWGR
Afu3g02600	Hypothetical protein (281 aa)	AQAALGSN*ALVSSTVVAASR

Table S1.iTRAQ quantified hemicellulases from *A. fumigatus* LF9 when different substrates were used as a main carbon sources.

Unused	Total	%Cov	Accession No	Name	Family	Peptide (95%)	Cellulose	Xylan	Starch
81.23±14.68	81.23±14.68	85.24±6.27	Afu1g14560	alpha-mannosidase	GH47	170±49	0.67±0.02	0.69±0.06	1.56±0.15
68.00±12.26	68.00±12.26	80.81±5.88	Afu3g00320	endo-1,4-beta-xylanase	GH11	135±44	1.96±0.17	6.90±0.51	0.66±0.05
44.21±8.10	44.21±8.10	67.13±5.85	Afu6g13610	endo-1,4-beta-xylanase, putative	GH10	84±28	0.75±0.05	7.07±1.30	0.78±0.08
38.48±5.65	38.48±5.65	70.20±6.46	Afu3g02090	beta-xylosidase	GH3	39±11	1.03±0.08	5.03±0.25	0.80±0.06
30.92±7.41	30.92±7.41	78.16±8.06	Afu4g09480	extracellular endo-1,4-beta-xylanase, putative	GH10	29±10	1.77±0.59	7.58±1.24	0.87±0.10
28.76±7.42	28.76±7.42	48.12±8.38	Afu1g10790	alpha-1,2-mannosidase family protein, putative	GH92	19±7	0.69±0.03	0.72±0.05	1.00±0.05
25.71±5.00	25.71±5.00	49.95±5.47	Afu6g14620	alpha-L-arabinofuranosidase	NA	28±10	2.59±0.25	6.18±0.54	0.71±0.09
24.93±7.90	24.93±7.90	62.80±8.13	Afu6g00770	extracellular arabinanase, putative	GH43	41±16	0.72±0.07	0.53±0.09	0.86±0.16
20.32±2.51	20.32±2.51	40.24±8.27	Afu4g03580	alpha-galactosidase	NA	20±5	0.93±0.08	6.17±0.72	0.77±0.07
19.27±4.90	19.27±4.90	51.25±5.68	Afu5g10520	alpha-1,2-mannosidase family protein	GH92	12±4	0.99±0.08	1.24±0.13	1.90±0.28
18.81±4.50	18.81±4.50	35.45±6.98	Afu1g14170	beta-galactosidase, putative	GH35	12±3	1.32±0.03	2.17±0.43	1.00±0.13
18.58±4.39	18.58±4.39	53.82±11.16	Afu8g04710	xylosidase	GH43	14±4	0.85±0.04	6.33±0.90	0.86±0.07
18.41±1.71	18.41±1.71	68.05±6.85	Afu4g03240	cell wall galactomannoprotein Mpl	NA	22±5	1.22±0.18	5.02±1.14	1.39±0.21
15.10±2.38	15.11±2.38	51.56±8.20	Afu6g02560	alpha-galactosidase	NA	8±1	1.11±0.11	1.92±0.32	1.68±0.33
12.72±2.29	12.72±2.29	63.91±11.09	Afu2g15430	L-xylulose reductase	NA	15±4	1.01±0.07	0.73±0.10	2.07±0.47
12.31±2.21	12.31±2.21	36.62±5.69	Afu6g13760	alpha-1,2-mannosidase, putative subfamily	GH92	8±2	1.37±0.37	2.42±0.57	1.89±0.42
11.38±1.91	11.38±1.91	37.44±5.95	Afu5g02130	alpha-galactosidase	NA	8±3	0.54±0.02	0.28±0.03	0.43±0.05
11.18±4.43	11.18±4.43	51.75±11.91	Afu5g09860	esterase, putative	NA	6±3	1.13±0.09	1.06±0.24	1.55±0.51
10.40±2.90	10.40±2.90	40.15±7.09	Afu3g00420	acetyl xylan esterase	NA	8±2	1.51±0.22	7.06±2.01	0.91±0.14

10.02±3.16	10.42±3.54	44.43±9.34	Afu2g12770	arabinofuranosidase	GH62	6±2	1.08±0.10	3.36±0.18	0.81±0.06
9.18±2.66	11.33±2.71	36.74±5.92	Afu3g00470	endo-1,4-beta-xylanase, putative	GH11	10±4	1.51±0.20	4.49±0.98	0.81±0.14
8.77±3.15	8.77±3.15	36.06±9.26	Afu1g06910	arabinogalactan endo-1,4-beta-galactosidase,	GH53	5±2	0.50±0.08	0.91±0.06	0.88±0.10
8.70±3.25	8.70±3.25	29.35±3.63	Afu1g09900	alpha-L-arabinofuranosidase	NA	5±2	1.04±0.02	4.24±0.98	0.76±0.06
7.66±1.86	7.66±1.86	36.59±10.54	Afu2g15160	alpha-L-arabinofuranosidase A	NA	5±1	1.00±0.08	4.09±0.80	0.86±0.07
7.00±3.31	7.00±3.31	58.48±12.31	Afu3g14510	rhamnogalacturonan acetyltransferase, putative	NA	4±2	0.51±0.06	0.26±0.02	0.56±0.11
6.60±2.24	6.60±2.24	27.08±2.94	Afu6g08840	beta-mannosidase	GH2	4±2	1.12±0.16	2.07±0.20	2.16±0.31
5.98±2.68	5.98±2.68	20.23±4.42	Afu2g04480	xylosidase/arabinosidase, putative	GH43	5±2	0.99±0.08	6.28±1.19	1.77±0.37
5.02±1.01	5.02±1.01	53.33±16.36	Afu1g04820	D-xylose reductase	NA	3±1	1.07±0.09	3.49±0.85	1.01±0.03
4.23±1.41	4.23±1.41	24.14±8.65	Afu2g00930	Possible beta-xylosidase, family 43 of glycosyl	GH43	2±1	0.82±0.11	2.14±0.62	0.54±0.08
3.88±1.33	3.88±1.33	28.47±5.03	Afu2g00650	arabinosidase	GH43, GH32, GH62, GH68	2±1	0.76±0.05	1.91±0.42	1.49±0.18
3.51±1.66	3.51±1.66	27.58±6.78	Afu1g17220	extracellular polygalacturonase, putative	GH28	2±1	8.53±2.94	0.36±0.04	0.29±0.05
3.51±1.67	3.51±1.67	28.23±10.04	Afu8g01970	extracellular endo-polygalacturonase, putative	GH28	2±1	0.83±0.01	0.44±0.02	0.72±0.02
2.63±1.51	2.63±1.51	27.38±6.56	Afu3g09070	carboxylesterase, putative	NA	2±1	1.08±0.18	2.69±0.90	0.68±0.05

NA: data not available

PL: Pectase lyase

Table S2. Lignin degrading enzymes, peptidases and proteases by *A. fumigatus* LF9 when different substrates were used as a main carbon sources.

Unused	Total	%Cov	Accession No.	Name	Peptide (95%)	Cellulose	Xylan	Starch
Lignin degrading enzymes								
13.95±3.54	13.95±3.54	32.25±5.81	Afu1g01180	isoamyl alcohol oxidase	9±3	1.17±0.08	1.24±0.23	1.45±0.21
20.69±4.39	20.69±4.39	50.48±7.37	Afu6g03620	isoamyl alcohol oxidase	31±10	0.80±0.01	1.09±0.10	1.29±0.11
42.64±5.93	42.64±5.93	73.38±7.29	Afu8g01670	bifunctional catalase-peroxidase Cat2	34±7	1.15±0.06	1.21±0.10	3.00±0.42
15.02±2.71	15.02±2.71	89.08±8.53	Afu5g09240	Cu,Zn superoxide dismutase	14±4	0.86±0.04	4.08±0.42	0.74±0.18
14.22±2.29	14.22±2.29	53.17±12.29	Afu2g14480	oxidoreductase, FAD-binding, putative	14±4	0.95±0.01	1.03±0.16	0.91±0.10
11.79±0.84	11.79±0.84	72.50±4.60	Afu1g14550	Mn superoxide dismutase MnSOD	9±2	1.08±0.10	0.75±0.08	2.17±0.24
10.00±0.73	10.00±0.73	57.17±8.98	Afu4g09110	cytochrome c peroxidase, putative	7±1	1.03±0.02	4.38±0.63	2.56±0.37
29.63±3.42	29.63±3.42	65.29±10.53	Afu3g00840	FAD-dependent oxygenase, putative	44±10	0.63±0.05	0.97±0.13	0.81±0.10
11.10±2.09	11.10±2.09	60.37±4.97	Afu4g11580	Mn superoxide dismutase	6±1	1.02±0.05	0.89±0.17	2.91±0.91
3.77±1.77	3.77±1.77	29.57±8.05	Afu1g15670	laccase	2±1	0.43±0.08	0.52±0.04	0.33±0.02
233.67±64.97	2.87±1.61	10.17±11.20	Afu7g05070	FAD dependent oxidoreductase, putative	2±0	1.09±0.00	2.67±1.52	7.27±5.87
10.53±4.52	10.53±4.52	39.87±8.66	Afu3g02270	mycelial catalase Cat1	5±2	1.83±0.05	2.83±0.39	0.74±0.11
2.01±0.00	227.51±25.50	18.48±6.47	Afu4g00610	aryl-alcohol dehydrogenase, putative	2±0	1.04±0.00	0.82±0.00	1.42±0.00
243.25±76.43	2.27±0.34	16.18±20.02	Afu1g15960	glutathione reductase	3±0	0.75±0.00	0.55±0.00	0.88±0.00
3.14±0.90	3.14±0.90	37.64±8.99	Afu1g15610	oxidoreductase,	2±1	1.54±0.12	1.00±0.19	1.05±0.19
2.98±0.73	2.98±1.73	34.70±17.35	Afu7g01010	alcohol dehydrogenase, putative	2±1	1.29±0.10	1.10±0.04	1.14±0.02

2.02±0.02	2.02±0.02	20.05±4.31	Afu2g04710	cytochrome b5, putative	2±0	2.13±0.64	4.18±1.44	1.06±0.29
Peptidases and Proteases								
58.93±12.65	58.93±12.65	62.59±8.26	Afu4g09030	aminopeptidase	48±14	1.01±0.04	1.47±0.16	3.31±0.31
36.78±11.98	36.78±11.98	60.05±5.90	Afu7g05930	metallopeptidase MepB	23±9	1.03±0.06	1.44±0.18	2.64±0.19
30.41±6.90	30.41±6.90	54.61±7.48	Afu4g06140	dipeptidyl peptidase III	22±6	1.05±0.06	1.35±0.15	2.27±0.25
29.95±5.32	29.95±5.32	71.09±4.54	Afu4g11800	alkaline serine protease Alp1	92±36	0.75±0.02	2.21±0.37	1.28±0.21
25.83±6.31	25.83±6.31	59.86±8.32	Afu5g08050	aminopeptidase P, putative	16±7	0.97±0.06	1.70±0.17	2.08±0.33
20.45±3.67	20.45±3.67	44.43±5.79	Afu5g03990	aspartyl aminopeptidase	14±4	1.63±0.32	2.07±0.21	1.74±0.32
26.25±6.38	26.26±6.38	50.00±4.73	Afu5g04330	aminopeptidase B	15±4	0.96±0.04	1.74±0.18	2.42±0.21
18.67±4.24	18.67±4.24	45.51±5.96	Afu5g13300	aspartic endopeptidase Pep1	12±4	1.13±0.06	0.52±0.03	0.64±0.05
21.43±5.53	21.43±5.53	42.09±6.52	Afu2g09030	secreted dipeptidyl peptidase	15±5	1.49±0.11	2.14±0.33	2.25±0.48
12.78±2.27	12.78±2.27	34.99±8.53	Afu5g01200	carboxypeptidase S1, putative	9±3	0.67±0.05	1.12±0.09	0.59±0.06
11.21±2.27	11.21±2.27	37.50±4.89	Afu4g03490	tripeptidyl-peptidase	7±2	0.32±0.00	0.09±0.01	0.09±0.01
19.49±1.57	19.49±1.57	38.71±2.39	Afu3g07030	glutaminase A	15±3	2.18±0.08	2.93±0.26	0.73±0.06
14.58±3.91	14.58±3.91	68.81±7.17	Afu3g08380	inorganic diphosphatase, putative	9±3	1.40±0.23	0.99±0.18	2.57±0.79
13.03±3.69	13.03±3.69	50.56±5.14	Afu7g04760	gamma-glutamyltranspeptidase	8±3	0.97±0.04	1.95±0.18	1.30±0.17
8.28±2.53	8.28±2.53	43.38±8.56	Afu3g08930	tripeptidyl peptidase A	4±1	0.62±0.06	0.61±0.02	0.57±0.03
11.75±2.83	11.75±2.83	40.30±7.49	Afu8g07080	elastinolytic metalloproteinase Mep	10±4	0.93±0.32	1.62±0.41	0.81±0.05
5.04±1.76	5.04±1.76	35.10±8.17	Afu2g01250	serine peptidase, family S28, putative	4±2	0.83±0.08	2.04±0.50	1.10±0.20
93.84±5.98	4.99±0.70	25.84±15.28	Afu6g06800	peptidase, putative	3±1	1.49±0.00	0.70±0.09	0.64±0.05
4.65±0.95	4.65±0.95	29.32±3.40	Afu6g00310	serine carboxypeptidase	3±1	1.15±0.09	0.51±0.05	0.38±0.09

3.43±0.80	3.43±0.80	23.88±5.62	Afu3g02970	aspergillopepsin, putative	5±2	5.89±0.06	0.27±0.03	0.23±0.01
7.09±4.24	7.09±4.24	36.11±8.78	Afu3g08290	aspartyl aminopeptidase	5±3	1.37±0.13	2.94±1.78	0.82±0.22
4.70±1.65	4.70±1.65	25.79±3.82	Afu8g04730	oligopeptidase family protein	2±1	0.82±0.24	4.83±1.71	1.07±0.38
2.19±1.70	2.38±0.27	10.82±12.06	Afu5g09210	alkaline serine protease Alp2	2±0	1.44±0.00	1.44±0.00	0.93±0.00
2.02±0.00	2.21±1.99	3.79±1.76	Afu4g14000	tripeptidyl peptidase A	2±0	1.29±0.00	0.79±0.00	0.66±0.00
2.70±1.65	4.06±1.67	24.29±12.92	Afu6g10250	alkaline serine protease AorO, putative	2±1	1.09±0.03	1.68±0.58	1.18±0.66
2.00±0.00	2.00±0.00	20.90±3.92	Afu8g04120	carboxypeptidase S1, putative	2±1	4.51±0.29	1.74±0.18	0.49±0.04

Table S3. iTRAQ quantified amylases, proteins involved in cell morphogenesis and hypothetical protein in different substrate conditions.

Unused	Total	%Cov	Accession No.	Name	Peptide (95%)	Cellulose	Xylan	Starch
Amylases								
18.87±4.26	18.88±4.26	31.86±3.83	Afu2g00710	alpha-amylase, putative	15±5	1.06±0.08	0.57±0.05	1.25±0.23
21.32±3.08	21.32±3.08	48.66±8.98	Afu3g00900	alpha-amylase AmyA	17±4	0.76±0.04	1.10±0.16	1.32±0.21
Lipases, Chitinases and phosphatases								
4.26±1.70	4.26±1.70	37.72±5.90	Afu2g03110	alkaline phosphatase	2±1	1.06±0.24	5.89±0.79	1.11±0.21
7.51±2.61	7.51±2.61	45.18±12.80	Afu1g17250	conidial hydrophobin RodB	4±1	0.52±0.09	0.27±0.04	0.31±0.05
5.20±0.88	5.21±0.88	31.61±5.88	Afu8g00930	chitosanase, putative	3±1	0.38±0.01	0.35±0.06	0.50±0.04
8.66±3.66	8.66±3.66	36.87±6.02	Afu5g02040	extracellular lipase, putative	14±6	0.72±0.11	4.10±1.80	0.60±0.07
7.22±1.40	7.22±1.40	29.41±5.98	Afu1g03570	acid phosphatase PHOa	5±1	0.67±0.04	1.13±0.19	1.16±0.13
11.15±2.04	11.15±2.04	45.06±5.65	Afu4g01290	endo-chitosanase, pseudogene	13±4	1.15±0.04	0.23±0.04	0.60±0.06
10.57±3.17	11.45±3.23	50.81±14.57	Afu3g11280	class V chitinase, putative	9±4	1.13±0.19	2.61±0.73	3.19±0.46
7.97±1.50	7.97±1.50	44.88±8.22	Afu1g16480	acid phosphatase, putative	4±1	1.09±0.08	1.56±0.98	1.65±0.20
14.15±1.92	14.15±1.92	43.12±4.98	Afu3g07160	class V chitinase, putative	12±4	0.74±0.05	2.19±0.58	1.53±0.33
9.05±2.56	9.17±2.50	40.63±7.73	Afu1g07440	Hsp70 chaperone	5±1	0.96±0.02	1.35±0.18	3.16±0.34
15.18±4.17	15.18±4.17	52.04±9.63	Afu3g01530	phosphatidylglycerol specific phospholipase C,	20±6	1.28±0.06	1.84±0.30	2.47±0.46
17.55±3.60	17.55±3.60	47.34±3.18	Afu4g03660	acid phosphatase, putative	25±7	0.81±0.03	0.58±0.09	1.63±0.24
39.47±5.47	39.47±5.47	72.40±1.89	Afu8g01410	class V chitinase ChiB1	83±25	1.78±0.46	2.63±0.62	2.64±0.57
29.09±3.50	29.09±3.50	80.11±11.16	Afu6g07940	lactoylglutathione lyase	23±4	1.02±0.05	0.80±0.08	2.88±0.18
7.53±3.43	7.53±3.43	32.94±6.59	Afu3g14680	lysophospholipase Plb3	4±2	0.70±0.05	1.67±0.12	1.83±0.14

2.80±0.66	2.80±0.66	9.46±0.81	Afu5g03760	class III chitinase ChiA1	3±1	0.96±0.13	5.23±1.10	0.76±0.07
Hypothetical proteins								
3.19±1.17	3.19±1.17	15.03±5.07	Afu8g02510	hypothetical protein	2±1	2.68±0.20	5.34±0.88	0.79±0.09
6.24±3.08	6.24±3.08	55.56±9.87	Afu2g05720	conserved hypothetical protein	6±3	0.82±0.13	0.70±0.04	2.18±0.77
40.90±6.62	40.90±6.62	68.56±3.40	Afu2g05240	conserved hypothetical protein	64±20	0.65±0.06	0.29±0.02	0.51±0.05
36.89±8.46	36.89±8.46	46.39±3.20	Afu6g03150	hypothetical protein	44±15	0.98±0.05	5.66±1.07	0.78±0.06
24.40±4.32	24.40±4.32	81.67±5.49	Afu3g01130	hypothetical protein	17±5	0.84±0.02	3.16±0.24	1.29±0.07
24.65±2.01	24.65±2.01	59.55±4.11	Afu3g02600	hypothetical protein	35±9	0.49±0.05	1.59±0.14	0.38±0.04
20.79±5.77	20.79±5.77	48.58±7.94	Afu7g04680	conserved hypothetical protein	11±3	1.14±0.04	3.65±0.58	0.80±0.08
13.75±5.95	13.75±5.95	45.61±7.20	Afu5g01190	conserved hypothetical protein	8±3	1.84±0.35	3.98±1.60	0.61±0.07
15.01±0.96	15.01±0.96	64.34±9.28	Afu5g01120	conserved hypothetical protein	10±1	0.55±0.05	0.43±0.03	0.67±0.06
13.52±2.90	13.52±2.90	32.18±8.18	Afu3g07870	conserved hypothetical protein	9±2	1.47±0.16	2.28±0.34	0.93±0.08
11.88±2.27	11.88±2.27	42.38±10.45	Afu5g10930	hypothetical protein	12±4	0.53±0.05	0.53±0.05	1.08±0.13
9.29±3.09	9.29±3.09	34.26±9.01	Afu3g12690	conserved hypothetical protein	6±3	1.02±0.06	1.15±0.10	1.78±0.50
11.04±1.54	11.04±1.54	59.75±11.75	Afu5g14560	hypothetical protein	7±2	1.41±0.05	2.15±0.27	1.06±0.09
9.30±2.48	9.30±2.48	32.11±4.55	Afu6g06560	conserved hypothetical protein	6±1	0.97±0.08	1.95±0.45	0.81±0.11
11.36±4.17	11.36±4.17	26.77±4.14	Afu4g13990	hypothetical protein	21±6	0.40±0.06	0.83±0.07	0.73±0.06
5.96±1.93	5.96±1.93	18.83±3.60	Afu7g05340	conserved hypothetical protein	3±1	0.68±0.06	0.84±0.11	0.95±0.31
8.53±0.87	8.53±0.87	56.57±6.12	Afu2g17630	hypothetical protein	7±2	0.92±0.06	4.88±0.91	0.87±0.08
6.51±1.65	6.51±1.65	46.59±4.10	Afu5g01420	conserved hypothetical protein	4±1	1.02±0.33	0.75±0.13	0.91±0.13
5.80±1.81	5.80±1.81	47.76±11.07	Afu5g03520	hypothetical protein	3±1	0.94±0.05	1.14±0.10	1.05±0.21
5.32±0.77	5.32±0.77	57.11±20.77	Afu6g00180	hypothetical protein	16±6	0.66±0.04	1.21±0.13	1.82±0.18

9.03±1.92	9.03±1.92	31.32±3.36	Afu5g02100	hypothetical protein	6±1	1.05±0.07	3.21±0.53	2.96±0.50
6.48±2.16	6.48±2.16	31.56±4.72	Afu1g16430	hypothetical protein	4±1	1.34±0.24	1.42±0.42	2.32±0.57
6.09±1.44	6.09±1.44	57.80±13.78	Afu8g00430	hypothetical protein	5±1	1.01±0.33	0.93±0.33	1.66±0.56
5.75±0.74	5.75±0.74	31.41±6.26	Afu6g14470	hypothetical protein	3±0	1.71±0.17	7.23±2.00	1.18±0.16
5.00±1.00	5.00±1.00	27.40±3.51	Afu1g09960	conserved hypothetical protein	4±2	1.21±0.37	0.85±0.06	2.50±0.28
7.38±2.20	6.07±1.62	18.82±9.49	Afu3g03610	hypothetical protein	3±1	0.94±0.04	1.06±0.27	1.14±0.28
4.15±1.26	4.15±1.26	50.30±9.53	Afu4g08450	conserved hypothetical protein	4±1	0.51±0.07	0.54±0.10	0.47±0.05
4.00±0.00	4.00±0.00	18.73±3.65	Afu6g02010	hypothetical protein	4±0	0.98±0.13	5.24±0.60	1.39±0.11
4.04±0.04	4.04±0.04	35.44±3.10	Afu3g00960	conserved hypothetical protein	3±1	0.96±0.09	2.07±0.34	2.30±0.45
4.04±1.39	4.04±1.39	57.18±12.84	Afu4g08310	conserved hypothetical protein	3±1	0.66±0.05	0.74±0.07	1.32±0.18
3.74±1.99	3.74±1.99	23.28±13.73	Afu3g10390	conserved hypothetical protein	2±1	1.13±0.22	1.05±0.15	1.58±0.55
5.00±1.73	5.00±1.73	18.99±3.91	Afu6g14400	conserved hypothetical protein	3±1	1.47±0.01	1.25±0.12	0.87±0.10
3.44±0.60	3.44±0.60	39.15±6.72	Afu1g02290	conserved hypothetical protein	4±2	0.98±0.10	1.19±0.23	1.52±0.26
5.00±1.00	5.00±1.00	62.05±11.48	Afu3g14940	hypothetical protein	3±1	0.94±0.14	1.05±0.10	0.90±0.09
9.28±3.30	9.28±3.30	50.87±12.89	Afu2g15770	conserved hypothetical protein	5±2	1.07±0.12	1.90±0.38	2.48±0.56
3.62±0.56	3.62±0.56	24.72±3.98	Afu2g15420	hypothetical protein	3±1	1.20±0.12	1.03±0.23	0.56±0.08
6.24±2.33	6.24±2.33	50.36±2.57	Afu4g02840	conserved hypothetical protein	3±2	1.02±0.15	0.71±0.14	1.57±0.31
8.50±34.70	3.67±0.85	12.16±5.44	Afu8g00630	conserved hypothetical protein, putative	2±1	1.57±0.64	4.12±2.52	3.07±2.72
3.10±1.02	3.10±1.02	32.34±8.54	Afu2g15710	hypothetical protein	2±1	1.02±0.10	1.79±0.50	1.91±0.64
2.76±0.44	2.00±0.00	11.30±13.15	Afu2g09140	conserved hypothetical protein	2±0	1.02±0.00	1.65±0.00	1.63±0.00
2.06±0.05	2.10±0.09	29.09±8.16	Afu2g01840	conserved hypothetical protein	2±0	1.62±0.50	1.40±0.36	3.10±0.87
3.50±0.87	3.50±0.87	28.33±2.92	Afu6g11720	hypothetical protein	2±0	1.03±0.25	1.20±0.15	1.37±0.21

2.12±0.41	2.12±0.41	14.00±6.15	Afu5g14420	hypothetical protein	2±0	0.49±0.08	0.65±0.08	1.51±0.13
3.41±1.41	3.41±1.41	20.87±5.92	Afu1g05270	conserved hypothetical protein	4±1	1.05±0.09	2.46±0.31	1.17±0.13
2.00±0.00	2.00±0.00	31.39±12.63	Afu6g10840	hypothetical protein	2±0	0.88±0.09	1.90±0.33	5.26±1.22
2.00±0.00	2.00±0.00	22.38±14.72	Afu2g12000	conserved hypothetical protein	2±1	0.51±0.01	0.31±0.04	0.55±0.07

Table S4. The band showing cellulolytic activity was cut, digested and analyzed by LC-MS/MS. The proteins identified in the band are listed below along with their mass, protein score, emPAI and Peptides detected.

Accession	GS	Name	Pro_Mass	Prot_score	emPAI	Peptides
Afu3g00320	Afu3g00320	endo-1,4-beta-xylanase (XlnA), putative (229 aa)	24536	1111	2.16	42
Afu7g06150	Afu7g06150	endoglucanase, putative (235 aa)	25883	742	2.37	33
Afu1g16190	Afu1g16190	cell wall glucanase Crf1 (396 aa)	40487	132	0.27	9
Afu3g02970	Afu3g02970	aspergillopepsin, putative (270 aa)	28225	52	0.25	3
Afu1g07300	Afu1g07300	tetracycline transporter-like protein, putative (461 aa)	49371	45	0.07	1
Afu1g09060	Afu1g09060	conserved hypothetical protein (513 aa)	57133	43	0.06	2
Afu2g03620	Afu2g03620	oxidoreductase, short-chain	53366	42	0.06	2
Afu3g12510	Afu3g12510	vesicular fusion ATPase, putative (771 aa)	84710	41	0.04	4
Afu4g02960	Afu4g02960	UDP-sugar hydrolase, putative (397 aa)	43996	39	0.08	1

Afu3g09300	Afu3g09300	conserved hypothetical protein (168 aa)	17248	29	0.2	3
Afu5g12560	Afu5g12560	hypothetical protein (733 aa)	83681	26	0.04	2
Afu1g15080	Afu1g15080	hypothetical protein (620 aa)	69179	26	0.05	1
Afu7g06780	Afu7g06780	hypothetical protein (235 aa)	26238	26	0.13	1
Afu4g03730	Afu4g03730	dienelactone hydrolase (493 aa)	54642	25	0.06	3
Afu5g00230	Afu5g00230	hypothetical protein (583 aa)	64230	25	0.05	2
Afu1g10610	Afu1g10610	hypothetical protein (175 aa)	19842	24	0.17	2
Afu6g12310	Afu6g12310	conserved hypothetical protein (300 aa)	33701	24	0.1	1
Afu3g12340	Afu3g12340	conserved hypothetical protein (693 aa)	75736	23	0.04	3
Afu1g06060	Afu1g06060	hypothetical protein (1503 aa)	164339	23	0.02	2
Afu2g04140	Afu2g04140	conserved hypothetical protein (1560 aa)	171043	23	0.02	4
Afu1g12200	Afu1g12200	conserved hypothetical protein (313 aa)	35110	23	0.09	1

Afu4g13050	Afu4g13050	hypothetical protein (485 aa)	54219	21	0.06	2
Afu4g11600	Afu4g11600	hypothetical protein (845 aa)	94168	21	0.03	2
Afu4g08030	Afu4g08030	conserved hypothetical protein (625 aa)	69539	19	0.05	2
Afu2g04920	Afu2g04920	conserved hypothetical protein (1467 aa)	154207	18	0.02	1
Afu2g05940	Afu2g05940	conserved hypothetical protein (370 aa)	40602	18	0.08	1
Afu6g07380	Afu6g07380	conserved hypothetical protein (634 aa)	65757	18	0.05	1
Afu2g10690	Afu2g10690	MFS phosphate transporter, putative (667 aa)	75128	17	0.04	1
Afu2g09270	Afu2g09270	hypothetical protein (409 aa)	45569	16	0.07	1
Afu2g14260	Afu2g14260	conserved hypothetical protein (960 aa)	102846	16	0.03	1