WRINKLED1 transcription factor : how much do we know about its regulatory mechanism?

Kong, Que; Ma, Wei

2018

Kong, Q., & Ma, W. (2018). WRINKLED1 transcription factor : how much do we know about its regulatory mechanism? Plant Science, 272, 153-156. doi:10.1016/j.plantsci.2018.04.013

<https://hdl.handle.net/10356/138303>

<https://doi.org/10.1016/j.plantsci.2018.04.013>

© 2018 Elsevier B.V. All rights reserved. This paper was published in Plant Science and is made available with permission of Elsevier B.V.

Downloaded on 26 Apr 2023 22:06:02 SGT

Title: WRINKLED1 transcription factor: how much do we know about its regulatory mechanism?

Authors: Que Kong¹ and Wei Ma^{2*}

Affiliation:

¹MSU-DOE Plant Research Laboratory, Michigan State University, East Lansing, MI 48824, USA.

²School of Biological Sciences, Nanyang Technological University, 637551, Singapore.

***Corresponding author:**

Wei Ma; Email: weima@ntu.edu.sg

Abstract

Many plant species produce and build up triacylglycerol (TAG) in their seeds as a main resource to provide carbon and energy during seedling development. Plant seed oils are important not only for human diets but also as renewable feedstock of industrial uses. WRINKLED1 (WRI1), an APETALA2 (AP2) transcription factor, plays an essential role in the transcriptional regulation of TAG biosynthesis as WRI1 regulates the expression of key genes in the glycolytic and fatty acid biosynthetic pathways. Recent work has identified intrinsic structural disorder in WRI1 that may affect the stability of the protein. Furthermore, WRI1 activity is modulated by post-translational modifications and interacting partners. These progresses shed new light on regulatory functions of WRI1 at the molecular levels, paving new paths to the use of WRI1 for bioengineering of TAG in plants.

Key words

Arabidopsis, transcription factor, plant oil biosynthesis, post-translational modifications, protein stability, protein–protein interaction.

Contents

- 1) Introduction**
- 2) WRI1 is a master transcriptional regulator of plant oil biosynthetic pathways**
- 3) Regulators that control *AtWRI1* expression**
- 4) *AtWRI1*' activity is mediated by posttranslational modifications and interacting partners**
- 5) Targets of *AtWRI1* in pathways other than oil biosynthesis**
- 6) *WRI1* orthologs in other plant species**
- 7) *WRI1* is important for translational research**
- 8) Future Perspectives**

Figure Legend

Acknowledgements

Reference

Introduction

Plant oils (mainly triacylglycerol, or TAG) are accumulated by numerous plant species in their seeds as the storage resource (carbon and energy) to support seedling development. Plant oils are also an essential supply for human diets and for industry as renewable feed stocks. It has been predicted that global demand for plant oils will double by 2030, necessitating technological breakthroughs [1].

TAG biosynthesis requires two main steps: fatty acid biosynthesis in the plastids and TAG assembly in the endoplasmic reticulum. After export from plastids, fatty acids are converted into acyl-CoA and assembled mainly via the eukaryotic phospholipid biosynthetic pathway [2]. In the last step of oil biosynthetic process, diacylglycerol is acylated using either acyl-CoA or phospholipids to form TAG, catalyzed by diacylglycerol acyltransferase (DGAT) or phosphatidylcholine:diacylglycerol acyltransferase (PDAT) [3-5]. The regulatory mechanism of TAG biosynthesis is complicated and not completely understood, hence the identification of gene regulators and studying their regulatory mechanisms are important for both understanding the oil biosynthetic process and designing strategies to efficiently increase plant oil production [6-7].

WRI1 is a master transcriptional regulator of plant oil biosynthetic pathways

WRINKLED1 (WRI1), a member of the APETALA2 (AP2) transcription factor family, is a key regulator of plant oil biosynthesis [8-9] (also see Figure 1A). The seed oil content in *wri1-1* (the *Arabidopsis* loss-of-function mutant of *AtWRI1*) is reduced by 80% compared to the wild-type (WT) [10]. The transcriptomic analysis of developing seeds of WT and *wri1-1* has shown that

the majority of the genes which display reduced expression in *wri1-1* are related to the glycolytic and fatty acid biosynthetic pathways [11]. Recent work has identified that many genes in glycolysis and fatty acid biosynthesis are targets of AtWRI1, such as genes encoding acetyl-CoA carboxylase, acyl carrier protein, ketoacyl-ACP synthase, a subunit of pyruvate kinase [12-14]. WRI1 is thus referred as a "master regulator" for transcriptional regulation of seed oil biosynthesis [1].

Regulators that control *AtWRI1* expression

Key regulators of seed maturation, including LEAFY COTYLEDON1 (LEC1) [15] and LEAFY COTYLEDON2 (LEC2) [12], seem to be responsible for activating *AtWRI1*. *LEC2* co-expresses with *AtWRI1*, leading to the activation of *AtWRI1* during Arabidopsis embryo development [12]. *AtWRI1* expression is increased in the gain-of-function mutant of *LEC1*, *tnp* [15], implicating a role for LEC1 in controlling *AtWRI1* [14, 16]. However, it should be noted that the gene promoter cis-elements, to which LEC1 and LEC2 recognize and bind, have yet to be identified (Figure 1B). How LEC1 and LEC2 regulate the expression of *AtWRI1* thus remain to be further elucidated [14]. In addition, the Arabidopsis transcription factor FUSCA3 (FUS3) is another positive regulator of *AtWRI1*. Comparative microarray analysis reveals that *AtWRI1* expression is reduced in the *fus3* mutant compared to WT [17]. Based on functional redundancy between LEC2 and FUS3, FUS3 has been speculated to regulate *AtWRI1* expression similar to that by LEC2 [17]. A recent ChIP-chip experiment [chromatin immunoprecipitation ("ChIP") followed by DNA microarray ("chip")] has confirmed that *AtWRI1* is a direct target of FUS3 [18] (Figure 1B). *AtWRI1* expression is induced by sucrose treatment [9]. AtWRI1 also activates some sugar-responsive genes, hence possibly playing a role in sugar signaling that mediates carbon flow to

storage oil [9]. *FUS3* expression does not influence the expression of *LEC1* or *LEC2*, *AtWRI1* activation in response to sucrose is thus suggested to be directly regulated by *FUS3* instead of indirectly through *LEC1* or *LEC2* [6]. It is highly possible that some plant species might have additional upstream *WRI1* regulator(s) other than those involved in seed maturation (e.g. *LEC1*, *LEC2*, and *FUS3*). In oil palm mesocarp, where *EgWRI1* is highly expressed, expression of key regulators for seed maturation remain low, indicating that *EgWRI1* is controlled by novel fruit-specific regulator(s) [19].

Recent work has identified that the transcription factor *MYB89* represses *AtWRI1* [20]. *AtWRI1* expression is elevated in *Arabidopsis* loss-of-function mutant, *myb89*, and upregulation of *MYB89* leads to repression of *AtWRI1* [20]. Further ChIP assay confirmed that *MYB89* directly binds the promoter of *AtWRI1*, indicating that *AtWRI1* is a direct target of *MYB89* [20] (Figure1B).

***AtWRI1*' activity is mediated by posttranslational modifications and interacting partners**

A number of mechanisms of both positive and negative transcriptional regulation of *WRI1* activities have been illustrated. Recent progress also has begun to shed light on *WRI1* function at the protein level. These efforts include *in silico* analysis of protein structural features, characterization of functional domains/motifs, and the identification of *WRI1* interacting partners. Using CULLIN3-based E3 ligase adaptor BTB/POZMATH 1 (*BPM1*) protein as a prey, *AtWRI1* was identified to interact with *BPM1* in yeast two-hybrid (Y2H) assay [21]. *AtWRI1* is able to interact with multiple other *BPMs*; and the degradation of *AtWRI1* is mediated by the 26S proteasome through interaction with E3 adaptor *BPMs* [21]. This work establishes a

connection between E3 ligase and plant lipid metabolism via degradation of the key transcriptional regulator, AtWRI1.

In silico analysis has revealed that AtWRI1 protein possesses intrinsic disorder features. Three intrinsically disordered regions (IDRs) are predicted using several computational programs [22]. A PEST motif, a well-characterized peptide signal sequence for protein degradation, is present in the IDR3 domain of AtWRI1 [22]. A transactivation domain (TAD) has been identified in AtWRI1 C-terminus, which does not overlap with the IDR3-PEST motif [22]. AtWRI1 has been engineered by removal of the IDR3-PEST motif or by mutation of putative phosphorylation residues in the IDR3-PEST motif. The engineered AtWRI1 proteins show enhanced stability, and overexpression of the engineered *AtWRI1* lead to increase of TAG biosynthesis in Arabidopsis compared to WT [22]. Phosphorylation is hence suggested as a possible mechanism that mediates AtWRI1 stability via modulation of the IDR3-PEST motif [22].

AtWRI1 has been recently identified as a new client of 14-3-3 proteins (a family of conserved regulatory proteins with ability to bind to phosphorylated proteins) [23]. AtWRI1 physically interacts with 14-3-3s in both yeast and plant cells [23]. Overexpression of a *14-3-3* enhances AtWRI1-mediated TAG biosynthesis, as well as transcriptional activity and stability of AtWRI1 [23]. The 14-3-3 binding motif in AtWRI1 overlaps with the binding motif for BPMs [23]. It is proposed that the interaction between AtWRI1 and 14-3-3 either prevents E3 ligase-BPMs from interacting with AtWRI1 or separates AtWRI1 from the bound E3 ligase-BPMs [23].

KIN10 (an essential SNF1-related protein kinase in the sugar signal transduction cascade) phosphorylates AtWRI1 through physical interaction[24]. The KIN10-mediated AtWRI1 phosphorylation is important for the AtWRI1 degradation. KIN10 phosphorylates AtWRI1 at

residues T70 and S166 [24]. The phosphorylation deficient mutant AtWRI1^{T70A/S166A} displays increased protein stability compared to WT AtWRI1 in the presence of KIN10 [24]. Interestingly, the 14-3-3 binding motif [23] and KIN10 phosphorylated residues [24] are close, suggesting that these two modulation processes might be overlapping probably via the proteasomal degradation pathway (Figure 1A).

In silico analysis also predicts that AtWRI1 is one of potential substrates of the kinase OST1 (SnRK2.6) [25]. SnRK2 kinases are essential regulators in ABA signal transduction [26-28]. The proposed OST1 phosphorylation motif in AtWRI1 protein is positioned in between the two AP2 domains [25]. SnRK2.6 mediates the stability of ABA-responsive-element Binding Factor 3 (ABF3) through phosphorylation [25]. Therefore, it is possible that SnRK2s are involved in phosphorylation of AtWRI1, which in turn affects the activity of AtWRI1 and AtWRI1-mediated oil biosynthesis.

The Mediator 15 (MED15), a subunit of the Arabidopsis mediator complex [29-30], physically interacts with AtWRI1 in both *in vitro* and *in vivo* assays [31]. *MED15* overexpression has resulted in the activation of a number of AtWRI1 target genes. ChIP assays suggest that MED15 binds the promoters of these AtWRI1 target genes [31]. However, expression of the AtWRI1 target genes remain elevated in transgenic *wri1* plants overexpressing *MED15*, indicating the presence of alternative transcription factors that interact with MED15 to regulate those genes [31].

Targets of AtWRI1 in pathways other than oil biosynthesis

While binding to the promoters of *PINs* at AW-boxes, AtWRI1 binds to the promoter of *GH3.3* (a gene involved in auxin degradation) at a novel binding motif other than AW-box, implicating

a role in mediating root auxin homeostasis [32]. AtWRI1 is also able to activate the expression of *Citrus sinensis* *Terpene Synthase 1* (*CsTPS1*) [33]. AtWRI1 is a homolog of the *C. sinensis* AP2 transcription factor, CitAP2.10, which stimulates the biosynthesis of (+)-valencene in orange [33]. Thus far, we do not fully understand how regulation of the alternative targets of AtWRI1 is connected to growth and development (e.g. root growth [32]). However, realizing the fact that WRI1 targets genes which might alter other physiological processes (e.g. hormone homeostasis [32]) is critical for the application of WRI1 in translational research to increase oil content.

WRI1 orthologs in other plant species

WRI1 orthologs have been identified from both monocot and dicot species, including *Brassica napus*, *Zea mays*, *Elaeis guineensis*, *Gossypium spp*, *Ricinus communis*, *Brachypodium distachyon*, *Camelina sativa*, and *Cocos nucifera* [34-39]. AtWRI1 and WRI1 orthologs are able to complement the phenotypes of *wri1* mutants [8, 35-36, 40]. Similar to AtWRI1, numerous WRI1 orthologs are found to express highly in developing seeds. However, oil palm WRI1 expresses highly in non-seed tissue, the oil accumulating mesocarp, and the transcript of *EgWRI1* increases as fruit ripening [19]. In addition, WRI1 orthologs from avocado [41] and nutsedge [42] were found to express highly in mesocarp and underground stem tuber, respectively. Structural features and functional motifs, including “VYL” [35], IDR, and PEST motif [22], are conserved features across the characterized WRI1 proteins [22, 35, 38, 40].

WRI1 is important for translational research

Overexpression of AtWRI1 or other WRI1 orthologs have resulted in increased seed oil content in transgenic plants [8, 34, 37-38, 43-44]. Ectopic expression of AtWRI1 and *BnWRI1* elevate oil

content in leaves of transgenic *Arabidopsis* and *Brachypodium* plants [38, 45]. *ZmWRII* overexpression, driven by the embryo-preferred *OLEOSIN* (*OLE*) promoter, has significantly increased seed oil accumulation in transgenic maize plants; however, little or no increase of oil content is resulted when the expression is driven by the *19 KD ZEIN* promoter (aimed to target the expression in starch endosperm) [37]. Using the *FUS3* promoter to drive *AtWRII* expression, in order to facilitate the extension of oil biosynthesis in the mid-phase of seed development, has efficiently increased seed oil content in transgenic *Arabidopsis* plants [46].

In tobacco leaves, transient overexpression of *AtWRII* and *WRII* orthologs significantly increase TAG accumulation [22, 42, 47]. Transient co-expression of *WRII* and *DGATI* has resulted in significantly higher oil content in tobacco leaves compared to expressing *WRII* alone, suggesting a synergistic effect between the two genes [47]. In transient expression assays using tobacco leaves, ectopic expression of stabilized forms of *WRII*, resulted from mutation at the phosphorylation sites in the IDR3-PEST motif, enhances oil production [22]. Transient co-expression of *AtWRII* and a *14-3-3* in tobacco leaves also enhances protein stability of *AtWRII*, leading to increased oil production [23].

In addition, overexpression of selected transcription factors that upregulate *WRII*, identified from plant species other than *Arabidopsis*, have proven to be useful for the increase of seed oil content. For instance, overexpression of soybean transcription factors, *GmZF351* and *GmDREBL*, enhance seed oil in transgenic *Arabidopsis* [48-49]. The two soybean transcription factors are able to activate *AtWRII* by binding to its promoter [48-49]. Overexpression in maize of *ZmLEC1*, an activator of *ZmWRII*, has increased seed oil up to 48% [37].

It should not be overlooked that overexpression of *WRI1* might cause abnormal growth or cell death in certain plant species [8, 14, 38]. Therefore, selection of suitable promoters of target plant species is important for the improvement of oil crops or development of new oil crops.

Future perspectives

To date, how *WRI1* expression is regulated still remains largely unknown. Also unclear is how LEC1, LEC2, and FUS3 regulate the expression of *WRI1*. It is likely that additional regulators also control the expression of *WRI1* during the developmental process or in response to environmental cues (e.g. sucrose). Addressing these questions may require detailed analyses of the *WRI1* promoters and the layered regulatory network.

Efforts have been made recently to bridge the structural features to functions of WRI1. Identification and characterization of the IDR [22], PEST motif [22], TAD [22], 14-3-3 and E3 ligase adaptor binding motif [23], and KIN10 kinase phosphorylation residues [24] in AtWRI1 protein have shed new lights onto the structure/function relationship of AtWRI1. Future work will focus on understanding the functions of individual domains and motifs, as well as how protein-protein interactions affect the dynamics of AtWRI1. Phosphorylation at T70/S166 of the IDR3-PEST motif has been shown to be important for AtWRI1 function [22, 24]. Given that AtWRI1 protein is enriched of putative phosphorylation sites [22], the fine characterization of AtWRI1 phosphorylation sites and identification of the new kinase(s) that phosphorylate AtWRI1 will be of great significance. Moreover, phosphorylation is proposed to play a dual role in mediating the stability and activity of AtWRI1 [23]. It is proposed that, in the early stage of embryo development, phosphorylation of AtWRI1 creates a binding site for 14-3-3, resulting in stabilization of AtWRI1 protein; however, during the late stage of embryo development,

phosphorylation of IDR3-PEST motif in AtWRI1 facilitates the AtWRI1 degradation [23] (Figure 1C). Thus, how AtWRI1 activity is dynamically modulated by the upstream kinases (e.g. SnRK2.6 [25]), in response to diverse environmental and developmental signals, requires attention.

Recent progress also indicates that AtWRI1 has novel targets other than genes in glycolysis and fatty acid biosynthesis via binding to AW-box or novel binding sites [32]. Understanding the effects of WRI1 on these genes unrelated to oil biosynthesis might be helpful to overcome the side effects of *WRI1* overexpression aimed for the improvement of oil crops [8, 14, 38]. Genome editing and protein engineering approaches can be applied to increase WRI1 binding specificity to the oil biosynthetic genes.

Figure Legend

Figure 1. Structural characteristics of AtWRI1 and a simplified model for the regulation of AtWRI1-mediated oil gene expression. A) Schematic representation of AtWRI1 showing the locations of the conserved functional motif of "VYL", three intrinsically disordered regions, the PEST motif, the transactivation domain (TAD), the 14-3-3 and BPM binding motifs, and the KIN10 kinase phosphorylation residues (T70 and S166). B) Regulation of *WRI1* expression. LEC1 and LEC2 have been suggested as key regulators mediating the expression of *WRI1*,

despite the lacking of evidence for LEC1/LEC2 direct binding to the *WRI1* promoter. FUS3 and MYB89 have been shown to directly bind to the *WRI1* promoter, acting as positive and negative regulators, respectively. C) A proposed model of possible regulatory mechanisms for AtWRI1. Post-translational modifications (such as phosphorylation and ubiquitination) of AtWRI1 occur in response to different environmental cues and developmental states, leading to altered protein stability and transcriptional activity, which in turn affects the expression of AtWRI1 target genes.

Acknowledgements

We are grateful to Dr. Ling Yuan (University of Kentucky) for critical reading and valuable suggestions on the manuscript. We apologize to all authors whose important work was not cited because of space limitations. The work on WRII in the Ma Lab is currently supported by a Nanyang Technological University Startup grant (to W.M.).

Reference

- [1] K.D. Chapman, J.B. Ohlrogge, Compartmentation of triacylglycerol accumulation in plants, *J Biol Chem*, 287 (2012) 2288-2294.
- [2] P.D. Bates, T.P. Durrett, J.B. Ohlrogge, M. Pollard, Analysis of acyl fluxes through multiple pathways of triacylglycerol synthesis in developing soybean embryos, *Plant Physiol*, 150 (2009) 55-72.
- [3] J. Zou, Y. Wei, C. Jako, A. Kumar, G. Selvaraj, D.C. Taylor, The *Arabidopsis thaliana* TAG1 mutant has a mutation in a diacylglycerol acyltransferase gene, *Plant J*, 19 (1999) 645-653.
- [4] M. Zhang, J. Fan, D.C. Taylor, J.B. Ohlrogge, DGAT1 and PDAT1 acyltransferases have overlapping functions in *Arabidopsis* triacylglycerol biosynthesis and are essential for normal pollen and seed development, *Plant Cell*, 21 (2009) 3885-3901.
- [5] A. Dahlqvist, U. Stahl, M. Lenman, A. Banas, M. Lee, L. Sandager, H. Ronne, S. Stymne, Phospholipid:diacylglycerol acyltransferase: an enzyme that catalyzes the acyl-CoA-independent formation of triacylglycerol in yeast and plants, *Proc Natl Acad Sci U S A*, 97 (2000) 6487-6492.
- [6] M. Zhang, X. Cao, Q. Jia, J. Ohlrogge, FUSCA3 activates triacylglycerol accumulation in *Arabidopsis* seedlings and tobacco BY2 cells, *Plant J*, 88 (2016) 95-107.
- [7] P.D. Bates, S. Stymne, J. Ohlrogge, Biochemical pathways in seed oil synthesis, *Current opinion in plant biology*, 16 (2013) 358-364.
- [8] A. Cernac, C. Benning, WRINKLED1 encodes an AP2/EREB domain protein involved in the control of storage compound biosynthesis in *Arabidopsis*, *Plant J*, 40 (2004) 575-585.
- [9] T. Masaki, N. Mitsui, H. Tsukagoshi, T. Nishii, A. Morikami, K. Nakamura, ACTIVATOR of Spomin::LUC1/WRINKLED1 of *Arabidopsis thaliana* transactivates sugar-inducible promoters, *Plant Cell Physiol*, 46 (2005) 547-556.
- [10] N. Focks, C. Benning, wrinkled1: A novel, low-seed-oil mutant of *Arabidopsis* with a deficiency in the seed-specific regulation of carbohydrate metabolism, *Plant Physiol*, 118 (1998) 91-101.
- [11] S.A. Ruuska, T. Girke, C. Benning, J.B. Ohlrogge, Contrapuntal networks of gene expression during *Arabidopsis* seed filling, *Plant Cell*, 14 (2002) 1191-1206.
- [12] S. Baud, M.S. Mendoza, A. To, E. Harscoet, L. Lepiniec, B. Dubreucq, WRINKLED1 specifies the regulatory action of LEAFY COTYLEDON2 towards fatty acid metabolism during seed maturation in *Arabidopsis*, *Plant J*, 50 (2007) 825-838.
- [13] K. Maeo, T. Tokuda, A. Ayame, N. Mitsui, T. Kawai, H. Tsukagoshi, S. Ishiguro, K. Nakamura, An AP2-type transcription factor, WRINKLED1, of *Arabidopsis thaliana* binds to the AW-box sequence conserved among proximal upstream regions of genes involved in fatty acid synthesis, *Plant J*, 60 (2009) 476-487.
- [14] C. Marchive, K. Nikovics, A. To, L. Lepiniec, S. Baud, Transcriptional regulation of fatty acid production in higher plants: molecular bases and biotechnological outcomes. , *Eur. J. Lipid Sci. Technol.* , 116 (2014) 1332-1343.
- [15] S.A. Casson, K. Lindsey, The turnip mutant of *Arabidopsis* reveals that LEAFY COTYLEDON1 expression mediates the effects of auxin and sugars to promote embryonic cell identity, *Plant Physiol*, 142 (2006) 526-541.
- [16] M. Santos-Mendoza, B. Dubreucq, S. Baud, F. Parcy, M. Caboche, L. Lepiniec, Deciphering gene regulatory networks that control seed development and maturation in *Arabidopsis*, *Plant J*, 54 (2008) 608-620.

- [17] A. Yamamoto, Y. Kagaya, H. Usui, T. Hobo, S. Takeda, T. Hattori, Diverse roles and mechanisms of gene regulation by the Arabidopsis seed maturation master regulator FUS3 revealed by microarray analysis, *Plant Cell Physiol*, 51 (2010) 2031-2046.
- [18] F. Wang, S.E. Perry, Identification of direct targets of FUSCA3, a key regulator of Arabidopsis seed development, *Plant Physiol*, 161 (2013) 1251-1264.
- [19] F. Bourgis, A. Kilaru, X. Cao, G.F. Ngando-Ebongue, N. Drira, J.B. Ohlrogge, V. Arondel, Comparative transcriptome and metabolite analysis of oil palm and date palm mesocarp that differ dramatically in carbon partitioning, *Proc Natl Acad Sci U S A*, 108 (2011) 12527-12532.
- [20] D. Li, C. Jin, S. Duan, Y. Zhu, S. Qi, K. Liu, C. Gao, H. Ma, M. Zhang, Y. Liao, M. Chen, MYB89 Transcription Factor Represses Seed Oil Accumulation, *Plant Physiol*, 173 (2017) 1211-1225.
- [21] L. Chen, J.H. Lee, H. Weber, T. Tohge, S. Witt, S. Roje, A.R. Fernie, H. Hellmann, Arabidopsis BPM Proteins Function as Substrate Adaptors to a CULLIN3-Based E3 Ligase to Affect Fatty Acid Metabolism in Plants, *Plant Cell*, 25 (2013) 2253-2264.
- [22] W. Ma, Q. Kong, M. Grix, J.J. Mantyla, Y. Yang, C. Benning, J.B. Ohlrogge, Deletion of a C-terminal intrinsically disordered region of WRINKLED1 affects its stability and enhances oil accumulation in Arabidopsis, *Plant J*, 83 (2015) 864-874.
- [23] W. Ma, Q. Kong, J.J. Mantyla, Y. Yang, J.B. Ohlrogge, C. Benning, 14-3-3 protein mediates plant seed oil biosynthesis through interaction with AtWRI1, *Plant J*, 88 (2016) 228-235.
- [24] Z. Zhai, H. Liu, J. Shanklin, Phosphorylation of WRINKLED1 by KIN10 Results in Its Proteasomal Degradation, Providing a Link between Energy Homeostasis and Lipid Biosynthesis, *Plant Cell*, 29 (2017) 871-889.
- [25] C. Sirichandra, M. Davanture, B.E. Turk, M. Zivy, B. Valot, J. Leung, S. Merlot, The Arabidopsis ABA-activated kinase OST1 phosphorylates the bZIP transcription factor ABF3 and creates a 14-3-3 binding site involved in its turnover, *PloS one*, 5 (2010) e13935.
- [26] H. Fujii, P.E. Verslues, J.K. Zhu, Arabidopsis decuple mutant reveals the importance of SnRK2 kinases in osmotic stress responses in vivo, *Proc Natl Acad Sci U S A*, 108 (2011) 1717-1722.
- [27] H. Fujii, J.K. Zhu, Arabidopsis mutant deficient in 3 abscisic acid-activated protein kinases reveals critical roles in growth, reproduction, and stress, *Proc Natl Acad Sci U S A*, 106 (2009) 8380-8385.
- [28] Y. Fujita, K. Nakashima, T. Yoshida, T. Katagiri, S. Kidokoro, N. Kanamori, T. Umezawa, M. Fujita, K. Maruyama, K. Ishiyama, M. Kobayashi, S. Nakasone, K. Yamada, T. Ito, K. Shinozaki, K. Yamaguchi-Shinozaki, Three SnRK2 protein kinases are the main positive regulators of abscisic acid signaling in response to water stress in Arabidopsis, *Plant Cell Physiol*, 50 (2009) 2123-2132.
- [29] S. Backstrom, N. Elfving, R. Nilsson, G. Wingsle, S. Bjorklund, Purification of a plant mediator from Arabidopsis thaliana identifies PFT1 as the Med25 subunit, *Molecular cell*, 26 (2007) 717-729.
- [30] S. Mathur, S. Vyas, S. Kapoor, A.K. Tyagi, The Mediator complex in plants: structure, phylogeny, and expression profiling of representative genes in a dicot (Arabidopsis) and a monocot (rice) during reproduction and abiotic stress, *Plant Physiol*, 157 (2011) 1609-1627.
- [31] M.J. Kim, I.C. Jang, N.H. Chua, The Mediator Complex MED15 Subunit Mediates Activation of Downstream Lipid-Related Genes by the WRINKLED1 Transcription Factor, *Plant Physiol*, 171 (2016) 1951-1964.

- [32] Q. Kong, W. Ma, H. Yang, G. Ma, J.J. Mantyla, C. Benning, The Arabidopsis WRINKLED1 transcription factor affects auxin homeostasis in roots, *Journal of experimental botany*, (2017) doi:10.1093/jxb/erx1275.
- [33] S.L. Shen, X.R. Yin, B. Zhang, X.L. Xie, Q. Jiang, D. Grierson, K.S. Chen, CitAP2.10 activation of the terpene synthase CsTPS1 is associated with the synthesis of (+)-valencene in 'Newhall' orange, *Journal of experimental botany*, 67 (2016) 4105-4115.
- [34] J. Liu, W. Hua, G. Zhan, F. Wei, X. Wang, G. Liu, H. Wang, Increasing seed mass and oil content in transgenic Arabidopsis by the overexpression of wr1-like gene from Brassica napus, *Plant physiology and biochemistry*, 48 (2010) 9-15.
- [35] W. Ma, Q. Kong, V. Arondel, A. Kilaru, P.D. Bates, N.A. Thrower, C. Benning, J.B. Ohlrogge, Wrinkled1, a ubiquitous regulator in oil accumulating tissues from Arabidopsis embryos to oil palm mesocarp, *PloS one*, 8 (2013) e68887.
- [36] B. Pouvreau, S. Baud, V. Vernoud, V. Morin, C. Py, G. Gendrot, J.P. Pichon, J. Rouster, W. Paul, P.M. Rogowsky, Duplicate maize Wrinkled1 transcription factors activate target genes involved in seed oil biosynthesis, *Plant Physiol*, 156 (2011) 674-686.
- [37] B. Shen, W.B. Allen, P. Zheng, C. Li, K. Glassman, J. Ranch, D. Nubel, M.C. Tarczynski, Expression of ZmLEC1 and ZmWRI1 increases seed oil production in maize, *Plant Physiol*, 153 (2010) 980-987.
- [38] Y. Yang, J. Munz, C. Cass, A. Zienkiewicz, Q. Kong, W. Ma, J. Sedbrook, C. Benning, Ectopic Expression of WRINKLED1 Affects Fatty Acid Homeostasis in Brachypodium distachyon Vegetative Tissues, *Plant Physiol*, 169 (2015) 1836-1847.
- [39] D. Tajima, A. Kaneko, M. Sakamoto, Y. Ito, N.T. Hue, M. Miyazaki, Y. Ishibashi, T. Yuasa, M. Iwaya-Inoue, Wrinkled 1 (WRI1) Homologs, AP2-Type Transcription Factors Involving Master Regulation of Seed Storage Oil Synthesis in Castor Bean (*Ricinus communis* L.), *American Journal of Plant Sciences*, 4 (2013) 333-339.
- [40] D. An, H. Kim, S. Ju, Y.S. Go, H.U. Kim, M.C. Suh, Expression of Camelina WRINKLED1 Isoforms Rescue the Seed Phenotype of the Arabidopsis wr1 Mutant and Increase the Triacylglycerol Content in Tobacco Leaves, *Frontiers in plant science*, 8 (2017) 34.
- [41] A. Kilaru, X. Cao, P.B. Dabbs, H.J. Sung, M.M. Rahman, N. Thrower, G. Zynda, R. Podicheti, E. Ibarra-Laclette, L. Herrera-Estrella, K. Mockaitis, J.B. Ohlrogge, Oil biosynthesis in a basal angiosperm: transcriptome analysis of Persea Americana mesocarp, *Bmc Plant Biol*, 15 (2015) 203.
- [42] A. Grimberg, A.S. Carlsson, S. Marttila, R. Bhalerao, P. Hofvander, Transcriptional transitions in Nicotiana benthamiana leaves upon induction of oil synthesis by WRINKLED1 homologs from diverse species and tissues, *Bmc Plant Biol*, 15 (2015) 192.
- [43] D. An, M.C. Suh, Overexpression of Arabidopsis WRI1 enhanced seed mass and storage oil content in Camelina sativa *Plant Biotechnol Rep*, 9 (2015) 137-148.
- [44] R. Sun, R. Ye, L. Gao, L. Zhang, R. Wang, T. Mao, Y. Zheng, D. Li, Y. Lin, Characterization and Ectopic Expression of CoWRI1, an AP2/EREBP Domain-Containing Transcription Factor from Coconut (*Cocos nucifera* L.) Endosperm, Changes the Seeds Oil Content in Transgenic Arabidopsis thaliana and Rice (*Oryza sativa* L.), *Frontiers in plant science*, 8 (2017) 63.
- [45] Sanjaya, T.P. Durrett, S.E. Weise, C. Benning, Increasing the energy density of vegetative tissues by diverting carbon from starch to oil biosynthesis in transgenic Arabidopsis, *Plant Biotechnol J*, 9 (2011) 874-883.

- [46] M. Kanai, S. Mano, M. Kondo, M. Hayashi, M. Nishimura, Extension of oil biosynthesis during the mid-phase of seed development enhances oil content in Arabidopsis seeds, *Plant Biotechnol J*, 14 (2016) 1241-1250.
- [47] T. Vanhercke, A. El Tahchy, P. Shrestha, X.R. Zhou, S.P. Singh, J.R. Petrie, Synergistic effect of WRI1 and DGAT1 coexpression on triacylglycerol biosynthesis in plants, *FEBS Lett*, 587 (2013) 364-369.
- [48] Y.Q. Zhang, X. Lu, F.Y. Zhao, Q.T. Li, S.L. Niu, W. Wei, W.K. Zhang, B. Ma, S.Y. Chen, J.S. Zhang, Soybean GmDREBL Increases Lipid Content in Seeds of Transgenic Arabidopsis, *Sci Rep*, 6 (2016) 34307.
- [49] Q.T. Li, X. Lu, Q.X. Song, H.W. Chen, W. Wei, J.J. Tao, X.H. Bian, M. Shen, B. Ma, W.K. Zhang, Y.D. Bi, W. Li, Y.C. Lai, S.M. Lam, G.H. Shui, S.Y. Chen, J.S. Zhang, Selection for a Zinc-Finger Protein Contributes to Seed Oil Increase during Soybean Domestication, *Plant Physiol*, 173 (2017) 2208-2224.