





# Complete Genome Sequence of *Citricoccus* sp. Strain SGAir0253, Isolated from Indoor Air in Singapore

Lakshmi Chandrasekaran,<sup>a</sup> Daniela I. Drautz-Moses,<sup>a</sup> Akira Uchida,<sup>a</sup>  Rikky W. Purbojati,<sup>a</sup> Anthony Wong,<sup>a</sup> Kavita K. Kushwaha,<sup>a</sup> Alexander Putra,<sup>a</sup> Nicolas E. Gaultier,<sup>a</sup> Balakrishnan N. V. Premkrishnan,<sup>a</sup> Cassie E. Heinle,<sup>a</sup> Vineeth Kodengil Vettath,<sup>a</sup>  Ana Carolina M. Junqueira,<sup>b</sup> Stephan C. Schuster<sup>a</sup>

<sup>a</sup>Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological University, Singapore

<sup>b</sup>Departamento de Genética, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

**ABSTRACT** *Citricoccus* sp. strain SGAir0253 was isolated from indoor air collected in Singapore. Its genome sequence was assembled using single-molecule real-time sequencing. It comprises one chromosome of 3.32 Mb and two plasmids of 137 kb and 99 kb. The genome consists of 2,950 protein-coding genes, 49 tRNAs, and 9 rRNAs.

Members of the bacterial genus *Citricoccus* are Gram-positive, nonmotile bacteria classified in the phylum *Actinobacteria*, family *Micrococcaceae* (1). The first reported draft genome sequence of a *Citricoccus* sp. was of an isolate from the Cuatro Ciénegas Basin (CCB) in Coahuila, Mexico (2). The members of this genus have been isolated and reported from various environments, such as marine macroalgae (3), mold-colonized walls (4), saline wastewater bioreactors (5), wastewater treatment plants (6), medieval wall paintings (1), desert soil (7), and human skin (8). There are no previous reports indicating the presence of *Citricoccus* spp. in air samples, and therefore, this work provides the first evidence that members of this genus can also be found in air.

The strain SGAir0253 was isolated from an indoor air sample collected in Singapore (global position system coordinates, 1.304842°N, 103.791614°E) using the Andersen single-stage impactor (SKC, USA). The air was impacted onto marine agar (Becton, Dickinson, USA), and further isolation of colonies was carried out by culturing onto Trypticase soy agar (TSA) at 30°C. A single colony was grown in lysogeny broth (Becton, Dickinson) at 30°C overnight and subjected to DNA extraction using the Wizard genomic DNA purification kit (Promega, USA), according to the manufacturer's protocol. Library preparation was performed with the SMRTbell template prep kit 1.0 (Pacific Biosciences), followed by single-molecule real-time (SMRT) sequencing on the Pacific Biosciences RS II platform.

For the following analysis, default parameters were used for all software unless stated otherwise. Quality control of raw reads was done with PreAssembler filter v1 within the Hierarchical Genome Assembly Process (HGAP) version 3 protocol (9) implemented in the PacBio SMRT Analysis 2.3.0 package. HGAP was also used for the *de novo* assembly of 78,591 subreads, followed by polishing with Quiver (9). The consensus assembly generated three contigs (Table 1). All three contigs were circularized using Circlator 1.1.4 (10).

Taxonomic identification was performed using average nucleotide identity (ANI) analysis (11) against a database of bacterial refseq genomes that was created using a text filter for "type, synonymtype, proxytype," which showed 82% identity with *Tepidibacter formicigenes*. Phyla-AMPHORA (12) was run using MarkerScanner.pl with an added -DNA flag and MarkerAlignTrim.pl with options -With reference and -output-Format phylip. Phylotyping.pl was run with default parameters, which showed 89.4%

**Citation** Chandrasekaran L, Drautz-Moses DI, Uchida A, Purbojati RW, Wong A, Kushwaha KK, Putra A, Gaultier NE, Premkrishnan BNV, Heinle CE, Vettath VK, Junqueira ACM, Schuster SC. 2019. Complete genome sequence of *Citricoccus* sp. strain SGAir0253, isolated from indoor air in Singapore. *Microbiol Resour Announc* 8:e00606-19. <https://doi.org/10.1128/MRA.00606-19>.

**Editor** Christina A. Cuomo, Broad Institute

**Copyright** © 2019 Chandrasekaran et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stephan C. Schuster, [scschuster@ntu.edu.sg](mailto:scschuster@ntu.edu.sg).

L.C. and D.I.D.-M. contributed equally to this work.

**Received** 30 May 2019

**Accepted** 23 August 2019

**Published** 12 September 2019

**TABLE 1** Assembly statistics for *Citricoccus* sp. strain SGAir0253

Contig	Length (bp)	Coverage (fold)	G+C content (%)	GenBank accession no.
Chromosome	3,323,969	202	73.8	CP039424
Plasmid 1	137,948	221	67.4	CP039425
Plasmid 2	99,421	175	68.4	CP039426

identity to *Micrococcus luteus*. As the identity was below the threshold for species-level identification for ANI results, we also performed 16S identification by extracting the full-length 16S rRNA gene sequence with Barnap 0.7 (13). The BLASTn (14) alignment of the 16S rRNA gene sequence to the Silva database resulted in 98.7% identity to *Citricoccus* sp. strain CH26A. Hence, only genus-level assignment was done for this strain.

The NCBI Prokaryotic Genome Annotation Pipeline (PGAP) version 4.2 (15) was used for genome annotation. A total of 3,134 genes were predicted, comprising 2,950 protein-coding genes (PCGs), 9 rRNA operons (3 each of 5S, 16S, and 23S rRNAs), 49 tRNAs, 3 noncoding RNAs, and 123 pseudogenes. Annotation for plasmid 1 predicted proteins carrying antibiotic resistance (class A beta-lactamase, major facilitator superfamily [MFS], and tetracycline [TetR]), as well as protein CopG, which is responsible for copy number regulation of the plasmid. Plasmid 2 contains metal resistance proteins (copper and arsenic). Both plasmids also contain the plasmid mobilization relaxosome protein MobC and a ParA family protein which is essential for plasmid partition.

**Data availability.** The complete genome sequences of *Citricoccus* sp. strain SGAir0253 and its plasmids are available in DDBJ/EMBL/GenBank under accession numbers CP039424, CP039425, and CP039426 and SRA accession number SRR9043820.

#### ACKNOWLEDGMENT

This work was supported by a Singapore Ministry of Education Academic Research Fund tier 3 grant (MOE2013-T3-1-013).

#### REFERENCES

- Altenburger P, Kampfer P, Schumann P, Steiner R, Lubitz W, Busse H-J. 2002. *Citricoccus muralis* gen. nov., sp. nov., a novel actinobacterium isolated from a medieval wall painting. *Int J Syst Evol Microbiol* 52: 2095–2100. <https://doi.org/10.1099/00207713-52-6-2095>.
- Hayano-Kanashiro C, López-Arredondo DL, Cruz-Morales P, Alcaraz L-D, Olmedo G, Barona-Gómez F, Herrera-Estrella L. 2011. First draft genome sequence of a strain from the genus *Citricoccus*. *J Bacteriol* 193: 6092–6093. <https://doi.org/10.1128/JB.06045-11>.
- Leiva S, Alvarado P, Huang Y, Wang J, Garrido I. 2015. Diversity of pigmented Gram-positive bacteria associated with marine macroalgae from Antarctica. *FEMS Microbiol Lett* 362:fnv206. <https://doi.org/10.1093/femsle/fnv206>.
- Schäfer J, Martin K, Kampfer P. 2010. *Citricoccus parietis* sp. nov., isolated from a mould-colonized wall and emended description of *Citricoccus alkalitolerans* Li et al. 2005. *Int J Syst Evol Microbiol* 60:271–274. <https://doi.org/10.1099/ijs.0.012567-0>.
- Meng F-X, Yang X-C, Yu P-S, Pan J-M, Wang C-S, Xu X-W, Wu M. 2010. *Citricoccus zhacaiensis* sp. nov., isolated from a bioreactor for saline wastewater treatment. *Int J Syst Evol Microbiol* 60:495–499. <https://doi.org/10.1099/ijs.0.011635-0>.
- Nielsen MB, Kjeldsen KU, Ingvorsen K. 2011. Description of *Citricoccus nitrophenolicus* sp. nov., a para-nitrophenol degrading actinobacterium isolated from a wastewater treatment plant and emended description of the genus *Citricoccus* Altenburger et al. 2002. *Antonie Van Leeuwenhoek* 99:489–499. <https://doi.org/10.1007/s10482-010-9513-6>.
- Li W-J, Chen H-H, Zhang Y-Q, Kim C-J, Park D-J, Lee J-C, Xu L-H, Jiang C-L. 2005. *Citricoccus alkalitolerans* sp. nov., a novel actinobacterium isolated from a desert soil in Egypt. *Int J Syst Evol Microbiol* 55:87–90. <https://doi.org/10.1099/ijs.0.63237-0>.
- Ndiaye C, Bassene H, Cadoret F, Raoult D, Lagier JC, Sokhna C. 2018. '*Citricoccus massiliensis*' sp. nov., a new bacterial species isolated from human skin by culturomics. *New Microbes New Infect* 23:83–85. <https://doi.org/10.1016/j.nmni.2018.02.004>.
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Shen L, Liu Y, Xu B, Wang N, Zhao H, Liu X, Liu F. 2017. Comparative genomic analysis reveals the environmental impacts on two Arctic-bacter strains including sixteen Sphingobacteriaceae species. *Sci Rep* 7:2055. <https://doi.org/10.1038/s41598-017-02191-4>.
- Wang Z, Wu M. 2013. A phylum-level bacterial phylogenetic marker database. *Mol Biol Evol* 30:1258–1262. <https://doi.org/10.1093/molbev/mst059>.
- Seemann T. 2013. Barnap 0.7: rapid ribosomal RNA prediction.
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402. <https://doi.org/10.1093/nar/25.17.3389>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.