Encapsulation of probiotics for increased thermal stability for storage and increased viability passing through the gastro-intestinal tract

Li, Aaron Zongwei

2020

Li, A. Z. (2020). Encapsulation of probiotics for increased thermal stability for storage and increased viability passing through the gastro-intestinal tract. Master's thesis, Nanyang Technological University, Singapore.

<https://hdl.handle.net/10356/143784>

<https://doi.org/10.32657/10356/143784>

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Downloaded on 26 Apr 2023 22:25:18 SGT

**NANYANG
TECHNOLOGICAL
UNIVERSITY**

SINGAPORE

**ENCAPSULATION OF PROBIOTICS FOR INCREASED THERMAL STABILITY
FOR STORAGE AND INCREASED VIABILITY PASSING THROUGH THE
GASTRO-INTESTINAL TRACT**

LI ZONGWEI AARON

SCHOOL OF CHEMICAL AND BIOMEDICAL ENGINEERING

2020

**ENCAPSULATION OF PROBIOTICS FOR INCREASED THERMAL STABILITY
FOR STORAGE AND INCREASED VIABILITY PASSING THROUGH THE
GASTRO-INTESTINAL TRACT**

LI ZONGWEI AARON

SCHOOL OF CHEMICAL AND BIOMEDICAL ENGINEERING

A thesis submitted to the Nanyang Technological University
in partial fulfilment of the requirement for the degree of
Master of Engineering

2020

Statement of Originality

I hereby certify that the work embodied in this thesis is the result of original research, is free of plagiarised materials, and has not been submitted for a higher degree to any other University or Institution.

07/07/20

Date

Li Zongwei Aaron

Supervisor Declaration Statement

I have reviewed the content and presentation style of this thesis and declare it is free of plagiarism and of sufficient grammatical clarity to be examined. To the best of my knowledge, the research and writing are those of the candidate except as acknowledged in the Author Attribution Statement. I confirm that the investigations were conducted in accord with the ethics policies and integrity standards of Nanyang Technological University and that the research data are presented honestly and without prejudice.

07 Jul 2020

.....
Date

C. W. Ning

.....
Professor Chen Wei Ning, William

Authorship Attribution Statement

Please select one of the following; *delete as appropriate:

*(A) This thesis **does not** contain any materials from papers published in peer-reviewed journals or from papers accepted at conferences in which I am listed as an author.

07/07/20

Date

Li Zongwei Aaron

Acknowledgement

I would like to take this opportunity to show my gratitude to my supervisor, Professor Chen Wei Ning, William for giving me the opportunity to undertake the Master of Engineering under his mentorship. Prof. William has always been very helpful and supportive to me and have provided me with many opportunities. He was always there during the entire duration of the course and would be there whenever I needed him. He also provided me with the much-appreciated guidance, nurturing me and allowing me to mature both in a personal and professional capacity. He is also always supportive of me, giving me the liberty to explore and drive the project and research in my own way, but also constantly guiding me to ensure that I am on the right track. He also given me many different opportunities and experiences which allowed me to constantly improve myself better and mould myself into a all-rounded researcher. I am also very grateful to Dr Jaslyn Lee for all her encouragement and guidance throughout the course of my studies. She was a wonderful mentor and a great colleague and friend whom I can approach if I needed any assistance regarding my project. I would also like to take this chance to thank my colleagues: Mr Mok Wai Kit, Ms Yeo Ying Tong, Mr Tan Yong Xing for all their support and help throughout my studies. Also, I thank my colleagues Salmah, Audrey and Serene from the admin department for their help and support as well.

List of Figures

Figure 1: Schematic Diagram of a Spray Dryer

Figure 2: Schematic diagram of Maillard Reaction

Figure 3: *L. Plantarum* incubated anaerobically in BD GasPak™ EZ container system

Figure 4: BUCHI Mini Spray Dryer B-290

Figure 5: Overview of simulated in vitro digestion system

Figure 6: Particle size distribution a: pectin, b: whey protein, c: WP2, d: WP1

Figure 7: Viability of *L. Plantarum* before and after spray drying

Figure 8: SEM image of encapsulated probiotics. a,b: Free *L. Plantarum*, c: WP1, d: WP2

Figure 9: SEM image of encapsulated probiotics at Week 8 of storage. a,b: WP1, c,d: WP2

Figure 10: Viability of *L. Plantarum* in storage (Room Temperature)

Figure 11: Viability of *L. Plantarum* after 15 and 30 min of pasteurization at 85°C

Figure 12: Viability of *L. Plantarum* before and after in vitro digestion

List of Tables

Table 1: Microorganisms used as probiotics

Table 2: Stock Solution Composition for Simulated Digestion Fluids

Table 3: Viable cell count of *L. Plantarum* before and after spray drying

Table of Contents

1. Introduction	1
1.1 Hypothesis/Problem Statement	1
1.2 Objective and Scope.....	2
2. Literature Review	2
2.1 Probiotics	2
2.1.1 Health Benefits of Probiotics	3
2.1.2 Probiotics Mechanism of Action	3
2.1.3 Lactic Acid Bacteria	4
2.1.4 Lactobacillus Plantarum	5
2.2 Gastrointestinal Tract	5
2.3 Encapsulation	7
2.3.1 Spray Drying.....	7
2.3.2 Encapsulation materials.....	9
2.3.3 Pectin	9
2.3.4 Whey Protein	10
2.4 Maillard Reaction	11
3. Materials and Methods.....	13
3.1 Culture of Probiotics	13
3.2 Enumeration of Viable Cells.....	13
3.3 Maillard Conjugation of Encapsulation materials.....	14
3.4 Conjugation Efficiency.....	14
3.5 Size distribution of whey protein-pectin conjugates.....	15
3.6 Encapsulation through spray-drying of <i>L.Plantarum</i>	15
3.7 Shelf-life testing of microencapsulated <i>L.Plantarum</i>	16
3.7 Scanning Electron Microscopy (SEM) imaging.....	16
3.8 Viability of <i>L.Plantarum</i> after pasteurization.....	16
3.9 In-vitro Gastrointestinal Digestion.....	16
3.10 Statistical Analysis.....	18
4. Results and Discussion	19
4.1 Characterization of Encapsulants	19
4.2 Effects of Spray Drying on viability of <i>L.Plantarum</i>	20
4.3 Viability of Encapsulated <i>L.Plantarum</i> during storage.....	23
4.4 Protective effects of encapsulation on <i>L.Plantarum</i> during pasteurization	24
4.5 Viability of Encapsulated <i>L.Plantarum</i> after in vitro digestion	24
5. Conclusion	25

6. References	26
---------------------	----

Abstract

Probiotics are defined as living bacteria that, when administered in adequate amounts, confer a health benefit on the host. (Hill, Guarner, Reid, Gibson, & Merenstein, 2014) Therefore, more and more people are beginning to explore the use of probiotics supplements for their proposed health benefits. However, probiotics being living bacteria, are susceptible to temperature and environmental fluctuations. In order to be able to provide a certain health benefit to the host, these live bacteria should maintain a high enough level of viability throughout the food processing procedures, during the storage duration and while passing through the gastrointestinal tract. (Chávarri, Marañón, & Villarán, 2012) One of the most common food processing methods for inactivation of harmful pathogens is through the use of heat. At present, there are few probiotics that are stable at high temperatures, thereby increasing the urgency to improve their heat resistance either through identification of new heat-stable strains or to develop novel delivery systems to protect them. (Solanki et al., 2013) Hence, much interest has been gathered on using encapsulation as a means of delivery and protection of the probiotics to able survive the food processing procedures and eventually be viable in the gastrointestinal tract.

In this study, the use of Maillard conjugation of whey protein with citrus pectin as encapsulants is explored for the encapsulation of *Lactobacillus plantarum* (DSM 12028) for increased thermal stability for storage and increased viability passing through the Gastrointestinal Tract. Viability of *L. Plantarum* encapsulated in 2% whey protein and 1% pectin (WP1) in room temperature (25°C) storage after 60 days was found to be 7 log units higher than free *L. Plantarum* cells. The encapsulation of WP1 also resulted in an increase in viability through the upper gastrointestinal tract with a log (CFU/g) of 8.15 compared to 4.90 in free unencapsulated *L. Plantarum* cells. WP1 also showed a high heat tolerance in pasteurization with only a 0.33 log units reduction after 30 minutes at 85°C.

1. Introduction

1.1 Hypothesis/Problem Statement

Probiotics have been consumed in many forms over the last several centuries across many different civilizations for their proposed health benefits. However, probiotics being living bacteria, are susceptible to temperature and environmental fluctuations. In order to be able to provide a certain health benefit to the host, these live bacteria should maintain a high enough level of viability throughout the food processing procedures, during the storage duration and while passing through the gastrointestinal tract. (Chávarri et al., 2012) One of the most common food processing methods for inactivation of harmful pathogens is through the use of heat. At present, there are few probiotics that are stable at high temperatures, thereby increasing the urgency to improve their heat resistance either through identification of new heat-stable strains or to develop novel delivery systems to protect them. (Solanki et al., 2013) Hence, much interest has been gathered on using encapsulation as a means of delivery and protection of the probiotics to able survive the food processing procedures and eventually be viable in the gastrointestinal tract. Many different methods of encapsulation of probiotics has since been explored; forming of probiotic emulsions using different encapsulants, (Qi, Liang, Yun, & Guo, 2019) forming microcapsules and microgels using extrusion method, (Heidebach, Först, & Kulozik, 2012) freezing drying of probiotic cells, (Chalat Santivarangkna, Kulozik, & Foerst, 2007) and spray drying of probiotic cells. (De Castro-Cislaghi, Carina Dos Reis, Fritzen-Freire, Lorenz, & Sant'Anna, 2012) Although much research has been done in the field of probiotic encapsulation many of the studies still present some inherent problems:

- The inability to overcome both the problem of thermal stability and navigation of gastrointestinal transit
- Difficulty for incorporation into existing food production due to the shape, size, form etc. of encapsulated particles
- Difficulty for scale up production for commercialization

Hence in this study, the use of Maillard conjugation of whey protein with citrus pectin as encapsulants is explored for the encapsulation of *Lactobacillus plantarum* (DSM 12028) for increased thermal stability for storage and increased viability passing through the Gastrointestinal Tract.

1.2 Objective and Scope

- Encapsulation of *Lactobacillus plantarum* (DSM 12028) using Maillard conjugation of whey protein and citrus pectin
- To test the conjugation efficiency of whey protein and citrus pectin conjugation
- To test the viability of *Lactobacillus plantarum* (DSM 12028) after encapsulation
- To test the viability of encapsulated *Lactobacillus plantarum* (DSM 12028) in storage at room temperature (25°C)
- To test the viability of *Lactobacillus plantarum* (DSM 12028) after pasteurization at (85°C)
- To test the viability of encapsulated *Lactobacillus plantarum* (DSM 12028) after passing through gastrointestinal tract

2. Literature Review

2.1 Probiotics

The term probiotics is developed from the Greek phrase “pro bios” which means “for life”. This would mean that the food or substance promotes improved health in the host. Since ancient times, people have been consuming probiotics in the form of fermented food across many different civilizations. Calpis (Japanese), Kaymak (Turkish), Kefir (Russian), Kimchi (Korean) and Tofu (Southeast Asian) are some examples for such food that have been consumed. (Bagchi, 2014) It was not until the early 1900s that Elie Metchnikoff hypothesized that lactic acid bacteria (LAB) can offer health benefits such as promote longevity through the modification of gut microbiota and replacing harmful gut microbes with useful ones. Following that, Henry Tissier postulated that oral administration of useful microbes (*Bifidobacteria*) to patients with diarrhoea can help improve the conditions through restoring of the gut flora. (Merenstein & Salminen, 2017) Since there, much more research has been done with regards to probiotics and in 2001, Food and Agriculture Organization of the United Nations (FAO) together with World Health Organization (WHO) came up with a definition that is widely used and recognized today: “Live microorganisms which when administered in adequate amounts confer a health benefit on the host”. (Morelli & Capurso, 2012) Through the years of research, scientist have found that probiotics have the potential to provide many different health benefits. Some of these health benefits could include : anti-pathogenic, anti-cancer, anti-inflammatory, anti-diabetic and even angiogenic activities. (Kerry et al., 2018) Many of these benefits however, are specific to the different genus, species and strains.

2.1.1 Health Benefits of Probiotics

Probiotics coming from fermented food such as kefir, kimchi, tofu, etc. have been ingested by humans for many years in the belief that they can provide health benefits. In recent years, as more research has been done into understanding our gut microbiota, many studies have postulated that probiotics are playing crucial roles in modulation of gastrointestinal immunological, and respiratory functions. (Floch et al., 2011) Due to the constant scientific evaluation in this area, there is now strong evidence that the use of probiotics can help in treating and preventing certain diseases. (Boyle, Robins-Browne, & Tang, 2006) Probiotics have been said to provide a number of health benefits which includes: a) prevention of gastrointestinal pathogens through the maintenance of a healthy intestinal flora, ; b) strengthening of the immune system; c) lowering of blood pressure and reducing blood cholesterol; d) having anti-carcinogenic properties; e) improved utilization of nutrients; f) prevention of inflammation; g) production of antimicrobial compounds; h) treatment and prevention of diarrhoea; i) improvement of lactose tolerance; j) treatment of urinary tract infections and respiratory infections have also been documented (D'Aimmo, Modesto, & Biavati, 2007; Kechagia et al., 2013; Marco, Pavan, & Kleerebezem, 2006; Parvez, Malik, Ah Kang, & Kim, 2006; Tripathi & Giri, 2014)

2.1.2 Probiotics Mechanism of Action

Over the years, there have been many studies done to try and understand the mechanisms of action of probiotics. However, the exact mechanisms by which how the effects are exerted on the host is not completely understood. (Harish & Varghese, 2006) According to (Hemaiswarya, Raja, Ravikumar, & Carvalho, 2013), how effective the probiotics is on the host depends strongly on their adherence and how well they can colonize of the gut in order to improve the host's immune system. Different species of probiotics differ significantly in their mechanism of action; None of the probiotic species found till date is able to exhibit all mechanisms of action to prevent or treat all the different diseases. (O'Hara & Shanahan, 2007; Oelschlaeger, 2010) Some of the proposed mechanisms of probiotic actions consist of: a) the production of bacteriocins which provides an antimicrobial effect; b) the production of lactic acid, and short-chain fatty acids which brings down the pH in the gut lumen to inhibit pathogens colonization; c) the activation of tight junction proteins for maintenance of the epithelial barrier to prevent the development of a leaky intestine; d) the prevention of inflammation and apoptosis of the lining intestinal epithelial cells (Harish & Varghese, 2006; Michail, 2005; Ng, Hart, Kamm, Stagg, & Knight, 2009; Sarowska, Choroszy-Król,

Regulska-Ilow, Frej-Madrzak, & Jama-Kmiecik, 2013; Sherman, Ossa, & Johnson - Henry, 2009)

2.1.3 Lactic Acid Bacteria

In order for the probiotic strains to be used, they are required to be identified by their genus, species and strain designation and deposited into an international culture collection. Till date, there have been many different probiotics strains including different types of bacteria and even yeast. (Anadón, Martínez-Larrañaga, Ares, & Martínez, 2016)

Probiotic Microorganisms	Species Involved	References
<i>Lactobacillus</i>	<i>L. Plantarum</i> , <i>L. paracasei</i> , <i>L. acidophilus</i> , <i>L. casei</i> , <i>L. rhamnosus</i> , <i>L. crispatus</i> , <i>L. gasseri</i> , <i>L. reuteri</i> , <i>L. bulgricus</i>	(Dixit, Wagle, & Vakil, 2016; Rault, Béal, Ghorbal, Ogier, & Bouix, 2007)
<i>Streptococcus</i>	<i>S. thermophilus</i> , <i>S. salivarius</i> , <i>S. sanguis</i> , <i>S. oralis</i> , <i>S. mitis</i>	(Arora, Singh, & Sharma, 2013)
<i>Bifidobacterium</i>	<i>B. longum</i> , <i>B. catenulatum</i> , <i>B. breve</i> , <i>B. animalis</i> , <i>B. bifidum</i>	(Kurmán & Rasic, 1991; Sakata et al., 2002)
<i>Bacillus</i>	<i>B. coagulans</i> , <i>B. subtilis</i> , <i>B. laterosporus</i>	(Nguyen et al., 2016)
<i>Lactococcus</i>	<i>L. lactis</i>	(Eid et al., 2016)
<i>Enterococcus</i>	<i>E. faecium</i>	(Lund, Adamsson, & Edlund, 2002)
<i>Pediococcus</i>	<i>P. acidilactici</i>	(Sornplang & Piyadeatsoontorn, 2016)
<i>Propionibacterium</i>	<i>P. jensenii</i> , <i>P. freudenreichii</i>	(Zárate, 2012)
<i>Peptostreptococcus</i>	<i>P. productus</i>	(Kontula et al., 1998)
<i>Escherichia coli</i>	<i>E. coli</i> Nissle	(Ukena et al., 2007)
<i>Saccharomyces</i>	<i>S. boulardii</i>	(Chen et al., 2013)

Table 1: Microorganisms used as probiotics

Presently, the most common and widely researched probiotic microbes are Lactic Acid Bacteria (LAB) which include *Lactobacillus*, *Streptococcus* and *Bifidobacterium*. (Vanderhoof, 2000) These bacteria produce Lactic Acid as one of the major end products. Amongst those, *Lactobacillus* is known to be the largest genus of the group. *Lactobacillus* comes from the phylum *Firmicutes*, class of *Bacilli*, order of *Lactobacillales* and family of *Lactobacillaceae*. They are rod shaped bacteria which are anaerobic, Gram positive and non-spore forming. (M. De Angelis, 2016)

2.1.4 Lactobacillus Plantarum

One of the most studied species of the genus would be *Lactobacillus plantarum* as many strains within the species have found to have distinct probiotic properties. They are said to produce many desirable health benefits including having antipathogenic properties, antiallergic properties, ability to reduce cholesterol and even improving brain function in autistic children. (Adlerberth et al., 1996; Kumar et al., 2016; Puertollano et al., 2008; Umbrello & Esposito, 2016; Zendeboodi, Khorshidian, Mortazavian, & da Cruz, 2020) Given the myriad of beneficial properties, this allows for *Lactobacillus plantarum* to be a promising candidate for use as probiotics. However, the guidelines from FAO/WHO emphasized the need for probiotic strains to be viable throughout the gastrointestinal tract to be able to confer the proposed health benefits upon entering the large intestine, regardless the delivery mode applied. (Bosnea, Moschakis, & Biliaderis, 2017; Morelli & Capurso, 2012)

2.2 Gastrointestinal Tract

As it is critical to ensure that probiotics are viable through to their destination of action in order to provide the proposed benefits, understanding of the barriers and harsh conditions they face through the gastrointestinal transit becomes critical. After being ingested, probiotics face many different harsh conditions throughout the Gastrointestinal Tract; which includes the mouth, the stomach as well as the small intestine.

In the mouth, probiotics are mainly affected by mineral salts, mucin and salivary amylase. However, as the pH within the mouth is relatively neutral, pH 6-7, which is similar to that of the colon where the probiotics are conditioned for. (Yeung, Arroyo-Maya, McClements, & Sela, 2016) The transit time within the mouth is also short, hence the harm caused to the probiotics is not as significant.

As they proceed into the stomach, the gastric fluids which are highly acidic, reduces the pH as low as pH 1-3 causing the conditions to be increasingly challenging. (Lankaputhra, 1995)

This low pH causes a disruption in the cytoplasmic pH of the probiotics and in turn reduces the glycolytic enzyme activity and reduces the survivability of many probiotics under these conditions. (Cotter & Hill, 2003; Sarao & Arora, 2017) Within the stomach, there are also other factors such as high salt content, enzymatic activity and mechanical churning which also reduces the viability of probiotics. (González-Ferrero, Irache, & González-Navarro, 2018; Surono, Verhoeven, Verbruggen, & Venema, 2018; Yeung et al., 2016)

Upon reaching the small intestine, probiotics are affected by the digestive enzymes and bile acids present. According to Hamner et. al (2013), (Hamner, McInnerney, Williamson, Franklin, & Ford, 2013) bile acids possess some antibacterial properties which could disrupt cell membranes and damage bacteria DNA. (Begley, Hill, & Gahan, 2006) mentioned however, that some probiotics could circumvent this through production of bile salt hydrolases (BSHs) to unconjugated bile acids. However, the exact mechanisms in which probiotics and bile acids interact remains unclear and further research needs to be done to better understand these interactions. (Jia & Xie, 2018)

According to the guidelines from FAO/WHO, even before the probiotics weather the conditions of the gastrointestinal tract, there should be a “minimum therapeutic” level of live probiotic microbes in a food product of at least 10^6 CFU/g of live cells during the entire shelf-life of the product. (Morelli & Capurso, 2012) This means to say that, to account for the loss in the gastrointestinal tract as well, the conditions for an effective probiotic product would be much stricter. Many reports have also mentioned, that probiotics survivability in many products which uses free probiotic cells is very poor. (de Vos, Faas, Spasojevic, & Sikkema, 2010) Some of these probiotic products in the market encapsulate them in a form of supplement such as tablet or capsules. However, although these formulations are large enough to encapsulate a large amount of probiotics, they are too large to pass directly through the pyloric sphincter which leads to the small intestine. Hence, when the encapsulated probiotics are released into the stomach, they become susceptible to the unforgiving environmental conditions there. (Yao et al., 2020) Hence, in order to ensure that all the necessary conditions for probiotic effectiveness are met, methods to protect the probiotic strains through the food processing procedures, storage conditions and the harsh environment of the gastrointestinal tract have been actively researched. One of the most common and widely researched methods would be to encapsulate the probiotics.

2.3 Encapsulation

Encapsulation is known as either a mechanical or physiochemical process in order to encase a substance, which could be a functional or bioactive compound, in a material in order to protect it, or for further delivery and release. (de Vos et al., 2010) These particles produced could have diameters ranging from a few nanometres to a few millimetres. These results in small particles that are capsules which form a shell or coating to contain active agents or core material. (Chávarri et al., 2012) This technique has seen increased research and interest, due to its applications in the pharmaceutical industry for drug and vaccines delivery. It is also been increasingly used in the food industry for the delivery of bioactive molecules (e.g. carotenoids, fatty acids, vitamins and antioxidants) or live cells such as probiotics. (Nedovic, Kalusevic, Manojlovic, Levic, & Bugarski, 2011) Encapsulation is a versatile technique that could be used for many purposes. It can be used to mask undesirable taste, odour or colouration which could affect the product performance. It can also be used to prevent interactions and reactions between the active food ingredients to be delivered and other ingredients or food components. Apart of that, it can also be used for controlled release of the food ingredient. (Sobel, Versic, & Gaonkar, 2014) In the case of probiotics, encapsulation is utilized to protect them from several factors:

- Processing conditions (temperature, oxidation, shear, etc.)
- Storage conditions (oxygen, moisture, temperature, packaging, etc.)
- Degradation by Gastrointestinal Tract (enzymes, low pH, bile acids, etc.)

This allows the probiotics to be released in a biologically active and viable state in the colon, (Picot & Lacroix, 2004) (Ding & Shah, 2007) while keeping crucial factors such as preserving the aroma and taste, maintaining the nutritional value, and keeping product appearance consistent. (Parra Huertas, 2010) There are several methods of encapsulation. These methods include extrusion, emulsification, coacervation and spray drying. (Rathore, Desai, Liew, Chan, & Heng, 2013) Among those, the most commonly used microencapsulation technique in the food industry due its flexibility, cost effectiveness and ability to be scaled up easily, would be spray drying. (Pu, Bankston, & Sathivel, 2011)

2.3.1 Spray Drying

Spray drying relies on the use of rapid dehydration to transform a feeding suspension, emulsion, solution, or dispersion into small dried particles by spraying the feed onto a heated surface or medium. Depending on the properties of the feed to be sprayed, the dried particles could appears as powders, granules or agglomerates. (Patel, Patel, & Suthar, 2009) Since the

1940s, pharmaceutical industry has successfully utilized spray drying for production of drug substances such as antacids, vitamins, antibiotics and analgesics. Then, from the late 1950s onwards, spray-drying was picked up by the food industry to encapsulate various oils to prevent them from degradation and oxidation and to produce powders from liquids for easier storage and extending shelf-life. (Phisut, 2012) Spray drying is presented as a viable method for drying of heat-sensitive active biological substances, such as live cells, proteins and even enzymes, with very minimal activity losses. (Idham, Muhamad, & Sarmidi, 2012) The process of microencapsulation by spray drying begins with the formation of an emulsion or suspension of coating or shell material with the core material. Then the emulsion or suspension is fed into the atomizer where it is sprayed out as tiny droplets into the drying chamber. This chamber is circulated with hot dry air which rapidly dehydrates and evaporate the droplets leaving the core material entrapped in the coating material as small dried particles. The dried particles are then sent through the cyclone and collected in a container. (Khuenpet, Charoenjarasrerk, Jaijit, Arayapoonpong, & Jittanit, 2016)

Figure 1: Schematic Diagram of a Spray Dryer (Khuenpet et al., 2016)

The utilization of spray drying for probiotic bacteria in the food industry has been around for decades and is used predominantly by the dairy industry. (Huang et al., 2017) The technique involves dispersing the probiotic cells into the encapsulating polymer solution and atomized into the spray drying chamber to evaporate the solvents to form the encapsulated particles. (De Castro-Cislaghi et al., 2012) In spray drying, bacterial cultures usually die from exposure

to heat, oxidative and osmotic stress, resulting from the high temperature during the dehydration. (Paéz et al., 2012) According to (Peighambaroust, Tafti, & Hesari, 2011), the main reasons for the loss of viability of probiotic cells is due to the combination of temperature and time, and that the inlet air temperature has little impact on probiotic death. (C Santivarangkna, Kulozik, & Foerst, 2008) proposed that outlet air temperature is the key parameter which affects the survivability of probiotic cultures during spray drying, and this parameter is dependent on the other parameters such as air flow rate, inlet air temperature, aspiration droplet size, and feed pump rates. (Chalat Santivarangkna et al., 2007) This claim that lower outlet air temperature results in an increased survival of probiotic cells is also supported by other researchers. (Ananta, Volkert, & Knorr, 2005; Desmond, Ross, O'callaghan, Fitzgerald, & Stanton, 2002) During a study done on several strains of probiotics isolated from kefir, (Golowczyc, Silva, Teixeira, De Antoni, & Abraham, 2011) found that *L.Plantarum* exhibited the highest survival rate across different outlet temperatures (70, 75, 80, 85°C). A similar results was found by (Barbosa et al., 2015; Iaconelli et al., 2015) and they concluded that sensitivity to different drying methods is bacterial strain specific.

2.3.2 Encapsulation materials

The selection of a wall coating material for microencapsulation is of paramount importance to obtain a stable and effective encapsulation. Encapsulation materials are typically selected based on the physicochemical properties such as melting/glass transition temperature, crystallinity, solubility, molecular weight and emulsifying properties. (Gharsallaoui, Roudaut, Chambin, Voilley, & Saurel, 2007) Many researchers such as (Agnihotri, Mishra, Goda, & Arora, 2012; Anandharamakrishnan & Ishwarya, 2015) emphasize the need to select appropriate wall materials as they determine the physical and chemical properties of the resulting microcapsules. Some of the materials commonly used to encapsulate probiotic cells include different polysaccharides (alginates, pectin, xanthan gum, gum arabic, carrageenan, chitosan, starch, etc.) oligosaccharides (maltodextrin, sucrose and corn syrup), proteins (soy protein, whey protein, casein, albumin and gluten), and lipids (waxes, paraffin, diglycerides, oils, fats, etc) (Das, Ray, Raychaudhuri, & Chakraborty, 2014; De Castro-Cislaghi et al., 2012; Serna-Cock & Vallejo-Castillo, 2013)

2.3.3 Pectin

Pectin is biopolymer which is increasingly used as an encapsulation material. This is due to several desirable properties it has, like superior gelling properties, great binding abilities the ability to stabilize emulsions, (Assadpour, Jafari, & Maghsoudlou, 2017; Gharehbeglou,

Jafari, Hamishekar, Homayouni, & Mirzaei, 2019) Given that it is typically obtained from fruits, it can also be considered as a food safe ingredient. (Colodel, das Graças Bagatin, Tavares, & de Oliveira Petkowicz, 2017) Different types of pectin exhibit different hydrophobicity. Unlike low methoxylated pectin which is usually hydrophilic, high methoxylated pectin is known to be very hydrophobic and interacts well with hydrophobic molecules. This enables pectin to play an important role in the incorporation of hydrophobic drugs, such as fluoroquinolone which is a type of antibiotics, into the encapsulation matrix to allow for controlled release. (Cacicedo et al., 2018) Other than that, pectin is also commonly used for the encapsulation of bioactive ingredients (such as polyphenols, carotenoids etc.) through several methods, such as spray drying, forming of emulsions and coacervation. (Fathi, Martin, & McClements, 2014)

2.3.4 Whey Protein

Whey proteins which are derived from whey, In the food industry, whey proteins are one of the most used food ingredients given their desirable functional properties. They are typically used as stabilizers, in gelling, texturizing and even as emulsifiers for various oil-in-water emulsions. (de Castro et al., 2017; Foegeding, Davis, Doucet, & McGuffey, 2002) The main components in whey protein, β -lactoglobulin and α -lactalbumin, are known to be able to form protective films (wall system), allowing them to be optimal wall material for encapsulation. Whey protein isolate (WPI) may also be formulated into nanoparticles, which could be used to as a medium of encapsulation for several different types of bioactive compounds, such as carotenoids, vitamins, and isoflavonoids etc. (Abbasi, Emam-Djomeh, Mousavi, & Davoodi, 2014; Flores, Singh, & Kong, 2014) (Schuck, Dolivet, Mejean, Hervé, & Jeantet, 2013) also found that through the use of sweet whey as the drying medium, probiotics can be spray dried with close to 100% survival rate. However, the use of whey protein also comes with some drawbacks. Within the food industry, the main limitations for the use of whey protein as a high value ingredient include: a) having poor protein solubility under low pH conditions, which causes phase separation and turbidity of the mixture (Akhtar & Dickinson, 2007); b) once the proteins are hydrolysed, they exhibit poor emulsification properties which limits its utility by causing challenges such as poor emulsion formation ability, instability, and becoming very sticky, resulting in difficulty in spray drying (Singh & Dalgleish, 1998); c) the proteins also tend to be physical instable, causing sedimentation, creaming and aggregation during processing, storage in highly charged environments as well as during thermal processing. (Yadav, Parris, Johnston, Onwulata, & Hicks, 2010) In order to prevent

undesirable effects such as whey protein aggregation, the conjugation of whey proteins to different polysaccharides was explored as a viable method.(Akhtar & Dickinson, 2007; Xu, Wang, Jiang, Yuan, & Gao, 2012; Zhu, Damodaran, & Lucey, 2010) Conjugation is able to successfully modify the functional properties of a myriad of milk protein/peptide-based ingredients, which includes whey protein ingredients as well. In particular, (Setiowati, Saeedi, Wijaya, & Van der Meeren, 2017) found that conjugation of whey protein isolate and low methoxyl pectin significantly improved heat stability.

2.4 Maillard Reaction

Over the years, researchers have explored means of altering properties and improving the functionality of proteins for different industrial applications. One such method is through the use of Maillard Reaction (MR). Maillard reaction has been widely researched for altering and improving the properties of whey protein significantly. (Spotti et al., 2013) Maillard Reaction is a reaction between proteins or amino acids and carbohydrates. The ϵ -amino groups in the proteins covalently bonds with the end carbonyl groups of polysaccharides under heated conditions. This covalent crosslinking of the proteins and carbohydrates results in highly desirable properties for food processing such as increased heat stability and better emulsifying properties. (Kato, 2002; Pellegrino, Van Boekel, Gruppen, Resmini, & Pagani, 1999) According to (Hodge, 1953), Maillard Reaction divided into three stages; namely the early, intermediate and advanced stages.

In the early stage, the free amino group from the protein is bonded to the reducing sugar group in the carbohydrate to form a Schiff base and produces a H_2O molecule. The Schiff base is unstable and would undergo spontaneous rearrangement to form either an Amadori (Aldose) product or a Heyn's (Ketose) product depending on the reducing sugar. (Wrodnigg & Eder, 2001) Then, during the intermediate stage, the Amadori/Heyn's product will undergo several reactions such as dehydration, isomerization, cyclization and condensations, which results in sugar degradation and fragmentation, and amino acids will undergo degradation as well. (Koubaa et al., 2018) The advanced stage on the other hand become much more complex. The reactions that occur become variable and highly dependent on environmental conditions. Although some studies mentioned that some of the products from advanced stage Maillard Reaction have beneficial properties such as antihypertensive antimicrobial and antioxidant properties, (Nooshkam, Babazadeh, & Jooyandeh, 2018; Nooshkam, Varidi, & Bashash, 2019; H.-Y. Wang, Qian, & Yao, 2011) others stated that products can lead to diseases like diabetes and Alzheimer.(Silván, Assar, Srey, Del Castillo, & Ames, 2011)

Figure 2: Schematic diagram of Maillard Reaction (O'Mahony, Drapala, Mulcahy, & Mulvihill, 2017)

Hence, it is advisable to use more superficial Maillard conjugation methods, which is sufficient to improve protein properties such as increased solubility and enhanced thermal stability, to prevent any complications. (de Oliveira, Coimbra, de Oliveira, Zuñiga, & Rojas, 2016) It can be categorized into two different processes; wet and dry heating, both having their own sets of advantages and disadvantages. During dry heating, the rate of reaction can be easily controlled owing to the lower moisture diffusion resulting from having a lower water content. (Nasirpour, Scher, & Desobry, 2006). However, dry heating is usually more tedious and needs a considerably long reaction time with harsher conditions. During the process, formation of insoluble substances might occur as well. Apart from relying on dry heating to conjugate whey protein and polysaccharides, researchers have also managed to form complexes through the wet heating method as well. (W.-q. Wang, Bao, & Chen, 2013) In the wet heating process, there is a retardation of the rate of reaction which is caused by the dilution of the water-soluble reactants given the higher water content. (Nasirpour et al., 2006) The wet heating process is also much faster as it is much less laborious. Similar to the dry heating process, it is also possible to control the reaction degree of wet heating effectively.

However, these might also result in addition interactions such as electrostatic or hydrophobic interactions. (Perusko, Al-Hanish, Velickovic, & Stanic-Vucinic, 2015)

3. Materials and Methods

3.1 Culture of Probiotics

Bacterial strain *Lactobacillus plantarum* (DSM 12028), obtained from the Leibniz Institute DSMZ German Collection of Microorganism and Cell was inoculated into 5mL MRS broth (Sigma Aldrich) and incubated at 37 °C for 48 h which serves as the stock culture. After which, the stock culture was inoculated into a 250mL Erlenmeyer Flask at an optical density of 0.2 at 600nm wavelength, which corresponds to 10^8 cells/mL (Ramos, Thorsen, Schwan, & Jespersen, 2013) and incubated at 37 °C for 48 h. When harvesting, cells were initially centrifuged at 5000g at 4°C for 10 minutes and washed thrice with milliQ water, then the cells were kept at 4°C to be used for encapsulation.

3.2 Enumeration of Viable Cells

Enumeration of *L. Plantarum* was done using the spread plate method. 100 µL of each cell suspension was taken and serially diluted in 900 µL of milliQ water. 100 µL of each dilution (10^{-6} – 10^{-9}) was added into sterile petri dishes which has 15 mL of MRS agar poured into each dish that was cooled and set. A L-shaped spreader is then used to spread the 100 µL suspension across the entire agar plate and incubated anaerobically at 37°C for 48 h in BD GasPak™ EZ container systems and the colonies were counted between 15 and 300 and the number of colony-forming units (CFU) in 1 mL of dilution was determined (CFU = no. of colonies × dilution factor).

Figure 3: *L. Plantarum* incubated anaerobically in BD GasPak™ EZ container system

3.3 Maillard Conjugation of Encapsulation materials

The citrus peel pectin was obtained from Sigma Aldrich, USA and the 15% protein content sweet whey powder was obtained from Jinan Meishubao Biotechnologies, China. The pectin powder was added to 500mL of milliQ water at 2% w/v and mixed for 30 minutes at 60°C until fully dissolved. The sweet whey powder was diluted in 500mL of milliQ water to obtain a 2% w/v whey protein solution. The method for wet heating was adapted from the method proposed by Gao F et. al, 2019 (Gao et al., 2019). Briefly, the whey protein and pectin solution were added together to obtain 2:1, 1:1 whey protein:pectin ratio solutions (WP2, WP1) respectively. Both solutions were adjusted to pH 7.0 and constantly stirred magnetically and maintained at a temperature of 85°C for 48h to allow Maillard conjugation to occur. The solutions were then rapidly cooled to room temperature for use in encapsulation.

3.4 Conjugation Efficiency

Using the O-phthaldialdehyde (OPA) method modified from (Davidov-Pardo, Joye, Espinal-Ruiz, & McClements, 2015), the conjugation efficiency was inversely determined through analysing the free amino groups. By determining the difference in free amino groups, we can determine the change in conjugation. The OPA reagent used was obtained from Sigma Aldrich, USA and the UV Spectrophotometer used was from the NanoDrop 2000 from Thermo Scientific. Briefly, 200 µL of sample solution was mixed with 4 mL of OPA reagent and then heated through a water bath at 35 °C for 2 min. Then, using the UV

Spectrophotometer, the absorbance of the reacted solution was taken at a wavelength of 340 nm. Conjugation Efficiency (CE) was back calculated from the change in the amount of unconjugated free amino groups using the following equation:

$$CE = (1 - A_{\text{After}}/A_{\text{Before}}) \times 100\%$$

Where A_{After} and A_{Before} were the amount of unconjugated amino groups found in the solution before and after conjugation respectively. A control group of 2% whey protein solution, was also included.

3.5 Size distribution of whey protein-pectin conjugates

Size distribution of the whey protein-pectin conjugates were measured through dynamic light scattering using the Malvern Zetasizer Nano SZ, UK. and the mean particle size was expressed as d50.

3.6 Encapsulation through spray-drying of *L. Plantarum*

200mL of both WP2 and WP1 solutions were inoculated with *L. Plantarum*, and stirred constantly for 30 minutes with the CFU in each at 7.6×10^9 CFU/mL of WP solution. A control group with 7.6×10^9 CFU/mL of *L. Plantarum* in 200mL of milliQ water was also added. BUCHI Mini Spray Dryer B-290, Switzerland was used to process samples at inlet temperature of 120°C, air flow rate of 35m³/h, compressed air pressure of 6 bar, feed rate of 7.5mL/min and outlet temperature of between 55-60°C. All the solutions were stirred constantly throughout the spray drying process to ensure even mixing.

Figure 4: BUCHI Mini Spray Dryer B-290

3.7 Shelf-life testing of microencapsulated *L.Plantarum*

The powder was collected after spray-drying and stored at 25°C for 60 days and enumerated fortnightly. To enumerate the surviving *L.Plantarum* bacteria, 200mg of encapsulated *L.Plantarum* powder was dissolved in 20mL of milliQ water and mixed for 30 minutes. The solution was then serially diluted and plated according to the enumeration method mentioned in 3.2 above.

3.7 Scanning Electron Microscopy (SEM) imaging

Scanning Electron Microscopy imaging of the encapsulated samples was done using the JEOL JSM 6701F scanning electron microscope from Japan. The accelerating voltage was 6.0kV. Trace amount of samples were added to a double sided adhesive carbon tape before sputtered coated with platinum layer using the JEC-3000FC Auto Fine Coater to inhibit charging.

3.8 Viability of *L.Plantarum* after pasteurization

The pasteurization testing method was done in accordance to the method proposed by (Mao, Pan, Hou, Yuan, & Gao, 2018). 200mg of encapsulated *L.Plantarum* powder was dissolved in 20mL of milliQ water and mixed for 30 minutes. Thereafter, the mixture was then heated in a water bath at 85 °C for 15 minutes. Then, the samples were cooled to room temperature and the surviving bacteria was enumerated according to the method described in 3.2.

3.9 In-vitro Gastrointestinal Digestion

All the reagents and enzymes used were obtained from Sigma Aldrich, USA. Stock solutions of Simulated Salivary Fluid, Simulated Gastric Fluid and Simulated Intestinal Fluid were prepared according to the table below:

Constituents	Stock Concentration		SSF (pH 7)		SGF (pH 3)		SIF (pH 7)	
			Vol of Stock (500mL)	Final Salt Conc.	Vol of Stock (500mL)	Final Salt Conc.	Vol of Stock (500mL)	Final Salt Conc.
	g/L	mol/L	mL	mmol/L	mL	mmol/L	mL	mmol/L
KCl	37.3	0.5	15.1	15.09	6.9	6.9	6.8	6.8
KH ₂ PO ₄	68	0.5	3.7	1.35	0.9	0.9	0.8	0.8
NaHCO ₃	84	1	6.8	13.68	12.5	25	42.5	85
NaCl	117	2	-	-	11.8	47.2	9.6	38.4
MgCl ₂ (H ₂ O) ₆	30.5	0.15	0.5	0.15	0.4	0.12	1.1	0.33
NH ₄ (CO ₃) ₂	48	0.5	0.06	0.06	0.5	0.5	-	-
For pH Adjustment								
NaOH		1	-	-	-	-	-	-
HCl		6	0.09	1.1	1.3	15.6	0.7	8.4
Added directly during digestion (Not added to simulated digestion fluids)								
CaCl ₂ (H ₂ O) ₂	44.1	0.3		1.5		0.15		0.6

Table 2: Stock Solution Composition for Simulated Digestion Fluids (Minekus et al., 2014)

Figure 5: Overview of simulated in vitro digestion system. Adapted from (Minekus et al., 2014)

The gastrointestinal digestion process comprises of 3 parts, namely the oral phase, gastric phase and the intestinal phase.

Oral Phase

Firstly, 1g of encapsulated *L. Plantarum* powder was added to a 250mL Erlenmeyer Flask and dissolved in 4 mL of milliQ water and mixed for 10 minutes to obtain 5g of slurry. Then, the slurry is added with 5mL of Simulated Salivary Fluid (SSF). The amylase concentration is ensured at 75 U/mL in the final mix. After which 25 μ L of 0.3M CaCl₂ is added to obtain a concentration of 0.75mM. Then, the pH is adjusted to pH 7. Once done, the mixture is placed in the shaker incubator to react for 2 minutes, shaking at 200rpm and 37°C.

Gastric phase

After that is done, 10mL of Simulated Gastric Fluid (SGF) is added at a ratio of 1: 1 with the oral phase mixture. Ensure that the pepsin concentration is 2000 U/mL in the final mix. (Added to the SGF beforehand) 5 μ L of 0.3M CaCl₂ is added to obtain a concentration of 0.075mM. Then, the pH is adjusted to pH 3 using 6M HCl. Once done, the mixture is placed back into the shaker incubator to react for 2 hours, shaking at 200rpm and 37°C.

Intestinal phase

Finally, 20mL of Simulated Intestinal Fluid is added at a ratio of 1: 1 with the gastric phase mixture. Ensure that the pancreatin concentration of 13.369 mg/mL and a bile salt concentration of 10mM in the final mix. (Added to SIF beforehand and stirred for 30 minutes at 37°C to fully dissolve) 40 μ L of 0.3M CaCl₂ is added to obtain a concentration of 0.3mM. Then, the pH is adjusted back to pH 7 using 1M NaOH. Once done, the mixture is placed back into the shaker incubator to react for 2 hours, shaking at 200rpm and 37°C. After the digestion is completed, the flasks were immersed into an ice bath to stop the digestion reaction. The surviving bacteria is then enumerated using the method described in 3.2.

3.10 Statistical Analysis

All experiments conducted were performed in triplicates and the values were reported as mean $\pm$ standard deviation. Significance was determined at $p \leq 0.05$ and analysed using ANOVA.

4. Results and Discussion

4.1 Characterization of Encapsulants

Figure 6: Particle size distribution a: pectin, b: whey protein, c: WP2, d: WP1

Comparing the size distributions in figure 6, it was found that the average size for whey protein is the smallest at about 100nm and pectin is the largest at about 2000nm. Upon

conjugation, the average size is reduced to 850-950nm. The change in particle size is indicative of the conjugation that has occurred. The conjugation efficiency test of WP1 and WP2 found a 16.25% and 19.17% respectively. According to (Davidov-Pardo et al., 2015), this low conjugation efficiency could be due to the reduced accessibility to the conjugation sites caused by steric hindrance from the larger polysaccharides, in this case the pectin.

4.2 Effects of Spray Drying on viability of *L.Plantarum*

Figure 7: Viability of *L.Plantarum* before and after spray drying

	Total Viable Cell Count	Spray dried powder obtained/ 200ml solution
Before Encapsulation	1.533×10^{12}	
Free <i>L.Plantarum</i>	1.044×10^{10}	101mg
WP1	6.035×10^{10}	4.28g
WP2	2.055×10^{10}	3.13g

Table 3: Viable cell count of *L.Plantarum* before and after spray drying

After the *L.Plantarum* was spray dried into powder, the viability of the microencapsulated *L.Plantarum* was compared. According to Fig. 3 above, it might seem that the free *L.Plantarum* viability is higher than those encapsulated in the Maillard conjugates. This is because of the density of the *L.Plantarum* bacteria. Although the free *L.Plantarum* has a higher CFU/ml, the amount of spray dried powder produced is much lesser compared to the encapsulated powder. There was 101mg of free *L.Plantarum* powder per 200ml of solution, compared to 4.28g and 3.13g per 200ml for WP1 and WP2 respectively. Comparing the cell count of the *L.Plantarum* inoculated, it was found that there was a reduction of about 2 log units decrease in all the spray dried powders. This is partly due to the amount of sample lost

from powder that was stuck onto the walls of the spray dryer. Although it is expected that both encapsulated powders had a higher viable cell count compared to the free *L.Plantarum* cells, it is interesting to note that the WP1 which has lesser pectin content had a viable cell count of which was 3 fold higher than that of WP2.

Figure 8: SEM image of encapsulated probiotics. a,b: Free *L.Plantarum*, c: WP1, d: WP2

Examining the SEM images of the particles (Figure 8), we can see that in a and b, the free *L.Plantarum* tend to aggregate together. (Janković, Frece, Abram, & Gobin, 2012) mentioned that aggregation by Lactobacilli is correlated to adhesion. This serves as one of the mechanisms used for colonization of the gut. It is also mentioned that there are two types of aggregation. Autoaggregation, which is the aggregation among the same strain and coaggregation, which is aggregation of different species. It is said that coaggregation is the mechanism that prevents colonization of harmful pathogens. (Kos et al., 2003) Images c and d shows that the encapsulated particles are generally between the size of 3-20 μm have relatively smooth surfaces and are spherical in shape. Comparing this with images from Figure 9, the surfaces of the particles which were kept in storage at room temperature for 8

weeks tend to appear rougher and have started deforming. This is indicative of the disintegration of the particles which results in the loss of viability. It is interesting to note however, that many of the particles have smaller particles within them, suggesting that the encapsulation is a matrix type encapsulation rather than a reservoir type one.

Figure 9: SEM image of encapsulated probiotics at Week 8 of storage. a,b: WP1, c,d: WP2

4.3 Viability of Encapsulated *L.Plantarum* during storage

Figure 10: Viability of *L.Plantarum* in storage (Room Temperature)

L.Plantarum encapsulated in WP1 and WP2 were found to have a relatively stable viability over the 8 weeks with an overall log reduction of 1.3 and 1.6 respectively. In comparison, free *L.Plantarum* showed a 9.7 log reduction over the course of 8 weeks. Something interesting to note is that for WP1 and WP2, there was a slight increase in viability during the course of the storage. This is consistent with a finding presented by (Mao et al., 2018) which suggested that this could be due to the prebiotic effects from the encapsulating materials. However, WP2 did not exhibit any significant increase compared to WP1 even though it has a higher amount of pectin, which is a prebiotic. Hence, as the effects are not consistent, more research must be done to identify the exact mechanism behind this occurrence.

4.4 Protective effects of encapsulation on *L.Plantarum* during pasteurization

Figure 11: Viability of *L.Plantarum* after 15 and 30 min of pasteurization at 85°C

As seen from figure 11, WP1 and WP2 were also found to have significant protective effect during pasteurization. There was only a 0.3 log reduction for WP1 and 1.2 log reduction for WP2 where viability of the free *L.Plantarum* became indeterminant after 15 minutes. The reason the decrease in viability in WP2 is more significant could be a result of a higher pectin content. According to (Einhorn-Stoll, Kastner, Urbisch, Kroh, & Drusch, 2019; Fraeye et al., 2007), the thermal degradation becomes more significant with increasing temperatures above 60°C. Hence, with a higher pectin content, WP2 is more susceptible to pectin thermal degradation and a lower viability of probiotic cells.

4.5 Viability of Encapsulated *L.Plantarum* after in vitro digestion

Figure 12: Viability of *L.Plantarum* before and after in vitro digestion

Figure 12 shows the comparison of viability of *L.Plantarum* before and after in vitro digestion. WP1, WP2 and free *L.Plantarum* cells showed a 2, 1.8 and 6.3 log reduction respectively. This difference between WP1 and WP2 could be attributed to the amount of pectin present in the encapsulating material as well. As pectin is known to be a dietary fiber and prebiotic, it is not well degraded by the enzymes present in the gastrointestinal tract. (Bang et al., 2018) Therefore, with a higher pectin content, the WP2 particles tend to be more resilient and offer better protection for the *L.Plantarum* bacteria.

5. Conclusion

In summary, from this study, it is shown that whey protein-pectin conjugate proves to be an ideal wall material for encapsulation of probiotics for increased heat stability in storage and surviving the harsh conditions of the gastrointestinal tract. However, more research has to be done in several areas. Optimizing the parameters for spray drying to increase the yield of viable probiotics is necessary to overcome the high percentage loss of viable cells. Other methods of encapsulation could also be explored for an increased yield. The composition of protein to carbohydrate ratio could also be tweaked to improve functionality of the particles. Addition coating such as chitosan could be added for improved functionality. In this study, materials used were commercial material which contributes to the cost. In order to reduce the cost of production further, materials from food side streams such as okara protein isolate or durian seed pectin could be considered to be used as encapsulating material.

6. References

- Abbasi, A., Emam-Djomeh, Z., Mousavi, M. A. E., & Davoodi, D. (2014). Stability of vitamin D3 encapsulated in nanoparticles of whey protein isolate. *Food Chemistry*, 143, 379-383.
- Adlerberth, I., Ahrne, S., Johansson, M.-L., Molin, G., Hanson, L. A., & Wold, A. E. (1996). A mannose-specific adherence mechanism in *Lactobacillus plantarum* conferring binding to the human colonic cell line HT-29. *Applied and Environmental Microbiology*, 62(7), 2244-2251.
- Agnihotri, N., Mishra, R., Goda, C., & Arora, M. (2012). Microencapsulation—a novel approach in drug delivery: a review. *Indo Global Journal of Pharmaceutical Sciences*, 2(1), 1-20.
- Akhtar, M., & Dickinson, E. (2007). Whey protein–maltodextrin conjugates as emulsifying agents: An alternative to gum arabic. *Food Hydrocolloids*, 21(4), 607-616.
- Anadón, A., Martínez-Larrañaga, M. R., Ares, I., & Martínez, M. A. (2016). Probiotics: safety and toxicity considerations. In *Nutraceuticals* (pp. 777-798): Elsevier.
- Anandharamakrishnan, C., & Ishwarya, S. P. (2015). Introduction to encapsulation of food ingredients. *Spray Drying Techniques for Food Ingredient Encapsulation*, 37-64.
- Ananta, E., Volkert, M., & Knorr, D. (2005). Cellular injuries and storage stability of spray-dried *Lactobacillus rhamnosus* GG. *International dairy journal*, 15(4), 399-409.
- Arora, T., Singh, S., & Sharma, R. K. (2013). Probiotics: interaction with gut microbiome and antiobesity potential. *Nutrition*, 29(4), 591-596.
- Assadpour, E., Jafari, S.-M., & Maghsoudlou, Y. (2017). Evaluation of folic acid release from spray dried powder particles of pectin-whey protein nano-capsules. *International journal of biological macromolecules*, 95, 238-247.
- Bagchi, T. (2014). Traditional food & modern lifestyle: Impact of probiotics. *The Indian journal of medical research*, 140(3), 333.
- Bang, S.-J., Kim, G., Lim, M. Y., Song, E.-J., Jung, D.-H., Kum, J.-S., . . . Seo, D.-H. (2018). The influence of in vitro pectin fermentation on the human fecal microbiome. *Amb Express*, 8(1), 1-9.
- Barbosa, J., Borges, S., Amorim, M., Pereira, M., Oliveira, A., Pintado, M., & Teixeira, P. (2015). Comparison of spray drying, freeze drying and convective hot air drying for the production of a probiotic orange powder. *Journal of Functional Foods*, 17, 340-351.

- Begley, M., Hill, C., & Gahan, C. G. (2006). Bile salt hydrolase activity in probiotics. *Applied and Environmental Microbiology*, 72(3), 1729-1738.
- Bosnea, L., Moschakis, T., & Biliaderis, C. (2017). Microencapsulated cells of *Lactobacillus paracasei* subsp. *paracasei* in biopolymer complex coacervates and their function in a yogurt matrix. *Food & function*, 8(2), 554-562.
- Boyle, R. J., Robins-Browne, R. M., & Tang, M. L. (2006). Probiotic use in clinical practice: what are the risks? *The American journal of clinical nutrition*, 83(6), 1256-1264.
- Cacicedo, M. L., Islan, G. A., Drachemberg, M. F., Alvarez, V. A., Bartel, L. C., Bolzán, A. D., & Castro, G. R. (2018). Hybrid bacterial cellulose–pectin films for delivery of bioactive molecules. *New Journal of Chemistry*, 42(9), 7457-7467.
- Chávarri, M., Marañón, I., & Villarán, M. C. (2012). Encapsulation technology to protect probiotic bacteria. In *Probiotics*: IntechOpen.
- Chen, X., Yang, G., Song, J.-H., Xu, H., Li, D., Goldsmith, J., . . . Kelly, C. P. (2013). Probiotic yeast inhibits VEGFR signaling and angiogenesis in intestinal inflammation. *PLoS One*, 8(5), e64227.
- Colodel, C., das Graças Bagatin, R. M., Tavares, T. M., & de Oliveira Petkowicz, C. L. (2017). Cell wall polysaccharides from pulp and peel of cubiu: A pectin-rich fruit. *Carbohydrate Polymers*, 174, 226-234.
- Cotter, P. D., & Hill, C. (2003). Surviving the acid test: responses of gram-positive bacteria to low pH. *Microbiology and molecular biology reviews*, 67(3), 429-453.
- D'Aimmo, M. R., Modesto, M., & Biavati, B. (2007). Antibiotic resistance of lactic acid bacteria and *Bifidobacterium* spp. isolated from dairy and pharmaceutical products. *International journal of food microbiology*, 115(1), 35-42.
- Das, A., Ray, S., Raychaudhuri, U., & Chakraborty, R. (2014). Microencapsulation of probiotic bacteria and its potential application in food technology. *International Journal of Agriculture, Environment and Biotechnology*, 7(1), 47-53.
- Davidov-Pardo, G., Joye, I. J., Espinal-Ruiz, M., & McClements, D. J. (2015). Effect of maillard conjugates on the physical stability of zein nanoparticles prepared by liquid antisolvent coprecipitation. *Journal of agricultural and food chemistry*, 63(38), 8510-8518.
- De Castro-Cislaghi, F. P., Carina Dos Reis, E. S., Fritzen-Freire, C. B., Lorenz, J. G., & Sant'Anna, E. S. (2012). *Bifidobacterium* Bb-12 microencapsulated by spray drying with whey: Survival under simulated gastrointestinal conditions, tolerance to NaCl, and viability during storage. *Journal of Food Engineering*, 113(2), 186-193.

- de Castro, R. J. S., Domingues, M. A. F., Ohara, A., Okuro, P. K., dos Santos, J. G., Brexó, R. P., & Sato, H. H. (2017). Whey protein as a key component in food systems: Physicochemical properties, production technologies and applications. *Food structure*, 14, 17-29.
- de Oliveira, F. C., Coimbra, J. S. d. R., de Oliveira, E. B., Zuñiga, A. D. G., & Rojas, E. E. G. (2016). Food protein-polysaccharide conjugates obtained via the Maillard reaction: A review. *Critical reviews in food science and nutrition*, 56(7), 1108-1125.
- de Vos, P., Faas, M. M., Spasojevic, M., & Sikkema, J. (2010). Encapsulation for preservation of functionality and targeted delivery of bioactive food components. *International dairy journal*, 20(4), 292-302.
- Desmond, C., Ross, R., O'callaghan, E., Fitzgerald, G., & Stanton, C. (2002). Improved survival of *Lactobacillus paracasei* NFBC 338 in spray - dried powders containing gum acacia. *Journal of applied microbiology*, 93(6), 1003-1011.
- Ding, W., & Shah, N. (2007). Acid, bile, and heat tolerance of free and microencapsulated probiotic bacteria. *Journal of food science*, 72(9), M446-M450.
- Dixit, Y., Wagle, A., & Vakil, B. (2016). Patents in the field of probiotics, prebiotics, synbiotics: A review. *Journal of Food: Microbiology Safety & Hygiene*, 01-02.
- Eid, R., Jakee, J., Rashidy, A., Asfour, H., Omara, S., Kandil, M., . . . Seida, A. (2016). Potential antimicrobial activities of probiotic *Lactobacillus* strains isolated from raw milk. *J Probiotics Health*, 4, 1e8.
- Einhorn-Stoll, U., Kastner, H., Urbisch, A., Kroh, L. W., & Drusch, S. (2019). Thermal degradation of citrus pectin in low-moisture environment-Influence of acidic and alkaline pre-treatment. *Food Hydrocolloids*, 86, 104-115.
- Fathi, M., Martin, A., & McClements, D. J. (2014). Nanoencapsulation of food ingredients using carbohydrate based delivery systems. *Trends in Food Science & Technology*, 39(1), 18-39.
- Floch, M. H., Walker, W. A., Madsen, K., Sanders, M. E., Macfarlane, G. T., Flint, H. J., . . . Kelly, C. P. (2011). Recommendations for probiotic use—2011 update. *Journal of clinical gastroenterology*, 45, S168-S171.
- Flores, F. P., Singh, R. K., & Kong, F. (2014). Physical and storage properties of spray-dried blueberry pomace extract with whey protein isolate as wall material. *Journal of Food Engineering*, 137, 1-6.

- Foegeding, E. A., Davis, J. P., Doucet, D., & McGuffey, M. K. (2002). Advances in modifying and understanding whey protein functionality. *Trends in Food Science & Technology*, 13(5), 151-159.
- Fraeye, I., De Roeck, A., Duvetter, T., Verlent, I., Hendrickx, M., & Van Loey, A. (2007). Influence of pectin properties and processing conditions on thermal pectin degradation. *Food Chemistry*, 105(2), 555-563.
- Gao, F., Zhang, X., Wang, H., Sun, X., Wang, J., & Wang, C. (2019). Comparison of dry-and wet-heat induced changes in physicochemical properties of whey protein in absence or presence of inulin. *Food science and biotechnology*, 28(5), 1367-1374.
- Gharehbeglou, P., Jafari, S. M., Hamishekar, H., Homayouni, A., & Mirzaei, H. (2019). Pectin-whey protein complexes vs. small molecule surfactants for stabilization of double nano-emulsions as novel bioactive delivery systems. *Journal of Food Engineering*, 245, 139-148.
- Gharsallaoui, A., Roudaut, G., Chambin, O., Voilley, A., & Saurel, R. (2007). Applications of spray-drying in microencapsulation of food ingredients: An overview. *Food Research International*, 40(9), 1107-1121.
- Golowczyc, M. A., Silva, J., Teixeira, P., De Antoni, G. L., & Abraham, A. G. (2011). Cellular injuries of spray-dried *Lactobacillus* spp. isolated from kefir and their impact on probiotic properties. *International journal of food microbiology*, 144(3), 556-560.
- González-Ferrero, C., Irache, J., & González-Navarro, C. (2018). Soybean protein-based microparticles for oral delivery of probiotics with improved stability during storage and gut resistance. *Food Chemistry*, 239, 879-888.
- Hamner, S., McInnerney, K., Williamson, K., Franklin, M. J., & Ford, T. E. (2013). Bile salts affect expression of *Escherichia coli* O157: H7 genes for virulence and iron acquisition, and promote growth under iron limiting conditions. *PLoS One*, 8(9), e74647.
- Harish, K., & Varghese, T. (2006). Probiotics in humans—evidence based review. *Calicut Med J*, 4(4), e3.
- Heidebach, T., Först, P., & Kulozik, U. (2012). Microencapsulation of probiotic cells for food applications. *Critical reviews in food science and nutrition*, 52(4), 291-311.
- Hemaiswarya, S., Raja, R., Ravikumar, R., & Carvalho, I. S. (2013). Mechanism of action of probiotics. *Brazilian archives of Biology and technology*, 56(1), 113-119.
- Hill, C., Guarner, F., Reid, G., Gibson, G., & Merenstein, D. (2014). Pot B6, Morelli L, Canani RB, Flint HJ, Salminen S, Calder PC, Sanders ME. Expert consensus

- document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol*, 11, 506-514.
- Hodge, J. E. (1953). Dehydrated foods, chemistry of browning reactions in model systems. *Journal of agricultural and food chemistry*, 1(15), 928-943.
- Huang, S., Vignolles, M.-L., Chen, X. D., Le Loir, Y., Jan, G., Schuck, P., & Jeantet, R. (2017). Spray drying of probiotics and other food-grade bacteria: A review. *Trends in Food Science & Technology*, 63, 1-17.
- Iaconelli, C., Lemetais, G., Kechaou, N., Chain, F., Bermúdez-Humarán, L. G., Langella, P., . . . Beney, L. (2015). Drying process strongly affects probiotics viability and functionalities. *Journal of biotechnology*, 214, 17-26.
- Janković, T., Frece, J., Abram, M., & Gobin, I. (2012). Aggregation ability of potential probiotic *Lactobacillus plantarum* strains. *Int J Sanit Eng Res*, 6(1), 19-24.
- Jia, W., & Xie, G. (2018). Probiotics, bile acids and gastrointestinal carcinogenesis. *Nature Reviews Gastroenterology & Hepatology*, 15(4), 205-205.
- Kato, A. (2002). Industrial applications of Maillard-type protein-polysaccharide conjugates. *Food Science and Technology Research*, 8(3), 193-199.
- Kechagia, M., Basoulis, D., Konstantopoulou, S., Dimitriadi, D., Gyftopoulou, K., Skarmoutsou, N., & Fakiri, E. M. (2013). Health benefits of probiotics: a review. *International Scholarly Research Notices*, 2013.
- Kerry, R. G., Patra, J. K., Gouda, S., Park, Y., Shin, H.-S., & Das, G. (2018). Benefaction of probiotics for human health: A review. *Journal of food and drug analysis*, 26(3), 927-939.
- Khuenpet, K., Charoenjarasrerk, N., Jaijit, S., Arayapoonpong, S., & Jittanit, W. (2016). Investigation of suitable spray drying conditions for sugarcane juice powder production with an energy consumption study. *Agriculture and Natural Resources*, 50(2), 139-145.
- Kontula, P., Jaskari, J., Nollet, L., De Smet, I., von Wright, A., Poutanen, K., & Mattila-Sandholm, T. (1998). The colonization of a simulator of the human intestinal microbial ecosystem by a probiotic strain fed on a fermented oat bran product: effects on the gastrointestinal microbiota. *Applied Microbiology and Biotechnology*, 50(2), 246-252.

- Kos, B., Šušković, J., Vuković, S., Šimpraga, M., Frece, J., & Matošić, S. (2003). Adhesion and aggregation ability of probiotic strain *Lactobacillus acidophilus* M92. *Journal of applied microbiology*, 94(6), 981-987.
- Koubaa, M., Roohinejad, S., Mungure, T. E., El-Din, B. A., Greiner, R., & Mallikarjunan, K. (2018). Effect of emerging processing technologies on Maillard reactions. In *Encyclopedia of Food Chemistry* (pp. 76-82): Elsevier.
- Kumar, M., Dhaka, P., Vijay, D., Vergis, J., Mohan, V., Kumar, A., . . . Rawool, D. B. (2016). Antimicrobial effects of *Lactobacillus plantarum* and *Lactobacillus acidophilus* against multidrug-resistant enteroaggregative *Escherichia coli*. *International journal of antimicrobial agents*, 48(3), 265-270.
- Kurmann, J., & Rasic, J. (1991). The Health Potential of Products Containing Bifidobacteria. Therapeutic Properties of Fermented Milks. Applied Food Sci. In: Elsevier London, UK.
- Lankaputhra, W. (1995). Survival of *Lactobacillus acidophilus* and *Bifidobacterium* spp. in the presence of acid and bile salt. *Cult. Dairy Prod. J.*, 30, 2-7.
- Lund, B., Adamsson, I., & Edlund, C. (2002). Gastrointestinal transit survival of an *Enterococcus faecium* probiotic strain administered with or without vancomycin. *International journal of food microbiology*, 77(1-2), 109-115.
- M. De Angelis, M. G. (2016). *Lactobacillus* SPP.: General Characteristics. *Reference Module in Food Science*. doi:<https://doi.org/10.1016/B978-0-08-100596-5.00851-9>
- Mao, L., Pan, Q., Hou, Z., Yuan, F., & Gao, Y. (2018). Development of soy protein isolate-carrageenan conjugates through Maillard reaction for the microencapsulation of *Bifidobacterium longum*. *Food Hydrocolloids*, 84, 489-497.
- Marco, M. L., Pavan, S., & Kleerebezem, M. (2006). Towards understanding molecular modes of probiotic action. *Current opinion in biotechnology*, 17(2), 204-210.
- Merenstein, D., & Salminen, S. (2017). Probiotics and prebiotics.
- Michail, S. (2005). The mechanism of action of probiotics. *Practic Gastroenterol*, 2, 29-47.
- Minekus, M., Alming, M., Alvito, P., Ballance, S., Bohn, T., Bourlieu, C., . . . Dupont, D. (2014). A standardised static in vitro digestion method suitable for food—an international consensus. *Food & function*, 5(6), 1113-1124.
- Morelli, L., & Capurso, L. (2012). FAO/WHO guidelines on probiotics: 10 years later. *Journal of clinical gastroenterology*, 46, S1-S2.
- Nasirpour, A., Scher, J., & Desobry, S. (2006). Baby foods: formulations and interactions (a review). *Critical reviews in food science and nutrition*, 46(8), 665-681.

- Nedovic, V., Kalusevic, A., Manojlovic, V., Levic, S., & Bugarski, B. (2011). An overview of encapsulation technologies for food applications. *Procedia Food Science*, 1, 1806-1815.
- Ng, S., Hart, A., Kamm, M., Stagg, A., & Knight, S. C. (2009). Mechanisms of action of probiotics: recent advances. *Inflammatory bowel diseases*, 15(2), 300-310.
- Nguyen, H.-T., Truong, D.-H., Kouhondé, S., Ly, S., Razafindralambo, H., & Delvigne, F. (2016). Biochemical engineering approaches for increasing viability and functionality of probiotic bacteria. *International journal of molecular sciences*, 17(6), 867.
- Nooshkam, M., Babazadeh, A., & Jooyandeh, H. (2018). Lactulose: Properties, techno-functional food applications, and food grade delivery system. *Trends in Food Science & Technology*, 80, 23-34.
- Nooshkam, M., Varidi, M., & Bashash, M. (2019). The Maillard reaction products as food-born antioxidant and antibrowning agents in model and real food systems. *Food Chemistry*, 275, 644-660.
- O'Hara, A. M., & Shanahan, F. (2007). Mechanisms of action of probiotics in intestinal diseases. *TheScientificWorldJournal*, 7.
- O'Mahony, J. A., Drapala, K. P., Mulcahy, E. M., & Mulvihill, D. M. (2017). Controlled glycation of milk proteins and peptides: Functional properties. *International dairy journal*, 67, 16-34.
- Oelschlaeger, T. A. (2010). Mechanisms of probiotic actions—a review. *International journal of medical microbiology*, 300(1), 57-62.
- Paéz, R., Lavari, L., Vinderola, G., Audero, G., Cuatrin, A., Zaritzky, N., & Reinheimer, J. (2012). Effect of heat treatment and spray drying on lactobacilli viability and resistance to simulated gastrointestinal digestion. *Food Research International*, 48(2), 748-754.
- Parra Huertas, R. A. (2010). Food microencapsulation: a review. *Revista Facultad Nacional de Agronomía Medellín*, 63(2), 5669-5684.
- Parvez, S., Malik, K. A., Ah Kang, S., & Kim, H. Y. (2006). Probiotics and their fermented food products are beneficial for health. *Journal of applied microbiology*, 100(6), 1171-1185.
- Peighambardoust, S., Tafti, A. G., & Hesari, J. (2011). Application of spray drying for preservation of lactic acid starter cultures: a review. *Trends in Food Science & Technology*, 22(5), 215-224.

- Pellegrino, L., Van Boekel, M., Gruppen, H., Resmini, P., & Pagani, M. (1999). Heat-induced aggregation and covalent linkages in β -casein model systems. *International dairy journal*, 9(3-6), 255-260.
- Perusko, M., Al-Hanish, A., Velickovic, T. C., & Stanic-Vucinic, D. (2015). Macromolecular crowding conditions enhance glycation and oxidation of whey proteins in ultrasound-induced Maillard reaction. *Food Chemistry*, 177, 248-257.
- Picot, A., & Lacroix, C. (2004). Encapsulation of bifidobacteria in whey protein-based microcapsules and survival in simulated gastrointestinal conditions and in yoghurt. *International dairy journal*, 14(6), 505-515.
- Pu, J., Bankston, J. D., & Sathivel, S. (2011). Developing microencapsulated flaxseed oil containing shrimp (*Litopenaeus setiferus*) astaxanthin using a pilot scale spray dryer. *Biosystems Engineering*, 108(2), 121-132.
- Puertollano, E., Puertollano, M. A., Cruz-Chamorro, L., de Cienfuegos, G. A., Ruiz-Bravo, A., & de Pablo, M. A. (2008). Orally administered *Lactobacillus plantarum* reduces pro-inflammatory interleukin secretion in sera from *Listeria monocytogenes* infected mice. *British Journal of Nutrition*, 99(4), 819-825.
- Qi, W., Liang, X., Yun, T., & Guo, W. (2019). Growth and survival of microencapsulated probiotics prepared by emulsion and internal gelation. *Journal of food science and technology*, 56(3), 1398-1404.
- Ramos, C. L., Thorsen, L., Schwan, R. F., & Jespersen, L. (2013). Strain-specific probiotics properties of *Lactobacillus fermentum*, *Lactobacillus plantarum* and *Lactobacillus brevis* isolates from Brazilian food products. *Food microbiology*, 36(1), 22-29.
- Rathore, S., Desai, P. M., Liew, C. V., Chan, L. W., & Heng, P. W. S. (2013). Microencapsulation of microbial cells. *Journal of Food Engineering*, 116(2), 369-381.
- Rault, A., Béal, C., Ghorbal, S., Ogier, J.-C., & Bouix, M. (2007). Multiparametric flow cytometry allows rapid assessment and comparison of lactic acid bacteria viability after freezing and during frozen storage. *Cryobiology*, 55(1), 35-43.
- Sakata, S., Kitahara, M., Sakamoto, M., Hayashi, H., Fukuyama, M., & Benno, Y. (2002). Unification of *Bifidobacterium infantis* and *Bifidobacterium suis* as *Bifidobacterium longum*. *International journal of systematic and evolutionary microbiology*, 52(6), 1945-1951.

- Santivarangkna, C., Kulozik, U., & Foerst, P. (2007). Alternative drying processes for the industrial preservation of lactic acid starter cultures. *Biotechnology progress*, 23(2), 302-315.
- Santivarangkna, C., Kulozik, U., & Foerst, P. (2008). Inactivation mechanisms of lactic acid starter cultures preserved by drying processes. *Journal of applied microbiology*, 105(1), 1-13.
- Sarao, L. K., & Arora, M. (2017). Probiotics, prebiotics, and microencapsulation: A review. *Critical reviews in food science and nutrition*, 57(2), 344-371.
- Sarowska, J., Choroszy-Król, I., Regulska-Ilow, B., Frej-Madrzak, M., & Jama-Kmiecik, A. (2013). The therapeutic effect of probiotic bacteria on gastrointestinal diseases. *Adv Clin Exp Med*, 22(5), 759-766.
- Schuck, P., Dolivet, A., Mejean, S., Hervé, C., & Jeantet, R. (2013). Spray drying of dairy bacteria: New opportunities to improve the viability of bacteria powders. *International dairy journal*, 31(1), 12-17.
- Serna-Cock, L., & Vallejo-Castillo, V. (2013). Probiotic encapsulation. *African Journal of Microbiology Research*, 7(40), 4743-4753.
- Setiowati, A. D., Saeedi, S., Wijaya, W., & Van der Meeren, P. (2017). Improved heat stability of whey protein isolate stabilized emulsions via dry heat treatment of WPI and low methoxyl pectin: Effect of pectin concentration, pH, and ionic strength. *Food Hydrocolloids*, 63, 716-726.
- Sherman, P. M., Ossa, J. C., & Johnson - Henry, K. (2009). Unraveling mechanisms of action of probiotics. *Nutrition in Clinical Practice*, 24(1), 10-14.
- Silván, J. M., Assar, S. H., Srey, C., Del Castillo, M. D., & Ames, J. M. (2011). Control of the Maillard reaction by ferulic acid. *Food Chemistry*, 128(1), 208-213.
- Singh, A., & Dalgleish, D. (1998). The emulsifying properties of hydrolyzates of whey proteins. *Journal of dairy science*, 81(4), 918-924.
- Sobel, R., Versic, R., & Gaonkar, A. G. (2014). Introduction to microencapsulation and controlled delivery in foods. In *Microencapsulation in the food industry* (pp. 3-12): Elsevier.
- Solanki, H. K., Pawar, D. D., Shah, D. A., Prajapati, V. D., Jani, G. K., Mulla, A. M., & Thakar, P. M. (2013). Development of microencapsulation delivery system for long-term preservation of probiotics as biotherapeutics agent. *BioMed research international*, 2013.

- Sornplang, P., & Piyadeatsoontorn, S. (2016). Probiotic isolates from unconventional sources: a review. *Journal of animal science and technology*, 58(1), 1-11.
- Spotti, M. J., Perduca, M. J., Piagentini, A., Santiago, L. G., Rubiolo, A. C., & Carrara, C. R. (2013). Gel mechanical properties of milk whey protein–dextran conjugates obtained by Maillard reaction. *Food Hydrocolloids*, 31(1), 26-32.
- Surono, I., Verhoeven, J., Verbruggen, S., & Venema, K. (2018). Microencapsulation increases survival of the probiotic *Lactobacillus plantarum* IS - 10506, but not *Enterococcus faecium* IS - 27526 in a dynamic, computer - controlled in vitro model of the upper gastrointestinal tract. *Journal of applied microbiology*, 124(6), 1604-1609.
- Tripathi, M. K., & Giri, S. K. (2014). Probiotic functional foods: Survival of probiotics during processing and storage. *Journal of Functional Foods*, 9, 225-241.
- Ukena, S. N., Singh, A., Dringenberg, U., Engelhardt, R., Seidler, U., Hansen, W., . . . Rogler, G. (2007). Probiotic *Escherichia coli* Nissle 1917 inhibits leaky gut by enhancing mucosal integrity. *PLoS One*, 2(12), e1308.
- Umbrello, G., & Esposito, S. (2016). Microbiota and neurologic diseases: potential effects of probiotics. *Journal of Translational Medicine*, 14(1), 298.
- Vanderhoof, J. A. (2000). Probiotics and intestinal inflammatory disorders in infants and children. *Journal of pediatric gastroenterology and nutrition*, 30, S34-S38.
- Wang, H.-Y., Qian, H., & Yao, W.-R. (2011). Melanoidins produced by the Maillard reaction: Structure and biological activity. *Food Chemistry*, 128(3), 573-584.
- Wang, W.-q., Bao, Y.-h., & Chen, Y. (2013). Characteristics and antioxidant activity of water-soluble Maillard reaction products from interactions in a whey protein isolate and sugars system. *Food Chemistry*, 139(1-4), 355-361.
- Wrodnigg, T. M., & Eder, B. (2001). The Amadori and Heyns rearrangements: landmarks in the history of carbohydrate chemistry or unrecognized synthetic opportunities? In *Glycoscience* (pp. 115-152): Springer.
- Xu, D., Wang, X., Jiang, J., Yuan, F., & Gao, Y. (2012). Impact of whey protein–Beet pectin conjugation on the physicochemical stability of β -carotene emulsions. *Food Hydrocolloids*, 28(2), 258-266.
- Yadav, M. P., Parris, N., Johnston, D. B., Onwulata, C. I., & Hicks, K. B. (2010). Corn fiber gum and milk protein conjugates with improved emulsion stability. *Carbohydrate Polymers*, 81(2), 476-483.

- Yao, M., Xie, J., Du, H., McClements, D. J., Xiao, H., & Li, L. (2020). Progress in microencapsulation of probiotics: A review. *Comprehensive Reviews in Food Science and Food Safety*, 19(2), 857-874.
- Yeung, T. W., Arroyo-Maya, I. J., McClements, D. J., & Sela, D. A. (2016). Microencapsulation of probiotics in hydrogel particles: enhancing *Lactococcus lactis* subsp. *cremoris* LM0230 viability using calcium alginate beads. *Food & function*, 7(4), 1797-1804.
- Zárate, G. (2012). Dairy Propionibacteria: Less conventional probiotics to improve the human and animal health. *Probiotic in Animals*, 153-202.
- Zendeboodi, F., Khorshidian, N., Mortazavian, A. M., & da Cruz, A. G. (2020). Probiotic: Conceptualization from a new approach. *Current Opinion in Food Science*.
- Zhu, D., Damodaran, S., & Lucey, J. A. (2010). Physicochemical and emulsifying properties of whey protein isolate (WPI)– dextran conjugates produced in aqueous solution. *Journal of agricultural and food chemistry*, 58(5), 2988-2994.