Climate-forced sea-level lowstands in the Indian Ocean during the last two millennia

Kench, Paul S.; McLean, Roger F.; Owen, Susan D.; Ryan, Emma; Morgan, Kyle Meredith; Ke, Lin; Wang, Xianfeng; Roy, Keven

2019

Kench, P. S., McLean, R. F., Owen, S. D., Ryan, E., Morgan, K. M., Ke, L., . . . Roy, K. (2020). Climate-forced sea-level lowstands in the Indian Ocean during the last two millennia. *Nature Geoscience*, 13(1), 61-64. doi:10.1038/s41561-019-0503-7

<https://hdl.handle.net/10356/146658>

<https://doi.org/10.1038/s41561-019-0503-7>

© The Author(s), under exclusive licence to Springer Nature Limited. All rights reserved.
This paper was published in *Nature Geoscience* and is made available with permission of The Author(s).

Downloaded on 26 Apr 2023 19:05:37 SGT

1 Climate-forced sea level lowstands in the Indian Ocean during the last two millennia

2 Paul S. Kench^{1*}, Roger F. McLean², Susan D. Owen³, Emma Ryan⁴, Kyle E. Morgan⁵, Lin Ke⁶,
3 Xianfeng Wang^{5,6}, Keven Roy^{5,7}

4 ¹*Department of Earth Sciences, Simon Fraser University, Burnaby, Canada*

5 ²*School of Science, University of New South Wales - Canberra, ACT, Australia*

6 ³*School of Resource and Environmental Management, Simon Fraser University, Burnaby, Canada*

7 ⁴*School of Environment, University of Auckland, Private Bag 92109, Auckland, New Zealand*

8 ⁵*Asian School of the Environment, Nanyang Technological University, Singapore 637459, Singapore*

9 ⁶*Earth Observatory of Singapore, Nanyang Technological University, Singapore 639798, Singapore*

10 ⁷*Hiscox Group, 1 Great St. Helen's, London, EC3A 6HX, UK*

11 Sea-level reconstructions over the past two millennia provide a pre-industrial context to assess
12 whether the magnitude and rate of modern sea-level change is unprecedented. Sea-level
13 records from the Indian Ocean over the past 2,000 years are sparse, while records from the
14 Atlantic and Pacific Oceans show variations less than 0.25 m and no significant negative
15 excursions. Here we present evidence of two low sea-level phases in the Maldives, Indian Ocean
16 based on fossil coral microatolls. Microatoll growth is constrained by low water levels and
17 consequently, they are robust recorders of past sea level. U-Th dating of the Maldivian corals
18 identified lowstands at 234-605 CE and 1481-1807 CE when sea-level fell to maximum depths of -
19 0.88 m and -0.89 m respectively. These lowstands are synchronous with reductions in radiative
20 forcing and sea surface temperature associated with the Late Antiquity Little Ice Age and Little
21 Ice Age. Our results provide the first high fidelity observations of lower sea levels during these
22 cool periods and show rates of change of up to 4.24 mm.yr⁻¹. Our data also confirms the
23 acceleration of relative sea-level rise over the past two centuries and suggests the current
24 magnitude and rate of sea-level rise is not unprecedented.

25 Resolving the precise behaviour of relative sea level (RSL) over the past two millennia, and particularly
26 prior to the industrial era, provides a robust basis to understand natural sea-level variability and
27 evaluate whether the scale and pace of modern sea-level rise is exceptional¹⁻³. Reconstruction of late
28 Holocene RSL also provides constraints for calibrating geophysical models of Earth's rheology and
29 glacio-isostatic adjustments (GIA)^{4,5}. An enhanced understanding of the relationships between sea
30 level, climate change and oceanic processes also provides the basis for improved projections of future
31 sea level⁶⁻⁸ and the impacts on coastal systems and societies⁹⁻¹¹.

32 Despite the importance of constructing pre-industrial sea-level records, knowledge of sea-level
33 variability over the Common Era (CE, the past 2,000 years) is limited and sea-level response to known
34 climatic variations such as the Medieval Climate Anomaly (MCA), Little Ice Age (LIA) and 20th Century
35 warming is poorly constrained^{4,6}. Furthermore, large differences in RSL behaviour exist between near-
36 field, intermediate-field and far-field locations and between ocean basins¹². Studies using salt-marsh
37 proxy records, have developed high resolution records spanning the past two millennia for the North
38 American Atlantic and Gulf coasts, which show sea level continually rising throughout the mid-late
39 Holocene to its present maximum^{3,13,14}. Several notable features characterize these records. First, RSL
40 appears to have not deviated significantly from its linear trend until the onset of modern sea-level rise in
41 the late 19th century. Second, low amplitude variations $< \pm 0.25$ m have been detected, corresponding
42 to shifts in global mean temperature. Third, there is little evidence of any marked sea-level excursions
43 below present level in the late Holocene.

44 In contrast to the Atlantic, sea-level dynamics over the past 2,000 years in the Indian and Pacific
45 Oceans remain poorly resolved. Broad-scale Holocene reconstructions from these ocean basins infer
46 sea level fell smoothly from a mid-Holocene highstand^{4,15,16}, although empirical data confirming this
47 trend are scarce^{5,17,18}. Notably, many records derived for the Indo-Pacific are constructed from coral-
48 based data in tropical far-field locations. However, such observational data are not continuous and are

restricted to time intervals and environmental settings that favour preservation of those records⁴. For example, existing coral-based sea-level reconstructions comprise a rich dataset (Supplementary Fig. 1) spanning the early to mid-Holocene transgression that terminated in a highstand, at which time reefs and corals grew above current sea level¹⁵⁻¹⁷. This widespread high sea-level signature is likely to have occluded post-highstand coral growth from emergent reef platforms, and with it any clear-cut evidence of RSL, rendering inferences of sea-level trends and variability over the last 2,000 years largely speculative.

Coral microatolls as paleo sea-level indicators

We present the first field evidence of post-highstand sea-level behaviour in the Maldives, central Indian Ocean (Fig. 1), that includes two periods of low sea level in the last 2,000 years. During this time both empirical paleo sea-level data from the region^{15,17,18} and geophysical model results¹⁹ depict a smoothly falling sea to its present position (Fig. 1D). Our new evidence, derived from high precision U-Th dating of 39 *in situ* fossil coral microatolls (*Porites* spp.) that have been preserved on the Mahutigalaa reef platform, Huvadhoo atoll, challenges this view (Fig. 1B, C, Supplementary Information). Such microatolls are the principal geological indicator of Holocene RSL changes used in far-field tropical mid-ocean settings^{4,5,20}. Details of the location, diameter, elevation and age of the fossil microatolls on Mahutigalaa are summarised in Supplementary Information.

Timing and elevation of low and variable sea levels

Two spatially and temporally discrete clusters of U-Th dated fossil microatolls (39 in total ranging in diameter from 1.0 to 4.95 m, with a mean of 2.66 m) identify an oscillating sea surface and two low RSL excursions following the mid-late Holocene highstand (+0.5 m above mean sea level (MSL)) of 4,000 to 2,100 years ago¹⁷ (Fig. 2A). The first lowstand occurred between 234 to 605 CE (371 years) and is defined by 11 microatolls. These microatolls cover a 0.24 m depth range and have a mean elevation of -0.77 ± 0.14 m below present height of living coral (HLC), with two microatolls defining the lowest level

73 of -0.88 ± 0.14 m below HLC. These elevations equate to mean and maximum depths of -1.34 and $-$
74 1.45 ± 0.14 m below mean sea level (MSL). The second lowstand occurred over eight centuries later
75 from 1481 to 1807 CE (326 years), as defined by 12 microatolls, and has a mean elevation of $-0.71 \pm$
76 0.14 m below HLC. Eleven of the microatolls fit within a narrow 185 yr window (1481 to 1666 CE), are
77 found across a 0.12 m depth range and have a mean elevation of -0.69 ± 0.14 m below HLC. A further
78 microatoll (sample 10) defines a lower sea-level elevation of -0.89 ± 0.14 m below HLC at 1807 CE.
79 Seven microatolls with diameters over 4.0 m attest to prolonged sea-level stability during both
80 lowstands.

81 A temporal hiatus exists in the dataset between 784 and 1471 CE in which no coral evidence has been
82 identified on Mahutigalaa. From the earliest identified lowstand, corals indicate an increase in sea level
83 between 653 and 784 CE to an elevation -0.18 m below HLC. Corals of similar elevation dated at 1471
84 CE also record the onset of the second fall in sea level. Consequently, while there is a lack of evidence
85 in the intervening 687 year period, microatolls close to present sea level constrain the beginning and
86 end of this gap and imply sea level was near present level, if not higher, during this period. Notably, our
87 data also reveal a marked increase in sea level over the past 210 years beginning from the lowstand
88 (1807 CE) with a magnitude of sea-level rise of 0.89 m.

89 Absence of low sea level evidence on coral reefs

90 Our Maldivian results provide the first observational data that identifies two RSL excursions that fell well
91 below the present level of the sea surface. Mean and maximum depths of >0.7 to 0.89 ± 0.14 m were
92 achieved and the temporal pattern clearly indicates large oscillations in sea-level behaviour over the
93 past two millennia in the Indian Ocean. To date, such oscillations have not been detected globally and
94 coral evidence across this timeframe is limited (SI Fig. 1). We believe this general lack of reefal
95 evidence is a legacy of the mid-Holocene high sea level which had two major, but generally
96 unrecognised, effects. First, the emergence of highstand reefs as sea-level fell occluded coral growth

from reef surfaces. Second, any potential accommodation space that was available has been overprinted by subsequent reef growth in the last two centuries. Preservation of coral records is therefore, rare, except in exceptional locations such as the lagoon platform reefs in the Maldives.

Significantly, GIA modelling in this far-field region is unable to resolve sea-level fluctuations of the scale observed¹⁹ (Fig. 1D), suggesting that the primary drivers of the two lowstands, and the timeframe of oscillations, cannot be explained by glacio-isostatic, gravitational and rotational effects. Instead, global climate fluctuations^{7,29} forcing variations in sea surface temperature (SST) are considered the most likely mechanism driving steric sea-level variability shown in our data²⁵.

Low sea levels and little ice ages

While SST records from the central Indian Ocean are scarce, regional and global proxy data show that both periods of lower sea level are coincident with intervals of substantial climate cooling (Fig. 2). The Late Antique Little Ice Age (LALIA) was a long-lasting, spatially synchronized period characterised by variable and colder climates that occurred from the second to the seventh centuries^{24,30-33}, (Fig. 2B). This interval had 13 of the coolest decades spanning the last 2,000 years²³. Of note, volcanism has been implicated in forcing the radiative cooling associated with the LALIA after which temperatures increased rapidly in the seventh century³¹. Consistent with this change is the progressive increase in elevation of microatolls in our coral data, leading toward the MCA, characterized by warmer temperatures and a gap in the global microatoll record (SI Fig. 1, Fig. 2A). The LIA is also known to have started abruptly 1275-1300 CE in response to increased volcanism and reinforced by ocean/sea-ice feedbacks³⁴ and a solar minimum²⁷ and extended through to the mid-19th century³⁵. Our physical records present the first evidence of falling and low sea levels in these climatic cool intervals with lower SSTs (Fig. 2C).

Interpreting modern sea level behaviour

Our results have profound implications for interpreting modern and future sea-level change. The magnitude of the sea-level oscillations, as revealed in our coral-based data, over the last two millennia, are larger (up to 0.89 m) than earlier empirical and model reconstructions suggest. Based on maximum and minimum elevation microatolls in our dataset, rates of sea-level rise and fall across the past 1800 years range from 4.24 to 2.80 mm.yr⁻¹ (Supplementary Table 4). Periods of sea-level fall occurred at rates of -2.80 (-92 to 401 CE) and -2.75 mm.yr⁻¹ (1521-1757 CE) whilst sea-level rose at a rate of 4.24 mm.yr⁻¹ between 401 and 717 CE. Consequently, the magnitude and rates of sea-level change currently observed in the Indian Ocean, at twice the global average³⁶, are not unprecedented over the past 2,000 years. Furthermore, coastal ecosystems and communities have clearly experienced sea-level oscillations that are of lower frequency but are much greater in magnitude than the El Niño Southern Oscillation (ENSO) seesaw that presently characterizes sea-level variability in the Indo-Pacific. Whilst ENSO-generated sea levels are expected to become more extreme with SLR in the future³⁷, the possibility of large positive and negative sea-level changes of the scale demonstrated here should also be considered. Additionally, we present the first coral-based record of the most recent acceleration in sea level in the Indian Ocean, that began in the early 19th century (Fig. 2A). While providing validity for the earlier sea-level excursions, both the magnitude and rate (4.24 mm.yr⁻¹) of this recent rise is consistent with the 25 year tide gauge record from Gan (Addu atoll) located 110 km south of our field site, which shows an increase in sea level of ~3.4 mm.yr⁻¹ over the past two decades. Collectively, our coral evidence suggests that rates of recent sea-level change are not unprecedented over the past two millennia. However, these rates of rise set a sea-level trajectory that will exceed the elevation of the late Holocene highstand in the Indo-Pacific in the next century.

References

1. Kemp, A.C. et al. Climate related sea-level variations over the past two millennia. *Proc. Natl. Acad. Sci. U.S.A.* 108, 11017-11022 (2011).

- 144 2. Church, J. et al. Sea Level Change. in *Climate Change 2013: The Physical Science Basis.*
145 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on*
146 *Climate Change* (eds. Stocker, T.F. et al.) 1137-1216 (Cambridge University Press, 2013).
- 147 3. Kopp, R.E. et al. Temperature-driven sea-level variability in the Common Era. *Proc. Natl. Acad.*
148 *Sci. U.S.A.* 113, 1434-1441 (2016).
- 149 4. Lambeck, K., Rouby, H., Purcell, A., Sun, Y. & Sambridge, M. Sea level and global ice volumes
150 from the last glacial maximum to the Holocene. *Proc. Natl. Acad. Sci. U.S.A.* 111, 43, 15296-
151 15303 (2014).
- 152 5. Hallmann, N. et al. Ice volume and climate changes from a 6000 year sea-level record in French
153 Polynesia. *Nat. Commun.* 9, 285 (2018).
- 154 6. Mann, M.E. et al. Global signatures and dynamical origins of the Little Ice Age and Medieval
155 Climate Anomaly. *Science* 326, 1256-1260 (2009).
- 156 7. Horton, B.P. et al. Mapping sea-level change in time, space, and probability. *Annu. Rev. Environ.*
157 *Resour.* 43, 481-521 (2018).
- 158 8. Dutton, A., Webster, J.M., Zwart, D., Lambeck, K. & Wohlfarth, B. Tropical tales of polar ice:
159 evidence of Last Interglacial polar ice sheet retreat recorded by fossil reefs of the granitic
160 Seychelles islands. *Quat. Sci. Rev.* 107, 182-196 (2015).
- 161 9. Nicholls, R.J. et al. Sea-level scenarios for evaluating coastal impacts. *WIREs Clim. Change*, 5,
162 129-150 (2014).
- 163 10. Schuerch, M. et al. Future response of global coastal wetlands to sea-level rise. *Nature* 561, 231-
164 234 (2018).
- 165 11. Hinkel, J. et al. The ability of societies to adapt to twenty-first-century sea level rise. *Nat. Clim.*
166 *Chang.* 8, 570-578 (2018).

- 167 12. Khan, N.S. et al. Holocene relative sea level changes from near-, intermediate-, and far-field
168 locations. *Curr. Clim. Change Rep.* **1**, 247-262 (2015).
- 169 13. Kemp, A.C. et al., Relative sea-level change in Newfoundland, Canada during the past ~3000
170 years. *Quat. Sci. Rev.* **201**, 89-110 (2018).
- 171 14. Gerlach, M.J. et al. Reconstructing Common Era relative sea-level change on the Gulf Coast of
172 Florida. *Mar. Geol.* **390**, 254-269 (2017).
- 173 15. Woodroffe, C.D. Late Quaternary sea-level highstands in the central and eastern Indian Ocean: a
174 review. *Glob. Planet Change* **49**, 121-138 (2005).
- 175 16. Woodroffe, C., McGregor, H.V., Lambeck, K., Smithers, S.G., & Fink, D. Mid-Pacific microatolls
176 record sea-level stability over the past 5000 yr. *Geology*, **40**, 10, 951-954 (2012).
- 177 17. Kench, P.S. et al. Holocene reef growth in the Maldives: evidence of a mid-Holocene sea level
178 highstand in the central Indian Ocean. *Geology*, **37**, 455-458 (2009).
- 179 18. Gischler, G., Hudson, J.H. & Pisera, A. Late Quaternary reef growth and sea level in the Maldives
180 (Indian Ocean). *Mar. Geol.* **250**, 104-113 (2008).
- 181 19. Yokohama, Y. et al. Holocene Indian Ocean sea level, Antarctic melting history and past tsunami
182 deposits inferred using sea level reconstructions from the Sri Lanka, Southeastern Indian and
183 Maldivian coasts. *Quat. Sci. Rev.* **206**, 150-161 (2019).
- 184 20. Meltzner, A.J. & Woodroffe, C.D. Coral microatolls, in *Handbook of Sea-Level Research* (eds
185 Shennan, I., Long, A.J., & Horton, B.P.) 125-145 (Wiley, 2015).
- 186 21. Peltier, W.R., Argus, D.F. & Drummond, R. Space geodesy constrains ice-age terminal
187 deglaciation: The global ICE-6G_C (VM5a) model. *J. Geophys. Res. Solid Earth* **120**, 450-487
188 (2015).

- 189 22. Roy, K. & Peltier, W.R. Space-geodetic and water level gauge constraints on continental uplift and
190 tilting over North America: Regional convergence of the ICE-6G_C (VM5a/VM6) models. *Geophys. J.*
191 *Int.* 210, 2, 1115-1142 (2017).
- 192 23. Büntgen, U. et al. 2500 years of European climate variability and human susceptibility. *Science*
193 331, 578-582 (2011).
- 194 24. Ljungqvist, F.C. A new reconstruction of temperature variability in the extra-tropical northern
195 hemisphere during the last two millennia. *Geogr. Ann. A* 92, 339-351 (2010).
- 196 25. Oppo, D.W., Rosenthal, Y. & Linsley, B.K. 2,000-year-long temperature and hydrology
197 reconstructions from the Indo-Pacific warm pool. *Nature* 460, 1113-1116, (2009).
- 198 26. Linsley, B.K., Rosenthal, Y. & Oppo, D.W. Holocene evolution of the Indonesian throughflow and
199 the western Pacific warm pool. *Nat. Geo.* 3, 578–583 (2010).
- 200 27. Steinhilber, F., Beer, J. & Fröhlich, C. Total solar irradiance during the Holocene. *Geophys. Res.*
201 *Lett.* 36, L19704 (2009).
- 202 28. Gautier, E. et al., 2600-years of stratospheric volcanism through sulfate isotopes, *Nat. Commun.*
203 10, 466 (2019).
- 204 29. Mann, M.E. et al. Proxy-based reconstructions of hemispheric and global surface temperature
205 variations over the past two millennia. *Proc. Natl. Acad. Sci. U.S.A.* 105, 13252-13257 (2008).
- 206 30. Sigl, M. et al. Timing and climate forcing of volcanic eruptions for the past 2,500 years. *Nature*
207 523, 543-549 (2015).
- 208 31. Büntgen, U. et al. Cooling and societal change during the Late Antique Little Ice Age from 536 to
209 around 660 AD. *Nat. Geosci.* 9, 231-236 (2016).
- 210 32. Neukom, R. et al. Consistent multidecadal variability in global temperature reconstructions and
211 simulations over the Common Era. *Nat. Geosci.* 12, 643-649 (2019).

33. Toohey, M. et al. Disproportionately strong climate forcing from extratropical explosive volcanic eruptions. *Nat. Geosci.* **12**, 100-107 (2019).
34. Miller, G.H. et al. Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks. *Geophys. Res. Lett.* **39**, L02708 (2012).
35. Gebble, G. & Huybers, P. The Little Ice Age and 20th Century deep Pacific cooling. *Science*, **363**, 70-74 (2019).
36. Thompson, P.R. Forcing of recent decadal variability in the Equatorial and Northern Indian Ocean. *J. Geophys. Res. Oceans* **121**, 6762-6778 (2016).
37. Widlansky, M.J. Timmermann, A. & Cai, W. Future extreme sea-level seesaws in the tropical Pacific. *Sci. Adv.* **1**, 500560 (2015).

Acknowledgements

We acknowledge LaMer Group and the Small Island Research Centre, Fares-Maathodaa, Huvadhoo atoll for logistical support, Government of the Maldives for research permission and Ana Vila-Concejo, Eddie Beetham and Tracey Turner for field assistance.

Corresponding Author

All correspondence and requests for materials should be addressed to Paul Kench, Department of Earth Science, Simon Fraser University, Burnaby, Canada, pkench@sfu.ca

Author contributions

PK conceived the project; PK, RM, SO, ER, KM undertook fieldwork; XW and LK performed U-Th dating; KR undertook GIA modelling of field location; PK, RM and SO led manuscript development and interpretation; all authors contributed to manuscript revision.

Competing interests

Authors declare no competing interests.

Figure captions

Figure 1. Field location and Holocene sea-level history of the Maldives archipelago, Indian Ocean. a, Location of Maldives archipelago, central Indian Ocean. b, Locality of the Mahutigalaa reef platform, Huvadhoo atoll (orange box), southern Maldives and location of emergent mid-late Holocene corals in central Maldives¹⁷ (orange triangle). Base data British Admiralty Chart 709. c, Oblique aerial photograph of the Mahutigalaa reef platform (source: P. Kench 2016). d, Summary Holocene sea-level curve for the Maldives based on published coral data^{17,18} and output from the glacial isostatic adjustment (GIA) model ICE-6G_C (VM5a) for Mahutigalaa^{21,22}, see Methods for details. msl is mean sea level. Years (CE) is Common Era and Cal. Years BP denotes calibrated years before 1950. No observational data indicates no physical coral evidence to support sea-level reconstructions, dated from the last 1800 years, has been reported from the Maldives.

Figure 2. Reconstructed sea level in the central Indian Ocean and global paleoclimate variability over the past 2500 years. a, elevation and age of fossil microatolls from Mahutigalaa reef platform, Maldives. Red boxes are two fossil microatolls from Funadhoo reef (central Maldives, Fig. 1A) that mark the end of the mid-late Holocene highstand and dated using AMS¹⁷; orange boxes are microatolls dated using U-Th method in this study (Supplementary Table 1), vertical error bars ± 0.14 m. Age error bars are smaller than the size of symbols and width of boxes reflect microatoll diameter. Dashed boxes define corals included in lowstand calculations. Cal. Years BP is calibrated years before 1950. b, Reconstructed northern hemisphere (European) June, July, August temperature anomalies (red line and left x-axis) relative to the 1901-2000 period (grey dashed line), error is ± 1 root mean square error (RMSE, red dashed lines)²³; and northern hemisphere decadal temperature variations (blue line, right x-axis) with 2σ error (blue shading)²⁴. c, 2,000-year sea surface temperature reconstruction in Makassar Strait, Indonesia²⁵ (blue line); SST reconstruction²⁶, for the West Pacific Warm Pool (red line) and southern Makassar Strait (orange line). d, Total solar irradiance²⁷ (blue line), occurrence of major volcanic events through the Holocene (green bars) and volcanic events making major contributions to stratospheric sulfate levels based on $\Delta^{17}\text{O}$ analyses (brown squares), error bars are 1 s.d.²⁸. Vertical grey shading denotes time periods of lower sea levels in the Mahutigalaa coral record and climatic cool periods. LIA = Little Ice Age, MCA = Medieval Climate Anomaly, LALIA = Late Antiquity Little Ice Age.

263

264 **Methods**

265 Glacio-isostatic adjustment modelling

266 Glacio-isostatic adjustment (GIA) modelling for the field location was undertaken using the ICE-6G_C
267 (VM5a) model²¹ by one of the authors (KR) and presented in Figure 1D. The model is global in nature,
268 takes into account a full suite of geophysical impacts related to GIA (self-gravitation, migrating
269 shorelines, rotational feedback, etc.), and provides fully consistent sea-level histories globally²¹. The
270 model is based upon iterative refinements against a global suite of constraints (e.g. relative sea level,
271 rotational observables, space-geodesy), and successive incremental iterations of the model have
272 shown the general convergence of the methodology^{21,22}.

273 Field sampling of corals

274 Field surveys and sampling were undertaken during three field campaigns in 2016-2018. Detailed
275 description of the field site and a discussion of the environmental context of the fossil microatolls and
276 causes of variability in elevation are presented in the Supplementary Information. A total of 49 fossil
277 microatolls were identified on the reef platform ranging in diameter from 0.9 to 4.95 m (Supplementary
278 Table 1, Supplementary Figs 2 and 3). All fossil microatolls were found in currently free draining
279 locations. The diameter of microatolls reflects the period of growth of each colony at its limiting water
280 level. Consequently, larger diameter microatolls reflect a longer period of sea-level stability at which
281 that colony grew. For example, the larger-sized microatolls (e.g. ~5.0 m) reflect growth over a period of
282 ~170 years assuming average skeletal extension of 1.5 cm/yr. For each specimen the location was
283 documented using a Trimble Global Positioning System (GPS) survey with a horizontal accuracy of
284 ±1.5 m and the diameter was recorded. Plug cores (3 cm x 6 cm) were extracted from the microatoll
285 surface using a Nemo V2 handheld underwater drill. Cores were extracted from the centre and outer
286 edge of each specimen to record the initial timing that the microatolls achieved their vertical growth limit

(centre) and the timing of colony death (outer edge). Cores were bathed in hydrogen peroxide for 24 hours to neutralise organic material, rinsed, dried and stored for dating.

The elevation of fossil microatolls was determined using a combination of Real Time Kinematic (RTK) GPS and laser level surveys. Three benchmarks were inserted into the vegetated cay and static surveys undertaken using RTK GPS over a 12-hour period to determine the elevation of each benchmark with respect to sea level. The vertical elevation error of these surveys was ± 0.02 m. As microatolls were found in water depths ranging up to 2.0 m below high tide, RTK GPS survey was not possible, thus laser level surveys (vertical error of ± 0.03 m) were used to connect the elevation of microatoll upper surfaces to island benchmarks. The elevation of each colony was determined at its central core and outer edge (mean of 5 measurements) and linked directly to the nearest land-based benchmark. Of note, surveys were repeated on three occasions (each field season) to resolve survey error and relative differences between fossil microatoll colonies.

A biological elevation reference (height of living coral, HLC) was established for the field site based on surveys of the elevation of the living upper surface of 30 live *Porites* spp. microatolls, within 50 m of the fossil coral locations. At each coral a minimum of five elevation points were recorded around each specimen to account for within colony variability. The elevation range of the upper surface of living *Porites* colonies was ± 0.095 m. The surveys were undertaken in each field season to examine consistency in elevation data. This sampling strategy provides a reproducible method and ensures a direct comparative basis between the growth elevation of fossil and modern microatolls. The HLC was calculated as the average elevation of living *Porites* spp. microatolls and records the contemporary upper growth limit for microatolls, which inversely represents the constraining water level for coral growth. The HLC for Mahutigalaa was calculated as 0.567 ± 0.095 m below mean sea level, which is close to Mean Low Water Spring (MLWS) tide level. The elevation of fossil microatolls is presented as both a difference from present MSL and HLC. The combined survey error in evaluating change in elevation between fossil and living microatolls is calculated as ± 0.14 m.

312 Uranium-Thorium dating method

313 A total of 49 ^{230}Th ages were determined for 37 corals (including 12 replicate analyses, Supplementary
314 Data 1). Samples were gently crushed into 1 to 3-mm³ segments. Clean and pristine pieces were
315 selected under microscope, and subsequently ultrasonically cleaned (three times) in superclean
316 water. For each sample, 30 to 60 mg was used for U-Th chemistry^{40,41}. Replicates were prepared for
317 six subsamples (1,12,13,14,16, and 18). All samples were dissolved in ultrapure nitric acids and a
318 ^{229}Th - ^{233}U - ^{236}U mixed spike was then added to the solutions. HClO_4 was also added to remove organic
319 compounds. Each sample solution was refluxed at 175°C for 2 hours and dried. Samples were re-
320 dissolved by adding 2N ultrapure HCl solution. Co-precipitation was achieved by adding FeCl_3 and
321 concentrated NH_4OH . Co-precipitates were re-dissolved and put through an anion-exchange resin in
322 columns to separate and purify uranium and thorium. U and Th fractions were then dissolved in
323 1% HNO_3 (+0.1% HF) for instrumental analysis. U and Th isotopic compositions were determined on a
324 multi-collector inductively coupled plasma mass spectrometer (MC-ICP-MS, a Neptune Plus from
325 Thermo Fisher Scientific) at the Earth Observatory of Singapore and Asian School of the Environment
326 (ASE), Nanyang Technological University, Singapore. The measurements were undertaken using a
327 recently improved SEM peak-jumping technique^{42,43}. Uncertainties in the U and Th isotopic data and
328 ^{230}Th dates are quoted at the 2σ level in this paper.

329 Estimating the growth age of microatolls

330 U-Th dating determined the age of the centre of each microatoll as this site represents the time at
331 which the coral was first constrained by water level. The diameter and flat surface of each microatoll
332 represents a period of time that sea level was stable and continued to constrain vertical growth. The
333 outer edge of each colony therefore, represents the terminal age of an individual coral at a specific
334 water level. This terminal age may also correspond to the end of a particular sea level stand.

An estimate of the age of the outer edge of each microatoll was calculated from the distance between the central sample (of known ^{230}Th age) and outer edge of the colony, divided by an average extension rate of 1.5 cm.yr^{-1} (range $0.5\text{-}2.5 \text{ cm.yr}^{-1}$), as previously reported for massive corals in the mid-Holocene^{44,45} and recently reported from contemporary growth estimates in the Maldives⁴⁶.

To validate these estimates accelerator mass spectrometry (AMS) radiocarbon dating of the outer edge of six microatolls (Supplementary Table 2) was compared with the ^{230}Th ages of the centre of the same coral (Supplementary Table 1). The difference in age across the distance between samples on each microatoll surface provides an estimate of average extension rates (Supplementary Table 3). Of note the estimated extension rates ranged from 1.23 to 1.99 cm.yr^{-1} , with an average of 1.47 cm.yr^{-1} , consistent with previous historic estimates and contemporary measurements from the Maldives⁴⁶ and supports the use of an average value of 1.5 cm.yr^{-1} in this study.

Determination of sea-level lowstands

Two measures were adopted to define the elevation of lower sea level from the coral evidence. First, the mean elevation of all microatolls defining a specific sea-level lowstand was calculated to account for variations in microatoll elevation. Of note, corals considered to be impacted by localised ponding were excluded from calculations of sea-level position, consistent with previous studies²⁹. Second, estimates of the maximum sea-level fall in each period were established based on the deepest microatoll documented in each lowstand period.

Determination of rates of sea-level change

Determination of rates of sea-level change from the coral records was based on age and depth constraints of corals and the growth age of corals, as reflected in the diameter of each colony (Supplementary Table 4). Both the highest and deepest microatolls were identified as markers of the maximum variations in sea level throughout the record. The onset of changes in sea level were determined based on the age of the outer edge of the highest or deepest microatoll. The stabilization of

sea level was denoted by the central ^{230}Th age of the deepest (sea-level fall) or highest (sea-level rise) corals. The selection of the central age is consistent with the time that coral colonies were constrained by sea level.

Data availability statement

The authors declare that the data supporting the findings of this study are available within the paper and its supplementary information files.

Methods References

40. Edwards, R. L., Chen, J. H., & Wasserburg, G. J. ^{238}U - ^{234}U - ^{230}Th - ^{232}Th systematics and the precise measurement of time over the past 500,000 years. *Earth Planet Sci Letters* **81**, 175-192 (1987).
41. Shen, C.C. et al. High-precision and high-resolution carbonate ^{230}Th dating by MC-ICP-MS with SEM protocols. *Geochim. Cosmochim. Ac.* **99**, 71-86 (2012).
42. Cheng, H. et al. Improvements in ^{230}Th dating, ^{230}Th and ^{234}U half-life values, and U-Th isotopic measurements by multi-collector inductively coupled plasma mass spectrometry. *Earth Planet. Sci Lett.* **371**, 82-91 (2013).
43. Chiang, H. W., Lu, Y., Wang, X., Lin, K., & Liu, X. Optimizing MC-ICP-MS with SEM protocols for determination of U and Th isotope ratios and ^{230}Th ages in carbonates. *Quat. Geochronol.* **50**, 75-90 (2019).
44. Zhao, J. & Collins, L. Uranium series dating. in *Encyclopedia of Modern Coral Reefs* (ed. Hopley, D.) 1128-1132 (Springer, 2011).
45. Bessat, F. & Buigues, D. Two centuries of variation in coral growth in a massive *Porites* colony from Moorea (French Polynesia): A response of ocean-atmosphere variability from south central Pacific. *Palaeo. Palaeo. Palaeo.* **175**, 381-392 (2001).

- 382 46. Morgan, K.M. & Kench, P.S. Skeletal extension and calcification of reef-building corals in the
383 central Indian Ocean. *Marine Env. Res.* **81**, 71-82 (2012).