Third cumulant stein approximation for Poisson stochastic integrals

Privault, Nicolas

2019

Privault, N. (2019). Third cumulant stein approximation for Poisson stochastic integrals. *Journal of Theoretical Probability*, 32(3), 1461-1481.
<https://dx.doi.org/10.1007/s10959-018-0817-1>

<https://hdl.handle.net/10356/148588>

<https://doi.org/10.1007/s10959-018-0817-1>

© 2018 Springer Science+Business Media. This is a post-peer-review, pre-copyedit version of an article published in *Journal of Theoretical Probability*. The final authenticated version is available online at: <http://dx.doi.org/10.1007/s10959-018-0817-1>.

Downloaded on 26 Apr 2023 16:11:09 SGT

Third cumulant Stein approximation for Poisson stochastic integrals

Nicolas Privault

Division of Mathematical Sciences

School of Physical and Mathematical Sciences

Nanyang Technological University

21 Nanyang Link

Singapore 637371

nprivault@ntu.edu.sg

August 25, 2018

Abstract

We derive Edgeworth-type expansions for Poisson stochastic integrals, based on cumulant operators defined by the Malliavin calculus. As a consequence we obtain Stein approximation bounds for stochastic integrals, which are based on third cumulants instead of the L^3 norm term found in the literature. The use of the third cumulant results into a convergence rate faster than the classical Berry-Esseen rate on certain examples.

Key words: Stein approximation; Malliavin calculus; Poisson stochastic integral; cumulants; Edgeworth expansions.

Mathematics Subject Classification: 60H07, 62E17, 60H05.

1 Introduction

Edgeworth expansions have been derived on the Wiener space in [9], [2], [4], using a construction of cumulant operators based on the inverse L^{-1} of the Ornstein-Uhlenbeck operator [11]. This approach extends the results of [10], [13] on Stein approximation, Berry-Esseen bounds and the fourth moment theorem. Related Edgeworth type expansions have also been derived for the Itô-Skorohod integral $\delta(u)$ of a

process u on the Wiener space in [18].

In this paper we derive Edgeworth type expansions of the form

$$E[\delta(u)g(\delta(u))] = E[\|u\|_{L^2(\mathbb{R}_+)}^2 g'(\delta(u))] + \sum_{k=2}^n E[g^{(k)}(\delta(u))\Gamma_{k+1}^u \mathbf{1}] + E[g^{(n+1)}(\delta(u))R_n^u] \quad (1.1)$$

for the compensated Poisson stochastic integral $\delta(u) = \int_0^\infty u_t d(N_t - t)$ of an adapted process $(u_t)_{t \in \mathbb{R}_+}$ with respect to a standard Poisson process $(N_t)_{t \in \mathbb{R}_+}$, where Γ_k^u and R_n^u are respectively a cumulant type operator and a remainder term defined using the derivation operators of the Malliavin calculus on the Poisson space, see Proposition 3.1.

From (1.1), in Proposition 4.1 and Corollary 4.2 we deduce Stein approximation bounds of the form

$$d(\delta(u), \mathcal{N}) \leq E[|1 - \|u\|_{L^2(\mathbb{R}_+)}^2|] + E\left[\left|\int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle\right|\right] + 2E[|R_1^u|],$$

and

$$\begin{aligned} d(\delta(u), \mathcal{N}) &\leq |1 - \text{Var}[\delta(u)]| + \sqrt{\text{Var}[\|u\|_H^2]} + E\left[\left|\int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle\right|\right] \\ &\quad + 2E[|R_1^u|], \end{aligned} \quad (1.2)$$

when the process u is adapted with respect to the Poisson filtration, where D is a gradient operator acting on Poisson functionals, $\mathcal{N} \simeq \mathcal{N}(0, 1)$ is a standard Gaussian random variable and

$$d(F, G) := \sup_{h \in \mathcal{L}} |E[h(F)] - E[h(G)]|$$

is the Wasserstein distance between the laws of two random variables F and G , where $\mathcal{L}$ denotes the class of 1-Lipschitz functions on $\mathbb{R}$.

In Section 5 we present examples of adapted processes $(u_t)_{t \in \mathbb{R}_+}$ for which (1.2) holds, see Proposition 5.1 and Corollary 5.2, in relation with classical examples such as the

normalized sequence $((T_k - k)/\sqrt{k})_{k \geq 1}$, where $(T_k)_{k \geq 1}$ is the sequence of jump times of the standard Poisson process $(N_t)_{t \in \mathbb{R}_+}$.

In particular, when f is a differentiable deterministic function we obtain bounds of the form

$$d\left(\int_0^\infty f(t)d(N_t - t), \mathcal{N}\right) \leq |1 - \|f\|_{L^2(\mathbb{R}_+)}^2| + \left|\int_0^\infty f^3(t)dt\right| + 2\|f\|_{L^2(\mathbb{R}_+)} \int_0^\infty t|f'(t)|^2 dt, \quad (1.3)$$

depending on the regularity of the function f , see Corollary 5.3. This alternative approach, which is based on derivation operators, replaces the $L^3(\mathbb{R}_+)$ norm of f in the classical Stein bound

$$d\left(\int_0^\infty f(t)d(N_t - t), \mathcal{N}\right) \leq |1 - \|f\|_{L^2(\mathbb{R}_+)}^2| + \int_0^\infty |f^3(t)|dt, \quad (1.4)$$

see Corollary 3.4 of [14], with the third cumulant $\kappa_3^f = \int_0^\infty f^3(t)dt$, by removing the inner absolute value in the integral.

The main reason for the appearance of an L^3 norm in (1.4) instead of the third cumulant $\kappa_3^f = \int_0^\infty f^3(t)dt$ lies with the use of finite difference operators and the replacement of the chain rule of derivation with a Taylor expansion bound, see Theorem 3.1 of [14] and § 4.2 of the recent survey [3]. In the present paper, the use of derivation operators allows us instead to use the third cumulant $\kappa_3^f = \int_0^\infty f^3(t)dt$ in (1.3).

Taking f_k of the form

$$f_k(t) := \frac{1}{\sqrt{k}}g(t/k), \quad k \geq 1,$$

where $g \in \mathcal{C}^1(\mathbb{R})$ is such that $\|g\|_{L^2(\mathbb{R}_+)} = 1$ and $\int_0^\infty g^3(t)dt = 0$, (1.3) shows that

$$d\left(\int_0^\infty f_k(t)d(N_t - t), \mathcal{N}\right) \leq \frac{2}{k} \int_0^\infty t|g'(t)|^2 dt, \quad k \geq 1,$$

see (5.4) below, while Corollary 3.4 of [14] only yields the standard Berry-Esseen rate, see (5.6). This similarly improves on the bound

$$d\left(\int_0^\infty f(t)d(N_t - t), \mathcal{N}\right) \leq |1 - \|f\|_{L^2(\mathbb{R}_+)}^2| + \sqrt{\int_0^\infty |f'(t)|^2 \left(\int_0^t f(s)ds\right)^2 dt} \quad (1.5)$$

obtained using derivation operators on the Poisson space in Corollary 4.4 of [20], which also yields the standard Berry-Esseen rate in this case, see (5.5) below.

In Section 2 we recall some background material on the Malliavin calculus and cumulant operators for stochastic integrals on the Poisson space. In Section 3 we derive Edgeworth type expansions, based on a family of cumulant operators that are associated to the process u and specially defined for the Skorohod integral operator δ . In Section 4 we derive Stein type approximation bounds for stochastic integrals, and in Section 5 we consider adapted and deterministic examples.

2 Malliavin operators

Let $(T_k)_{k \geq 1}$ denote the sequence of jump times of a standard Poisson process $(N_t)_{t \in \mathbb{R}_+}$ with unit intensity and generating the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ on a probability space $(\Omega, \mathcal{F}_\infty, P)$, with $T_0 := 0$. The gradient operator D defined on random functionals

$$F \in \mathcal{S} := \{F = f(T_1, \dots, T_n) : f \in \mathcal{C}_b^1(\mathbb{R}^n)\},$$

as

$$D_t F = - \sum_{k=1}^n \mathbf{1}_{[0, T_k]}(t) \frac{\partial f}{\partial x_k}(T_1, \dots, T_n),$$

has the derivation property, cf. [7], [5], [15]. The operator D defines the Sobolev spaces $\mathbb{D}_{p,1}$ with the Sobolev norms

$$\|F\|_{\mathbb{D}_{p,1}} := \|F\|_{L^p(\Omega)} + \|DF\|_{L^p(\Omega, H)}, \quad F \in \mathcal{S}.$$

$p > 1$, where $H := L^2(\mathbb{R}_+)$.

Covariant derivative

In addition to the operator D , we will also need the following notion of covariant derivative, see [19] and references therein. In the sequel we let $W_{2,1}(\mathbb{R}_+)$ denote the Sobolev space of weakly differentiable functions on $\mathbb{R}_+$ such that

$$\|f\|_{W_{2,1}}^2 := \int_0^\infty |f(t)|^2 dt + \int_0^\infty t |f'(t)|^2 dt < \infty.$$

Definition 2.1 Let the operator $\tilde{\nabla}$ be defined on

$$u \in \mathcal{U} := \left\{ \sum_{i=1}^n h_i F_i : F_i \in \mathcal{S}, h_i \in W_{2,1}(\mathbb{R}_+), i = 1, \dots, n, n \geq 1 \right\},$$

as

$$\tilde{\nabla}_s u_t := D_s u_t - \dot{u}_t \mathbf{1}_{[0,t]}(s), \quad s, t \in \mathbb{R}_+,$$

where $\dot{u}_t$ denotes the time derivative of $t \mapsto u_t$ with respect to t .

By closability, the operator $\tilde{\nabla}$ extends to the Sobolev spaces $\tilde{\mathbb{D}}_{p,1}(H)$ of processes, $p \geq 1$, by the Sobolev norms

$$\|u\|_{\tilde{\mathbb{D}}_{p,1}(H)} := \|u\|_{L^p(\Omega, W_{2,1}(\mathbb{R}_+))} + \|Du\|_{L^p(\Omega, H \otimes H)}, \quad u \in \mathcal{U}.$$

We note that when a process $u \in \tilde{\mathbb{D}}_{p,1}(H)$ is $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted we have

$$\tilde{\nabla}_s u_t = D_s u_t = 0, \quad s > t. \quad (2.1)$$

Definition 2.2 Given $k \geq 1$ and $u \in \tilde{\mathbb{D}}_{2,1}(H)$ we define the operator power $(\tilde{\nabla}u)^k$ in the sense of matrix powers with continuous indices, as

$$(\tilde{\nabla}u)^k h_s = \int_0^\infty \cdots \int_0^\infty (\tilde{\nabla}_{t_k} u_s \tilde{\nabla}_{t_{k-1}} u_{t_k} \cdots \tilde{\nabla}_{t_1} u_{t_2}) h_{t_1} dt_1 \cdots dt_k, \quad s \in \mathbb{R}_+, \quad h \in H.$$

In particular, for $h \in W_{2,1}(\mathbb{R}_+)$ and $v \in \mathcal{U}$ we have

$$(\tilde{\nabla}h)v_s = \int_0^\infty v_t \tilde{\nabla}_t h(s) dt = -\dot{h}(s) \int_0^\infty v_t \mathbf{1}_{[0,s]}(t) dt = -\dot{h}(s) \int_0^s v_t dt, \quad s \in \mathbb{R}_+. \quad (2.2)$$

We note that when $h \in W_{2,1}(\mathbb{R}_+)$ is a deterministic function, (2.2) can be iterated to show that

$$\begin{aligned} (\tilde{\nabla}h)^k f_s &= \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_k} h(s) \tilde{\nabla}_{t_{k-1}} h(t_k) \cdots \tilde{\nabla}_{t_1} h(t_2) f(t_1) dt_1 \cdots dt_k \\ &= (-1)^k \dot{h}(s) \int_0^\infty \cdots \int_0^\infty \mathbf{1}_{[0,s]}(t_k) \mathbf{1}_{[0,t_k]}(t_{k-1}) \dot{h}(t_k) \cdots \mathbf{1}_{[0,t_2]}(t_1) \dot{h}(t_2) f(t_1) dt_1 \cdots dt_k \\ &= (-1)^k \dot{h}(s) \int_0^s \dot{h}(t_k) \int_0^{t_k} \cdots \dot{h}(t_2) \int_0^{t_2} f(t_1) dt_1 \cdots dt_k, \quad s \in \mathbb{R}_+, \quad k \geq 1, \end{aligned}$$

with the bound

$$\|(\tilde{\nabla}h)^k f\|_{L^2(\mathbb{R}_+)} \leq \|\tilde{\nabla}h\|_{L^2(\mathbb{R}_+^2)}^k \|f\|_{L^2(\mathbb{R}_+)} = \left(\int_0^\infty t |h'(t)|^2 dt \right)^{k/2} \|f\|_{L^2(\mathbb{R}_+)}.$$

This also yields

$$\begin{aligned}
\langle (\tilde{\nabla} h)^k f, h \rangle &= (-1)^k \int_0^\infty h(s) \dot{h}(s) \int_0^s \int_0^{t_k} \cdots \int_0^{t_2} \dot{h}(t_k) \cdots \dot{h}(t_2) f(t_1) dt_1 \cdots dt_k \\
&= \frac{1}{2} (-1)^{k-1} \int_0^\infty h^2(t_k) \dot{h}(t_k) \int_0^{t_k} \cdots \int_0^{t_2} \dot{h}(t_{k-1}) \cdots \dot{h}(t_2) f(t_1) dt_1 \cdots dt_k \\
&= \frac{1}{(k+1)!} \int_0^\infty h^{k+1}(t_1) f(t_1) dt_1, \quad f \in L^2(\mathbb{R}_+), \quad k \geq 1,
\end{aligned}$$

and in particular, for $h \in W_{2,1}(\mathbb{R}_+)$ we have $h \in \bigcap_{p=2}^\infty L^p(\mathbb{R}_+)$, with

$$\int_0^\infty |h^k(t)| dt \leq (k-1)! \langle |(\tilde{\nabla}|h|)^{k-2} h|, |h| \rangle \leq (k-1)! \|h\|_{W_{2,1}}^k, \quad k \geq 2.$$

Note that due to (2.1), when the process $u \in \tilde{\mathbb{D}}_{p,1}(H)$ is $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted, we have

$$(\tilde{\nabla} u)^k v_s = \int_0^s \int_0^{t_k} \cdots \int_0^{t_2} (\tilde{\nabla}_{t_k} u_s \tilde{\nabla}_{t_{k-1}} u_{t_k} \cdots \tilde{\nabla}_{t_1} u_{t_2}) v_{t_1} dt_1 \cdots dt_k, \quad s \in \mathbb{R}_+,$$

and as a consequence, the process $((\tilde{\nabla} u)^k v_s)_{s \in \mathbb{R}_+}$ is also $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted. In the sequel we simply denote $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_H = \langle \cdot, \cdot \rangle_{L^2(\mathbb{R}_+)}$.

Lemma 2.3 *Letting $n \geq 1$ and $u \in \tilde{\mathbb{D}}_{2,1}(H)$, we have*

$$\langle (\tilde{\nabla} u)^n u, u \rangle = \frac{1}{(n+1)!} \int_0^\infty u_t^{n+2} dt + \sum_{k=2}^{n+1} \frac{1}{k!} \left\langle (\tilde{\nabla} u)^{n+1-k} u, D \int_0^\infty u_t^k dt \right\rangle. \quad (2.3)$$

Proof. Using the adjoint $\tilde{\nabla}^* u$ of $\tilde{\nabla} u$ on H given by

$$(\tilde{\nabla}^* u) v_s := \int_0^\infty (\tilde{\nabla}_s u_t) v_t dt, \quad s \in \mathbb{R}_+, \quad v \in H,$$

with the duality relation

$$\langle v, (\tilde{\nabla}^* u) h \rangle = \langle (\tilde{\nabla} u) v, h \rangle, \quad h, v \in H,$$

we will show by induction on $1 \leq k \leq n+1$ that

$$\begin{aligned}
(\tilde{\nabla}^* u)^n u_{t_0} &= \int_0^\infty \cdots \int_0^\infty u_{t_n} \tilde{\nabla}_{t_0} u_{t_1} \tilde{\nabla}_{t_1} u_{t_2} \cdots \tilde{\nabla}_{t_{n-1}} u_{t_n} dt_1 \cdots dt_n \\
&= \sum_{i=2}^k \frac{1}{i!} \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-i}} u_{t_{n+1-i}} D_{t_{n+1-i}} u_{t_{n+2-i}}^i dt_1 \cdots dt_{n-i}
\end{aligned}$$

$$+\frac{1}{k!} \int_0^\infty \cdots \int_0^\infty u_{t_{n+1-k}}^k \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-k}} u_{t_{n+1-k}} dt_1 \cdots dt_{n+1-k}. \quad (2.4)$$

This relation holds for $k = 1$. Next, assuming that the identity (2.4) holds for some $k \in \{1, \dots, n\}$, and using the relation

$$\tilde{\nabla}_{t_{n-k}} u_{t_{n+1-k}} = D_{t_{n-k}} u_{t_{n+1-k}} - \mathbf{1}_{[0, t_{n+1-k}]}(t_{n-k}) \dot{u}_{t_{n+1-k}}, \quad t_{n-k}, t_{n+1-k} \in \mathbb{R}_+,$$

we have

$$\begin{aligned} (\tilde{\nabla}^* u)^n u_{t_0} &= \sum_{i=2}^k \frac{1}{i!} \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-i}} u_{t_{n+1-i}} D_{t_{n+1-i}} u_{t_{n+2-i}}^i dt_1 \cdots dt_{n+2-i} \\ &\quad + \frac{1}{k!} \int_0^\infty \cdots \int_0^\infty u_{t_{n+1-k}}^k \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-k}} u_{t_{n+1-k}} dt_1 \cdots dt_{n+1-k} \\ &= \sum_{i=2}^k \frac{1}{i!} \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-i}} u_{t_{n+1-i}} D_{t_{n+1-i}} u_{t_{n+2-i}}^i dt_1 \cdots dt_{n+2-i} \\ &\quad + \frac{1}{k!} \int_0^\infty \cdots \int_0^\infty u_{t_{n+1-k}}^k \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-k-1}} u_{t_{n-k}} D_{t_{n-k}} u_{t_{n+1-k}} dt_1 \cdots dt_{n+1-k} \\ &\quad - \frac{1}{k!} \int_0^\infty \cdots \int_0^\infty \int_{t_{n-k}}^\infty \dot{u}_{t_{n+1-k}} u_{t_{n+1-k}}^k \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-1-k}} u_{t_{n-k}} dt_1 \cdots dt_{n+1-k} \\ &= \sum_{i=2}^k \frac{1}{i!} \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-i}} u_{t_{n+1-i}} D_{t_{n+1-i}} u_{t_{n+2-i}}^i dt_1 \cdots dt_{n+2-i} \\ &\quad + \frac{1}{(k+1)!} \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-k-1}} u_{t_{n-k}} D_{t_{n-k}} u_{t_{n+1-k}}^{k+1} dt_1 \cdots dt_{n+1-k} \\ &\quad - \frac{1}{(k+1)!} \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-k-1}} u_{t_{n-k}} \int_{t_{n-k}}^\infty (u_t^{k+1})' dt dt_1 \cdots dt_{n-k} \\ &= \sum_{i=2}^{k+1} \frac{1}{i!} \int_0^\infty \cdots \int_0^\infty \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-i}} u_{t_{n+1-i}} D_{t_{n+1-i}} u_{t_{n+2-i}}^i dt_1 \cdots dt_{n+2-i} \\ &\quad + \frac{1}{(k+1)!} \int_0^\infty \cdots \int_0^\infty u_{t_{n-k}}^{k+1} \tilde{\nabla}_{t_0} u_{t_1} \cdots \tilde{\nabla}_{t_{n-k-1}} u_{t_{n-k}} dt_1 \cdots dt_{n-k} \\ &= \sum_{i=2}^{k+1} \frac{1}{i!} (\tilde{\nabla}^* u)^{n+1-i} D_{t_0} \int_0^\infty u_s^i ds + \frac{1}{(k+1)!} (\tilde{\nabla}^* u)^{n-k} u_{t_0}^{k+1}, \end{aligned}$$

which shows by induction that (2.4) holds for $k = 1, \dots, n$. In particular, for $k = n$ we have

$$(\tilde{\nabla}^* u)^n u_t = \frac{1}{(n+1)!} u_t^{n+1} + \sum_{i=2}^{n+1} \frac{1}{i!} (\tilde{\nabla}^* u)^{n+1-i} D_t \int_0^\infty u_s^i ds, \quad t \in \mathbb{R}_+,$$

which yields (2.3) by integration with respect to $t \in \mathbb{R}_+$ and duality. $\square$

Cumulant operators

We recall the definition of the cumulant operators introduced in § 6 of [17] on the Poisson space. Given $k \geq 2$ and $u \in \widetilde{\mathbb{D}}_{2,1}(H)$ an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process we define the cumulant operators

$$\Gamma_k^u : \mathbb{D}_{2,1} \longrightarrow L^2(\Omega), \quad k \geq 2,$$

by

$$\Gamma_k^u F := F \langle (\widetilde{\nabla} u)^{k-2} u, u \rangle + \langle (\widetilde{\nabla} u)^{k-1} u, DF \rangle, \quad k \geq 2.$$

As a consequence of Lemma 2.3 we have

$$\Gamma_k^u \mathbf{1} = \frac{1}{(k-1)!} \int_0^\infty u_t^k dt + \sum_{i=2}^{k-1} \frac{1}{i!} \left\langle (\widetilde{\nabla} u)^{k-1-i} u, D \int_0^\infty u_t^i dt \right\rangle,$$

and $|\Gamma_k^u \mathbf{1}| \leq \|u\|_{W_{2,1}(\mathbb{R}_+)}^k$, a.s., $k \geq 2$.

When $h \in W_{2,1}(\mathbb{R}_+)$ is a deterministic function we find

$$\Gamma_k^h \mathbf{1} = \frac{1}{(k-1)!} \int_0^\infty h^k(t) dt = \frac{1}{(k-1)!} \kappa_k(h), \quad k \geq 2,$$

where $\kappa_k(h) = \int_0^\infty h^k(s) ds$ is the cumulant of order $k \geq 2$ of the compensated Poisson stochastic integral $\int_0^\infty h(t) d(N_t - t)$.

Poisson-Skorohod integral

In addition, D has a closable adjoint operator δ with domain $\text{Dom}(\delta)$, that satisfies the duality relation

$$E \left[\int_0^\infty u_t D_t F dt \right] = E[F \delta(u)], \quad F \in \mathbb{D}_{2,1}, \quad u \in \text{Dom}(\delta), \quad (2.5)$$

and coincides with the compensated Poisson stochastic integral on square-integrable processes $(u_t)_{t \in \mathbb{R}_+}$ adapted to the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ generated by $(N_t)_{t \in \mathbb{R}_+}$, i.e., we have

$$\delta(u) = \int_0^\infty u_t d(N_t - t).$$

In addition, the operators $\tilde{\nabla}$, δ and D satisfy the commutation relation

$$D_t \delta(u) = u_t + \delta(\tilde{\nabla}_t^* u), \quad (2.6)$$

for $u \in \tilde{\mathbb{D}}_{2,1}(H)$ an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process, cf. Lemma 4.5 in [17] and Relations (2.16), (2.19) and Lemma 2.4 in [19].

Recall that when $(u_t)_{t \in \mathbb{R}_+}$ is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process, $\delta(u)$ coincides with the Itô integral of u .

3 Edgeworth type expansions

Classical Edgeworth expansions are used in particular as asymptotic expansions around the Gaussian cumulative distribution function $\Phi(x)$ for the cumulative distribution function $P(F \leq x)$ of a centered random variable F with $E[F^2] = 1$, as

$$\Phi(x) + c_1 \phi(x) H_1(x) + \cdots + c_m \phi(x) H_m(x) + \cdots,$$

where $\phi(x)$, $x \in \mathbb{R}$, is the standard Gaussian density, $H_k(x)$ is the Hermite polynomial of degree $k \geq 1$, and c_k is a coefficient depending on the sequence of cumulants $(\kappa_n)_{n \geq 1}$ of a random variable F , cf. Chapter 5 of [8] and § A.4 of [12].

Edgeworth type expansions of the form

$$E[Fg(F)] = \sum_{l=1}^n \frac{\kappa_{l+1}}{l!} E[g^{(l)}(F)] + E[g^{(n+1)}(F) \Gamma_{n+1} F], \quad n \geq 1,$$

have been obtained by the Malliavin calculus in [9], [2], [4], written here for F a centered random variable, where Γ_{n+1} is a cumulant type operator on the Wiener space such that $n!E[\Gamma_n F]$ coincides with the cumulant κ_{n+1} of order $n+1$, $n \in \mathbb{N}$, cf. [11], extending results of [1].

An infinite Edgeworth type expansion for the compensated Poisson stochastic integral of a deterministic function $f \in \bigcap_{p=2}^{\infty} L^p(\mathbb{R}_+)$ can be written as

$$E[\delta(f)g(\delta(f))] = E \left[\int_0^\infty f(s)(g(\delta(f) + f(s)) - g(\delta(f))) ds \right] \quad (3.1)$$

$$\begin{aligned}
&= \sum_{k=1}^{\infty} \frac{1}{k!} \int_0^{\infty} f^{k+1}(s) ds E[g^{(k)}(\delta(f))] \\
&= \sum_{k=1}^{\infty} \frac{1}{k!} \kappa_{k+1}(f) E[g^{(k)}(\delta(f))], \quad g \in \mathcal{C}_b^{\infty}(\mathbb{R}),
\end{aligned}$$

using a standard integration by parts for finite difference operators on the Poisson space.

In this section we establish an Edgeworth type expansion of any finite order with an explicit remainder term for the compensated Poisson stochastic integral $\delta(u)$ of an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process u .

Before proceeding to the statement of general expansions in Proposition 3.1, we derive an expansion of order one for a deterministic integrand $f \in W_{2,1}(\mathbb{R}_+)$. In the sequel we let $\mathcal{C}_b^n(\mathbb{R}_+)$ denote the space of $\mathcal{C}_b^n$ functions bounded together with their derivatives on $\mathbb{R}_+$, $n \geq 1$.

By the duality relation (2.5) between D and δ , the chain rule of derivation for D and the commutation relation (2.6) we get, for $g \in \mathcal{C}_b^2(\mathbb{R})$,

$$\begin{aligned}
E[\delta(f)g(\delta(f))] &= E[g'(\delta(f))\langle f, D\delta(f) \rangle] \\
&= E[g'(\delta(f))\langle f, f \rangle] + E[g'(\delta(f))\langle f, \delta(\tilde{\nabla} * f) \rangle] \\
&= E[g'(\delta(f))\langle f, f \rangle] + E[\langle \tilde{\nabla}^* f, D(g'(\delta(f))f) \rangle] \\
&= E[g'(\delta(f))\langle f, f \rangle] + E[g''(\delta(f))\langle (\tilde{\nabla} f)f, D\delta(f) \rangle] \\
&= E[g'(\delta(f))\langle f, f \rangle] + \frac{1}{2} \int_0^{\infty} f^3(t) dt E[g''(\delta(f))] + E[g''(\delta(f))\langle (\tilde{\nabla} f)f, \delta(\tilde{\nabla}^* f) \rangle] \\
&= \kappa_2(f) E[g'(\delta(f))] + \frac{1}{2} \kappa_3(f) E[g''(\delta(f))] + E[g''(\delta(f))\langle (\tilde{\nabla} f)f, \delta(\tilde{\nabla}^* f) \rangle],
\end{aligned}$$

since by Lemma 2.3 we have

$$\langle (\tilde{\nabla} f)f, f \rangle = - \int_0^{\infty} f(t) f'(t) \int_0^t f(s) ds dt = \frac{1}{2} \int_0^{\infty} f^3(t) dt = \frac{1}{2} \kappa_3^f.$$

In the next proposition we derive general Edgeworth type expansions for adapted integrands processes $(u_t)_{t \in \mathbb{R}_+}$.

Proposition 3.1 *Let $n \geq 0$ and assume that $u \in \widetilde{\mathbb{D}}_{n+1,1}(H)$ is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process. Then for all $g \in \mathcal{C}_b^{n+1}(\mathbb{R})$ and bounded $G \in \mathbb{D}_{2,1}$ we have*

$$\begin{aligned} E[G\delta(u)g(\delta(u))] &= E[g(\delta(u))\langle u, DG \rangle] + \sum_{k=1}^n E[g^{(k)}(\delta(u))\Gamma_{k+1}^u G] \\ &+ E \left[Gg^{(n+1)}(\delta(u)) \left(\int_0^\infty \frac{u_s^{n+2}}{(n+1)!} ds + \sum_{k=2}^{n+1} \frac{1}{k!} \left\langle (\widetilde{\nabla} u)^{n+1-k} u, D \int_0^\infty u_t^k dt \right\rangle + \langle (\widetilde{\nabla} u)^n u, \delta(\widetilde{\nabla}^* u) \rangle \right) \right]. \end{aligned}$$

Proof. By the duality relation (2.5) between D and δ , the chain rule of derivation for D and the commutation relation (2.6), we get

$$\begin{aligned} &E[Gg(\delta(u))\langle (\widetilde{\nabla} u)^k u, D\delta(u) \rangle] - E[Gg'(\delta(u))\langle (\widetilde{\nabla} u)^{k+1} u, D\delta(u) \rangle] \\ &= E[Gg(\delta(u))\langle (\widetilde{\nabla} u)^k u, u \rangle] + E[Gg(\delta(u))\langle (\widetilde{\nabla} u)^k u, \delta(\widetilde{\nabla}^* u) \rangle] - E[Gg'(\delta(u))\langle (\widetilde{\nabla} u)^{k+1} u, D\delta(u) \rangle] \\ &= E[Gg(\delta(u))\langle (\widetilde{\nabla} u)^k u, u \rangle] + E[\langle \widetilde{\nabla}^* u, D(Gg(\delta(u))(\widetilde{\nabla} u)^k u) \rangle] - E[Gg'(\delta(u))\langle (\widetilde{\nabla} u)^{k+1} u, D\delta(u) \rangle] \\ &= E[Gg(\delta(u))\langle (\widetilde{\nabla} u)^k u, u \rangle] + E[g(\delta(u))\langle (\widetilde{\nabla} u)^{k+1} u, DG \rangle] + E[Gg(\delta(u))\langle \widetilde{\nabla}^* u, D((\widetilde{\nabla} u)^k u) \rangle] \\ &= E[g(\delta(u))\Gamma_{k+2}^u G], \end{aligned}$$

since $\langle \widetilde{\nabla}^* u, D((\widetilde{\nabla} u)^k u) \rangle = 0$ as u is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted process, see Lemma 4.4 of [16]. Therefore, we have

$$\begin{aligned} E[G\delta(u)g(\delta(u))] &= E[Gg'(\delta(u))\langle u, D\delta(u) \rangle] + E[g(\delta(u))\langle u, DG \rangle] \\ &= E[g(\delta(u))\langle u, DG \rangle] + E[Gg^{(n+1)}(\delta(u))\langle (\widetilde{\nabla} u)^n u, D\delta(u) \rangle] \\ &\quad + \sum_{k=0}^{n-1} \left(E[Gg^{(k+1)}(\delta(u))\langle (\widetilde{\nabla} u)^k u, D\delta(u) \rangle] - E[Gg^{(k+2)}(\delta(u))\langle (\widetilde{\nabla} u)^{k+1} u, D\delta(u) \rangle] \right) \\ &= E[g(\delta(u))\langle u, DG \rangle] + \sum_{k=1}^n E[g^{(k)}(\delta(u))\Gamma_{k+1}^u G] + E[Gg^{(n+1)}(\delta(u))\langle (\widetilde{\nabla} u)^n u, D\delta(u) \rangle] \\ &= E[g(\delta(u))\langle u, DG \rangle] + \sum_{k=1}^n E[g^{(k)}(\delta(u))\Gamma_{k+1}^u G] \\ &\quad + E[Gg^{(n+1)}(\delta(u))\langle (\widetilde{\nabla} u)^n u, u \rangle] + E[Gg^{(n+1)}(\delta(u))\langle (\widetilde{\nabla} u)^n u, \delta(\widetilde{\nabla}^* u) \rangle], \end{aligned}$$

and we conclude by Lemma 2.3. $\square$

When $G = 1$, Proposition 3.1 shows that

$$E[\delta(u)g(\delta(u))] = \sum_{k=1}^{n+1} \frac{1}{k!} E \left[\int_0^\infty u_s^{k+1} ds g^{(k)}(\delta(u)) \right]$$

$$\begin{aligned}
& + \sum_{k=2}^{n+1} \sum_{i=2}^k \frac{1}{i!} E \left[\left\langle (\tilde{\nabla} u)^{k-i} u, D \int_0^\infty u_t^i dt \right\rangle g^{(k)}(\delta(u)) \right] \\
& + E[g^{(n+1)}(\delta(u)) \langle (\tilde{\nabla} u)^n u, \delta(\tilde{\nabla}^* u) \rangle], \quad n \geq 0.
\end{aligned}$$

On the other hand, when $f \in W_{2,1}(\mathbb{R}_+)$ is a deterministic function and $g \in \mathcal{C}_b^\infty(\mathbb{R})$ we find

$$\begin{aligned}
E[\delta(f)g(\delta(f))] &= \sum_{k=1}^{n+1} \frac{1}{k!} \int_0^\infty f^{k+1}(s) ds E[g^{(k)}(\delta(f))] + E[g^{(n+1)}(\delta(f)) \langle (\tilde{\nabla} f)^n f, \delta(\tilde{\nabla}^* f) \rangle] \\
&= \sum_{k=1}^{n+1} \frac{1}{k!} \kappa_{k+1}(f) E[g^{(k)}(\delta(f))] + E[g^{(n+1)}(\delta(f)) \langle (\tilde{\nabla} f)^n f, \delta(\tilde{\nabla}^* f) \rangle],
\end{aligned}$$

$n \geq 0$, showing, as n tends to $+\infty$, that

$$E[\delta(f)g(\delta(f))] = \sum_{k=1}^{\infty} \int_0^\infty \frac{f^{k+1}(s)}{k!} ds E[g^{(k)}(\delta(f))] = E \left[\int_0^\infty f(s) (g(\delta(f) + f(s)) - g(\delta(f))) ds \right],$$

which recovers (3.1) and the standard integration by parts identity for finite difference operators on the Poisson space.

4 Stein approximation

We let $\mathcal{N} \simeq \mathcal{N}(0, 1)$ denote a standard Gaussian random variable. In comparison with the results of [2], our bounds apply to a different stochastic integral representation.

In the case $n = 0$, Proposition 3.1 reads

$$E[g'(\delta(u)) \langle u, u \rangle - \delta(u)g(\delta(u))] = -E[g'(\delta(u)) \langle u, \delta(\tilde{\nabla}^* u) \rangle],$$

for $u \in \tilde{\mathbb{D}}_{2,1}(H)$ and $g \in \mathcal{C}_b^1(\mathbb{R})$. Applying this relation to the solution g_x of the Stein equation

$$\mathbf{1}_{(-\infty, x]}(z) - \Phi(z) = g'_x(z) - z g_x(z), \quad z \in \mathbb{R},$$

which satisfies $\|g_x\|_\infty \leq \sqrt{2\pi}/4$ and $\|g'_x\|_\infty \leq 1$, cf. Lemma 2.2-(v) of [6], yields the expansion

$$P(\delta(u) \leq x) - \Phi(x) = E[(1 - \langle u, u \rangle) g'_x(\delta(u))] - E[\langle u, \delta(\tilde{\nabla} u) \rangle g'_x(\delta(u))], \quad x \in \mathbb{R},$$

around the Gaussian cumulative distribution function $\Phi(x)$, with $u \in \widetilde{\mathbb{D}}_{2,1}(H)$.

On the other hand, given $h : \mathbb{R} \rightarrow \mathbb{R}$ an absolutely continuous function with bounded derivative, the functional equation

$$h(z) - E[h(\mathcal{N})] = g'(z) - zg(z), \quad z \in \mathbb{R}, \quad (4.1)$$

has a solution $g_h \in \mathcal{C}_b^1(\mathbb{R})$ which is twice differentiable and satisfies the bounds

$$\|g'_h\|_\infty \leq \|h'\|_\infty \quad \text{and} \quad \|g''_h\|_\infty \leq 2\|h'\|_\infty, \quad x \in \mathbb{R},$$

cf. Lemma 1.2-(v) of [10] and references therein.

Proposition 4.1 *Let $u \in \widetilde{\mathbb{D}}_{2,1}(H)$ be adapted with respect to the Poisson filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$. We have*

$$\begin{aligned} d(\delta(u), \mathcal{N}) &\leq E[|1 - \langle u, u \rangle|] + E \left[\left| \int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle \right| \right] \\ &\quad + 2E[|\langle (\widetilde{\nabla} u)u, \delta(\widetilde{\nabla}^* u) \rangle|]. \end{aligned} \quad (4.2)$$

Proof. For $n = 1$ and $G = 1$, Proposition 3.1 shows that

$$\begin{aligned} E[\delta(u)g(\delta(u))] &= E[g'(\delta(u))\langle u, u \rangle] + \frac{1}{2}E \left[g''(\delta(u)) \left(\int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle \right) \right] \\ &\quad + E[g''(\delta(u))\langle (\widetilde{\nabla} u)u, \delta(\widetilde{\nabla}^* u) \rangle], \end{aligned}$$

hence for any absolutely continuous function $h : \mathbb{R} \rightarrow \mathbb{R}$ with bounded derivative, denoting by g_h the solution to (4.1) we have

$$\begin{aligned} E[h(\delta(u))] - E[h(\mathcal{N})] &= E[\delta(u)g_h(\delta(u)) - g'_h(\delta(u))] \\ &= E[g'_h(\delta(u))(\langle u, u \rangle - 1)] + \frac{1}{2}E \left[g''(\delta(u)) \left(\int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle \right) \right] \\ &\quad + 2E[g''_h(\delta(u))\langle (\widetilde{\nabla} u)u, \delta(\widetilde{\nabla}^* u) \rangle], \end{aligned}$$

hence

$$\begin{aligned} |E[\delta(u)h(\delta(u))] - E[h(\mathcal{N})]| &\leq \|h'\|_\infty E[|1 - \langle u, u \rangle|] \\ &\quad + \|h'\|_\infty E \left[\left| \int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle \right| \right] \\ &\quad + 2\|h'\|_\infty E[|\langle (\widetilde{\nabla} u)u, \delta(\widetilde{\nabla}^* u) \rangle|], \end{aligned}$$

which yields (4.2). □

As a consequence of Proposition 4.1 and the Itô isometry we have the following corollary.

Corollary 4.2 *Let $u \in \widetilde{\mathbb{D}}_{2,1}(H)$ be adapted with respect to the Poisson filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$. We have*

$$\begin{aligned} d(\delta(u), \mathcal{N}) \leq & |1 - \text{Var}[\delta(u)]| + \sqrt{\text{Var}[\|u\|_H^2]} + E \left[\left| \int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle \right| \right] \\ & + 2E[|\langle (\widetilde{\nabla} u)u, \delta(\widetilde{\nabla}^* u) \rangle|]. \end{aligned} \quad (4.3)$$

Proof. By the Itô isometry we have

$$\text{Var}[\delta(u)] = E \left[\left(\int_0^\infty u_t d(N_t - t) \right)^2 \right] = E[\langle u, u \rangle],$$

hence

$$\begin{aligned} E[|1 - \langle u, u \rangle|] &\leq E[|1 - E[\langle u, u \rangle]|] + E[|\langle u, u \rangle - E[\langle u, u \rangle]|] \\ &= |1 - \text{Var}[\delta(u)]| + \sqrt{E[(\langle u, u \rangle - E[\langle u, u \rangle])^2]} \\ &= |1 - \text{Var}[\delta(u)]| + \sqrt{\text{Var}[\|u\|_H^2]}. \end{aligned}$$

□

In particular, when $\text{Var}[\delta(u)] = 1$, (4.3) shows that

$$d(\delta(u), \mathcal{N}) \leq \sqrt{\text{Var}[\|u\|_H^2]} + E \left[\left| \int_0^\infty u_s^3 ds + \left\langle u, D \int_0^\infty u_t^2 dt \right\rangle \right| \right] + 2E[|\langle (\widetilde{\nabla} u)u, \delta(\widetilde{\nabla}^* u) \rangle|].$$

5 Examples

Adapted integrands

Although the present approach does not apply directly to the classical normalized sequence

$$F_k := \frac{T_k - k}{\sqrt{k}} = -\frac{1}{\sqrt{k}} \int_0^\infty \mathbf{1}_{[0, T_k]}(t) d(N_t - t), \quad k \geq 1,$$

due to the lack of time differentiability of the adapted integrand $t \mapsto \mathbf{1}_{[0, T_k]}(t)$, examples of this form can be treated via smoothed processes u_k as in (5.1) below, see also Corollary 5.2.

Proposition 5.1 *Let*

$$u_k(t) := g(t)\mathbf{1}_{[0,T_k]}(t) + g(T_k)\mathbf{1}_{(T_k,\infty)}(t)f(t - T_k), \quad t \in \mathbb{R}_+, \quad (5.1)$$

$k \in \mathbb{N}$, where g is a Lipschitz function on $\mathbb{R}_+$ and $f \in W_{2,1}(\mathbb{R}_+)$ satisfies $f(0) = 1$.

Then for every $k \in \mathbb{N}$ we have

$$\begin{aligned} & d(\delta(u_k), \mathcal{N}) \\ & \leq E \left[\left| 1 - \int_0^{T_k} g^2(t) dt - \|f\|_{L^2(\mathbb{R}_+)}^2 g^2(T_k) \right| \right] + E \left[\left| \int_0^{T_k} (g^2(s) - g^2(T_k)) g(s) ds \right| \right] \\ & \quad + E \left[|g^3(T_k)| \right] \left| \int_0^\infty f^3(t) dt \right| + 2\|f\|_{L^2(\mathbb{R}_+)}^2 E \left[\left| g(T_k) g'(T_k) \int_0^{T_k} g(t) dt \right| \right] \\ & \quad + 2E \left[\left| \int_0^{T_k} g'(t) \int_0^t g(s) ds \int_t^{T_k} g'(s) d(N_s - s) dt \right| \right] \\ & \quad + 2\|f\|_{L^2(\mathbb{R}_+)} E \left[\left| g'(T_k) \int_0^{T_k} g'(t) \int_0^t g(s) ds dt \right| \right] \\ & \quad + 2\|f\|_{L^2(\mathbb{R}_+)} E \left[\left| g(T_k) g'(T_k) \int_0^{T_k} g(s) ds \right| \right] \sqrt{\int_0^\infty t |f'(t)|^2 dt} \\ & \quad + 2\|f\|_{L^2(\mathbb{R}_+)} E \left[|g^3(T_k)| \right] \int_0^\infty t |f'(t)|^2 dt. \end{aligned} \quad (5.2)$$

Proof. We have

$$\delta(u_k) = \int_0^\infty u_k(t) d(N_t - t) = \int_0^{T_k} g(t) d(N_t - t) + g(T_k) \int_{T_k}^\infty f(t - T_k) d(N_t - t),$$

and

$$\text{Var}[\delta(u_k)] = E \left[\int_0^{T_k} g^2(t) dt \right] + E[g^2(T_k)] \int_0^\infty f^2(t) dt.$$

On the other hand, we have

$$\begin{aligned} \tilde{\nabla}_s u_k(t) &= D_s u_k(t) - (g'(t)\mathbf{1}_{[0,T_k]}(t) + g(T_k)f'(t - T_k)\mathbf{1}_{(T_k,\infty)}(t))\mathbf{1}_{[0,t]}(s) \\ &= -(g'(T_k)\mathbf{1}_{(T_k,\infty)}(t)f(t - T_k) - g(T_k)\mathbf{1}_{(T_k,\infty)}(t)f'(t - T_k))\mathbf{1}_{[0,T_k]}(s) \\ &\quad - (g'(t)\mathbf{1}_{[0,T_k]}(t) + g(T_k)f'(t - T_k)\mathbf{1}_{(T_k,\infty)}(t))\mathbf{1}_{[0,t]}(s) \\ &= -(g'(t)\mathbf{1}_{[0,T_k]}(t) + g'(T_k)\mathbf{1}_{(T_k,\infty)}(t)f(t - T_k))\mathbf{1}_{[0,t]}(s)\mathbf{1}_{[0,T_k]}(s) - g(T_k)\mathbf{1}_{\{T_k < s < t\}}f'(t - T_k), \end{aligned}$$

and

$$(\tilde{\nabla} u_k)u_k(t) = - \int_0^\infty g(s)(g'(t)\mathbf{1}_{[0,T_k]}(t) + g'(T_k)\mathbf{1}_{(T_k,\infty)}(t)f(t - T_k))\mathbf{1}_{[0,t]}(s)\mathbf{1}_{[0,T_k]}(s) ds$$

$$\begin{aligned}
& -g^2(T_k)f'(t-T_k) \int_0^\infty f(s-T_k) \mathbf{1}_{\{T_k < s < t\}} ds \\
= & -\mathbf{1}_{[0, T_k]}(t)g'(t) \int_0^t g(s)ds - \mathbf{1}_{(T_k, \infty)}(t)g'(T_k)f(t-T_k) \int_0^{T_k} g(s)ds \\
& + \mathbf{1}_{(T_k, \infty)}(t)g^2(T_k)f'(t-T_k) \int_{T_k}^t f(s-T_k)ds.
\end{aligned}$$

Given that

$$\begin{aligned}
\delta(\tilde{\nabla}_s^* u_k) &= - \left(\int_s^{T_k} g'(t)d(N_t - t) + g'(T_k) \int_{T_k}^\infty f(t-T_k)d(N_t - t) \right) \mathbf{1}_{[0, T_k]}(s) \\
&+ \mathbf{1}_{\{T_k < s\}}g(T_k) \int_s^\infty f'(t-T_k)d(N_t - t),
\end{aligned}$$

this yields

$$\begin{aligned}
& \langle (\tilde{\nabla} u_k)u_k, \delta(\tilde{\nabla}^* u_k) \rangle \\
= & \int_0^{T_k} \left(\int_t^{T_k} g'(s)d(N_s - s) + g'(T_k) \int_{T_k}^\infty f(s-T_k)d(N_s - s) \right) g'(t) \int_0^t g(s)dsdt \\
& - g(T_k) \int_{T_k}^\infty \int_t^\infty f'(s-T_k)d(N_s - s)g'(T_k)f(t-T_k) \int_0^{T_k} g(s)dsdt \\
& + g^3(T_k) \int_{T_k}^\infty \int_t^\infty f'(s-T_k)d(N_s - s)f'(t-T_k) \int_{T_k}^t f(s-T_k)dsdt \\
\cong & \int_0^{T_k} g'(t) \int_t^{T_k} g'(s)d(\hat{N}_s - s) \int_0^t g(s)dsdt \\
& + g'(T_k) \int_0^\infty f(s)d(\hat{N}_s - s) \int_0^{T_k} g'(t) \int_0^t g(s)dsdt \\
& - g(T_k)g'(T_k) \int_0^{T_k} g(s)ds \int_0^\infty f(t) \int_t^\infty f'(s)d(\hat{N}_s - s)dt \\
& + g^3(T_k) \int_0^\infty f'(t) \int_t^\infty f'(s)d(\hat{N}_s - s) \int_0^t f(s)dsdt,
\end{aligned}$$

where “ $\cong$ ” denotes equality in distribution and $(\hat{N}_t)_{t \in \mathbb{R}_+}$ is a standard Poisson process independent of $(T_k)_{k \geq 1}$. We also have

$$\begin{aligned}
D_s \int_0^\infty u_k^2(t)dt &= D_s \left(\int_0^{T_k} g^2(t)dt + g^2(T_k) \int_{T_k}^\infty f^2(t-T_k)dt \right) \\
&= D_s \left(\int_0^{T_k} g^2(t)dt + g^2(T_k) \int_0^\infty f^2(t)dt \right)
\end{aligned}$$

$$\begin{aligned}
&= g^2(T_k)D_s T_k + 2g(T_k)g'(T_k) \int_0^\infty f^2(t)dt D_s T_k \\
&= - \left(g^2(T_k) + 2g(T_k)g'(T_k) \int_0^\infty f^2(t)dt \right) \mathbf{1}_{[0, T_k]}(s),
\end{aligned}$$

consequently we have

$$\left\langle u_k, D \int_0^\infty u_k^2(t)dt \right\rangle = - \left(g^2(T_k) + 2g(T_k)g'(T_k) \int_0^\infty f^2(t)dt \right) \int_0^{T_k} g(t)dt,$$

and

$$\begin{aligned}
&\int_0^\infty u_k^3(s)ds + \left\langle u_k, D \int_0^\infty u_k^2(t)dt \right\rangle \\
&= \int_0^{T_k} (g^2(s) - g^2(T_k))g(s)ds + g^3(T_k) \int_0^\infty f^3(t)dt - 2g(T_k)g'(T_k) \int_0^\infty f^2(t)dt \int_0^{T_k} g(t)dt.
\end{aligned}$$

Hence by Corollary 4.2 we find

$$\begin{aligned}
&d(\delta(u_k), \mathcal{N}) \\
&\leq |1 - \text{Var}[\delta(u_k)]| + \sqrt{\text{Var}[\|u_k\|_H^2]} + E \left[\left| \int_0^\infty u_k^3(s)ds + \left\langle u_k, D \int_0^\infty u_k^2(t)dt \right\rangle \right| \right] \\
&\quad + 2E[| \langle (\tilde{\nabla} u_k)u_k, \delta(\tilde{\nabla}^* u_k) \rangle |] \\
&\leq E \left[\left| 1 - \int_0^{T_k} g^2(t)dt - g^2(T_k) \int_0^\infty f^2(t)dt \right| \right] \\
&\quad + E \left[\left| \int_0^{T_k} (g^2(s) - g^2(T_k))g(s)ds \right| \right] \\
&\quad + E[|g^3(T_k)|] \left| \int_0^\infty f^3(t)dt \right| + 2E \left[\left| g(T_k)g'(T_k) \int_0^{T_k} g(t)dt \right| \right] \int_0^\infty f^2(t)dt \\
&\quad + 2E \left[\left| \int_0^{T_k} \int_t^{T_k} g'(s)d(N_s - s)g'(t) \int_0^t g(s)dsdt \right| \right] \\
&\quad + 2E \left[\left| g'(T_k) \int_0^{T_k} g'(t) \int_0^t g(s)dsdt \right| \right] E \left[\left| \int_0^\infty f(s)d(N_s - s) \right| \right] \\
&\quad + 2E \left[\left| g(T_k)g'(T_k) \int_0^{T_k} g(s)ds \right| \right] E \left[\left| \int_0^\infty f(t) \int_t^\infty f'(s)d(N_s - s)dt \right| \right] \\
&\quad + 2E[|g^3(T_k)|] E \left[\left| \int_0^\infty f'(t) \int_t^\infty f'(s)d(N_s - s) \int_0^t f(s)dsdt \right| \right].
\end{aligned}$$

Finally, we note that

$$E \left[\left| \int_0^\infty f(t) \int_t^\infty f'(s)d(N_s - s)dt \right| \right] = E[| \langle f, \delta(\tilde{\nabla}^* f) \rangle |]$$

$$\begin{aligned}
&\leq \|f\|_{L^2(\mathbb{R}_+)} \sqrt{E[\|\delta(\tilde{\nabla}^* f)\|_{L^2(\mathbb{R}_+)}^2]} \\
&\leq \|f\|_{L^2(\mathbb{R}_+)} \sqrt{E\left[\int_0^\infty |\delta(\tilde{\nabla}_t^* f)|^2 dt\right]} \\
&= \|f\|_{L^2(\mathbb{R}_+)} \sqrt{\int_0^\infty \int_0^\infty |\tilde{\nabla}_t f(s)|^2 ds dt} \\
&= \|f\|_{L^2(\mathbb{R}_+)} \sqrt{\int_0^\infty \int_0^\infty |f'(s)|^2 \mathbf{1}_{[0,s]}(t) ds dt} \\
&= \|f\|_{L^2(\mathbb{R}_+)} \sqrt{\int_0^\infty s |f'(s)|^2 ds},
\end{aligned}$$

and

$$\begin{aligned}
E\left[\left|\int_0^\infty f'(t) \int_t^\infty f'(s) d(N_s - s) \int_0^t f(s) ds dt\right|\right] &= E[|\langle (\tilde{\nabla} f) f, \delta(\tilde{\nabla}^* f) \rangle|] \\
&\leq \|(\tilde{\nabla} f) f\|_{L^2(\mathbb{R}_+)} \sqrt{E\left[\int_0^\infty |\delta(\tilde{\nabla}_t^* f)|^2 dt\right]} \\
&\leq \|f\|_{L^2(\mathbb{R}_+)} \|\tilde{\nabla} f\|_{L^2(\mathbb{R}_+^2)}^2 = \|f\|_{L^2(\mathbb{R}_+)} \int_0^\infty s |f'(s)|^2 ds.
\end{aligned}$$

The above bound can also be obtained as

$$\begin{aligned}
E[|\langle (\tilde{\nabla} f) f, \delta(\tilde{\nabla}^* f) \rangle|] &= E[|\delta((\tilde{\nabla} f)^2 f)|] \leq \sqrt{E[|\delta((\tilde{\nabla} f)^2 f)|^2]} \\
&= \|(\tilde{\nabla} f)^2 f\|_{L^2(\mathbb{R}_+)} \leq \|f\|_{L^2(\mathbb{R}_+)} \|\tilde{\nabla} f\|_{L^2(\mathbb{R}_+^2)}^2 \\
&= \|f\|_{L^2(\mathbb{R}_+)} \int_0^\infty s |f'(s)|^2 ds.
\end{aligned}$$

□

When u_k is an adapted process of the form

$$u_k(t) := g_k(t) \mathbf{1}_{[0, T_k]}(t), \quad t \in \mathbb{R}_+,$$

where g_k is Lipschitz, and e.g. $f(t) := e^{-nt}$, Proposition 5.2 shows, by letting n tend to infinity, that

$$d\left(\int_0^{T_k} g_k(t) d(N_t - t), \mathcal{N}\right) \leq E\left[\left|1 - \int_0^{T_k} g_k^2(t) dt\right|\right] + E\left[\left|\int_0^{T_k} (g_k^2(s) - g_k^2(T_k)) g_k(s) ds\right|\right]$$

$$+2E \left[\left| \int_0^{T_k} g'_k(t) \int_t^{T_k} g'_k(s) d(N_s - s) \int_0^t g_k(s) ds dt \right| \right].$$

As a consequence, similarly to Corollary 4.2 we have the following result.

Corollary 5.2 *Let g_k be a Lipschitz function on $\mathbb{R}_+$ for $k \geq 0$. We have*

$$\begin{aligned} d \left(\int_0^{T_k} g_k(t) d(N_t - t), \mathcal{N} \right) &\leq E \left[\left| 1 - E \left[\int_0^{T_k} g_k^2(t) dt \right] \right| \right] + \sqrt{\text{Var} \left[\int_0^{T_k} g_k^2(t) dt \right]} \\ &+ E \left[\left| \int_0^{T_k} (g_k^2(s) - g_k^2(T_k)) g_k(s) ds \right| \right] \\ &+ 2E \left[\left| \int_0^{T_k} g'_k(t) \int_t^{T_k} g'_k(s) d(N_s - s) \int_0^t g_k(s) ds dt \right| \right]. \end{aligned}$$

For example, when $g_k(t) = 1/\sqrt{k}$ is constant, Corollary 5.2 recovers the standard Berry-Esseen rate

$$d \left(\frac{T_k - k}{\sqrt{k}}, \mathcal{N} \right) \leq \frac{\sqrt{\text{Var}[T_k]}}{k} = \frac{1}{\sqrt{k}}, \quad k \geq 1.$$

Deterministic integrands

Taking $k = 0$ and $g_0(0) = 1$ in Proposition 5.1, we also find the following corollary.

Corollary 5.3 *Let $f \in W_{2,1}(\mathbb{R}_+)$ be a deterministic function. We have*

$$d \left(\int_0^\infty f(t) d(N_t - t), \mathcal{N} \right) \leq \left| 1 - \int_0^\infty f^2(t) dt \right| + \left| \int_0^\infty f^3(t) dt \right| + 2\|f\|_{L^2(\mathbb{R}_+)} \int_0^\infty t |f'(t)|^2 dt.$$

Considering for example $f_k(t)$ of the form

$$f_k(t) := \frac{1}{\sqrt{k}} g(t/k), \quad k \geq 1,$$

where $g \in \mathcal{C}^1(\mathbb{R})$ is such that $\|g\|_{L^2(\mathbb{R}_+)} = \|f_k\|_{L^2(\mathbb{R}_+)} = 1$, we have

$$\begin{aligned} d \left(\int_0^\infty f_k(t) d(N_t - t), \mathcal{N} \right) &\leq \left| \int_0^\infty f_k^3(s) ds \right| + 2 \int_0^\infty t |f'_k(t)|^2 dt \\ &= \frac{1}{\sqrt{k}} \left| \int_0^\infty g^3(t) dt \right| + \frac{2}{k} \int_0^\infty t |g'(t)|^2 dt, \quad (5.3) \end{aligned}$$

$k \geq 1$. When g has constant sign, e.g. $g(x) := \sqrt{2b}e^{-bt}$ with $b > 0$ and $f_k(t) = \sqrt{2b/k}e^{-bt/k}$, (5.3) does not improve on the standard Berry-Esseen convergence rate

$$d\left(\int_0^\infty g_k(t)d(N_t - t), \mathcal{N}\right) \leq \frac{2}{3}\sqrt{\frac{2b}{k}}, \quad \text{and} \quad d\left(\int_0^\infty g_k(t)d(N_t - t), \mathcal{N}\right) \leq \sqrt{\frac{b}{3k}},$$

respectively obtained from (1.4) and (1.5), see § 4.2 of [20]. On the other hand, choosing g such that

$$\int_0^\infty g^3(t)dt = 0,$$

we find the bound

$$d\left(\int_0^\infty f_k(t)d(N_t - t), \mathcal{N}\right) \leq \frac{2}{k} \int_0^\infty t|g'(t)|^2 dt, \quad k \geq 1, \quad (5.4)$$

while Corollary 4.4 of [20] and Corollary 3.4 of [14] only yield the Berry-Esseen rates

$$\begin{aligned} d\left(\int_0^\infty f_k(t)d(N_t - t), \mathcal{N}\right) &\leq |1 - \|f_k\|_{L^2(\mathbb{R}_+)}^2| + \|f_k\|_{L^2(\mathbb{R}_+)} \sqrt{\int_0^\infty t|f'_k(t)|^2 dt} \\ &= \frac{1}{\sqrt{k}} \sqrt{\int_0^\infty t|g'(t)|^2 dt}, \quad k \geq 1, \end{aligned} \quad (5.5)$$

by (1.5), and

$$\begin{aligned} d\left(\int_0^\infty f_k(t)d(N_t - t), \mathcal{N}\right) &\leq |1 - \|f_k\|_{L^2(\mathbb{R}_+)}^2| + \int_0^\infty |f_k^3(s)| ds \\ &= \frac{1}{\sqrt{k}} \int_0^\infty |g(t)|^3 dt, \end{aligned} \quad (5.6)$$

by (1.4). For another example, choosing

$$f_k(t) := \frac{1}{C\sqrt{k}} \sum_{i=1}^m a_i e^{-b_i t/k}, \quad k \geq 1,$$

with $\|f_k\|_{L^2(\mathbb{R}_+)} = 1$, for $b_i > 0$, $a_i \in \mathbb{R}$, $i = 1, \dots, m$, such that

$$\sum_{1 \leq i, j, l \leq m} \frac{a_i a_j a_l}{b_i + b_j + b_l} = 0,$$

(5.4) yields the bound

$$d\left(\int_0^\infty f_k(t)d(N_t - t), \mathcal{N}\right) \leq \frac{1}{C^2 k} \sum_{1 \leq i, j \leq m} \frac{a_i a_j b_i b_j}{(b_i + b_j)^2}, \quad k \geq 1.$$

In particular, when $f_k(t) := k^{-1/2}(e^{-t/k} - ae^{-bt/k})/C$ with $a, b > 0$ and

$$C^2 := \frac{1}{2} - \frac{2a}{1+b} + \frac{a^2}{2b} = \frac{b(b+1) - 4ab + a^2(1+b)}{2b(b+1)} > 0,$$

for any $b > 0$ we can choose $a > 0$ satisfying the equation

$$\int_0^\infty f_k^3(t)dt = \frac{1}{C^3\sqrt{k}} \left(\frac{1}{3} - \frac{3a}{2+b} + \frac{3a^2}{1+2b} - \frac{a^3}{3b} \right) = 0,$$

which yields the bound

$$d \left(\int_0^\infty f_k(t)d(N_t - t), \mathcal{N} \right) \leq \frac{c(a, b)}{k}, \quad k \geq 1,$$

where $c(a, b)$ depends only on $a, b > 0$.

References

- [1] A.D. Barbour. Asymptotic expansions based on smooth functions in the central limit theorem. *Probab. Theory Relat. Fields*, 72(2):289–303, 1986.
- [2] H. Biermé, A. Bonami, I. Nourdin, and G. Peccati. Optimal Berry-Esseen rates on the Wiener space: the barrier of third and fourth cumulants. *ALEA Lat. Am. J. Probab. Math. Stat.*, 9(2):473–500, 2012.
- [3] S. Bourguin and G. Peccati. The Malliavin-Stein method on the Poisson space. In *Stochastic analysis for Poisson point processes*, volume 7 of *Bocconi & Springer Series*, pages 185–228. Springer-Verlag, 2016.
- [4] S. Campese. Optimal convergence rates and one-term Edgeworth expansions for multidimensional functionals of Gaussian fields. *ALEA Lat. Am. J. Probab. Math. Stat.*, 10(2):881–919, 2013.
- [5] E. Carlen and É. Pardoux. Differential calculus and integration by parts on Poisson space. In S. Albeverio, Ph. Blanchard, and D. Testard, editors, *Stochastics, Algebra and Analysis in Classical and Quantum Dynamics (Marseille, 1988)*, volume 59 of *Math. Appl.*, pages 63–73. Kluwer Acad. Publ., Dordrecht, 1990.
- [6] L.H.Y. Chen, L. Goldstein, and Q.-M. Shao. *Normal approximation by Stein's method*. Probability and its Applications (New York). Springer, Heidelberg, 2011.
- [7] R.J. Elliott and A.H. Tsoi. Integration by parts for Poisson processes. *J. Multivariate Anal.*, 44(2):179–190, 1993.
- [8] P. McCullagh. *Tensor methods in statistics*. Monographs on Statistics and Applied Probability. Chapman & Hall, London, 1987.
- [9] I. Nourdin and G. Peccati. Stein's method and exact Berry-Esseen asymptotics for functionals of Gaussian fields. *Ann. Probab.*, 37(6):2231–2261, 2009.
- [10] I. Nourdin and G. Peccati. Stein's method on Wiener chaos. *Probab. Theory Related Fields*, 145(1-2):75–118, 2009.

- [11] I. Nourdin and G. Peccati. Cumulants on the Wiener space. *J. Funct. Anal.*, 258(11):3775–3791, 2010.
- [12] I. Nourdin and G. Peccati. *Normal approximations with Malliavin calculus*, volume 192 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2012.
- [13] I. Nourdin and G. Peccati. The optimal fourth moment theorem. *Proc. Amer. Math. Soc.*, 143:3123–3133, 2015.
- [14] G. Peccati, J. L. Solé, M. S. Taqqu, and F. Utzet. Stein’s method and normal approximation of Poisson functionals. *Ann. Probab.*, 38(2):443–478, 2010.
- [15] N. Privault. Chaotic and variational calculus in discrete and continuous time for the Poisson process. *Stochastics and Stochastics Reports*, 51:83–109, 1994.
- [16] N. Privault. Laplace transform identities and measure-preserving transformations on the Lie-Wiener-Poisson spaces. *J. Funct. Anal.*, 263:2993–3023, 2012.
- [17] N. Privault. Cumulant operators for Lie-Wiener-Itô-Poisson stochastic integrals. *J. Theoret. Probab.*, 28(1):269–298, 2015.
- [18] N. Privault. Stein approximation for Itô and Skorohod integrals by Edgeworth type expansions. *Electron. Comm. Probab.*, 20:Article 35, 2015.
- [19] N. Privault. De Rham-Hodge decomposition and vanishing of harmonic forms by derivation operators on the Poisson space. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 19(2):1–34, 2016.
- [20] N. Privault and G.L. Torrisi. Probability approximation by Clark-Ocone covariance representation. *Electron. J. Probab.*, 18:1–25, 2013.