Carbene-catalyzed reaction of indolyl methylenemalononitriles and enals for access to complex tetrahydrocarbazoles

Mou, Chengli; Zhou, Liejin; Song, Runjiang; Chai, Huifang; Hao, Lin; Chi, Robin Yonggui

2020

Mou, C., Zhou, L., Song, R., Chai, H., Hao, L. & Chi, R. Y. (2020). Carbene-catalyzed reaction of indolyl methylenemalononitriles and enals for access to complex tetrahydrocarbazoles. *Organic Letters*, 22(7), 2542-2547.
<https://dx.doi.org/10.1021/acs.orglett.0c00418>

<https://hdl.handle.net/10356/148641>

<https://doi.org/10.1021/acs.orglett.0c00418>

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Organic Letters*, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <https://doi.org/10.1021/acs.orglett.0c00418>

Downloaded on 26 Apr 2023 21:24:59 SGT

Carbene-Catalyzed Reaction of Indolyl Methylenemalononitriles and Enals for Access to Complex Tetrahydrocarbazoles

Chengli Mou,^{||} Liejin Zhou,^{||} Runjiang Song,^{||} Huifang Chai,^{*} Lin Hao, and Yonggui Robin Chi^{*}

Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: A carbene-catalyzed enantioselective cascade reaction of substituted methylenemalononitriles and α -bromo enals is disclosed. Key steps of this cascade process include a formal [4 + 2] cycloaddition, aldol reaction, and intramolecular lactonization. Our reaction offers streamlined and highly stereoselective access to complex tetrahydrocarbazole derivatives, with simultaneous formation of four chemical bonds and four chiral centers.

Chiral hydrocarbazoles are polycyclic indole architectures found in numerous natural products such as alkaloids that exhibit important bioactivities (Figure 1A).¹ For example, (+)-vincadifformine and its analogues display remarkable

cytotoxicity in vitro against a total of 60 human tumor cell lines derived from nine cancer types.² (+)-Aristoloteline has been used as anti-inflammation and pain-reducing drugs.³ Uleine has been proven to be the major antimalarial indole alkaloid isolated from *Aspidosperma parvifolium*.⁴ In recent years, enantioselective catalytic access to chiral hydrocarbazoles and their derivatives via transition-metal-free approaches have received considerable attention.⁵ Most of the success in this regard relies on the use of chiral amine catalysts to activate one of the reaction partners via iminium⁶ or enamine⁷ catalytic pathways, as disclosed by the groups of MacMillan,⁸ Melchiorre,⁹ Chen,¹⁰ Zanardi,¹¹ and others.¹² A potentially efficient approach to make this class of molecules is functionalization of indole derivatives.¹³ In 2013, we reported that indole-3-carboxaldehydes could be activated via N-heterocyclic carbene (NHC) organic catalysis¹⁴ to form *ortho*-quinodimethane intermediates for formal [4 + 2] reactions with reactive ketones¹⁵ (Figure 1B). Based on studies from us and others, we postulate that reacting similar indole *ortho*-quinodimethane intermediates¹⁶ with NHC-bound dienophiles¹⁷ can effectively provide hydrocarbazoles bearing all-carbon rings fused with the indole skeleton (Figure 1C). Key steps in our approach (Figure 1C) include a formal [4 + 2] cycloaddition, aldol reaction, and intramolecular

Figure 1. Rapid access to complex hydrocarbazole derivatives.

lactonization (see [Supporting Information](#) for a complete pathway). The catalytic reaction efficiently builds two new rings and four chiral centers. The reaction products contain multiple functional groups and can be readily transformed to diverse hydrocarbazole derivatives. Our catalytic strategy can be extended from indole-based substrates to other allylidene malononitriles, providing quick access to multicyclic hydro-pyranone products.

We started by using methylenemalononitrile **1a** and α -bromoaldehyde **2a** as model substrates to search for suitable conditions ([Table 1](#)). NHC **A** gave the desired tetrahydro-

enantioselectivities but in lower yields (e.g., entries 11–14). Finally, we established the optimal reaction conditions for this reaction: using NHC **A** as catalyst, **2a** (3.0 equiv), Et₃N (3.0 equiv) as the base, and THF as the solvent. In this case, the desired product was obtained in a good yield with excellent enantioselectivity. As a technical note, the desired product **3a** we obtained in these cases are single diastereomers (dr >20:1). With acceptable conditions in hand ([Table 1](#), entry 6), we first examined various substituted methylenemalononitriles **1** for this transformation ([Scheme 1](#)). Different substituents on

Table 1. Optimization of Reaction Conditions^a

entry	NHC	base	solvent	yield (%) ^b	ee ^c
1	A	Et ₃ N	THF	30	99
2	B	Et ₃ N	THF	22	99
3	C	Et ₃ N	THF	21	99
4	D	Et ₃ N	THF	21	99
5 ^d	A	Et ₃ N	THF	54	99
6 ^e	A	Et ₃ N	THF	73	99
7 ^f	A	Et ₃ N	THF	70	99
8 ^g	A	Et ₃ N	THF	69	99
9 ^e	A	DBU	THF	—	—
10 ^e	A	Cs ₂ CO ₃	THF	—	—
11 ^e	A	Et ₃ N	DCM	45	99
12 ^e	A	Et ₃ N	DCE	45	99
13 ^e	A	Et ₃ N	PhCH ₃	44	99
14 ^e	A	Et ₃ N	1,4-dioxane	33	99

^aGeneral conditions (unless otherwise specified): **1a** (0.05 mmol), **2a** (0.10 mmol), base (0.10 mmol), NHC (0.01 mmol), 4 Å MS (50 mg), THF (1.0 mL), rt, 24 h. Dr values for all the products (>20:1) were determined by crude ¹H NMR. ^bIsolated yield of **3a**. ^cEe was determined via HPLC on chiral stationary phase. ^d**1a** (0.05 mmol), **2a** (0.10 mmol), base (0.15 mmol). ^e**1a** (0.05 mmol), **2a** (0.15 mmol), base (0.15 mmol). ^f**2a** (0.2 mmol), base (0.15 mmol) was used. ^g**2a** (0.25 mmol), base (0.15 mmol) was used.

carbazole product **3a** in a promising yield and excellent ee (entry 1). Other NHC catalysts bearing different *N*-substituents could also provide the target products with excellent ee values, albeit with lower yields (entries 2–4). By increasing the amount of Et₃N to 3.0 equiv, the product could be obtained in a moderate yield (54% yield, entry 5). Then the amount of **2a** was examined (entries 6–8). To our delight, the yield of **3a** could be improved to 73% with 3 equiv of **2a** used (entry 6). A further increase of the amount of **2a** led to a slight drop of the reaction yield (entries 7–8). Strong organic or inorganic bases (e.g., entries 9, 10) could not be used for this reaction. A variety of organic solvents could be used in this reaction, and the product could be obtained in excellent

Scheme 1. Scope of Methylenemalononitriles **1**^a

^aReaction conditions as stated in [Table 1](#), entry 6. Yields are isolated yields after purification via SiO₂ column chromatography. Dr values for all the products (>20:1) were determined by crude ¹H NMR. Ee values were determined via HPLC on chiral stationary phase.

the indole core of **1** were well tolerated. Substitution at the 5- or 6-positions with electron-withdrawing or electron-donating groups gave the products **3b** to **3g** with moderate to good yields and excellent enantioselectivities. The *N*-Boc group of **1a** could be switched to an *N*-Ts group, with the desired product **3h** afforded in a moderate yield and excellent optical purity. Electron-donating *N*-substituents were not effective for this transformation. Benzofuranmalononitrile could also work in this cascade cycloaddition reaction, with the corresponding product **3i** afforded in a moderate yield with excellent enantioselectivity. We have also examined the benzothiophene-derived malononitrile substrate, but the reaction yield was very low. Replacement of the dicyano unit on the substrate **1a** with a cyanoester unit also led to no product formation.

We then examined the scope of various substituted α -bromoaldehydes **2** ([Scheme 2](#)). Electron-withdrawing and -donating groups were well tolerated on the benzene rings of the substrate **2**, with all the desired multicyclic tetrahydrocarbazoles **3** afforded in good yields and excellent ee values as single

Scheme 2. Scope of α -Bromoaldehydes 2^a

^aReaction conditions as stated in Table 1, entry 6. Yields are isolated yields after purification via SiO₂ column chromatography. Dr values for all the products (>20:1) were determined by crude ¹H NMR. Ee values were determined via HPLC on chiral stationary phase.

diastereomers (3j to 3o). In addition, the β -phenyl group of the α -bromoaldehyde 2a could be replaced with a 1-naphthyl group, with the corresponding product 3p afforded in a moderate yield and excellent enantioselectivity. Moreover, the β -phenyl group of the α -bromoaldehyde 2a could also be switched to a furanyl group or even an alkenyl group, although the optically pure products 3q and 3r could only be afforded in low yields under the current catalytic conditions.

To our delight, cinnamaldehyde could also be used instead of the α -bromoaldehydes 2 as the reaction substrates in this transformation, although the enantiomerically pure product 3s could only be afforded in 33% yield (Figure 2a). The application of other electrophiles such as imines or electron-deficient ketones in this cascade cyclization process remained unsuccessful (Figure 2b).

The versatility of the cascade reaction could be further demonstrated by using allylidene malononitriles 4 instead of 1 as the reaction substrates (Scheme 3). 1-Tetralone-derived allylidene malononitriles worked well under the current reaction conditions and gave the desired products 5a and 5b in good yields with excellent enantioselectivities. However, 1-

a) reaction using cinnamaldehyde instead of the α -Br enals 2

b) reactions using other electrophiles to replace half amount of 2a

Figure 2. Examination of various electrophiles.

Scheme 3. Reaction of Allylidene Malononitriles with α -Bromoaldehydes^a

^aReaction conditions as stated in Table 1, entry 6. Yields are isolated yields after purification via SiO₂ column chromatography. Dr values for all the products (>20:1) were determined by crude ¹H NMR. Ee values were determined via HPLC on chiral stationary phase.

indanone-derived allylidene malononitrile only gave the desired product 5c in 54% yield with 75% ee at this moment. β -Methylcinnamaldehyde-derived allylidene malononitrile could also be used in this transformation, with the desired product 5d afforded in a good yield and excellent enantioselectivity.

The chiral tetrahydrocarbazoles products can be used in various transformations (Scheme 4). The Boc group in 3a could be removed by acid and gave 6 in a good yield. 3a could be efficiently oxidized to carbazole derivative 7 with retention of the optical purity. Furthermore, the C–C bond of 3a could be hydrogenated and debrominated to give 8 in a good yield. Chiral alkynes 9 and 10 could be efficiently afforded from 3a through the Sonogashira coupling process without obvious erosion of the optical purities. The C–O bond of the lactone motif could be broken under mild basic conditions and give the heavily substituted product 11 and 12 in good yields and stereoselectivities. The malononitrile moiety of 3a could be smoothly converted to a carboxylate group, with compound 13 afforded in a good yield.

Scheme 4. Synthetic Transformations of Chiral Products 3a

^aDCM/TFA (*v* = 1:1), 40 °C, 2 h. ^bDDQ, PhCl, reflux, 12 h. ^cPt/C, H₂, EtOH/THF (*v* = 1:1), rt, 5 h. ^dCuI, PdCl₂(PPh₃)₂, Et₃N, THF, rt, 24 h. ^eFor 11: K₂CO₃, MeOH, rt, 0.5 h; for 12: BnNH₂, THF, rt, 48 h. ^fmCBA, K₂CO₃, MeOH, 0 °C, 0.5 h.

In summary, we have developed a new NHC-catalyzed cascade reaction for efficient synthesis of tetrahydrocarbazole derivatives. The optically pure products bear multiple cycles with four chiral centers. The afforded products could be used in various transformations and gave a variety of chiral functional molecules in good yields.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at

Experimental procedures and spectral data for all new compounds (PDF)

Accession Codes

CCDC 1921695 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Authors

Huifang Chai – Guizhou University of Traditional Chinese Medicine, Guizhou, Huaxi District, Guiyang 550025, China; Email: hfchai@126.com

Yonggui Robin Chi – Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371 Singapore; orcid.org/0000-0003-0573-257X; Email: robinchi@ntu.edu.sg

Authors

Chengli Mou – Guizhou University of Traditional Chinese Medicine, Guizhou, Huaxi District, Guiyang 550025, China;

Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371 Singapore

Liejun Zhou – Key Laboratory of the Ministry of Education for Advanced Catalysis Materials, Institute of Physical Chemistry, Zhejiang Normal University, Jinhua 321004, China; Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371 Singapore

Runjiang Song – Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371 Singapore

Lin Hao – Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371 Singapore

Complete contact information is available at:

Author Contributions

[†]C.M., L.Z., and R.S. contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Dr. Yongxin Li (NTU) for assistance with X-ray structure analysis. We acknowledge financial support by the Singapore National Research Foundation (NRF-NRFI2016-06), the Ministry of Education of Singapore (RG 1/18; RG 5/19; MOE2018-T3-1-003), an A*STAR Individual Research Grant (A1783c0008; A1783c0010), GSK-EDB Trust Fund, a Nanyang Research Award Grant, and Nanyang Technological University; the Guizhou Province First-Class Disciplines Project (Yiliu Xueke Jianshe Xiangmu)-GNYL(2017)008, Guizhou University of Traditional Chinese Medicine (China), QMY [2017]101 (China); the National Natural Science Foundation of China (Nos. 21772029, 21901233), the National Key Technologies R&D Program (No. 2014BAD23B01), the “Thousand Talent Plan”, the 10 Talent Plan (Shicengci) of Guizhou Province [2016]5649, and Guizhou University, Department of Science and Technology of Guizhou Province [2019]1020, Undergraduate Innovation and Pioneer Training Program, Guizhou University of Traditional Chinese Medicine 2018(31), Program of Introducing Talents of Discipline to Universities of China (111 Program, D20023).

REFERENCES

- (1) For selected reviews, see: (a) Bonjoch, J.; Solé, D. Synthesis of Strychnine. *Chem. Rev.* 2000, 100, 3455. (b) Knölker, H.-J.; Reddy, K. R. Isolation and Synthesis of Biologically Active Carbazole Alkaloids. *Chem. Rev.* 2002, 102, 4303. (c) Schmidt, A. W.; Reddy, K. R.; Knölker, H.-J. Occurrence, Biogenesis, and Synthesis of Biologically Active Carbazole Alkaloids. *Chem. Rev.* 2012, 112, 3193. (d) Ishikawa, H. G.; Elliott, I.; Velcicky, J.; Choi, Y.; Boger, D. L. Total Synthesis of (–)- and *ent*-(+)-Vindoline and Related Alkaloids. *J. Am. Chem. Soc.* 2006, 128, 10596. (e) Nikitenko, A.; Evrard, D.; Sabb, A. L.; Vogel, R. L.; Stack, G.; Young, M.; Lin, M.; Harrison, B. L.; Potoski, J. R. First Scale-Up: Problems and Resolutions on the Synthesis of WAY-253752, a Novel, Dual-Acting SSRI/5HT_{1A} Antagonist. *Org. Process Res. Dev.* 2008, 12, 76. (f) Kato, D.; Sasaki, Y.; Boger, D. L. Asymmetric Total Synthesis of Vindoline. *J. Am. Chem. Soc.* 2010, 132, 3685. (g) Tan, F.; Cheng, H.-G. Catalytic asymmetric synthesis of tetrahydrocarbazoles. *Chem. Commun.* 2019, 55, 6151.

- (2) Hollingshead, M. G.; Alley, M. C.; Camalier, R. F.; Abbott, B. J.; Mayo, J. G.; Malspeis, L.; Grever, M. R. In vivo cultivation of tumor cells in hollow fibers. *Life Sci.* 1995, 57, 131.
- (3) (a) Stoermer, D.; Heathcock, C. H. Total synthesis of alloaristoteline, (–)-serratoline, and (+)-aristoteline. *J. Org. Chem.* 1993, 58, 564. (b) Muñoz, O.; Christen, P.; Cretton, S.; Backhouse, N.; Torres, V.; Correa, O.; Costa, E.; Miranda, H.; Delporte, C. Chemical study and anti-inflammatory, analgesic and antioxidant activities of the leaves of *Aristolelia chilensis* (Mol.) Stuntz, Elaeocarpaceae. *J. Pharm. Pharmacol.* 2011, 63, 849.
- (4) Dolabela, M. F.; Pova, M. M.; Brandão, G. C.; Rocha, F. D.; Soares, L. F.; de Paula, R. C.; de Oliveira, A. B. *Aspidosperma* species as sources of anti-malarials: uleine is the major anti-malarial indole alkaloid from *Aspidosperma parvifolium* (Apocynaceae). *Malar. J.* 2015, 14, 498.
- (5) (a) Nicolaou, K. C.; Reingruber, R.; Sarlah, D.; Braš, S. Enantioselective Intramolecular Friedel–Crafts α -Arylation of Aldehydes. *J. Am. Chem. Soc.* 2009, 131, 2086. (b) Conrad, J. C.; Kong, J.; Laforteza, B. N.; MacMillan, D. W. C. Enantioselective α -Arylation of Aldehydes via Organo-SOMO Catalysis. An Ortho-Selective Arylation Reaction Based on an Open-Shell Pathway. *J. Am. Chem. Soc.* 2009, 131, 11640. (c) Gioia, C.; Hauville, A.; Bernardi, L.; Fini, F.; Ricci, A. Organocatalytic Asymmetric Diels–Alder Reactions of 3-Vinylindoles. *Angew. Chem., Int. Ed.* 2008, 47, 9236. (d) Tan, B.; Hernández-Torres, G.; Barbas, C. F., III Highly Efficient Hydrogen-Bonding Catalysis of the Diels–Alder Reaction of 3-Vinylindoles and Methyleneindolinones Provides Carbazolespirooxindole Skeletons. *J. Am. Chem. Soc.* 2011, 133, 12354. (e) Müller, S.; Webber, M. J.; List, B. The Catalytic Asymmetric Fischer Indolization. *J. Am. Chem. Soc.* 2011, 133, 18534. (f) Loh, C. C. J.; Raabe, G.; Enders, D. Enantioselective Synthesis of Tetrahydrocarbazoles through a Michael Addition/Ciamician–Plancher Rearrangement Sequence: Asymmetric Synthesis of a Potent Constrained Analogue of MS-245. *Chem. - Eur. J.* 2012, 18, 13250. (g) Tian, X.; Hofmann, N.; Melchiorre, P. Asymmetric Vinylogous Diels–Alder Reactions Catalyzed by a Chiral Phosphoric Acid. *Angew. Chem., Int. Ed.* 2014, 53, 2997. (h) Hansen, C. L.; Ohm, R. G.; Olsen, L. B.; Ascic, E.; Tanner, D.; Nielsen, T. E. Catalytic Enantioselective Synthesis of Tetrahydrocarbazoles and Exocyclic Pictet–Spengler-Type Reactions. *Org. Lett.* 2016, 18, 5990.
- (6) (a) Jen, W. S.; Wiener, J. J. M.; MacMillan, D. W. C. New Strategies for Organic Catalysis: The First Enantioselective Organocatalytic 1,3-Dipolar Cycloaddition. *J. Am. Chem. Soc.* 2000, 122, 9874. (b) Ahrendt, K. A.; Borths, C. J.; MacMillan, D. W. C. New Strategies for Organic Catalysis: The First Highly Enantioselective Organocatalytic Diels–Alder Reaction. *J. Am. Chem. Soc.* 2000, 122, 4243.
- (7) (a) List, B.; Lerner, R. A.; Barbas, C. F. Proline-Catalyzed Direct Asymmetric Aldol Reactions. *J. Am. Chem. Soc.* 2000, 122, 2395. (b) List, B. The Direct Catalytic Asymmetric Three-Component Mannich Reaction. *J. Am. Chem. Soc.* 2000, 122, 9336. (c) List, B. Direct Catalytic Asymmetric α -Amination of Aldehydes. *J. Am. Chem. Soc.* 2002, 124, 5656. (d) Yang, J. W.; Chandler, C.; Stadler, M.; Kampen, D.; List, B. *Nature* 2008, 452, 453.
- (8) (a) Jones, S. B.; Simmons, B.; MacMillan, D. W. C. Nine-Step Enantioselective Total Synthesis of (+)-Minfiensine. *J. Am. Chem. Soc.* 2009, 131, 13606. (b) Laforteza, B. N.; Pickworth, M.; MacMillan, D. W. C. Enantioselective Total Synthesis of (–)-Minovincine in Nine Chemical Steps: An Approach to Ketone Activation in Cascade Catalysis. *Angew. Chem., Int. Ed.* 2013, 52, 11269. (c) Horning, B. D.; MacMillan, D. W. C. Nine-Step Enantioselective Total Synthesis of (–)-Vincorine. *J. Am. Chem. Soc.* 2013, 135, 6442. (d) Jones, S. B.; Simmons, B.; Mastracchio, A.; MacMillan, D. W. C. Collective synthesis of natural products by means of organocascade catalysis. *Nature* 2011, 475, 183.
- (9) (a) Liu, Y.; Nappi, M.; Arceo, E.; Vera, S.; Melchiorre, P. Asymmetric Catalysis of Diels–Alder Reactions with in Situ Generated Heterocyclic *ortho*-Quinodimethanes. *J. Am. Chem. Soc.* 2011, 133, 15212. (b) Liu, Y.; Nappi, M.; Escudero-Adán, E. C.; Melchiorre, P. Multicatalytic Asymmetric Synthesis of Complex Tetrahydrocarbazoles via a Diels–Alder/Benzoin Reaction. *Org. Lett.* 2012, 14, 1310.
- (10) Xiao, Y.-C.; Zhou, Q.-Q.; Dong, L.; Liu, T.-Y.; Chen, Y.-C. Asymmetric Diels–Alder Reaction of 2-Methyl-3-indolylmethanols via in Situ Generation of *o*-Quinodimethanes. *Org. Lett.* 2012, 14, 5940.
- (11) Rassu, G.; Curti, C.; Zambrano, V.; Pinna, L.; Brindani, N.; Pelosi, G.; Zanardi, F. Exploiting the Distal Reactivity of Indolyl Methylenealononitriles: An Asymmetric Organocatalyzed [4 + 2] Cycloaddition with Enals Enables the Assembly of Elusive Dihydrocarbazoles. *Chem. - Eur. J.* 2016, 22, 12637.
- (12) (a) Zheng, C.; Lu, Y.; Zhang, J.; Chen, X.; Chai, Z.; Ma, W.; Zhao, G. The enantioselective, organocatalyzed Diels–Alder reaction of 2-vinylindoles with α,β -unsaturated aldehydes: an efficient route to functionalized tetrahydrocarbazoles. *Chem. - Eur. J.* 2010, 16, 5853. (b) Wang, X.-F.; Chen, J.-R.; Cao, Y.-J.; Cheng, H.-G.; Xiao, W.-J. An Enantioselective Approach to Highly Substituted Tetrahydrocarbazoles through Hydrogen Bonding-Catalyzed Cascade Reactions. *Org. Lett.* 2010, 12, 1140. (c) Zhu, X.-Y.; An, X.-L.; Li, C.-F.; Zhang, F.-G.; Hua, Q.-L.; Chen, J.-R.; Xiao, W.-J. A Practical and Enantioselective Approach to Tetrahydrocarbazoles by Asymmetric Organocatalysis. *ChemCatChem* 2011, 3, 679. (d) Cao, Y.-J.; Cheng, H.-G.; Lu, L.-Q.; Zhang, J.-J.; Cheng, Y.; Chen, J.-R.; Xiao, W.-J. Organocatalytic Multiple Cascade Reactions: A New Strategy for the Construction of Enantioenriched Tetrahydrocarbazoles. *Adv. Synth. Catal.* 2011, 353, 617. (e) Wu, X.; Zhu, H.-J.; Zhao, S.-B.; Chen, S.-S.; Luo, Y.-F.; Li, Y.-G. An Asymmetric Dehydrogenative Diels–Alder Reaction for the Synthesis of Chiral Tetrahydrocarbazole Derivatives. *Org. Lett.* 2018, 20, 32.
- (13) (a) Liu, C.; Han, X.; Wang, X.; Widenhofer, R. A. Platinum-Catalyzed Intramolecular Alkylation of Indoles with Unactivated Olefins. *J. Am. Chem. Soc.* 2004, 126, 3700. (b) Huang, H.-X.; Peters, R. A highly strained planar-chiral platinumacycle for catalytic activation of internal olefins in the Friedel–Crafts alkylation of indoles. *Angew. Chem., Int. Ed.* 2009, 48, 604. (c) Bandini, M.; Eichholzer, A. Enantioselective Gold-Catalyzed Allylic Alkylation of Indoles with Alcohols: An Efficient Route to Functionalized Tetrahydrocarbazoles. *Angew. Chem., Int. Ed.* 2009, 48, 9533. (d) Wu, Q.-F.; Zheng, C.; You, S.-L. Enantioselective Synthesis of Spiro Cyclopentane-1,3'-indoles and 2,3,4,9-Tetrahydro-1*H*-carbazoles by Iridium-Catalyzed Allylic Dearomatization and Stereospecific Migration. *Angew. Chem., Int. Ed.* 2012, 51, 1680. (e) Kawano, M.; Kiuchi, T.; Negishi, S.; Tanaka, H.; Hoshikawa, T.; Matsuo, J.-I.; Ishibashi, H. Regioselective Inter- and Intramolecular Formal [4 + 2] Cycloaddition of Cyclobutanones with Indoles and Total Synthesis of ($\pm$)-Aspidospermidine. *Angew. Chem., Int. Ed.* 2013, 52, 906. (f) Wang, Y.-D.; Zhang, P.-C.; Liu, Y.; Xia, F.; Zhang, J.-L. Enantioselective gold-catalyzed intermolecular [2 + 2] versus [4 + 2]-cycloadditions of 3-styrylindoles with *N*-allenamides: observation of interesting substituent effects. *Chem. Sci.* 2015, 6, 5564. (g) Liu, Q.-J.; Yan, W.-G.; Wang, L.; Zhang, X. P.; Tang, Y. One-Pot Catalytic Asymmetric Synthesis of Tetrahydrocarbazoles. *Org. Lett.* 2015, 17, 4014.
- (14) For selected recent reviews, see: (a) Ryan, S. J.; Candish, L.; Lupton, D. W. Acyl anion free *N*-heterocyclic carbene organocatalysis. *Chem. Soc. Rev.* 2013, 42, 4906. (b) Hopkinson, M. N.; Richter, C.; Schedler, M.; Glorius, F. An overview of *N*-heterocyclic carbenes. *Nature* 2014, 510, 485. (c) Mahatthananchai, J.; Bode, J. W. On the Mechanism of *N*-Heterocyclic Carbene-Catalyzed Reactions Involving Acyl Azoliums. *Acc. Chem. Res.* 2014, 47, 696. (d) Flanagan, D. M.; Romanov-Michailidis, F.; White, N. A.; Rovis, T. Organocatalytic Reactions Enabled by *N*-Heterocyclic Carbenes. *Chem. Rev.* 2015, 115, 9307. (e) Menon, R. S.; Biju, A. T.; Nair, V. Recent advances in employing homoenolates generated by *N*-heterocyclic carbene (NHC) catalysis in carbon–carbon bond-forming reactions. *Chem. Soc. Rev.* 2015, 44, 5040. (f) Chen, X.-Y.; Liu, Q.; Chauhan, P.; Enders, D. *N*-Heterocyclic Carbene Catalysis via Azolium Dienolates: An Efficient Strategy for Remote Enantioselective Functionalizations. *Angew. Chem., Int. Ed.* 2018, 57, 3862. (g) Murauski, K. J. R.; Jaworski, A. A.; Scheidt, K. A. A continuing challenge: *N*-heterocyclic

carbene-catalyzed syntheses of γ -butyrolactones. *Chem. Soc. Rev.* 2018, 47, 1773.

(15) Chen, X. K.; Yang, S.; Song, B.-A.; Chi, Y. R. Functionalization of Benzylic C(sp³)H Bonds of Heteroaryl Aldehydes through N-Heterocyclic Carbene Organocatalysis. *Angew. Chem., Int. Ed.* 2013, 52, 11134.

(16) For two reviews of indole-based oQDMs, see: (a) Magnus, P.; Gallagher, T.; Brown, P.; Pappalardo, P. The indole 2–3-quinodimethane strategy for the synthesis of indole alkaloids. *Acc. Chem. Res.* 1984, 17, 35. (b) Pindur, U.; Erfanian-Abdoust, H. Indolo-2,3-quinodimethanes and stable cyclic analogs for regio- and stereo-controlled syntheses of [b]-annelated indoles. *Chem. Rev.* 1989, 89, 1681. For recent examples, see: (c) Segura, J. L.; Martin, N. o-Quinodimethanes: Efficient Intermediates in Organic Synthesis. *Chem. Rev.* 1999, 99, 3199. (d) Kuroda, N.; Takahashi, Y.; Yoshinaga, K.; Mukai, C. A Novel Generation of Indole-2,3-quinodimethanes. *Org. Lett.* 2006, 8, 1843. (e) Liu, Y.; Nappi, M.; Arceo, E.; Vera, S.; Melchiorre, P. Asymmetric Catalysis of Diels–Alder Reactions with in Situ Generated Heterocyclic *ortho*-Quinodimethanes. *J. Am. Chem. Soc.* 2011, 133, 15212. (f) Zhou, L.; Xu, B.; Zhang, J. Metal-Free Dehydrogenative Diels–Alder Reactions of 2-Methyl-3-Alkylindoles with Dienophiles: Rapid Access to Tetrahydrocarbazoles, Carbazoles, and Heteroacenes. *Angew. Chem., Int. Ed.* 2015, 54, 9092. (g) Gu, B.-Q.; Zhang, H.; Su, R.-H.; Deng, W.-P. Organocatalytic asymmetric synthesis of dihydrocarbazoles via a formal [4 + 2] cycloaddition of in situ generated o-quinodimethanes with enals. *Tetrahedron* 2016, 72, 6595. (h) Yang, B.; Gao, S. Recent advances in the application of Diels–Alder reactions involving o-quinodimethanes, aza-o-quinone methides and o-quinone methides in natural product total synthesis. *Chem. Soc. Rev.* 2018, 47, 7926.

(17) For selected reviews about unsaturated acyl azolium, see:

(a) Zeidler, K. Stereoselective Synthesis of (*E*)- α,β -Unsaturated Esters via Carbene-Catalyzed Redox Esterification. *Org. Lett.* 2006, 8, 637. (b) Guin, J. S.; De Sarkar, D.; Grimme, S.; Studer, A. Biomimetic Carbene-Catalyzed Oxidations of Aldehydes Using TEMPO. *Angew. Chem., Int. Ed.* 2008, 47, 8727. (c) Kaebamrungs, J.; Mahatthananchai, J.; Zheng, P.; Bode, J. W. An Enantioselective Claisen Rearrangement Catalyzed by N-Heterocyclic Carbenes. *J. Am. Chem. Soc.* 2010, 132, 8810. (d) De Sarkar, S.; Studer, A. NHC-Catalyzed Michael Addition to α,β -Unsaturated Aldehydes by Redox Activation. *Angew. Chem., Int. Ed.* 2010, 49, 9266. (e) De Sarkar, S.; Grimme, S.; Studer, A. NHC Catalyzed Oxidations of Aldehydes to Esters: Chemoselective Acylation of Alcohols in Presence of Amines. *J. Am. Chem. Soc.* 2010, 132, 1190. (f) Ryan, S. J.; Candish, L.; Lupton, D. W. N-Heterocyclic Carbene-Catalyzed (4 + 2) Cycloaddition/Decarboxylation of Silyl Dienol Ethers with α,β -Unsaturated Acid Fluorides. *J. Am. Chem. Soc.* 2011, 133, 4694. (g) Kravina, A. G.; Mahatthananchai, J.; Bode, J. W. Enantioselective, NHC-Catalyzed Annulations of Trisubstituted Enals and Cyclic *N*-Sulfonylimines via α,β -Unsaturated Acyl Azoliums. *Angew. Chem., Int. Ed.* 2012, 51, 9433. (h) De Sarkar, S.; Biswas, A.; Samanta, R. C.; Studer, A. Catalysis with N-Heterocyclic Carbenes under Oxidative Conditions. *Chem. - Eur. J.* 2013, 19, 4664. (i) Mo, J.; Shen, L.; Chi, Y. R. Direct β -Activation of Saturated Aldehydes to Michael Acceptors through Oxidative NHC Catalysis. *Angew. Chem., Int. Ed.* 2013, 52, 8588.

(j) Chen, X.-Y.; Gao, Z.-H.; Song, C.-Y.; Zhang, C.-L.; Wang, Z.-X.; Ye, S. N-heterocyclic carbene catalyzed cyclocondensation of α,β -unsaturated carboxylic acids: enantioselective synthesis of pyrrolidinone and dihydropyridinone derivatives. *Angew. Chem., Int. Ed.* 2014, 53, 11611. (k) Li, G.-T.; Gu, Q.; You, S.-L. Enantioselective annulation of enals with 2-naphthols by triazolium salts derived from l-phenylalanine. *Chem. Sci.* 2015, 6, 4273. (l) Yetra, S. R.; Mondal, S.; Mukherjee, S.; Gonnade, R. G.; Biju, A. T. Enantioselective Synthesis of Spirocyclohexadienones by NHC-Catalyzed Formal [3 + 3] Annulation Reaction of Enals. *Angew. Chem., Int. Ed.* 2016, 55, 268. (m) Levens, A.; Ametovski, A.; Lupton, D. W. Enantioselective (4 + 2) Annulation of Donor-Acceptor Cyclobutanes by N-Heterocyclic Carbene Catalysis. *Angew. Chem., Int. Ed.* 2016, 55, 16136. (n) Wu, X.; Liu, B.; Zhang, Y.; Jeret, M.;

Wang, H.; Zheng, P.; Yang, S.; Song, B.-A.; Chi, Y. R. Enantioselective Nucleophilic β -Carbon-Atom Amination of Enals: Carbene-Catalyzed Formal [3 + 2] Reactions. *Angew. Chem., Int. Ed.* 2016, 55, 12280.

(o) Zhang, C.; Hooper, J. F.; Lupton, D. W. N-Heterocyclic Carbene Catalysis via the α,β -Unsaturated Acyl Azolium. *ACS Catal.* 2017, 7, 2583. (p) Chen, X.-Y.; Liu, Q.; Chauhan, P.; Li, S.; Peuronen, A.; Rissanen, K.; Jafari, E.; Enders, D. N-Heterocyclic Carbene Catalyzed [4 + 2] Annulation of Enals via a Double Vinylogous Michael Addition: Asymmetric Synthesis of 3,5-Diaryl Cyclohexenones. *Angew. Chem., Int. Ed.* 2017, 56, 6241. (q) Wu, X.; Hao, L.; Zhang, Y.; Rakesh, M.; Reddi, R.; Yang, S.; Song, B.-A.; Chi, Y. R. Construction of Fused Pyrrolidines and β -Lactones by Carbene-Catalyzed C–N, C–C, and C–O Bond Formations. *Angew. Chem., Int. Ed.* 2017, 56, 4201. (r) Chen, X.-Y.; Li, S.; Vetica, F.; Kumar, M.; Enders, D. N-Heterocyclic-Carbene-Catalyzed Domino Reactions via Two or More Activation Modes. *iScience* 2018, 2, 1. (s) Mukherjee, S.; Ghosh, A.; Marelli, U. K.; Biju, A. T. N-Heterocyclic Carbene-Catalyzed Michael–Michael–Lactonization Cascade for the Enantioselective Synthesis of Tricyclic δ -Lactones. *Org. Lett.* 2018, 20, 2952. (t) Mondal, S.; Yetra, S. R.; Mukherjee, S.; Biju, A. T. NHC-Catalyzed Generation of α,β -Unsaturated Acylazoliums for the Enantioselective Synthesis of Heterocycles and Carbocycles. *Acc. Chem. Res.* 2019, 52, 425. (u) Wu, X.; Zhou, L.; Maiti, R.; Mou, C.; Pan, L.; Chi, Y. R. Sulfinate and Carbene Co-catalyzed Rauhut–Currier Reaction for Enantioselective Access to Azepino[1,2-*a*]indole. *Angew. Chem., Int. Ed.* 2019, 58, 477.