

Wearable activity trackers for promoting physical activity : a systematic meta-analytic review

Li, Caining; Chen, Xiaoyu; Bi, Xinhua

2021

Li, C., Chen, X. & Bi, X. (2021). Wearable activity trackers for promoting physical activity : a systematic meta-analytic review. *International Journal of Medical Informatics*, 152, 104487-. <https://dx.doi.org/10.1016/j.ijmedinf.2021.104487>

<https://hdl.handle.net/10356/151480>

<https://doi.org/10.1016/j.ijmedinf.2021.104487>

© 2021 Elsevier B.V. All rights reserved. This paper was published in *International Journal of Medical Informatics* and is made available with permission of Elsevier B.V.

Downloaded on 25 Mar 2023 06:18:34 SGT

Wearable activity trackers for promoting physical activity:

A systematic meta-analytic review

Caining Li ^a, Xiaoyu Chen ^b, and Xinhua Bi ^a

a. School of Management, Jilin University, Changchun, China

b. Wee Kim Wee School of Communication and Information

Nanyang Technological University, Singapore

1st corresponding author: Xinhua Bi

Email address: bxhpaper@163.com

Postal address: School of Management, Jilin University, Changchun City, Jilin Province, 130025, China

2nd corresponding author: Xiaoyu Chen

Email address: xiaoyu001@e.ntu.edu.sg

Postal address: Nanyang Technological University, WKWSCI Building, 31 Nanyang Link, #05-16, Singapore 637718

Please cite as: Li, C., Chen, X., & Bi, X. (2021). Wearable activity trackers for promoting physical activity: A systematic meta-analytic review. *International Journal of Medical Informatics*, 152(8), 104487. <https://doi.org/10.1016/j.ijmedinf.2021.104487>

Highlights

- We conduct a systematic meta-analytic review to synthesize the existing evidence on the effectiveness of wearable activity trackers (WATs) for promoting physical activity (PA)
- Interventions with WATs significantly increased daily steps and weekly moderate-to-vigorous physical activity (MVPA) but had no impact on light physical activity (LPA) or sedentary behavior
- Two PA outcomes (daily steps and weekly MVPA) were associated with participants' characteristics (i.e., gender, age, health status, and baseline PA level) and intervention features (i.e., technology features, types of expert support, and intervention length)

Summary points

What already known on the topic?

- Wearable activity tracker (WAT) is a widely-accepted technology for nudging people toward a more active lifestyle.
- Inconsistent findings exist in the literature regarding the effectiveness of WATs for promoting physical activities.

What this review article adds?

- The interventions with WATs have small-to-medium effects on daily steps and weekly moderate-to-vigorous physical activities, whereas having no impact on light physical activities and sedentary behaviors.
- The effect size of daily steps and weekly MVPA, as outcomes of WATs-based interventions, is contingent on users' characteristics and interventions' features.

Wearable activity trackers for promoting physical activity:

A systematic meta-analytic review

Abstract

Purpose: Although wearable activity trackers hold a promise of nudging people toward a more active lifestyle, current research reveals inconsistent findings regarding their effectiveness. The objectives of this paper are two-fold: (1) to synthesize evidence on the effects of wearable activity trackers for improving physical activities, and (2) to identify potential moderators of effect size.

Methods: A systematic meta-analytic review was conducted. Forty-eight eligible papers based on forty-four distinct trials were identified through a systematic literature search process. Two authors independently extracted information from each study based on predefined data fields. Random-effects meta-analysis, subgroup analysis, and meta-regression analysis were employed.

Results: First, interventions with wearable activity trackers significantly increased daily steps and weekly moderate-to-vigorous physical activity but had no impact on light physical activity or sedentary behavior. Second, daily steps and weekly moderate-to-vigorous physical activity were associated with participants' characteristics (i.e., gender, age, medical condition, and baseline physical activity level) and intervention features (i.e., sensors, modes of expert support, and intervention duration). The identified factors explained 53% of the total variance for weekly moderate-to-vigorous physical activity.

Conclusions: The use of wearable activity trackers effectively improves conscious exercise behavior, including daily steps and weekly moderate-to-vigorous physical activity, but not effective for modifying habitual behavior, such as light physical activity and sedentary behavior. We also explicitly show that the extent to which the interventions with wearable activity trackers help users is contingent on the type of users and the design and delivery of interventions. Future studies are called to validate the findings and to offer theoretical explanations.

Keywords: Physical activity interventions; Wearable activity trackers; Effectiveness; Meta-analysis; Moderators

Paper type: Review article

Wearable activity trackers for promoting physical activity:

A systematic meta-analytic review

1. Introduction

1.1 Research background

In modern society, many people are habitually inactive during commuting, work, and leisure [1]. Insufficient physical activity (PA) is a leading risk factor for chronic diseases, mental health problems, and quality of life [2]. For substantial health benefits, PA guidelines recommend that adults do at least 150 minutes of moderate-to-vigorous physical activity (MVPA) per week, accumulate 10,000 steps per day, and reduce sedentary behavior [3, 4]. Nevertheless, according to a global survey across 168 countries [2], over one-quarter of people do not meet the PA level recommendations.

Wearable activity trackers (WATs) have the potential to nudge people toward a more active lifestyle. By integrating sensors, algorithms, and human-computer interaction designs, WATs could benefit users in two ways. First, WATs support self-monitoring of PA, such as steps taken, timing and intensity of PA, distance walked, and calories burned. Second, WATs help build self-regulatory skills, including goal-setting, action plans, behavior control, and goal achievement [5, 6].

1.2 Problem statement and research objectives

Despite the increasing popularity of WATs (e.g., Fitbit, Mi Band, Apple Watch, Garmin) in the consumer market, formal adoption in public health practice is scarce [7]. One exception was a national public health project carried out by Fitbit and Singapore's Health Promotion Board in 2019. Some empirical studies have been conducted to assess the effectiveness of WATs, albeit with mixed findings (e.g., [8, 9]). The extent to which WATs are able to promote PA remains unclear in the literature. Hence, a meta-analysis method has

been employed by several review studies. This approach can overcome the equivocation in the existing literature by calculating the weighted average value [10]. Although several relevant meta-analytic papers have come out in recent years [11-24], there are still two research gaps.

First, previous review articles did not examine all types of PA. Existing meta-analytic reviews found that WATs are effective for improving PA by assessing step counts [11, 12, 14, 16, 17, 19, 22-24], MVPA [12, 14, 17, 19, 22-24], or overall PA [13, 15, 16, 18, 20, 21]. However, there is a lack of in-depth analysis of light PA (LPA) and sedentary behavior, leading to an incomplete understanding of the WATs' effectiveness. To fill this gap, we propose the first research objective.

RO1: To synthesize evidence on the effects of WATs for promoting PA in terms of steps, LPA, MVPA, and sedentary behavior

Second, although large heterogeneity of WATs' effect size was observed, potential moderators were not systematically investigated. Specifically, by comparing between reviews with a focus on at-risk populations (e.g., chronic disease patients, older adults) [13, 15-19] and those without a specific population focus [21-24], we noticed that the former often reported larger PA effect sizes (reflected by *standard mean difference, SMD*) (Table 1).

Table 1. Comparison between reviews focused on at-risk populations and reviews without a specific population focus

| PA | Reviews focused on at-risk populations | Reviews without a specific population focus |
|--------------------|---|---|
| Step counts | [19] <i>SMD</i> = 1.23, 95% <i>CI</i> 0.75-1.70 [16] <i>SMD</i> = 0.55, 95% <i>CI</i> 0.40-0.70 [17] <i>SMD</i> = 0.54, 95% <i>CI</i> 0.34-0.73 | [23] <i>SMD</i> = 0.25, 95% <i>CI</i> 0.17-0.32 [22] <i>SMD</i> = 0.24, 95% <i>CI</i> 0.16-0.33 |
| MVPA | [17] <i>SMD</i> = 0.34, 95% <i>CI</i> 0.15-0.52 [19] <i>SMD</i> = 0.22, 95% <i>CI</i> -0.62-1.06 | [22] <i>SMD</i> = 0.27, 95% <i>CI</i> 0.15-0.39 [23] <i>SMD</i> = 0.01, 95% <i>CI</i> -0.15-0.13 |
| Overall PA | [18] <i>SMD</i> = 0.43, 95% <i>CI</i> 0.19-0.68 [13] <i>SMD</i> = 0.30, 95% <i>CI</i> 0.16-0.44 [15] <i>SMD</i> = 0.34, 95% <i>CI</i> 0.23-0.44 | [21] <i>SMD</i> = 0.26, 95% <i>CI</i> 0.04-0.49 |

Note: *SMD* = standard mean difference; *CI* = confidence interval

In addition to health status, past research suggested that participants' gender, age, and baseline PA level were associated with PA outcomes of using WATs [11, 25, 26]. Such findings indicated that different target populations could respond diversely to WATs. Notably, previous articles mainly focused on a particular population, lacking a comparison between multiple groups.

Even if focusing on a particular target population, previous meta-analytic reviews reported large heterogeneity of effect size across trials [11, 12, 14, 16, 19, 20]. However, the sources of the heterogeneity were not well explained. The heterogeneity of effect size could stem from intervention features, such as functionalities of WATs, intervention duration, and intervention delivery [23, 27]. Unfortunately, there is a lack of comprehensive evidence on how intervention features could moderate the effects of WATs.

Therefore, we propose the second research objective.

RO2: To explore potential moderators of WATs' effects through examining participants' characteristics and intervention features

This article is important in three ways. First, by considering the PA outcome at various intensity levels (i.e., steps, MVPA, LPA, and sedentary behavior), we analyze the influence of WATs use on PA thoroughly. This helps to understand the role of WATs from a fine-grained perspective. Second, by reporting eligible studies on a wide range of populations (see TableA.1 for details), we provide a holistic view of the effectiveness of WATs. This allows for comparing the role of WATs in promoting PA across different target user groups. Third, by analyzing the moderating role of participants' characteristics and intervention features, we fill a common gap in previous review articles: lacking a systematic investigation into the cross-studies heterogeneity.

2. Methods

We conducted and reported this review according to *the Cochrane Handbook for Systematic Reviews of Interventions* [28] and *the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines* [29]. The details of the applied methods were described as following.

2.1 Eligibility criteria

We assessed the eligibility of studies based on article type, participants, interventions, comparators, outcomes, and study design (Table 2).

Article type. Eligible studies were restricted to English-language publications. To ensure the credibility of the sources, only peer-reviewed journal papers were considered. Other publications, including book sections, conference abstracts, generics, and theses, were excluded.

Participants. One research objective of this review is to compare the effects of WATs across different populations, so we did not apply any population restrictions at first. However, we excluded participants with impaired movement ability from analysis as their activity patterns were non-comparable to other populations.

Interventions. WATs in this review refer to pedometer- or accelerometer-based devices, which are attached to the body to collect PA information [13]. Studies were included if they applied WATs alone or in combination with other intervention components to promote PA. Thus, technology design studies, validation tests, and studies that only used WATs as measurement tools were excluded because of irrelevance.

Comparators. The meta-analytic review must choose studies that involving a non-WATs (or blinded-WATs) comparator group. Studies provided any unblinded activity monitors to comparator group participants were excluded from analyses.

Outcomes. Studies were included if they reported objectively measured PA outcomes including step counts, MVPA, LPA, and sedentary behavior. Studies were excluded if they did not report any PA outcomes or only provided self-reported measures.

Study design. Only randomized controlled trials were included, while other study designs such as qualitative studies, cross-sectional surveys, pre-post or literature reviews were excluded.

Table 2. Inclusion and exclusion criteria

| Domain | Inclusion/exclusion criteria |
|----------------------|--|
| Article type | <ul style="list-style-type: none">• Inclusion- English-language, peer-reviewed journal papers;○ Exclusion- Non-English records; Article types other than peer-reviewed journal articles (e.g., book sections, conference abstracts, generics, theses); |
| Participants | <ul style="list-style-type: none">• Inclusion- A broad range of populations were eligible to be included;○ Exclusion- Participants with impaired motor ability, or for whom it was unsafe to be physically active without health professional supervision (e.g., patients diagnosed with a stroke, people immediately after surgery); |
| Interventions | <ul style="list-style-type: none">• Inclusion- Studies conducted a PA intervention by using WATs;○ Exclusion- Studies did not investigate WATs as an intervention tool for promoting PA (e.g., technology design studies, validation tests, WATs were solely used as a measurement tool in the studies); |
| Comparators | <ul style="list-style-type: none">• Inclusion- Studies involved a comparator group that did not provide unblinded WATs to participants;○ Exclusion- Studies lacked a non-WATs/blinded-WATs comparator group; |
| Outcomes | <ul style="list-style-type: none">• Inclusion- Studies reported objective measures of PA such as step counts, LPA, MVPA, and sedentary periods;○ Exclusion- Studies did not report PA outcomes or only provided self-reported PA; |
| Study design | <ul style="list-style-type: none">• Inclusion- Studies designed a randomized controlled trial (RCT) or cluster RCT with at least one intervention group and one control group;○ Exclusion- Studies did not apply RCTs (e.g., editorials, qualitative studies, literature reviews, conceptual papers, cross-sectional studies, pre-post study design). |

2.2. Information sources and search

Six major electronic databases were searched on March 25, 2020, including PubMed, PsycINFO, EMBASE, CINAHL, Science Direct, and Web of Science. Also, we updated our search results on March 24, 2021, to include the most recent research.

We used a combination of free-text terms and medical subject headings (MeSH) to develop the search query (Table 3). Two sets of keywords were used to identify papers involving WATs and PA outcomes of interest. We applied the search query in the field of title, abstract and keywords. The search query was adapted for each database as necessary. Furthermore, reference searching was performed to identify additional studies that the database search might have missed.

Table 3. The development of search query

| Purposes | Search query |
|---|---|
| Identifying studies involving WATs | #1: wearable tracker* OR wearable monitor* OR wearable device* OR wearable technolog* OR “activity tracker*” (MeSH) OR activity monitor* OR activity device* OR activity technolog* OR fitness tracker* (MeSH) OR fitness monitor* OR fitness device* OR fitness technolog* |
| Identifying studies involving PA | #2: physical activit* (MeSH) OR behavio*r change OR exercis* (MeSH) OR sedentary behavio*r (MeSH) OR sitting OR step* |
| For this review | #1 AND #2 |

2.3. Study selection

Retrieved records were screened in a three-step process. First, all records were imported into Endnote 20 (Clarivate, Pennsylvania), and duplicates were removed automatically. Second, the article type, title and abstract of each record were examined. Any records that did not meet the inclusion criteria were excluded. Third, all papers identified as requiring full-text review were examined. The primary author initially performed the three-step data selection process. To avoid bias, another author then performed the third step to verify the included/excluded studies. Any differences were resolved by consensus. The study selection process was similar to methods applied by Ringeval, Wagner [24].

2.4. Data collection and coding process

Data collection was informed by the Oxford Implementation Index, a common tool for incorporating implementation data into systematic reviews and meta-analysis [30]. We developed a data extraction sheet containing the following data items: citation; country/region; participants’ characteristics of recruitment settings, gender distribution, mean age, average baseline PA level, and medical condition; intervention descriptors of the model of WATs, sensor, functionalities, intervention duration, and combined intervention components; retention rate, sample size; and statistic findings (mean and SD of PA outcomes for the intervention and comparison groups).

Further, to address the second research objective (i.e., to explore potential moderators of effect size), four covariates related to participants’ characteristics and three covariates related to intervention features were coded dichotomy or categorical variables. Table 4 shows the coding rules.

Table 4. Coding rules of covariates

| Category | Covariates | Coding rules | Rationale |
|--------------------------------------|----------------------------------|--|---|
| Participants’ characteristics | Gender | “both male and female” = 0, “only male” = 1, “only female” = 2 | Biological properties |
| | Age | “mean age < 60 years” = 1, “mean age ≥ 60 years” = 2 | The definition of elderly [31] |
| | Baseline PA level | “below guidelines” = 1, “above guidelines” = 2 | PA guidelines [3, 4] |
| | Medical condition | “without medical conditions” = 1, “with a medical condition” = 2 | With or without clinical confirmed diseases (e.g., [32]) |
| Intervention features | PA sensor | “pedometer” = 1, “accelerometer” = 2 | Types of activity sensors [13] |
| | Combined intervention components | “null” = 0, “expert support delivered through one-way communication” = 1, “expert support delivered through two-way communication” = 2 | Communication types (i.e., with or without feedback) [33] |
| | Intervention duration | “≤ 12 weeks” = 1, “> 12 weeks” = 2 | Short/long-term WATs-based interventions [16] |

Data extraction and coding were performed independently by two authors. The inter-coder reliability was acceptable based on the agreement percentage (92%). The third author resolved all disagreements.

2.5. Risk of bias assessment

The *Cochrane Risk of Bias Tools* [34] was adapted to assess the risk of bias (ROB) for each study. The domain of *blinding participants* was excluded because the nature of the WATs-based intervention rendered blinding not feasible [13]. The domain of *blinding outcome assessment* was also excluded because of the use of objective PA measurements

[24]. Therefore, five ROB domains were assessed in this review, including *random sequence generation, allocation concealment, incomplete outcome data, selective reporting, and other sources of bias*. Each domain was scored as low, unclear, or high ROB following the criteria provided by Brickwood, Watson [22]. Two authors assessed the ROB of each selected study independently. The inter-coder reliability presented by the agreement percent was acceptable (94%). The third author resolved all disagreements.

2.6. Data synthesis, analysis, and interpretation

Empirical studies that investigated the effectiveness of WATs constitute the potential targets of our meta-analytic review. Meta-analysis can be conducted on two or more samples. However, to ensure robustness and practicability, the number of analyzable empirical studies must neither be too small nor too large. As a most recent review on this topic obtained a sample of 37 studies [24], we expect that the potential sample size of this review is suitable.

This review presented PA outcomes as steps/day, sedentary minutes/day, LPA minutes/day, and MVPA minutes/week to facilitate greater transferability for current PA guidelines [3, 4]. The mean and standard deviation (*SD*) of each PA outcome were derived for meta-analysis. Data presented as confidence interval level, standard error, p-values, and t-values were converted to *SD* based on statisticians' suggestions [28, 35]. We used R-3.6.2 to do random-effects meta-analyses.

For the first research objective, we calculated the mean difference (*MD*) and the standardized mean difference (*SMD*). The *SMD* was interpreted according to the Cohen rule of interpretation (small $0.2 \leq SMD < 0.5$, medium $0.5 \leq SMD < 0.8$, large $0.8 \leq SMD$) [36]. Heterogeneity across studies was assessed using I^2 , with per Higgins values of 25%, 50%, 75% were used as boundary limits for low, moderate, or high heterogeneity [37]. Funnel plots and Egger's test were used to examine potential publication bias. The trim-and-fill method

was used to adjust for potential missing studies. The fail-safe N was calculated to reflect the robustness of the results.

For the second research objective, subgroup meta-analysis and meta-regression analysis were conducted. First, interventions were sub-grouped according to each of the seven covariates (Table 4). Random-effects meta-analyses were conducted for each subgroup. T-tests were conducted to reflect the statistical significance of the intergroup difference. Second, the meta-regression analyses that including all covariates were calculated, and the model estimates were reported.

3. Results

3.1 Study selection

The initial search on the six databases resulted in 13,791 records. Of them, 6,298 were remained after removing duplicates. After a screening of article types, titles, and abstracts, 6,131 ineligible records were excluded. The remaining 167 articles were assessed based on the full-texts, resulting in 35 articles that met the eligibility criteria. Besides, 9 eligible articles were retrieved from reference searching. Also, 4 studies were identified through database updating (from March 25, 2020, to March 24, 2021). Finally, the formal dataset for coding and analysis comprised 48 articles (based on 43 distinct trials). Figure 1 outlines the process of dataset selection.




Figure 1. Flowchart of dataset selection

3.2. Characteristics of included studies

Most of the studies were published in the last 5 years ($n = 37$, 86%). A total of 5,808 participants were involved, with 3,218 in intervention groups and 2,679 in control groups. Descriptive statistics of participants, interventions, comparators, outcomes, and study design were stated below.

Participants. Table 5 presented a summary of participants' characteristics of gender, age, baseline PA level, medical condition, and recruitment settings. For detailed descriptions, one can refer to Table A.1 in the Appendix.

In summary, all trials were conducted in developed countries with an absence of literature in developing countries. Relatively small sample size was used in many trials ($n = 30$, 70%).

Many trials covered both male and female ($n = 33$, 77%), with a few focused solely on male ($n = 4$, 9%) or female ($n = 6$, 14%). About one-third of trials reached the elder (≥ 60 years old) ($n = 14$, 33%), leaving two-thirds involved younger populations ($n = 29$, 67%). More trials involved inactive participants ($n = 25$, 58%) than those involved with active participants (18, 42%).

About half trials recruited participants with a medical condition ($n = 21$, 49%), including diabetes [38, 39], cardiovascular disease [40, 41], peripheral artery disease [42], chronic obstructive pulmonary disease [43, 44], early knee osteoarthritis [45, 46], rheumatoid arthritis [47], systemic lupus erythematosus [47], and cancer [25, 32, 48-55]. The other half ($n = 22$, 51%) did not restrict participants according to medical conditions. Similarly, about half of the trials recruited participants mainly from clinics ($n = 22$, 51%). Other recruitment settings were communities ($n = 9$, 21%), workplaces ($n = 7$, 16%), social media (2, 5%), colleges courses (2, 5%), and newspapers (1, 2%).

Table 5. Summary of participants' characteristics

| Characteristics | No. trials (%) | Characteristics | No. trials (%) |
|------------------------|-----------------------|-----------------------------|-----------------------|
| Countries | | Baseline PA levels | |
| developed countries | 43 (100%) | below PA guidelines | 25 (58%) |
| developing countries | 0 (0%) | above PA guidelines | 18 (42%) |
| Sample size | | Medical condition | |
| < 100 | 30 (70%) | with medical conditions | 21 (49%) |
| ≥ 100 | 13 (30%) | without medical conditions | 22 (51%) |
| Gender | | Recruitment settings | |
| both genders | 33 (77%) | clinics | 21 (51%) |
| only female | 6 (14%) | communities | 9 (21%) |
| only male | 4 (9%) | workplaces | 7 (16%) |
| Age | | social Media | 2 (5%) |
| < 60 years old | 29 (67%) | college courses | 2 (5%) |
| ≥ 60 years old | 14 (33%) | newspaper | 1 (2%) |

Note: numbers in parentheses are percentages

Interventions. A detailed description of each studies' intervention features was listed in Table A.2 in the Appendix. Table 6 summarized the intervention features of all trials,

including the model of WATs, sensor, combined intervention components, and intervention duration.

The included trials utilized a wide range of WATs models. Fitbit devices were the most popular products ($n = 20$, 47%). The applied Fitbit models include Fitbit Alta [52], Fitbit Charge [40, 50, 51, 55], Fitbit Flex [45-47, 49, 53], Fitbit One [25, 32, 56-58], and Fitbit Zip [59-63]. Besides, Jawbone UP 24 [64-66] ($n = 3$, 7%), Polar (Polar M400 [54], Polar Active [67]) ($n = 2$, 5%), Garmin (Garmin Forerunner 210 [68], Garmin Vivofit [48]) ($n = 2$, 5%) and Fitbug [41, 44] ($n = 2$, 5%) were applied in more than one trials. Others ($n = 13$, 30%) adopted Activator [69], Active Link [70], Body Media FIT Core [71], LOMO Back[72], Misfit Flash [9], My Wellness Key [73], Sitfit [74], and study-specific WATs [38, 39, 43, 75, 76].

Two-fifths of trials applied pedometer-based WATs ($n = 17$, 40%), which could only monitor steps. The remaining three-fifths used accelerometer-based WATs ($n = 26$, 60%), capable of monitoring steps and timing and intensity of PA in detail.

About one-fifth of trials did not combine any intervention components other than WATs ($n = 8$, 19%). Others combined expert support ($n = 34$, 79%) and financial incentives ($n = 1$, 2%) into the PA interventions with WATs. Specifically, two delivery modes of expert support were identified: two-way communication ($n = 26$, 60%), including face-to-face interviews, phone calls, and group sessions; and one-way communication ($n = 8$, 19%) such as standard text messages, emails, and posts on social media.

Regarding intervention duration, the average intervention duration was 20-week. Short-term trials occupied two-thirds ($n = 29$, 67%), and one-third were long-term trials ($n = 14$, 33%).

Most trials had a high retention rate of over 80% ($n = 33, 77\%$), indicating low attendance bias.

Table 6. Summary of intervention features

| Intervention features | No. trials (%) | Intervention features | No. trials (%) |
|---------------------------------------|----------------|---|----------------|
| Model of WATs | | Combined intervention components | |
| Fitbit | 20 (47%) | expert support | 8 (19%) |
| Jawbone UP 24 | 3 (7%) | one-way communication | 26 (60%) |
| Polar | 3 (7%) | two-way communication | 1 (2%) |
| Garmin | 2 (5%) | monetary rewards | 8 (19%) |
| Fitbug | 2 (5%) | NA | 29 (67%) |
| Others | 13 (30%) | | 14 (33%) |
| Sensor pedometer accelerometer | | Intervention duration | |
| | 17 (40%) | ≤ 12 weeks | 10 (23%) |
| | 26 (60%) | > 12 weeks | 33 (77%) |
| | | Retention rate | |
| | | < 80% | |
| | | ≥ 80% | |

Comparators, outcomes, and study design. Included trials designed blinded WATs ($n = 4, 9\%$), wait-list control ($n = 8, 19\%$), standard practice or usual care ($n = 28, 65\%$), and blank control group ($n = 3, 7\%$) as comparators. All comparators were treated the same in the analysis, according to Hodkinson, Kontopantelis [13].

Trials reported at least one PA outcome of interest, such as step counts, MVPA, LPA, and sedentary behaviors. MVPA was the most frequently reported outcomes ($n = 31, 72\%$), followed by step counts ($n = 28, 65\%$), sedentary behavior ($n = 19, 44\%$), and the least reported was LPA ($n = 9, 21\%$).

All included trials conducted RCTs. Individual-based ($n = 38, 88\%$), paired ($n = 1, 2\%$), and cluster-based ($n = 4, 9\%$) randomization were applied. All randomization methods were treated the same in the analysis.

3.3. Risk of bias

According to the ROB assessment results (refers to Table A.3 in the Appendix), most trials had moderate-to-high-quality methodological design ($n = 39, 91\%$). In comparison, a few of them might at high risk of bias as they met only one or two criteria ($n = 4, 9\%$).

In detail, one trial was assessed as a high risk of selection bias because the group assignment was not based on randomization [59]. The management of incomplete data was assessed as high risk in nine trials as those omitted missing data regardless of the impact, for instance, Kenfield *et al.*, [32]. Two trials [71, 77] declared potential conflicts of interest. Thus high risks of other sources of bias were detected from them.

3.4. Overall effects of WATs

Table 7 summarized the results of meta-analyses on daily steps, weekly MVPA, LPA, and sedentary behavior. Overall, there were a significant moderate increase in daily steps and a small increase in weekly MVPA. However, the effect on LPA and sedentary behavior was small and nonsignificant. Detailed information was described in the following.

Table 7. Summary of overall effects meta-analyses

| Outcomes | k | Inverse-Variance Random Effects | | |
|-------------------|-----|--------------------------------------|--------|------------------------------|
| | | MD (95% CI) | I^2 | SMD (95% CI) |
| Steps/day | 32 | 1078.53 *** (772.18, 1384.87) | 78%*** | 0.50 *** (0.36, 0.63) |
| MVPA min/week | 34 | 42.99 *** (28.07, 57.92) | 84%*** | 0.33 *** (0.16, 0.51) |
| LPA min/day | 10 | -0.11 (-12.27, 12.06) | 88%*** | 0.09 (-0.28, 0.45) |
| Sedentary min/day | 21 | -0.24 (-15.83, 15.35) | 98%*** | 0.01 (-0.46, 0.48) |

Note: significant estimates are bolded; CI = confidence interval; significant codes include 0 '****' 0.001 '***' 0.01 '**' 0.05 '.' 0.1

Steps. A total of 32 records coded from 28 trials were analyzed to estimate the effects on daily steps (see Figure A.1). WATs-based interventions were associated with a significant increase in daily steps compared with the control groups ($MD = 1078.53, 95\% CI 772.18-$

1384.87). The average effect size reached medium level ($SMD = 0.50$, 95% CI 0.36-0.63). High heterogeneity ($I^2 = 78\%$) of the effect sizes across trials was observed.

Publication bias was detected by visual inspection of the funnel plot (Figure A.2) and Eggers' test ($p = 0.05$). As recommended [78], we applied the trim-and-fill method to examine the sensitivity of the results to potential publication bias. It turned out that the adjusted effect was still significant at $p < 0.001$, suggesting the result was less likely to be impacted by publication bias. Besides, the fail-safe N was 2063, further suggesting the robustness of the result.

MVPA. The meta-analysis for MVPA was based on 34 records coded from 31 trials (eFigure 3). There was a significant increase in weekly MVPA when comparing intervention groups versus controls ($MD = 42.99$, 95% CI 28.07-57.92). The SMD indicated that the effect size was small ($SMD = 0.33$, 95% CI 0.16-0.51). The heterogeneity was significantly large in terms of the magnitude of effect sizes across trials ($I^2 = 84\%$). Publication bias was not observed by visual inspection of the funnel plots (Figure A.4) and the Egger test ($p = 0.19$). The fail-safe N for the meta-analysis of MVPA was 712, indicating the robustness of the results.

LPA and sedentary behavior. Meta-analysis results indicated small and nonsignificant effect on LPA ($MD = -0.11$ min/day, 95% CI -12.27-12.06; $SMD = 0.09$, 95% CI -0.28-0.45) and sedentary behavior ($MD = -0.24$ min/day, 95% CI -15.83-15.35; $SMD = 0.01$; 95% CI -0.46-0.48). Thus, WATs-based intervention seemed to have no impact on the two PA outcomes, therefore they were not included in further analysis.

3.5. Analysis of Moderators

Subgroup analyses. Table 8 shown the subgroup analyses results according to participants' characteristics (i.e., gender, age, medical condition, baseline PA level) and

intervention features (sensor, modes of expert support, and intervention duration). All the subgroups revealed statistically significant effects on daily steps and weekly MVPA, indicating the effectiveness of WATs across different populations and diverse intervention designs.

Participants' characteristics as moderators. Gender, age, and medical condition were significantly associated with PA outcomes of daily steps and weekly MVPA. Baseline PA level was associated with daily steps but not associated with weekly MVPA.

Specifically, the subgroup analysis for gender suggested that the male population achieved significantly more increments than females in daily steps ($MD_{male} = 1324.52$, 95% CI 942.15-1706.90; $MD_{female} = 669.84$, 95% CI 80.79-1318.90; $t = 3.22$) and weekly MVPA ($MD_{male} = 108.46$, 95% CI 44.60-172.33; $MD_{female} = 69.44$, 95% CI 39.97-98.90; $t = 1.97$).

Subgroups for age indicated the pooled effects on daily steps ($MD_{older} = 1245.21$, 95% CI 602.77-1887.66; $MD_{younger} = 985.08$, 95% CI 657.74-1312.42; $t = 2.47$) and weekly MVPA ($MD_{older} = 57.63$, 95% CI 43.62-71.64; $MD_{younger} = 40.01$, 95% CI 18.96-61.05; $t = 6.41$) were significantly larger in older populations than in younger populations.

Subgroup analysis by medical condition suggested a significant difference between participants with versus without a medical condition in favor of the former for increment in daily steps ($MD_{with a medical condition} = 1342.30$, 95% CI 829.60-1854.99; $MD_{without a medical condition} = 818.32$, 95% CI 514.97-1121.67; $t = 6.90$) and weekly MVPA ($MD_{with a medical condition} = 56.48$, 95% CI 40.00-73.97; $MD_{without a medical condition} = 30.59$, 95% CI 8.76-52.41; $t = 7.42$)

Sub-grouping by baseline PA level suggested that inactive participants significantly increased more daily steps ($MD_{below guidelines} = 1158.22$, 95% CI 708.86-1607.58;

$MD_{above\ guidelines} = 968.24$, 95% CI 627.38-1309.09; $t = 2.58$) than those already active at baseline. However, there is no difference regarding weekly MVPA.

Intervention features as moderators. The type of sensors and intervention duration significantly impacted the effect size of WATs-based interventions on daily steps and weekly MVPA. The delivery modes of expert support were associated with daily steps but not weekly MVPA.

Subgroup analysis by sensors suggested that accelerometer-based WATs had a significantly larger effect than pedometer-based WATs for daily steps ($MD_{accelerometer} = 1153.83$, 95% CI 723.94-1583.72; $MD_{pedometer} = 1009.85$, 95% CI 554.06-1465.64; $t = 1.78$) and weekly MVPA ($MD_{accelerometer} = 51.94$, 95% CI 31.39-72.50; $MD_{pedometer} = 24.00$, 95% CI 14.77-33.23; $t = 10.72$)

According to tests of expert support types, one-way communication had more effects for daily steps ($MD_{one-way\ communication} = 1556.94$, 95% CI 746.91-2366.98; $MD_{two-way\ communication} = 1075.87$, 95% CI 645.51-1506.23; $t = 3.23$) than two-way communication. The between-group difference for weekly MVPA was small and nonsignificant.

As for subgroup analysis by intervention duration, short-term interventions reported more effects than long-term interventions for improving daily steps ($MD_{\leq 12-week} = 1402.21$, 95% CI 950.00-1766.17; $MD_{> 12-week} = 528.12$, 95% CI 258.15-785.24; $t = 13.4$) and weekly MVPA ($MD_{\leq 12-week} = 56.46$, 95% CI 34.83-68.31; $MD_{> 12-week} = 30.89$, 95% CI 4.99-56.79; $t = 4.89$).

Table 8. Subgroup analyses of potential moderators: participants' characteristics and intervention features.

| Sub-group | Daily steps | | | | MVPA minutes/week | | | |
|------------------------------|-------------|----------------------------|-----------------------|----------------------------------|-------------------|----------------------------|-----------------------|----------------------------------|
| | <i>k</i> | <i>MD</i> (95% <i>CI</i>) | <i>p</i> ^a | t-test (<i>p</i> ^b) | <i>k</i> | <i>MD</i> (95% <i>CI</i>) | <i>p</i> ^a | t-test (<i>p</i> ^b) |
| Gender | | | | 3.22 (0.02) | | | | 1.97 (0.08) |
| male | 3 | 1324.52 (942.15-1706.90) | <0.001 | | 3 | 108.46 (44.60-172.33) | 0.001 | |
| female | 4 | 699.84 (80.79-1318.90) | 0.03 | | 6 | 69.44 (39.97-98.90) | <0.001 | |
| Age | | | | 2.47 (0.02) | | | | 6.41 (<0.001) |
| <60-year | 21 | 985.08 (657.74-1312.42) | <0.001 | | 23 | 40.01 (18.96-61.05) | <0.001 | |
| ≥60-year | 11 | 1245.21 (602.77-1887.66) | <0.001 | | 11 | 57.63 (43.62-71.64) | <0.001 | |
| Medical condition | | | | 6.90 (<0.001) | | | | 7.42 (<0.001) |
| without | 16 | 818.32 (514.97-1121.67) | <0.001 | | 16 | 30.59 (8.76-52.41) | 0.006 | |
| with | 16 | 1342.30 (829.60-1854.99) | <0.001 | | 18 | 56.48 (40.00-73.97) | <0.001 | |
| Baseline PA level | | | | 2.58 (0.01) | | | | -0.65 (0.52) |
| below guidelines | 22 | 1158.22 (708.86-1607.58) | <0.001 | | 12 | 40.98 (19.77-62.19) | <0.001 | |
| above guidelines | 10 | 968.24 (627.38-1309.09) | <0.001 | | 22 | 43.47 (23.24-63.70) | <0.001 | |
| Sensors | | | | 1.78 (0.08) | | | | 10.72 (<0.001) |
| pedometers | 14 | 1009.85 (554.06-1465.64) | <0.001 | | 10 | 24.00 (14.77-33.23) | <0.001 | |
| accelerometers | 18 | 1153.83 (723.94-1583.72) | <0.001 | | 24 | 51.94 (31.39-72.50) | <0.001 | |
| Expert support | | | | 3.23 (0.004) | | | | -0.39(0.69) |
| one-way communication | 9 | 1556.94 (746.91-2366.98) | <0.001 | | 7 | 43.96 (14.55-73.37) | 0.003 | |
| two-way communication | 15 | 1075.87 (645.51-1506.23) | <0.001 | | 19 | 41.58 (21.99-61.16) | <0.001 | |
| Duration | | | | 13.4 (<0.001) | | | | 4.89 (<0.001) |
| ≤12-week | 23 | 1358.08 (950.00-1766.17) | <0.001 | | 22 | 51.58 (34.83-68.31) | <0.001 | |
| >12-week | 9 | 521.70 (258.15-785.24) | <0.001 | | 12 | 30.89 (4.99-56.79) | 0.02 | |

Note: statistically significant between-group difference are bolded; CI = confidence interval;

Meta-regression analyses. Mixed-effect meta-regression was applied to examine the explanatory power of the identified participants' characteristics and intervention features for WATs' effects on daily steps and weekly MVPA.

Meta-regression analysis for daily steps. Results of the meta-regression model for daily steps were presented in Table 9. The current model slightly reduced the amount of unexplained heterogeneity from $I^2 = 78\%$ to 71% (Table 6), but the residual heterogeneity was still significant ($QE = 65.14, p < 0.01$). The amount of variation accounted for was $R^2 = 6\%$, whereas, the model was not statistically significant ($F = 1.34, p = 0.27$).

Among the tested factors, expert support modes and intervention duration were statistically significant at $p < 0.1$. Specifically, interventions with expert support delivered through one-way communication exerted more improvements in daily steps ($\beta = 1209.74, p = 0.05$) than those delivered through two-way communication ($\beta = 293.79, p = 0.53$). The coefficient estimates of duration indicated that long-term interventions reported fewer increments of daily steps ($\beta = -907.97, p = 0.08$) than short-term interventions. These findings were consistent with the results of subgroup analyses (Table 8). Although significant differences were observed in subgroup analyses, gender, age, medical condition, PA level, and sensors turned nonsignificant in the model with all the seven factors.

Table 9. Results of the meta-regression model for daily steps

| Model | Estimates: β | |
|---|---------------------------------------|------------------|
| $F = 1.34, p = 0.27;$ $R^2 = 6\%;$ $I^2 = 71\%;$ $QE = 65.14, p < 0.01$ | intercept | 1250.14 * |
| | Gender: male | -219.91 |
| | Gender: female | -160.77 |
| | Age: ≥ 60 years | 216.84 |
| | Medical condition: with | -72.76 |
| | PA level: above | -463.03 |
| | Sensor: accelerometer | -368.59 |
| | Expert support: one-way communication | 1209.74 * |
| | Expert support: two-way communication | 293.79 |
| | Duration: >12-week | -907.97^ |

Note: significant estimates of covariates are bolded; significant codes include 0 '****' 0.001 '***' 0.01 '**' 0.05 '.' 0.1

Meta-regression analysis for MVPA. Table 10 presented the results of the meta-regression model for weekly MVPA. The multivariable model was significant ($F = 2.77, p = 0.02$) and explained a large part of the total variance ($R^2 = 53\%$). The identified covariates reduced unexplained heterogeneity across studies from $I^2 = 84\%$ (Table 6) to 57%. However, the residual heterogeneity was still significant ($QE = 31.12, p < 0.01$).

According to the coefficients, weekly MVPA was larger when interventions focused on males ($\beta = 70.41, p = 0.01$). The finding of gender was in line with subgroup analyses (Table 7). However, the impact of other covariates turned nonsignificant in the meta-regression analysis.

Table 10. Results of the meta-regression analysis for weekly MVPA

| Model | Estimates: β (p) | |
|--|---|----------------------------|
| $F = 2.77, p = 0.02;$ $R^2 = 53\%;$ $I^2 = 57\%;$ $QE = 31.12, p < 0.01$ | Intercept | 36.87[^] |
| | Gender: male | 70.41 ^{**} |
| | Gender: female | 36.06[^] |
| | Age: ≥ 60 years | 1.77 |
| | Medical condition: with | 23.74 |
| | PA level: above | 3.61 |
| | Sensor: accelerometer | 12.49 |
| | Expert support: one-way communication | -31.88 |
| | Expert support: two-way communication | -30.06 |
| | Duration | -15.28 |

Note: significant estimates of covariates are bolded; significant codes include 0 '****' 0.001 '***' 0.01 '**' 0.05 '^' 0.1

4. Discussion

4.1. Key findings

ROI: Overall, we found that interventions with WATs had a medium effect on daily steps and a small effect on MVPA. The positive findings were consistent with some previous reviews [13, 17, 20, 22]. By involving studies on a broad range of populations, this review provides evidence with greater generality in supporting the use of WATs for improving steps and MVPA.

However, this paper could not support the use of WATs for improving LPA or reducing sedentary behavior. Comparison of findings of LPA was not possible due to a lack of quantitative assessment in previous review papers. As for findings on sedentary behavior, similar results were reported by Ringeval, Wagner [24], who found that Fitbit-based interventions did not improve sedentary behaviors. We assume that the null effect for LPA and sedentary behavior might be because: existing WATs-based intervention mainly targeted at motivating individuals' conscious effort to exercise (i.e., to reach weekly MVPA goals or daily steps goals) rather than altering habitual behaviors (i.e., to change LPA or sedentary behavior). It is known that regulatory processes of conscious behaviors differed from habitual activities [79]. Regulatory strategies for improving conscious exercises, such as monitoring, goal settings, feedbacks, and rewards, are often applied in existing WATs designs (e.g., [60, 64]). However, habit reversal or formation strategies, such as automatic association and priming effects, are not common in existing WATs-based PA intervention designs.

RO2: Daily steps and weekly MVPA were associated with participants' characteristics and intervention features in similar patterns (see Table 7 and Figure 2), which was reasonable as the two measurements reflecting the same construct of PA levels. The identified moderators explained a large part of the variance of weekly MVPA (Table 10) but failed to explain the variance of daily steps across studies (Table 9).




Figure 2. Pooled MD for daily steps and weekly MVPA from subgroup meta-analysis

Participants' characteristics. We found that the effect size of WATs-based interventions for PA promotion varied across different populations (see Table 7 and Figure 2). Specifically, subgroup analyses indicated that the effects of WATs-based interventions tended to be larger when involved males, older, or participants with medical conditions, for both steps and MVPA. Also, inactive participants achieved more daily steps than active participants, but there was no difference for weekly MVPA.

Franssen et al. [11] investigated the impact of participants' gender and age on the effects of consumer WATs for improving PA in a meta-analytic review. Their findings of gender were consistent with our findings (i.e., male participants benefit more than females from WATs-based PA interventions). However, their study reported that younger users had a higher increase in PA, which conflicted with ours. The impacts of participants' characteristics of with/without medical condition and baseline PA level were not examined in previous review papers. Because of lacking enough evidence in the literature, findings of WATs'

effect among different populations from this review were not conclusive and need further verification in the future.

Intervention features. Interventions that applied accelerometer-based WATs reported a larger effect size than pedometer-based trackers. The finding indicated that monitoring PA information in detail instead of reporting simple step counts made WATs more useful for users. In contrast, Hodkinson, Kontopantelis [13] reported that pedometers demonstrated more increase in PA than accelerometers ($SMD_{pedometers} = 0.52$, 95% *CI* 0.32-0.72; $SMD_{accelerometers} = 0.30$, 95% *CI* 0.16-0.44). However, we noticed that Hodkinson, Kontopantelis [13] did not assess steps and MVPA separately but compared pedometers' impact on daily steps with accelerometers' impact on MVPA/total PA [13]. As we pooled a larger effect on daily steps than MVPA (Table 7) from both accelerometer- and pedometer-based WATs, we assume that the effect size of pedometers might be exaggerated compared to accelerometers in Hodkinson's study [13].

When comparing the two delivery modes of expert support, we found human-dependent two-way communication (e.g., face-to-face communication) was no better than computer-facilitated one-way communication (e.g., automated message push). A similar conclusion was drawn from an experimental study, which found that combined interviewing into a Fitbit Charge group failed to achieve more benefits [80]. Such findings promise to transform traditional human-dependent PA interventions into more accessible and cost-effective computer-delivered interventions during health practices.

The investigation of intervention duration suggested that shorter interventions reported more increment for both PA outcomes than interventions with longer periods. Along with existing studies [1, 12], this indicated that the effects of WATs lessened over time.

The explanation power of the identified covariates. The seven factors' meta-regression analysis was significant and explained a large part of the effect-size variation ($R^2 = 53\%$) for weekly MVPA. However, it was not effective for explaining the cross-study variance of daily steps. Predictors of daily steps resulting from WATs-based intervention remain to be explored, and the explanation power of them for weekly MVPA needs to be verified in future studies.

4.2. Implications for literature

Based on the key findings discussed above, four theoretical contributions are summarized and implications for relevant literature are proposed.

First, we provided a more comprehensive and fine-grained understanding of the effectiveness of WATs by including steps, MVPA, LPA, and sedentary behavior as PA outcomes. The effectiveness of WATs depends on the applied PA measurements (i.e., medium effect in steps, small effect in MVPA, null effect in LPA, and sedentary behavior). As related empirical studies are still accumulating, future work is suggested to look into different PA outcome measurements to present a broader picture.

Second, we found that WATs had no impact on LPA and sedentary behaviors. We speculate that existing WATs-based PA intervention strategies did not help in modifying habitual activities [79]. Future trials and conceptual studies are warranted to improve WATs technologies and interventions to support the regulation of habitual activities.

Third, we found that the degree of effectiveness was affected by participants' characteristics and intervention features. There is a trend that the increment of PA tends to be larger when interventions involved males, older, or participants with a medical condition. Simultaneously, accelerometer-based, short-term interventions reported larger PA outcomes. However, these findings were not conclusive and more evidence is needed in future research.

Besides, it is unclear why the increment in PA was different across populations and how WATs-based intervention features influenced PA outcomes. For a deeper understanding of the role of WATs in nudging users' behavior change, we suggest more theory-driven studies to investigate the psycho-social mediators linking the technological interventions to behavioral outcomes.

Fourth, the findings of this review might be biased to developed countries because none of the included studies were conducted in developing countries. The research and application of WATs in developing countries may be hindered by the relatively high cost and technical barriers. Future studies in developing countries are expected as societies move up the economic ladder as well as the advancement of technologies.

4.3. Implications for health practices

The findings have several implications for health practitioners, including WATs developers and health management practitioners.

First, we noticed that steps and MVPA, which are mainly featured by conscious activities, could be improved by WATs-based interventions. Also, the effectiveness of WATs for promoting steps and MVPA is reliable because the effect size reached statistical significance regardless of participants' characteristics and intervention features. Thus, we recommend that health management practitioners consider WATs a reliable intervention tool for improving steps and MVPA.

Second, no evidence was found to support the use of WATs for improving LPA or sedentary behaviors, which were mainly accumulated by an individual's routine/habitual PA. The effectiveness of WATs for increasing steps and MVPA lessened over time, indicating the importance of developing an internal exercise habit rather than depending on external interventions. More efforts, such as the adaptation of environmental factors, are needed to

alter habitual PA, in addition to WATs. For instance, the introduction of a flexible sit-stand workstation combined with WATs may have substantial benefits for reducing the sitting time of office-based employees [81].

Third, related practitioners should be aware that different types of populations might respond differently to WATs-based interventions. As the female, younger, and healthy populations benefit less than other populations, there needs more effort to focus on these populations. For instance, designers could help those populations by developing their self-determined motivations, as suggested by James et al. [5].

Fourth, accelerometer-based WATs, which provide in detail PA information [4], including step counts, and timing and intensity level of PA, are more effective for promoting PA than pedometers [82]. Therefore, we suggest future PA interventions apply accelerometer-based PA trackers rather than pedometers.

Fifth, the results showed that human-dependent two-way communication was no better than computer-facilitated one-way communication for delivering WATs-based PA interventions. As human resources are generally high cost and with limited access [70], designers must consider computer-facilitated communication as an alternative way of delivering WATs-based PA interventions.

5. Conclusion

This systematic meta-analytic review analyzed 48 papers (based on 43 distinct trials) involving 5,808 participants. We found promising evidence on the effects of WATs for promoting daily steps and weekly MVPA. The extent to which the WATs help was contextual—that is, the effect of WATs for promoting PA was contingent on the type of users and the design and delivery of interventions. As the emerging of more empirical studies, future work is needed to examine the validity of the findings from the current study. Theory-

driven studies are warranted to deeply understand the psycho-social mechanisms underlying the relationship between the use of WAT and users' PA. Also, we found that interventions with WATs had no impact on LPA and sedentary behavior. Thus, the development of strategies specifically for modifying habitual PA, such as sedentary behavior and LPA, is one of the directions for advancing design for advancing future WAT technology design.

Appendix

Appendix was provided as a supplement document.

Acknowledgments

The first author (C. Li) would like to thank the assistance in data collection from the Wee Kim Wee School of Communication and Information, Nanyang Technological University, Singapore, where she worked as a visiting PhD student from 2019 to 2020. This work is partly supported by the China Scholarship Council (CSC No.: 201906170120).

CRedit authorship contribution statement

Caining Li: Conceptualization, Methodology, Data Curation, Formal Analysis, Writing-original draft, Writing-review & editing.

Xiaoyu Chen: Conceptualization, Data Curation, Writing-original draft, Writing-review & editing.

Xinhua Bi: Supervision, Writing-review & editing.

References

- [1] Stephenson A, McDonough SM, Murphy MH, Nugent CD, Mair JL. Using computer, mobile and wearable technology enhanced interventions to reduce sedentary behaviour: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* 2017;14:105. doi: 10.1186/s12966-017-0561-4.
- [2] Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million participants. *The Lancet Global Health.* 2018;6(10):e1077-e1086. doi: 10.1016/s2214-109x(18)30357-7.
- [3] WHO. Global recommendations on physical activity for health. Geneva: World Health Organization; 2010.
- [4] Yao J, Tan CS, Chen C, Tan J, Lim N, Muller-Riemenschneider F. Bright spots, physical activity investments that work: National Steps Challenge, Singapore: a nationwide mHealth physical activity programme. *Br J Sports Med.* 2020;54(17):1047-1048. doi: 10.1136/bjsports-2019-101662.
- [5] James TL, Wallace L, Deane JK. Using organismic integration theory to explore the associations between users' exercise motivations and fitness technology feature set use. *Management Information Systems Quarterly.* 2019;43(1):287-312. doi: 10.25300/misq/2019/14128.
- [6] Shin G, Jarrahi MH, Fei Y, Karami A, Gafinowitz N, Byun A, et al. Wearable activity trackers, accuracy, adoption, acceptance and health impact: A systematic literature review. *J Biomed Inform.* 2019;93:103153. doi: 10.1016/j.jbi.2019.103153.
- [7] Li C, Lin SH, Chib A. The state of wearable health technologies: a transdisciplinary literature review. *Mobile Media & Communication.* 2020. doi: 10.1177/2050157920966023.
- [8] Pope ZC, Barr-Anderson DJ, Lewis BA, Pereira MA, Gao Z. Use of wearable technology and social media to improve physical activity and dietary behaviors among college students: a 12-week randomized pilot study. *Int J Environ Res Public Health.* 2019;16(19):3579. doi: 10.3390/ijerph16193579.
- [9] Kim Y, Lumpkin A, Lochbaum M, Stegemeier S, Kitten K. Promoting physical activity using a wearable activity tracker in college students: A cluster randomized controlled trial. *J Sports Sci.* 2018;36(16):1889-1896. doi: 10.1080/02640414.2018.1423886.
- [10] Borenstein M, Hedges LV, Higgins JP, Rothstein HR. *Introduction to Meta-analysis.* Chichester (UK): John Wiley & Sons; 2011.
- [11] Franssen WMA, Franssen GHLM, Spaas J, Solmi F, Eijnde BO. Can consumer wearable activity tracker-based interventions improve physical activity and cardiometabolic health in patients with chronic diseases? A systematic review and meta-analysis of randomised controlled trials. *International Journal of Behavioral Nutrition and Physical Activity.* 2020;17(1):57. doi: 10.1186/s12966-020-00955-2.
- [12] Kirk MA, Amiri M, Pirbaglou M, Ritvo P. Wearable technology and physical activity behavior change in adults with chronic cardiometabolic disease: a systematic review and meta-analysis. *American Journal of Health Promotion.* 2019;33(5):778-791. doi: 10.1177/0890117118816278.
- [13] Hodkinson A, Kontopantelis E, Adeniji C, van Marwijk H, McMillan B, Bower P, et al. Accelerometer- and pedometer-based physical activity interventions among adults with cardiometabolic conditions: a systematic review and meta-analysis. *JAMA Netw Open.* 2019;2(10):e1912895. doi: 10.1001/jamanetworkopen.2019.12895.
- [14] Davergne T, Pallot A, Dechartres A, Fautrel B, Gossec L. Use of wearable activity trackers to improve physical activity behavior in patients with rheumatic and musculoskeletal diseases: a systematic review and meta-analysis. *Arthritis Care Res (Hoboken).* 2019;71(6):758-767. doi: 10.1002/acr.23752.
- [15] Braakhuis HEM, Berger MAM, Bussmann JBJ. Effectiveness of healthcare interventions using objective feedback on physical activity: A systematic review and meta-analysis. *J Rehabil Med.* 2019;51(3):151-159. doi: 10.2340/16501977-2522.
- [16] J SO, Sherrington C, E RYZ, Franco MR, Tiedemann A, Puhakka S, et al. Effect of interventions using physical activity trackers on physical activity in people aged 60 years and over: a systematic review and meta-analysis. *Br J Sports Med.* 2020;54(20):1188-1194. doi: 10.1136/bjsports-2018-100324.
- [17] Larsen RT, Christensen J, Juhl CB, Andersen HB, Langberg H. Physical activity monitors to enhance amount of physical activity in older adults - a systematic review and meta-analysis. *Eur Rev Aging Phys Act.* 2019;16:7. doi: 10.1186/s11556-019-0213-6.
- [18] Cooper C, Gross A, Brinkman C, Pope R, Allen K, Hastings S, et al. The impact of wearable motion sensing technology on physical activity in older adults. *Exp Gerontol.* 2018;112(1):9-19. doi: 10.1016/j.exger.2018.08.002.
- [19] Liu JY-W, Kor PP-K, Chan CP-Y, Kwan RY-C, Sze-Ki D. The effectiveness of a wearable activity tracker (WAT)-based intervention to improve physical activity levels in sedentary older adults: A systematic review and meta-analysis. *Archives of Gerontology and Geriatrics.* 2020;91:104211. doi: 10.1016/j.archger.2020.104211.
- [20] Tang MSS, Moore K, McGavigan A, Clark RA, Ganesan AN. Effectiveness of wearable trackers on physical activity in healthy adults: systematic review and meta-analysis of randomized controlled trials. *JMIR Mhealth Uhealth.* 2020;8(7):e15576. doi: 10.1139/apnm-2019-0129.
- [21] Goode A, Hall K, Batch B, Huffman K, Hastings S, Allen K, et al. The impact of interventions that integrate

- accelerometers on physical activity and weight loss: a systematic review. *Annals of Behavioral Medicine*. 2017;51(1):79-93. doi: 10.1007/s12160-016-9829-1.
- [22] Brickwood KJ, Watson G, O'Brien J, Williams AD. Consumer-based wearable activity trackers increase physical activity participation: systematic review and meta-analysis. *JMIR Mhealth Uhealth*. 2019;7(4):e11819. doi: 10.2196/11819.
- [23] Lynch C, Bird S, Lythgo N, Selva-Raj I. Changing the physical activity behavior of adults with fitness trackers: a systematic review and meta-analysis. *American Journal of Health Promotion*. 2020;34(4):418-430. doi: 10.1177/0890117119895204.
- [24] Ringeval M, Wagner G, Denford J, Paré G, Kitsiou S. Fitbit-based interventions for healthy lifestyle outcomes: systematic review and meta-analysis. *J Med Internet Res*. 2020;22(10):e23954. doi: 10.1186/s12955-020-01478-x.
- [25] Weiner LS, Takemoto M, Godbole S, Nelson SH, Natarajan L, Sears DD, et al. Breast cancer survivors reduce accelerometer-measured sedentary time in an exercise intervention. *Journal of Cancer Survivorship*. 2019;13(3):468-476. doi: 10.1007/s11764-019-00768-8.
- [26] Patel M, O'Kane AA, Assoc Comp M. Contextual Influences on the Use and Non-Use of Digital Technology While Exercising at the Gym2015. 2923-2932 p.
- [27] Maxwell-Smith CM, Cohen PA, Platell C, Tan P, Tan J, Salama P, et al. Increasing physical activity in cancer survivors at cardiovascular risk using Fitbits: Randomized controlled trial. *Journal of Clinical Oncology*. 2018;36(7):93.
- [28] Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane handbook for systematic reviews of interventions*. Chichester (UK): John Wiley & Sons; 2019.
- [29] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *Bmj*. 2009;62(10):e1-e34. doi: 10.1136/bmj.b2700.
- [30] Montgomery P, Underhill K, Gardner F, Operario D, Mayo-Wilson E. The Oxford Implementation Index: a new tool for incorporating implementation data into systematic reviews and meta-analyses. *J Clin Epidemiol*. 2013;66(8):874-882. doi: 10.1016/j.jclinepi.2013.03.006.
- [31] WHO. Health care in the elderly: report of the Technical Group on Use of Medicaments by the Elderly. World Health Organization. *Drugs*. 1981;22(4):279-294.
- [32] Kenfield SA, Van Blarigan EL, Ameli N, Lavaki E, Cedars B, Paciorek AT, et al. Feasibility, acceptability, and behavioral outcomes from a technology-enhanced behavioral change intervention (prostate 8): a pilot randomized controlled trial in men with prostate cancer. *Eur Urol*. 2019;75(6):950-958. doi: 10.1016/j.eururo.2018.12.040.
- [33] Cooper R, DeJong DV, Forsythe R, Ross TW. Communication in Coordination Games. *The Quarterly Journal of Economics*. 1992;107(2):739-771. doi: 10.2307/2118488.
- [34] Higgins J, Altman D, Gotzsche P, Juni P, Moher D, Oxman A, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *Bmj*. 2011;343(7829):d5928-d5928. doi: 10.1136/bmj.d5928.
- [35] Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol*. 2014;14:135.
- [36] Cohen J. A power primer. *Psychological bulletin*. 1992;112(1):155.
- [37] Higgins J, Thompson S, Deeks J, Altman D. Measuring inconsistency in meta-analyses. *Bmj*. 2003;327(7414):557-560.
- [38] Van Dyck D, De Greef K, Deforche B, Ruige J, Tudor-Locke CE, Kaufman J-M, et al. Mediators of physical activity change in a behavioral modification program for type 2 diabetes patients. *Int J Behav Nutr Phys Act*. 2011;8. doi: 10.1186/1479-5868-8-105.
- [39] van der Weegen S, Verwey R, Spreeuwenberg M, Tange H, van der Weijden T, de Witte L. It's LiFe! Mobile and web-based monitoring and feedback tool embedded in primary care increases physical activity: a cluster randomized controlled trial. *J Med Internet Res*. 2015;17(7):e184. doi: 10.2196/jmir.4579.
- [40] Duscha BD, Piner LW, Patel MP, Craig KP, Brady M, IimcGarrah RW, et al. Effects of a 12-week mHealth program on peak VO₂ and physical activity patterns after completing cardiac rehabilitation: A randomized controlled trial. *Am Heart J*. 2018;199:105-114. doi: 10.1016/j.ahj.2018.02.001.
- [41] Martin SS, Feldman DI, Blumenthal RS, Jones SR, Post WS, McKibben RA, et al. mActive: A randomized clinical trial of an automated mHealth intervention for physical activity promotion. *J Am Heart Assoc*. 2015;4(11). doi: 10.1161/jaha.115.002239.
- [42] Duscha BD, Piner LW, Patel MP, Crawford LE, Jones WS, Patel MR, et al. Effects of a 12-week mhealth program on functional capacity and physical activity in patients with peripheral artery disease. *Am J Cardiol*. 2018;122(5):879-884. doi: 10.1016/j.amjcard.2018.05.018.
- [43] Varas AB, Córdoba S, Rodríguez-Andonaegui I, Rueda MR, García-Juez S, Vilaró J. Effectiveness of a community-based exercise training programme to increase physical activity level in patients with chronic obstructive pulmonary disease: a randomized controlled trial. 2018;23(4).

- [44] Armstrong M, Hume E, McNeillie L, Chambers F, Wakenshaw L, Burns G, et al. Behavioural modification interventions alongside pulmonary rehabilitation improve COPD patients' experiences of physical activity. *Respir Med.* 2021;180:106353. doi: 10.1016/j.rmed.2021.106353.
- [45] Li LC, Sayre EC, Xie H, Clayton C, Feehan LM. A community-based physical activity counselling program for people with knee osteoarthritis: feasibility and preliminary efficacy of the Track-OA study. *JMIR Mhealth Uhealth.* 2017;5(6):e86. doi: 10.2196/mhealth.7863.
- [46] Li LC, Feehan LM, Xie H, Lu N, Shaw CD, Gromala D, et al. Effects of a 12-week multifaceted wearable-based program for people with knee osteoarthritis: randomized controlled trial. *JMIR Mhealth Uhealth.* 2020;8(7). doi: 10.2196/19116.
- [47] Li LC, Feehan LM, Xie H, Lu N, Shaw C, Gromala D, et al. Efficacy of a physical activity counselling program with use of wearable tracker in people with inflammatory arthritis: a randomized controlled trial. *Arthritis Care Res (Hoboken).* 2020. doi: 10.1002/acr.24199.
- [48] Lynch BM, Nguyen NH, Moore MM, Reeves MM, Rosenberg DE, Boyle T, et al. A randomized controlled trial of a wearable technology-based intervention for increasing moderate to vigorous physical activity and reducing sedentary behavior in breast cancer survivors: The ACTIVATE Trial. *Cancer.* 2019;125(16):2846-2855. doi: 10.1002/cncr.32143.
- [49] Van Blarigan EL, Chan H, Van Loon K, Kenfield SA, Chan JM, Mitchell E, et al. Self-monitoring and reminder text messages to increase physical activity in colorectal cancer survivors (Smart Pace): a pilot randomized controlled trial. 2019;19(1).
- [50] Cadmus-Bertram L, Tevaarwerk AJ, Sesto ME, Gangnon R, Van Remortel B, Date P. Building a physical activity intervention into clinical care for breast and colorectal cancer survivors in Wisconsin: a randomized controlled pilot trial. *J Cancer Surviv.* 2019;13(4):593-602. doi: 10.1007/s11764-019-00778-6.
- [51] Devine KA, Viola A, Levonyan-Radloff K, Mackowski N, Bozzini B, Chandler A, et al. Feasibility of FitSurvivor: A technology-enhanced group-based fitness intervention for adolescent and young adult survivors of childhood cancer. *Pediatr Blood Cancer.* 2020;67(9). doi: 10.1002/pbc.28530.
- [52] Maxwell-Smith C, Hince D, Cohen PA, Bulsara MK, Boyle T, Platell C, et al. A randomized controlled trial of WATAAP to promote physical activity in colorectal and endometrial cancer survivors. *Psychooncology.* 2019;28(7):1420-1429. doi: 10.1002/pon.5090.
- [53] Mendoza JA, Baker KS, Moreno MA, Whitlock K, Abbey-Lambertz M, Waite A, et al. A Fitbit and Facebook mHealth intervention for promoting physical activity among adolescent and young adult childhood cancer survivors: A pilot study. *Pediatric Blood and Cancer.* 2017;64(12). doi: 10.1002/pbc.26660.
- [54] Pope ZC, Zeng N, Zhang R, Lee HY, Gao Z. Effectiveness of combined smartwatch and social media intervention on breast cancer survivor health outcomes: a 10-week pilot randomized trial. *J Clin Med.* 2018;7(6). doi: 10.3390/jcm7060140.
- [55] Singh B, Spence RR, Sandler CX, Tanner J, Hayes SC. Feasibility and effect of a physical activity counselling session with or without provision of an activity tracker on maintenance of physical activity in women with breast cancer - A randomised controlled trial. *J Sci Med Sport.* 2020;23(3):283-290. doi: 10.1016/j.jsams.2019.09.019.
- [56] Hartman SJ, Nelson SH, Cadmus-Bertram LA, Patterson RE, Parker BA, Pierce JP. Technology- and Phone-Based Weight Loss Intervention. *Am J Prev Med.* 2016;51(5):714-721. doi: 10.1016/j.amepre.2016.06.024.
- [57] Cadmus-Bertram L, Marcus BH, Patterson RE, Parker BA, Morey BL. Randomized trial of a Fitbit-based physical activity intervention for women. *Am J Prev Med.* 2015;49(3):414-418.
- [58] Thorndike AN, Mills S, Sonnenberg L, Palakshappa D, Gao T, Pau CT, et al. Activity monitor intervention to promote physical activity of physicians-in-training: randomized controlled trial. *PLoS One.* 2014;9(6):e100251. doi: 10.1371/journal.pone.0100251.
- [59] Rote AE. Physical activity intervention using Fitbits in an introductory college health course. *Health Education Journal.* 2017;76(3):337-348. doi: 10.1177/0017896916674505.
- [60] Finkelstein EA, Haaland BA, Bilger M, Sahasranaman A, Sloan RA, Nang EEK, et al. Effectiveness of activity trackers with and without incentives to increase physical activity (TRIPPA): a randomised controlled trial. *Lancet Diabetes Endocrinol.* 2016;4(12):983-995. doi: 10.1016/s2213-8587(16)30284-4.
- [61] Muellmann S, Buck C, Voelcker-Rehage C, Bragina I, Lippke S, Meyer J, et al. Effects of two web-based interventions promoting physical activity among older adults compared to a delayed intervention control group in Northwestern Germany: Results of the PROMOTE community-based intervention trial. *Prev Med Rep.* 2019;15:100958. doi: 10.1016/j.pmedr.2019.100958.
- [62] Sloan RA, Kim Y, Sahasranaman A, Müller-Riemenschneider F, Biddle SJH, Finkelstein EA. The influence of a consumer-wearable activity tracker on sedentary time and prolonged sedentary bouts: secondary analysis of a randomized controlled trial. *BMC Res Notes.* 2018;11:189. doi: 10.1186/s13104-018-3306-9.
- [63] Nanda S, Hurt RT, Croghan IT, Mundi MS, Gifford SL, Schroeder DR, et al. Improving physical activity and body composition in a medical workplace using brief goal setting. *Mayo Clin Proc Innov Qual Outcomes.* 2019;3(4):495-505. doi: 10.1016/j.mayocpiqo.2019.07.002.
- [64] Ashton LM, Morgan PJ, Hutchesson MJ, Rollo ME, Collins CE. Feasibility and preliminary efficacy of the

- 'HEYMAN' healthy lifestyle program for young men: a pilot randomised controlled trial. *Nutr J*. 2017;16(1):2. doi: 10.1186/s12937-017-0227-8.
- [65] Brickwood K-J, Ahuja KDK, Watson G, O'Brien JA, Williams AD. Effects of activity tracker use with health professional support or telephone counseling on maintenance of physical activity and health outcomes in older adults: randomized controlled trial. *JMIR Mhealth Uhealth*. 2021;9(1). doi: 10.2196/18686.
- [66] Lyons EJ, Swartz MC, Lewis ZH, Martinez E, Jennings K. Feasibility and acceptability of a wearable technology physical activity intervention with telephone counseling for mid-aged and older adults: a randomized controlled pilot trial. *JMIR Mhealth Uhealth*. 2017;5(3):e28. doi: 10.2196/mhealth.6967.
- [67] Leinonen AM, Pyky R, Ahola R, Kangas M, Siirtola P, Luoto T, et al. Feasibility of gamified mobile service aimed at physical activation in young men: population-based randomized controlled study (MOPO). *JMIR Mhealth Uhealth*. 2017;5(10):e146. doi: 10.2196/mhealth.6675.
- [68] Obling KH, Overgaard K, Juul L, Maindal HT. Effects of a motivational, individual and locally anchored exercise intervention (MILE) on cardiorespiratory fitness: a community-based randomised controlled trial. *BMC Public Health*. 2019;19(1):239. doi: 10.1186/s12889-019-6556-0.
- [69] Renaud LR, Jelsma JGM, Huysmans MA, van Nassau F, Lakerveld J, Spekle EM, et al. Effectiveness of the multi-component dynamic work intervention to reduce sitting time in office workers- Results from a pragmatic cluster randomised controlled trial. 2020;84.
- [70] Thomas JG, Raynor HA, Bond DS, Luke AK, Cardoso CC, Foster GD, et al. Weight loss in weight watchers online with and without an activity tracking device compared to control: a randomized trial. *Obesity (Silver Spring)*. 2017;25(6):1014-1021. doi: 10.1002/oby.21846.
- [71] Jakicic JM, Davis KK, Rogers RJ, King WC, Marcus MD, Helsel D, et al. Effect of wearable technology combined with a lifestyle intervention on long-term weight loss: the IDEA randomized clinical trial. *Journal of American Medical Association*. 2016;316(11):1161-1171. doi: 10.1001/jama.2016.12858.
- [72] Brakenridge CL, Fjeldsoe BS, Young DC, Winkler EAH, Dunstan DW, Straker LM, et al. Evaluating the effectiveness of organisational-level strategies with or without an activity tracker to reduce office workers' sitting time: a cluster-randomised trial. *International Journal of Behavioral Nutrition and Physical Activity*. 2016;13. doi: 10.1186/s12966-016-0441-3.
- [73] Maselli M, Gobbi E, Carraro A. Effectiveness of individual counseling and activity monitors to promote physical activity among university students. *J Sports Med Phys Fitness*. 2019;59(1):132-140. doi: 10.23736/s0022-4707.17.07981-6.
- [74] Wyke S, Bunn C, Gray CM, Macaulay L, Loudon DW, Maxwell DJ, et al. The effect of a programme to improve men's sedentary time and physical activity: The European Fans in Training (EuroFIT) randomised controlled trial. *PLoS Med*. 2019;16(2):1-25. doi: 10.1371/journal.pmed.1002736.
- [75] Yang Y-P, Wang C-J, Wang J-J, Lin C-W, Yang Y-TC, Wang J-s, et al. The effects of an activity promotion system on active living in overweight subjects with metabolic abnormalities. *Obes Res Clin Pract*. 2017;11(6):718-727. doi: 10.1016/j.orcp.2017.06.002.
- [76] Nakata Y, Sasai H, Tsujimoto T, Hashimoto K, Kobayashi H. Web-based intervention to promote weight-loss maintenance using an activity monitor: A randomized controlled trial. *Prev Med Rep*. 2019;14:100839. doi: 10.1016/j.pmedr.2019.100839.
- [77] Poirier J, Bennett WL, Jerome GJ, Shah NG, Lazo M, Yeh HC, et al. Effectiveness of an activity tracker- and internet-based adaptive walking program for adults: a randomized controlled trial. *J Med Internet Res*. 2016;18(2):e34. doi: 10.2196/jmir.5295.
- [78] Duval S, Tweedie R. A nonparametric "trim and fill" method of accounting for publication bias in meta-analysis. *Journal of the American Statistical Association*. 2000;95(449):89-98. doi: 10.1080/01621459.2000.10473905.
- [79] Rebar AL, Dimmock JA, Jackson B, Rhodes RE, Kates A, Starling J, et al. A systematic review of the effects of non-conscious regulatory processes in physical activity. *Health psychology review*. 2016;10(4):395-407. doi: 10.1080/17437199.2016.1183505.
- [80] Ellingson LD, Lansing JE, DeShaw KJ, Peyer KL, Bai Y, Perez M, et al. Evaluating Motivational Interviewing and Habit Formation to Enhance the Effect of Activity Trackers on Healthy Adults' Activity Levels: Randomized Intervention. *JMIR Mhealth Uhealth*. 2019;7(2):e10988. doi: 10.2196/10988.
- [81] Coffeng JK, van der Ploeg HP, Castellano JM, Fernandez-Alvira JM, Ibanez B, Garcia-Lunar I, et al. A 30-month worksite-based lifestyle program to promote cardiovascular health in middle-aged bank employees: Design of the TANSNIP-PESA randomized controlled trial. *Am Heart J*. 2017;184. doi: 10.1016/j.ahj.2016.11.002.
- [82] Esakia A, McCrickard S, Harden S, Horning M, Ramalingam N. Using smartwatches to facilitate a group dynamics-based statewide physical activity intervention. *International Journal of Human-Computer Studies*. 2020;142. doi: 10.1016/j.ijhcs.2020.102501.