Observation of dislocation-induced topological modes in a three-dimensional acoustic topological insulator

Xue, Haoran; Jia, Ding; Ge, Yong; Guan, Yi-Jun; Wang, Qiang; Yuan, Shou-Qi; Sun, Hong-Xiang; Chong, Yidong; Zhang, Baile

2021

Xue, H., Jia, D., Ge, Y., Guan, Y., Wang, Q., Yuan, S., Sun, H., Chong, Y. & Zhang, B. (2021). Observation of dislocation-induced topological modes in a three-dimensional acoustic topological insulator. *Physical Review Letters*, 127(21), 214301-.
<https://dx.doi.org/10.1103/PhysRevLett.127.214301>

<https://hdl.handle.net/10356/154803>

<https://doi.org/10.1103/PhysRevLett.127.214301>

© 2021 American Physical Society. All rights reserved. This paper was published in *Physical Review Letters* and is made available with permission of American Physical Society.

Downloaded on 26 Apr 2023 22:52:46 SGT

Observation of Dislocation-Induced Topological Modes in a Three-Dimensional Acoustic Topological Insulator

Haoran Xue^{1,*}, Ding Jia,^{2,*} Yong Ge,² Yi-jun Guan,² Qiang Wang¹, Shou-qi Yuan,²

Hong-xiang Sun^{2,†}, Y. D. Chong^{1,3,‡} and Baile Zhang^{1,3,§}

¹*Division of Physics and Applied Physics, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore*

²*Research Center of Fluid Machinery Engineering and Technology, School of Physics and Electronic Engineering, Jiangsu University, Zhenjiang 212013, China*

³*Centre for Disruptive Photonic Technologies, Nanyang Technological University, Singapore 637371, Singapore*

(Received 6 May 2021; accepted 20 October 2021; published 16 November 2021)

The interplay between real-space topological lattice defects and the reciprocal-space topology of energy bands can give rise to novel phenomena, such as one-dimensional topological modes bound to screw dislocations in three-dimensional topological insulators. We obtain direct experimental observations of dislocation-induced helical modes in an acoustic analog of a weak three-dimensional topological insulator. The spatial distribution of the helical modes is found through spin-resolved field mapping, and verified numerically by tight-binding and finite-element calculations. These one-dimensional helical channels can serve as robust waveguides in three-dimensional media. Our experiment paves the way to studying novel physical modes and functionalities enabled by topological lattice defects in three-dimensional classical topological materials.

DOI: [10.1103/PhysRevLett.127.214301](https://doi.org/10.1103/PhysRevLett.127.214301)

There are two distinct types of topological phenomena that commonly occur in crystalline materials. The first is band topology defined in reciprocal space, which gives rise to topologically protected boundary modes in topological insulators and related materials [1,2]. The other involves topological lattice defects defined in real space, which are protected against local perturbations [3,4]. These two types of topological phenomena are usually studied separately, but when they are simultaneously present in a material, their interplay can give rise to many unique phenomena [5–19]. For example, it has been theoretically predicted that a screw dislocation in a three-dimensional (3D) topological insulator (TI) can host one-dimensional (1D) topologically protected helical modes [5]. The existence of helical modes is determined by the nonzero value of $\mathbf{B} \cdot \mathbf{G}_\nu / 2\pi$, where $\mathbf{B}$ is the Burgers vector of the dislocation (defined in real space), and $\mathbf{G}_\nu$ is a reciprocal vector defined as $\mathbf{G}_\nu = \sum_{i=1}^3 \nu_i \mathbf{b}_i$, where $\mathbf{b}_i$ are the primitive reciprocal lattice vectors and ν_i are the three weak topological indices [20]. This “bulk-dislocation correspondence” is a modification of the more well-known bulk-boundary correspondence principle, taking the topology of the lattice dislocation into account. Similar phenomena have been theoretically studied in the context of strong and weak 3D electronic TIs [5–7,11], 3D photonic Chern crystals [21], and weak 3D photonic TIs with a synthetic dimension [22]. Dislocation-induced topological modes have technological possibilities as a way to embed robust 1D waveguides in 3D TIs. Compared to other topological waveguide designs

such as hinge modes in 3D higher-order TIs [23–25], dislocation-induced topological modes can have larger bandwidth (they span the entire bulk gap, unlike hinge modes which usually exist inside a surface gap), and can easily support multimode transport (by increasing the Burgers vector or the spin Chern number).

Thus far, however, there have been few experimental observations of dislocation-induced topological modes. Some signatures of these modes have been found in real materials through conductivity measurements [26] and local density of states measurements [27]. Classical wave systems such as photonic crystals and acoustic crystals have been used to demonstrate localized modes bound to dislocations in 2D photonic Chern insulators [28] and weak 2D magnetomechanical TIs [29], disclination waveguiding in 2D valley photonic crystals [17], and fractional mode density at lattice defects in 2D topological crystalline insulators [30,31]. However, the topological helical modes induced by a screw dislocation in a 3D TI [5] have not previously been realized.

In this Letter, we present direct experimental observations of dislocation-induced topological helical modes in a 3D acoustic TI. The sample consists of stacked 2D acoustic TIs implementing a weak 3D acoustic TI with $\mathbf{G}_\nu = (0, 0, 1)2\pi/a_z$. A screw dislocation with $\mathbf{B} = (0, 0, a_z)$ is fabricated into the center of the sample. With this setup, the bulk-dislocation correspondence predicts a pair of gapless helical modes inside the topological band gap [5]. The existence of the helical modes is

FIG. 1. Construction of a 3D weak TI with a screw dislocation. (a) A single 2D TI with helical edge modes. (b) 2D TIs stacked along z to form a weak 3D TI. In the limit of zero interlayer couplings, the surface modes of the weak 3D TI are simply the helical edge modes of the decoupled 2D layers. (c) A screw dislocation is introduced by a cut-and-glue procedure; the helical edge modes on the bottom and top layers connect to modes running along the dislocation.

determined experimentally with field-mapping measurements, resolved over the effective spin and spanning the entire sample, both on the surface and in the bulk. The experimental results are corroborated by tight-binding calculations and full-wave simulations.

The design process is schematically illustrated in Fig. 1. We start with a 2D TI that hosts a pair of topologically protected helical modes on its edges [32,33], as shown in Fig. 1(a). A weak 3D TI can be constructed by stacking the 2D TIs in the z direction. The bulk band gap is maximized by letting the couplings between adjacent layers vanish. In that case, the gapless surface states on the side surfaces of the weak 3D TI are intrinsically 2D: they propagate only in the xy plane of each decoupled layer [Fig. 1(b)]. A screw dislocation along z is then introduced by a “cut and glue” procedure, as shown in Fig. 1(c). In tight-binding models as well as the real experimental sample, this can be accomplished by modifying the couplings between certain sites near the center of each layer. The introduction of the dislocation gives rise to a pair of dislocation-induced helical modes, which are connected to the helical edge modes on the bottom and top layers, as shown in Fig. 1(c).

To design the structure, we begin with a tight-binding model for a 2D TI described by a Hamiltonian of the form

$$H = \begin{bmatrix} H_{\uparrow}(\phi) & 0 \\ 0 & H_{\downarrow}(\phi) \end{bmatrix}, \quad (1)$$

where the submatrices $H_{\uparrow,\downarrow}$ describe 2D Chern insulators consisting of square lattices with complex couplings, as depicted in Fig. 2(a). Each is a copy of the Hofstadter model [34] with magnetic flux $\phi = \pm\pi/2$ (thus $H_{\downarrow} = H_{\uparrow}^*$) [35,36]. The design can be further simplified through the transformation $[\psi_A, \psi_B]^T = u^{-1}[\psi_{\uparrow}, \psi_{\downarrow}]^T$, where [37,38]

$$u = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & i \\ 1 & -i \end{bmatrix}. \quad (2)$$

This replaces the original complex couplings by real couplings of either positive or negative sign, as shown

in Fig. 2(b). Such a lattice can be realized in acoustics using acoustic resonators coupled by properly designed channels [39–41]. Figure 2(c) shows the spectrum of the resulting 2D TI, with open boundary condition along x and periodic boundary condition along y . There are two nontrivial band gaps traversed by helical edge modes, whose mode profiles are shown in Fig. 2(e). Finally, following the procedure depicted in Fig. 1, we stack the 2D TI to form a weak 3D TI, and introduce a central screw dislocation. This involves replacing the intralayer couplings along the line segment $y = 0, x > 0$ in each layer [where $(x, y) = (0, 0)$ denotes the center of the layer] with interlayer couplings implementing the “spiral staircase” pattern shown in Fig. 1(c).

Figure 2(d) shows the calculated spectrum for a lattice of size 16×16 , with periodic boundary conditions imposed along x and y to remove the surface modes on the sides. With this simulation geometry, there is an extra dislocation at $(8, 0)$ in addition to the dislocation at $(0, 0)$. In the plot, dislocation-induced helical modes can clearly be seen within the bulk band gap. Each dispersion curve of dislocation modes is two-fold degenerate, corresponding to the two modes at the two dislocations; red (blue) denote spin $\uparrow$ ($\downarrow$) for the modes evaluated at $(0, 0)$. The mode profiles are shown in Fig. 2(f).

We now introduce an acoustic lattice that realizes the above tight-binding model. It consists of two types of building blocks: acoustic resonators and coupling tubes; similar designs have previously been used to create acoustic realizations of other tight-binding models [39–47]. A photograph of a one-layer sample (without a dislocation) is shown in Fig. 3(a), and a schematic of part of it is shown in Fig. 3(b). Each cuboid resonator supports a dipolar mode at around 2144 Hz, and is connected to its neighbors through a series of coupling tubes. The coupling tubes shown in red (blue) in Fig. 3(b) correspond to positive (negative) couplings. Details about the acoustic structure are given in the Supplemental Material [48].

Prior to dealing with the full 3D lattice, we verify the features of the 2D lattice using simulations and experiments. We consider a finite acoustic lattice of size 8×8

FIG. 2. Lattice model design. (a) Square lattices with magnetic flux $\phi = \pi/2$ (left) and $\phi = -\pi/2$ (right) per plaquette. (b) A lattice model for a 2D TI obtained by combining and transforming the two lattice models in (a). Here, all couplings are real and have the same amplitude, and red (blue) colors denote positive (negative) couplings. (c) Edge spectrum for the lattice shown in (b), calculated using periodic boundary conditions in y and open boundary conditions in x . Black curves correspond to bulk bands and red and blue curves denote spin-up and spin-down modes on the left edge. (d) Dislocation spectrum for a weak 3D TI built up by stacking the 2D TI in (b) with vanishing interlayer couplings. The calculation uses lattice size 16×16 and periodic boundary conditions in both the x and y directions. The couplings at $y = 0$ and $x > 0$ are all replaced by interlayer couplings. This setup leads to two dislocations at $(0,0)$ and $(8,0)$. (e) Profile of the edge mode marked by the red circle in (c). (f) Profile of the dislocation mode marked by the red circle in (d).

[Fig. 3(a)]. The simulated eigenfrequencies are plotted in Fig. 3(c), with the two bulk band gaps shaded in gray. The modes inside the band gaps are indeed edge modes, as can be seen from the eigenmode profile plotted in Fig. 3(d). Throughout this work, we will focus on the second band gap at ~ 2200 Hz, which is larger.

To verify the existence of the edge modes experimentally, we put a speaker at the left edge [yellow star in Figs. 3(e) and 3(f)] and measure the acoustic field (both amplitude and phase) in each resonator. Details of the experimental procedure are given in the Supplemental Material [48]. The spin-resolved fields are obtained as $[\psi_{\uparrow}, \psi_{\downarrow}]^T = u[\psi_A, \psi_B]^T$.

FIG. 3. Design and demonstration of a 2D acoustic TI. (a) Photo of the fabricated sample for the 2D acoustic TI. (b) Unit cell for the acoustic lattice, with the red and blue coupling tubes enabling positive and negative couplings. (c) Simulated eigenfrequencies for the acoustic lattice shown in (a). The shaded gray regions denote the bulk band gap. (d) Eigenmode profile for one of the in-gap edge modes. (e) and (f) Measured field distributions at 2165 Hz for spin-up mode (e) and spin-down mode (f).

Figures 3(e) and 3(f) show the measured intensity distributions at 2165 Hz for spin-up and spin-down, respectively. Evidently, the two fields counterpropagate along the edge, as expected of spin-polarized helical edge modes.

Building on this 2D acoustic TI design, we construct a weak 3D acoustic TI with a screw dislocation according to the procedure in Fig. 1. The fabricated sample is shown in Fig. 4(a). It consists of four layers of the 2D acoustic TI with a central screw dislocation, with Burgers vector $\mathbf{B} = (0, 0, a_z)$. A close-up view of the dislocation is shown in Fig. 4(d). Figure 4(b) shows the simulated dispersion for the 3D acoustic structure, which agrees well with the dispersion obtained using the tight-binding model [Fig. 2(d)]. We can clearly see the emergence of the dislocation-induced helical modes, with a typical mode profile shown in Fig. 4(c).

To probe the dislocation-induced helical modes experimentally, we inject sound into one of the resonators at the center of the top layer; the source position is indicated by a

FIG. 4. Observation of dislocation-induced topological helical modes. (a) Photo of the fabricated sample for the weak 3D acoustic TI with dislocation. (b) Simulated band structure for the lattice shown in (a). Here, gray dots correspond to bulk modes and red and blue dots represent spin-up and spin-down dislocation modes. (c) Eigenmode profile for one of the dislocation modes. (d) An enlarged photo of the sample center area. The Burgers vector $\mathbf{B}$ is denoted by the red arrow. (e) Measured acoustic pressure spectra when the excitation is at the center of the first layer and the detection is at the second layer. Red and black curves are results obtained at one resonator next to the center and away from the center, respectively. (f) Measured acoustic intensity distributions at 2165 Hz in the three layers counted from the top layer. Red and blue colors represent spin-up and spin-down states, respectively. The subplots within the same column (i.e., the same layer) are normalized by the same value, while the subplots in different columns are normalized by different values.

yellow star in Fig. 4(f). We then scan the acoustic fields inside the whole sample [48]. Figure 4(e) plots the acoustic pressure spectra measured at two different resonators on the second layer (here the second layer refers to the layer right below the top layer). The red curve comes from measurements taken at a resonator next to the dislocation, which has strong peaks at frequencies corresponding to the predicted bulk band gaps, where the dislocation modes are expected to occur. The black curve is derived from measurements taken at a resonator that is $2\sqrt{2}a$ away from the dislocation (a is the spacing between adjacent resonators), and has less pronounced peaks at those frequencies. This indicates that the in-gap modes are localized around the dislocation.

Figure 4(f) shows site- and spin-resolved acoustic intensity distributions measured at 2165 Hz. As can be seen, the distributions for the two spins are very different.

For spin-up (upper row), the measured intensity is always localized around the center. For spin-down (lower row), the intensities in the second and the third layers are negligible, indicating that only spin-up modes can propagate downwards along the dislocation. In contrast, when the source is placed at the center of the bottom layer, the situation is reversed (i.e., only spin-down modes can propagate upwards along the dislocation), as shown in the Supplemental Material [48]. Thus, the direction of propagation along the dislocation is locked to the spin. The observed spatial profile and helical characteristics of the dislocation-induced topological modes agree well with the simulation results shown in Fig. 4(b).

To sum up, we have observed a pair of dislocation-induced helical modes in an acoustic analog of a weak 3D TI, thereby validating the long-predicted phenomenon of

dislocation-induced 1D topological modes in 3D TIs [5,6]. Our results, together with other recent related experimental efforts [18,49–51], coincide with growing interest in the possibilities of combining band topology with lattice defects [17,28–31]. One promising direction for future research is to extend the current design beyond the tight-binding regime. It has previously been shown that a smooth helical modulation can generate chiral modes in a magnetic Weyl crystal [21]; similarly, it may be possible to generate helical modes in a time-reversal invariant phononic or photonic crystal with four Weyl points, which would be helpful for implementing waveguides with high bandwidth, low loss, and multimode features. Moreover, although we have focused on screw dislocations, there are other types of lattice defects that may host topological modes with different characteristics, which could be similarly studied using acoustic structures. Finally, it would be interesting to study non-Hermitian band topological effects [52–62], which can have interesting interactions with topological lattice defects [63–66], by using passive [57,58] or active [59] methods to introduce non-Hermiticity into acoustic structures.

H. X., Q. W., Y. C., and B. Z. acknowledge support from the Singapore Ministry of Education Academic Research Fund under Tier 3 Grant No. MOE2016-T3-1-006 and Tier 2 Grant No. MOE2019-T2-2-085. D. J., Y. G., Y. G., S. Y., and H. S. acknowledge support from the National Natural Science Foundation of China under Grants No. 11774137, No. 12174159, and No. 51779107, the China Postdoctoral Science Foundation under Grant No. 2020M671351, and the State Key Laboratory of Acoustics, Chinese Academy of Science under Grant No. SKLA202115.

*These authors contributed equally to this work.

[†]jsdxshx@ujs.edu.cn

[‡]yidong@ntu.edu.sg

[§]blzhang@ntu.edu.sg

- [1] M. Z. Hasan and C. L. Kane, Colloquium: Topological insulators, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [2] X.-L. Qi and S.-C. Zhang, Topological insulators and superconductors, *Rev. Mod. Phys.* **83**, 1057 (2011).
- [3] N. D. Mermin, The topological theory of defects in ordered media, *Rev. Mod. Phys.* **51**, 591 (1979).
- [4] J. M. Kosterlitz, Nobel lecture: Topological defects and phase transitions, *Rev. Mod. Phys.* **89**, 040501 (2017).
- [5] Y. Ran, Y. Zhang, and A. Vishwanath, One-dimensional topologically protected modes in topological insulators with lattice dislocations, *Nat. Phys.* **5**, 298 (2009).
- [6] J. C. Y. Teo and C. L. Kane, Topological defects and gapless modes in insulators and superconductors, *Phys. Rev. B* **82**, 115120 (2010).
- [7] K.-I. Imura, Y. Takane, and A. Tanaka, Weak topological insulator with protected gapless helical states, *Phys. Rev. B* **84**, 035443 (2011).
- [8] V. Juričić, A. Mesaros, R.-J. Slager, and J. Zaanen, Universal Probes of Two-Dimensional Topological Insulators: Dislocation and π Flux, *Phys. Rev. Lett.* **108**, 106403 (2012).
- [9] A. Rüegg and C. Lin, Bound States of Conical Singularities in Graphene-Based Topological Insulators, *Phys. Rev. Lett.* **110**, 046401 (2013).
- [10] K. Shiozaki and M. Sato, Topology of crystalline insulators and superconductors, *Phys. Rev. B* **90**, 165114 (2014).
- [11] R.-J. Slager, A. Mesaros, V. Juričić, and J. Zaanen, Interplay between electronic topology and crystal symmetry: Dislocation-line modes in topological band insulators, *Phys. Rev. B* **90**, 241403(R) (2014).
- [12] H. Sumiyoshi and S. Fujimoto, Torsional Chiral Magnetic Effect in a Weyl Semimetal with a Topological Defect, *Phys. Rev. Lett.* **116**, 166601 (2016).
- [13] E. N. Lima, T. M. Schmidt, and R. W. Nunes, Topologically protected metallic states induced by a one-dimensional extended defect in the bulk of a 2D topological insulator, *Nano Lett.* **16**, 4025 (2016).
- [14] J. Liu and L. Balents, Anomalous Hall Effect and Topological Defects in Antiferromagnetic Weyl Semimetals: $\text{Mn}_3\text{Sn}/\text{Ge}$, *Phys. Rev. Lett.* **119**, 087202 (2017).
- [15] R. Queiroz, I. C. Fulga, N. Avraham, H. Beidenkopf, and J. Cano, Partial Lattice Defects in Higher-Order Topological Insulators, *Phys. Rev. Lett.* **123**, 266802 (2019).
- [16] R.-J. Slager, The translational side of topological band insulators, *J. Phys. Chem. Solids* **128**, 24 (2019).
- [17] Q. Wang, H. Xue, B. Zhang, and Y. D. Chong, Observation of Protected Photonic Edge States Induced by Real-Space Topological Lattice Defects, *Phys. Rev. Lett.* **124**, 243602 (2020).
- [18] Q. Wang, Y. Ge, H.-x. Sun, H. Xue, D. Jia, Y.-j. Guan, S.-q. Yuan, B. Zhang, and Y. Chong, Vortex states in an acoustic Weyl crystal with a topological lattice defect, *Nat. Commun.* **12**, 3654 (2021).
- [19] T. Li, P. Zhu, W. A. Benalcazar, and T. L. Hughes, Fractional disclination charge in two-dimensional C_n -symmetric topological crystalline insulators, *Phys. Rev. B* **101**, 115115 (2020).
- [20] L. Fu, C. L. Kane, and E. J. Mele, Topological Insulators in Three Dimensions, *Phys. Rev. Lett.* **98**, 106803 (2007).
- [21] L. Lu, H. Gao, and Z. Wang, Topological one-way fiber of second Chern number, *Nat. Commun.* **9**, 5384 (2018).
- [22] Q. Lin, X.-Q. Sun, M. Xiao, S.-C. Zhang, and S. Fan, A three-dimensional photonic topological insulator using a two-dimensional ring resonator lattice with a synthetic frequency dimension, *Sci. Adv.* **4**, 2774 (2018).
- [23] F. Schindler, A. M. Cook, M. G. Vergniory, Z. Wang, S. S. Parkin, B. A. Bernevig, and T. Neupert, Higher-order topological insulators, *Sci. Adv.* **4**, eaat0346 (2018).
- [24] J. Langbehn, Y. Peng, L. Trifunovic, F. von Oppen, and P. W. Brouwer, Reflection-Symmetric Second-Order Topological Insulators and Superconductors, *Phys. Rev. Lett.* **119**, 246401 (2017).
- [25] Z. Song, Z. Fang, and C. Fang, (d-2)-Dimensional Edge States of Rotation Symmetry Protected Topological States, *Phys. Rev. Lett.* **119**, 246402 (2017).
- [26] H. Hamasaki, Y. Tokumoto, and K. Edagawa, Dislocation conduction in Bi-Sb topological insulators, *Appl. Phys. Lett.* **110**, 092105 (2017).

- [27] A. K. Nayak, J. Reiner, R. Queiroz, H. Fu, C. Shekhar, B. Yan, C. Felser, N. Avraham, and H. Beidenkopf, Resolving the topological classification of bismuth with topological defects, *Sci. Adv.* **5**, 6996 (2019).
- [28] F.-F. Li, H.-X. Wang, Z. Xiong, Q. Lou, P. Chen, R.-X. Wu, Y. Poo, J.-H. Jiang, and S. John, Topological light-trapping on a dislocation, *Nat. Commun.* **9**, 2462 (2018).
- [29] I. H. Grinberg, M. Lin, W. A. Benalcazar, T. L. Hughes, and G. Bahl, Trapped State at a Dislocation in a Weak Magneto-mechanical Topological Insulator, *Phys. Rev. Applied* **14**, 064042 (2020).
- [30] C. W. Peterson, T. Li, W. Jiang, T. L. Hughes, and G. Bahl, Trapped fractional charges at bulk defects in topological insulators, *Nature (London)* **589**, 376 (2021).
- [31] Y. Liu, S. Leung, F.-F. Li, Z.-K. Lin, X. Tao, Y. Poo, and J.-H. Jiang, Bulk–disclination correspondence in topological crystalline insulators, *Nature (London)* **589**, 381 (2021).
- [32] C. L. Kane and E. J. Mele, Quantum Spin Hall Effect in Graphene, *Phys. Rev. Lett.* **95**, 226801 (2005).
- [33] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, Quantum spin Hall effect and topological phase transition in HgTe quantum wells, *Science* **314**, 1757 (2006).
- [34] D. R. Hofstadter, Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields, *Phys. Rev. B* **14**, 2239 (1976).
- [35] M. Hafezi, E. A. Demler, M. D. Lukin, and J. M. Taylor, Robust optical delay lines with topological protection, *Nat. Phys.* **7**, 907 (2011).
- [36] M. Hafezi, S. Mittal, J. Fan, A. Migdall, and J. Taylor, Imaging topological edge states in silicon photonics, *Nat. Photonics* **7**, 1001 (2013).
- [37] R. Süssstrunk and S. D. Huber, Observation of phononic helical edge states in a mechanical topological insulator, *Science* **349**, 47 (2015).
- [38] J. Ningyuan, C. Owens, A. Sommer, D. Schuster, and J. Simon, Time-and Site-Resolved Dynamics in a Topological Circuit, *Phys. Rev. X* **5**, 021031 (2015).
- [39] H. Xue, Y. Ge, H.-X. Sun, Q. Wang, D. Jia, Y.-J. Guan, S.-Q. Yuan, Y. Chong, and B. Zhang, Observation of an acoustic octupole topological insulator, *Nat. Commun.* **11**, 2442 (2020).
- [40] X. Ni, M. Li, M. Weiner, A. Alù, and A. B. Khanikaev, Demonstration of a quantized acoustic octupole topological insulator, *Nat. Commun.* **11**, 2108 (2020).
- [41] Y. Qi, C. Qiu, M. Xiao, H. He, M. Ke, and Z. Liu, Acoustic Realization of Quadrupole Topological Insulators, *Phys. Rev. Lett.* **124**, 206601 (2020).
- [42] M. Xiao, W.-J. Chen, W.-Y. He, and C. T. Chan, Synthetic gauge flux and Weyl points in acoustic systems, *Nat. Phys.* **11**, 920 (2015).
- [43] Y.-X. Xiao, G. Ma, Z.-Q. Zhang, and C. T. Chan, Topological Subspace-Induced Bound State in the Continuum, *Phys. Rev. Lett.* **118**, 166803 (2017).
- [44] H. Xue, Y. Yang, F. Gao, Y. Chong, and B. Zhang, Acoustic higher-order topological insulator on a kagome lattice, *Nat. Mater.* **18**, 108 (2019).
- [45] X. Ni, M. Weiner, A. Alù, and A. B. Khanikaev, Observation of higher-order topological acoustic states protected by generalized chiral symmetry, *Nat. Mater.* **18**, 113 (2019).
- [46] H. Xue, Y. Yang, G. Liu, F. Gao, Y. Chong, and B. Zhang, Realization of an Acoustic Third-Order Topological Insulator, *Phys. Rev. Lett.* **122**, 244301 (2019).
- [47] M. Weiner, X. Ni, M. Li, A. Alù, and A. B. Khanikaev, Demonstration of a third-order hierarchy of topological states in a three-dimensional acoustic metamaterial, *Sci. Adv.* **6**, eaay4166 (2020).
- [48] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.127.214301> for more numerical and experimental details.
- [49] L. Ye, C. Qiu, M. Xiao, T. Li, J. Du, M. Ke, and Z. Liu, Topological dislocation modes in three-dimensional acoustic topological insulators, *arXiv:2104.04172*.
- [50] S. S. Yamada, T. Li, M. Lin, C. W. Peterson, T. L. Hughes, and G. Bahl, Bound states at partial dislocation defects in multipole higher-order topological insulators, *arXiv:2105.01050*.
- [51] Z.-K. Lin, Y. Wu, B. Jiang, Y. Liu, S. Wu, and J.-H. Jiang, Single-plaquette gauge flux as a probe of topological phases on lattices, *arXiv:2105.02070*.
- [52] C. Poli, M. Bellec, U. Kuhl, F. Mortessagne, and H. Schomerus, Selective enhancement of topologically induced interface states in a dielectric resonator chain, *Nat. Commun.* **6**, 6710 (2015).
- [53] S. Weimann, M. Kremer, Y. Plotnik, Y. Lumer, S. Nolte, K. G. Makris, M. Segev, M. C. Rechtsman, and A. Szameit, Topologically protected bound states in photonic parity–time-symmetric crystals, *Nat. Mater.* **16**, 433 (2017).
- [54] S. Weidemann, M. Kremer, T. Helbig, T. Hofmann, A. Stegmaier, M. Greiter, R. Thomale, and A. Szameit, Topological funneling of light, *Science* **368**, 311 (2020).
- [55] H. Zhao, X. Qiao, T. Wu, B. Midya, S. Longhi, and L. Feng, Non-Hermitian topological light steering, *Science* **365**, 1163 (2019).
- [56] W. Zhu, X. Fang, D. Li, Y. Sun, Y. Li, Y. Jing, and H. Chen, Simultaneous Observation of a Topological Edge State and Exceptional Point in an Open and Non-Hermitian Acoustic System, *Phys. Rev. Lett.* **121**, 124501 (2018).
- [57] H. Gao, H. Xue, Q. Wang, Z. Gu, T. Liu, J. Zhu, and B. Zhang, Observation of topological edge states induced solely by non-Hermiticity in an acoustic crystal, *Phys. Rev. B* **101**, 180303(R) (2020).
- [58] H. Gao, H. Xue, Z. Gu, T. Liu, J. Zhu, and B. Zhang, Non-Hermitian route to higher-order topology in an acoustic crystal, *Nat. Commun.* **12**, 1888 (2021).
- [59] L. Zhang, Y. Yang, Y. Ge, Y.-j. Guan, Q. Chen, Q. Yan, F. Chen, R. Xi, Y. Li, D. Jia *et al.*, Acoustic non-Hermitian skin effect from twisted winding topology, *arXiv:2104.08844*.
- [60] T. Helbig, T. Hofmann, S. Imhof, M. Abdelghany, T. Kiessling, L. Molenkamp, C. Lee, A. Szameit, M. Greiter, and R. Thomale, Generalized bulk–boundary correspondence in non-Hermitian topoelectrical circuits, *Nat. Phys.* **16**, 747 (2020).
- [61] S. Liu, S. Ma, C. Yang, L. Zhang, W. Gao, Y. J. Xiang, T. J. Cui, and S. Zhang, Gain-and Loss-Induced Topological Insulating Phase in a Non-Hermitian Electrical Circuit, *Phys. Rev. Applied* **13**, 014047 (2020).
- [62] L. Xiao, T. Deng, K. Wang, G. Zhu, Z. Wang, W. Yi, and P. Xue, Non-Hermitian bulk–boundary correspondence in quantum dynamics, *Nat. Phys.* **16**, 761 (2020).

- [63] X.-Q. Sun, P. Zhu, and T. L. Hughes, Geometric Response and Disclination-Induced Skin Effects in Non-Hermitian Systems, *Phys. Rev. Lett.* **127**, 066401 (2021).
- [64] A. Panigrahi, R. Moessner, and B. Roy, Non-Hermitian dislocation modes: Stability and melting across exceptional points, [arXiv:2105.05244](#).
- [65] F. Schindler and A. Prem, Dislocation non-Hermitian skin effect, *Phys. Rev. B* **104**, L161106 (2021).
- [66] B. A. Bhargava, I. C. Fulga, J. v. d. Brink, and A. G. Moghaddam, Non-Hermitian skin effect of dislocations and its topological origin, [arXiv:2106.04567](#).