Anti-Markovnikov hydrofunctionalization of alkenes : use of a benzyl group as a traceless redox-active hydrogen donor

Lonca, Geoffroy Hervé; Ong, Derek Yiren; Tran, Thi Mai Huong; Tejo, Ciputra; Chiba, Shunsuke; Gagosz, Fabien

2017

Lonca, G. H., Ong, D. Y., Tran, T. M. H., Tejo, C., Chiba, S. & Gagosz, F. (2017). Anti-Markovnikov hydrofunctionalization of alkenes : use of a benzyl group as a traceless redox-active hydrogen donor. *Angewandte Chemie International Edition*, 129(38), 11598-11602. <https://dx.doi.org/10.1002/anie.201705368>

<https://hdl.handle.net/10356/154855>

<https://doi.org/10.1002/anie.201705368>

This is the peer reviewed version of the following article: Lonca, G. H., Ong, D. Y., Tran, T. M. H., Tejo, C., Chiba, S. & Gagosz, F. (2017). Anti-Markovnikov hydrofunctionalization of alkenes : use of a benzyl group as a traceless redox-active hydrogen donor. *Angewandte Chemie International Edition*, 129(38), 11598-11602, which has been published in final form at <https://doi.org/10.1002/ange.201705368>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Anti-Markovnikov Hydrofunctionalization of Alkenes - Use of a Benzyl Group as a Traceless Redox-active Hydrogen Donor

Geoffroy Hervé Lonca,^a Derek Yiren Ong,^b Thi Mai Huong Tran,^b Ciputra Tejo,^b Shunsuke Chiba,^{*b} and Fabien Gagosz^{*a,c}

Abstract: A protocol for the *anti*-Markovnikov hydrofunctionalization of alkenes, that utilizes a benzyl group as a traceless redox-active hydrogen donor, has been developed. Under copper catalysis and in the presence of CF₃- or N₃-containing hypervalent iodine reagents, a series of homoallylic alcohol derivatives could be regioselectivity hydrofunctionalized. A similar principle was also applied to the hydrofunctionalization of alkenols.

The ability in hydrofunctionalization of alkenes in a regioselective manner is of prime importance for the development of chemical processes to produce high value fine chemicals. Despite the recent remarkable progress in this area,^[1] a number of challenges still remain for the *anti*-Markovnikov hydrofunctionalization of non-activated alkenes.^[2] One of the most reliable ways to achieve the *anti*-Markovnikov hydrofunctionalization involves the use of carbon- or heteroatom centered radicals which can indeed add on the less hindered/substituted side of an unactivated alkene, thus generating a new C-C or C-heteroatom bond in an *anti*-Markovnikov manner (Scheme 1A). Subsequent delivery of a H atom onto the resulting highly reactive C-centered radical is then required to complete the transformation. This last step often appears to be a key issue to realize an efficient hydrofunctionalization.^[3] In this context, we wondered if a simple benzyl ether moiety, that would be introduced in an alkenyl substrate **I** possessing a hydroxyl group at a suitable position (Scheme 1B, **I**→**II**), could be used after the radical functionalizing event (**II**→**III**) for the precise intramolecular H-delivery *via* a remote H radical shift (**III**→**IV**).^[4,5] We reasoned that this H-radical shift could be driven by the formation of a more stable α -oxy benzyl radical **IV**, which could be, if the suitable redox reaction conditions are set, readily converted into an oxonium cation **V** in a subsequent single-electron-oxidation step (**IV**→**V**). The cationic intermediate **V** could finally be degraded by hydrolysis/solvolysis to release the original hydroxyl group with formation of benzaldehyde as a co-product

(**V**→**VI**). Overall, the temporarily introduced benzyl moiety would function both as a redox active traceless H-donor and as a protecting group for the hydroxy moiety. Herein, we demonstrate the feasibility of this concept with the Cu-catalyzed hydrotrifluoromethylation^[6] and hydroazidation^[7,8] of readily accessible homoallylic alcohols (Scheme 1C). The design, optimization, and substrate scope of these processes are described herein.

A. *anti*-Markovnikov Hydrofunctionalization of alkenes with radicals

B. Use of Bn ether for the redox active hydrogen radical source

C. Cu-catalyzed hydrotrifluoromethylation and -azidation of homoallylic alcohols (this work)

Scheme 1. Anti-Markovnikov hydrofunctionalization of alkenes

We initiated our investigations by studying the possibility to perform the hydrotrifluoromethylation of the readily accessible homoallylic alcohol benzyl ether **1a** (Scheme 2A). The use of Togni's reagent **2**^[9] as a source of trifluoromethyl radical along with a copper salt as a redox catalyst was considered as a potentially suitable set of reagents to achieve the desired transformation.^[10] An extensive screening of the reaction conditions (see SI for details) revealed that the *anti*-Markovnikov hydrotrifluoromethylation of **1a** could be efficiently performed in the presence of CuI (10 mol%) and Togni's reagent **2** (1.2 equiv) in dioxane-MeOH (9:1) solvent system at 60 °C. Under these conditions, **1a** was completely consumed after 12 h of reaction,

- [a] G. H. Lonca, Prof. F. Gagosz
Département de Chimie, UMR 7652 and 7653 CNRS, Ecole Polytechnique, 91128 Palaiseau, France.
- [b] D. Y. Ong, T. M. H. Tran, Dr. C. Tejo, Prof. S. Chiba
Division of Chemistry and Biological Chemistry
School of Physical and Mathematical Sciences
Nanyang Technological University, Singapore 637371 (Singapore)
E-mail: shunsuke@ntu.edu.sg
- [c] Prof. F. Gagosz
Department of Chemistry and Biomolecular Sciences, University of Ottawa, K1N 6N5, Ottawa, Canada.

Supporting information for this article is given via a link at the end of the document.

and subsequent treatment of the reaction mixture with TsOH (20 mol%) in MeOH for 1 h led to isolation of the hydrotrifluoromethylated alcohol **3a** in 92% yield (Scheme 2A).^[11]

As shown in Scheme 2B, the catalytic process is initiated by a single-electron reduction of Togni's reagent **2** by the Cu(I) species. This allows for the generation of a CF₃-radical along with a higher-valent Cu(II) species and a 2-iodobenzoate ion. The CF₃-radical then undergoes an *anti*-Markovnikov addition onto the alkene moiety of **1a** to produce a secondary carbon centered radical **VII**. A subsequent 1,5-H radical shift leads to the formation of an α -oxy benzyl radical **VIII**,^[12,13] which is then oxidized by the Cu(II) species, thus producing the oxonium carbocation **IX** while maintaining the redox catalytic turnover with regeneration of the active Cu(I) species. Trapping of oxonium **IX** with MeOH produces acetal **X**,^[14] which is finally degraded in the presence of TsOH in MeOH to provide **3a**.

A. Optimized reaction conditions: hydrotrifluoromethylation of **1a**

B. Proposed catalytic cycle

Scheme 2. Cu-catalyzed hydrotrifluoromethylation of **1a** with Togni's reagent **2**.

With optimized reaction conditions in hands, we next investigated the scope and limitation of this *anti*-Markovnikov hydrotrifluoromethylation. As shown in Scheme 3, the transformation could be applied to a variety of homoallylic alcohols **1b-k** with good efficiency in general. Various functional groups such as silyl ether (**3c**), cyclopropyl (**3d**), phosphonate (**3e**), and thienyl (**3g**) moieties were tolerated. A quaternary carbon center could be introduced at position α to the alcohol with high efficiency (**3h**). In addition to secondary alcohols **3b-h**, tertiary alcohols **3i** and **3j** could be afforded in good yields, while the yield of **3k**, derived from camphor, was moderate. Moreover,

the present method was found applicable to the synthesis of primary alcohol such as **3l**. It could also be employed for the hydrotrifluoromethylation of *N*-benzylamine derivatives **1m** and **1n** to deliver amines **3m** and **3n** in 59% and 49% yields, respectively.

Scheme 3. Substrate scope for the Cu-catalyzed hydrotrifluoromethylation. [a] The reactions were conducted using 0.2–0.3 mmol of **1** with 10 mol% of CuI in 1,4-dioxane-MeOH (0.3 M) for 12–24 h followed by treatment with TsOH (20 mol%) in MeOH for solvolysis and isolated yields were noted above unless otherwise stated. [b] ¹H NMR yield based on the internal standard (**3d** was volatile and **3m** was not very stable). [c] 20 mol% of CuI was used. TBDPS = *tert*-butyldiphenylsilyl; PhthN = phthalimido; PMP = *para*-methoxyphenyl.

The method was not limited to the introduction of a CF₃ group onto alkenes. As proof of the synthetic potential of our approach, hydroazidation of alkenes **1** with hypervalent iodine reagent **4** as a source of the azido radical was developed by slightly modifying the reaction conditions (Scheme 4).^[15] Treatment of alkene **1a** with **4** (1.5 equiv) in the presence of CuBr₂^[16] (20 mol%) as the catalyst in 1,2-dichloroethane at 60 °C delivered, after the acidic work-up, the hydroazidation product **5a** in 57% isolated yield. Notably, this product is a precursor of 1,5-diamino-2-pentanol in which the two amino functionalities are orthogonally protected. The hydroazidation protocol could be applied to other homoallylic alcohol benzyl ethers **1**, which were converted into the corresponding azidoalcohols **5** in moderate to good yields. Given the versatile

reactivity of organic azides,^[17] the products obtained using this protocol would be of high value in various applications.

We also sought for the possibility to apply the current protocol to 1,1-disubstituted alkenes. As shown in Scheme 5A, the Cu(I)-catalyzed hydrotrifluoromethylation of methyl substituted alkene **1o** delivered the desired adduct **3o** in 72% yield but with poor diastereoselectivity (60:40). The replacement of the methyl group by a bromine (for **1p**), a chlorine (for **1q**) or a fluorine (for **1r**) atom did not alter the reactivity, affording **3p**, **3q**, and **3r** in 73%, 65%, and 55% (65% brsm) yields, respectively, and more fascinatingly, we observed an enhancement of the diastereoselectivity (84:16 for **3p**, 85:15 for **3q**, and 85:15 for **3r**). A similar level of selectivity was obtained in the hydroazidation of bromoalkene **1p**, that afforded **5p** in decent 47% yield with a good diastereoisomeric ratio (80:20). The origin of the diastereoselectivity likely stems from the transition states in the 1,5-H shift process.^[18] In the possible chair-like transition states **XI** and **XII** (Scheme 5B), both the phenyl group of the benzyl moiety and the phthalimidomethyl group would preferentially adopt a pseudo-equatorial position. Considering that a trifluoroethyl group or an azidomethyl group would be more sterically demanding than a halogen atom,^[19] the reaction would more favourably proceed via transition state **XII**, in which the halogen atom stands in a pseudo-axial position. As a result, the major isomer from **1p-r** would be the one having the resulting hydroxyl group and the halogen atom in a relative 1,3-*anti*-configuration (see Scheme 5A). We took advantage of the high density of functional groups present in **3p** and **5p** to derivatize them. They could be readily converted into the valuable pyrrolidines **6p** and **7p** through a sequence of hydroxyl group protection as a TBS ether, removal of the phthalimide moiety with hydrazine followed by spontaneous intramolecular substitution of the secondary bromide by the resulting primary amine (Scheme 5C).

A. diastereoselective hydrofunctionalization of 1o-1q

B. Origin of the diastereoselectivity

C. Derivatization of 3p and 5p to pyrrolidines 6p and 7p

Scheme 5. Hydrofunctionalization of 1,1-disubstituted alkenes. [a] Diastereomeric ratio was determined by ¹H NMR spectroscopy. [b] Diastereomeric ratio was determined by ¹⁹F NMR spectroscopy. [c] 65% yield based on recovery of **1r** (brsm). [d] i) TBSCl, 2,6-lutidine, CH₂Cl₂, 0 °C, 95% yield; ii) aq. NH₂NH₂, EtOH, 60 °C, 97% yield of **6p** as an inseparable mixture of two diastereomers. [e] i) TBSCl, imidazole, DMF, 23 °C, 68% yield of the single diastereomer derived from the major isomer of **5p**; ii) aq. NH₂NH₂, EtOH, 60 °C, 96% yield of **7p** as the single diastereomer.

We also envisaged using the present concept to perform the hydrotrifluoromethylation of a simple alkenol **XIII** (Scheme 6). In this case, a suitably located alcohol moiety in **XIII** would play the role of a redox-active hydrogen donor. We considered that the 1,n-H shift step (**XIV**→**XV**) would proceed favourably from the α-H of the hydroxyl group to the reactive C-centered radical, while the resulting hydroxy-substituted C-centered radical intermediate **XV** could be subsequently oxidized into ketone **XVI**. Overall, this transformation would be a combination of *anti*-Markovnikov hydrotrifluoromethylation and alcohol oxidation.

Scheme 6. Cu-catalyzed hydrotrifluoromethylation of alkenes using alcohols as redox active hydrogen donor groups

Gratifyingly, we found that the treatment of 1-aryl-5-hexenols **8a-k** with **2** (2 equiv) in the presence of Cu(OAc)₂ (10 mol%)^[16] and 2,2'-bipyridine (10 mol%) in 1,4-dioxane at 60 °C resulted in the formation of the hydrotrifluoromethylated ketones **9a-k** in generally good yields (Scheme 7). Notably, the cyclopropyl moiety in **9k** could be kept intact during the redox radical process probably due to a higher oxidation rate of the α-hydroxy radical **XV** to ketone **9k** than the radical ring-opening. The method could be extended to the synthesis of benzene-tethered carbonyl compounds **9l** and **9m** as well as aliphatic ketones such as **9n**. Interestingly, the reaction of 1-phenyl-6-heptenol **8o**, which has a linker with an additional carbon unit (as compared to **8a-n**), proceeded smoothly via a probable 1,6-hydrogen shift to furnish **9o** in 66% yield. The method also allowed for the use of substituted alkenes. The reaction of 1,1-disubstituted alkene **8p** was shown to be highly efficient affording the desired hydrotrifluoromethylated ketone **9p** in 89% yield. The reaction of **8q** possessing a styryl moiety (as an example of internal alkene substrate) provided the γ-trifluoromethyl ketone **9q** in 57% yield via benzylic radical **XVII** intermediately formed by a regioselective addition of the CF₃ radical onto the styryl moiety.

In summary, this work demonstrated that a simple benzyl protecting group can be employed as a traceless redox-active hydrogen radical donor for the *anti*-Markovnikov alkene hydrofunctionalization of homoallylic alcohols and their derivatives. The hydrotrifluoromethylation and azidation developed herein are easy to set up, relatively efficient and exhibit a high functional group tolerance. This concept has been successfully extended to the hydrotrifluoromethylation of 5-hexenol and 6-heptenol derivatives. Application of this general concept to develop other types of catalytic redox molecular transformations is the subject of our ongoing investigations.

Scheme 7. Cu-catalyzed sequence of hydrotrifluoromethylation/oxidation of alkenols. [a] The reactions were conducted using 0.5 mmol of **8** in 1,4-dioxane (0.1 M) at 60 °C for 2–61 h and isolated yields were noted above. [b] The reaction was run at 80 °C. [c] brsm = based on recovery of the starting material.

Acknowledgements

This work was supported by funding from Ecole Polytechnique (for F.G.), Nanyang Technological University (for S.C.) and the Singapore Ministry of Education (Academic Research Fund Tier 1: RG2/15 for S.C.). F.G. and S.C. are grateful to PHC Merlion grant (Project 5.02.15) for the support of this collaboration project.

Keywords: *anti*-Markovnikov alkene functionalization • trifluoromethylation • azidation • radicals • copper catalysis

- [1] For reviews, see: a) M. Beller, J. Seayad, A. Tillack, H. Jiao, *Angew. Chem. Int. Ed.* **2004**, 43, 3368; *Angew. Chem.* **2004**, 116, 3448; b) L.-B. Han, M. Tanaka, *Chem. Commun.* **1999**, 395.
- [2] K. A. Margrey, D. A. Nicewicz, *Acc. Chem. Res.* **2016**, 49, 1997.
- [3] For a seminal work on radical mediated *anti*-Markovnikov hydroamination by Studer *et al.*, see: a) J. Guin, C. Mück-Lichtenfeld, S. Grimme, A. Studer, *J. Am. Chem. Soc.* **2007**, 129, 4498; b) J. Kemper, A. Studer, *Angew. Chem. Int. Ed.* **2005**, 44, 4914; *Angew. Chem.* **2005**, 117, 4993.

- [4] For selected reviews, see: a) S. Chiba, H. Chen, *Org. Biomol. Chem.* **2014**, *12*, 4051. b) F. Dénès, F. Beaufils, P. Renaud, *Synlett*, **2008**, 2389; c) J. Robertson, J. Pillai, R. K. Lush, *Chem. Soc. Rev.* **2001**, *30*, 94; d) G. Majetich, K. Wheelless, *Tetrahedron* **1995**, *51*, 7095.
- [5] Liu and Tan developed elegant strategies on remote aliphatic C-H functionalization triggered by radical trifluoromethylation of alkenes, see: a) P. Yu, S.-C. Zheng, N.-Y. Yang, B. Tan, X.-Y. Liu, *Angew. Chem. Int. Ed.* **2015**, *54*, 4041; *Angew. Chem.* **2015**, *127*, 4113; b) P. Yu, J.-S. Lin, L. Li, S.-C. Zheng, Y.-P. Xiong, L.-J. Zhao, B. Tan, X.-Y. Liu, *Angew. Chem. Int. Ed.* **2014**, *53*, 11890; *Angew. Chem.* **2014**, *126*, 12084; c) L. Huang, S.-C. Zheng, B. Tan, X.-Y. Liu, *Org. Lett.* **2015**, *17*, 1589; d) L. Huang, S.-C. Zheng, B. Tan, X.-Y. Liu, *Chem. Eur. J.* **2015**, *21*, 6718; e) L. Huang, J.-S. Lin, B. Tan, X.-Y. Liu, *ACS Catal.* **2015**, *5*, 2826.
- [6] For recent reports on *anti*-Markovnikov hydrotrifluoromethylation of alkenes, see: a) L. Zhu, L.-S. Wang, B. Li, B. Fu, C.-P. Zhang, W. Li, *Chem. Commun.* **2016**, *52*, 6371; b) Y. Cheng, S. Yu, *Org. Lett.* **2016**, *18*, 2962; c) H. Egami, Y. Usui, S. Kawamura, S. Nagashima, M. Sodeoka, *Chem. Asian J.* **2015**, *10*, 2190; d) S. Choi, Y. J. Kim, S. M. Kim, J. W. Yang, S. W. Kim, E. J. Cho, *Nature Commun.* **2014**, *5*, 4881; e) D. J. Wilger, N. J. Gesmundo, D. A. Nicewicz, *Chem. Sci.* **2013**, *4*, 3160; f) X. Wu, L. Chu, F.-L. Qing, *Angew. Chem. Int. Ed.* **2013**, *52*, 2198; *Angew. Chem.* **2013**, *125*, 2254; g) S. Mizuta, S. Verhoog, K. M. Engle, T. Khotavivattana, M. O'Duill, K. Wheelhouse, G. Rassias, M. Médebielle and V. Gouverneur, *J. Am. Chem. Soc.* **2013**, *135*, 2505.
- [7] For a report on *anti*-Markovnikov hydroazidation of alkenes, see: A. Kapat, A. König, F. Montermini, P. Renaud, *J. Am. Chem. Soc.* **2011**, *133*, 13890.
- [8] For reports on Markovnikov hydroazidation of alkenes, see: a) J. Waser, H. Nambu, E. M. Carreira, *J. Am. Chem. Soc.* **2005**, *127*, 8294; b) J. Waser, B. Gaspar, H. Nambu, E. M. Carreira, *J. Am. Chem. Soc.* **2006**, *128*, 11693; c) B. Gaspar, J. Waser, E. M. Carreira, *Synthesis*, **2007**, 3839.
- [9] For a review, see: J. Charpentier, N. Früh, A. Togni, *Chem. Rev.* **2015**, *115*, 650.
- [10] For the mechanistic study on the reactions of Togni reagents with Cu(I) complex, see: a) F. Wang, D. Wang, X. Wan, L. Wu, P. Chen, G. Liu, *J. Am. Chem. Soc.* **2016**, *138*, 15547; b) S. Kawamura, H. Egami, M. Sodeoka, *J. Am. Chem. Soc.* **2015**, *137*, 4865.
- [11] Use of MOM, Et, and PMB ethers was not optimal for this transformation (see the SI for more details).
- [12] The selective hydrogen atom delivery from the benzylic position (IV→VIII) was supported by the reaction of deuterated substrate **D₂-1a** which, under identical reaction conditions, was converted into mono-deuterated compound **D-3a**.
-
- [13] Beau reported regioselective de-O-benzoylation on carbohydrate templates using xanthate-mediated 1,7-H radical shift, see: A. Attouche, D. Urban, J.-M. Beau, *Angew. Chem. Int. Ed.* **2013**, *52*, 9572; *Angew. Chem.* **2013**, *125*, 9751.
- [14] Formation of acetal **X** was confirmed prior to treatment with TsOH in MeOH.
- [15] To the best of our knowledge, this is the first example of hydroazidation of non-activated alkenes with azido iodine(III) reagent **4**. For reports on the difunctionalization of alkenes with **4**, see: a) Y.-A. Yuan, D.-F. Lu, Y.-R. Chen, H. Xu, *Angew. Chem. Int. Ed.* **2016**, *55*, 534; *Angew. Chem.* **2016**, *128*, 544; b) G. Fumagalli, P. T. G. Rabet, S. Boyd, M. F. Greaney, *Angew. Chem. Int. Ed.* **2015**, *54*, 11481; *Angew. Chem.* **2015**, *127*, 11643; c) M.-Z. Lu, C.-Q. Wang, T.-P. Loh, *Org. Lett.* **2015**, *17*, 6110; d) B. Zhang, A. Studer, *Org. Lett.* **2013**, *15*, 4548.
- [16] Cu(II) species could be readily reduced to Cu(I) by various factors to initiate the catalytic cycle. For several relevant literature precedents, see: (a) S. Chiba, *Chem. Lett.* **2012**, *41*, 1554; b) Y.-F. Wang, K. K. Toh, J.-Y. Lee, S. Chiba, *Angew. Chem. Int. Ed.* **2011**, *50*, 5927; *Angew. Chem.* **2011**, *123*, 6049; c) J. Kim, S. Chang, *J. Am. Chem. Soc.* **2010**, *132*, 10272; d) A. Y. S. Malkhasian, M. E. Finch, B. Nikolovski, A. Menon, B. E. Kucera, F. A. Chavez, *Inorg. Chem.* **2007**, *46*, 2950; e) J. J. Teo, Y. Chang, H. C. Zeng, *Langmuir* **2006**, *22*, 7369.
- [17] a) S. Bräse, C. Gil, K. Knepper, V. Zimmermann, *Angew. Chem. Int. Ed.* **2005**, *44*, 5188; *Angew. Chem.* **2005**, *117*, 5320; b) *Organic Azides: Syntheses and Applications*; S. Bräse, K. Banert, Eds.; Wiley: Chichester, U.K., 2010.
- [18] A. E. Dorigo, M. A. McCarrick, R. J. Loncharich, K. N. Houk, *J. Am. Chem. Soc.* **1990**, *112*, 7508.
- [19] The relative steric demands are evaluated on the basis of the A-values for a bromine atom (0.48–0.67 kcal/mol), a chlorine atom (0.53–0.64 kcal/mol), a fluorine atom (0.25–0.42 kcal/mol), and an ethyl group (1.79 kcal/mol) whose value should be close to that of a trifluoroethyl or azidomethyl group. For A-values of various substituents, see: E. L. Eliel, S. H. Wilen, M. P. Doyle, *Basic Organic Stereochemistry*, Wiley, 2001, p. 443.

COMMUNICATION

Geoffroy Hervé Lonca, Derek Yiren Ong,
Thi Mai Huong Tran, Ciputra Tejo,
Shunsuke Chiba* and Fabien Gagosz*

Page No. – Page No.

**Anti-Markovnikov
Hydrofunctionalization of Alkenes -
Use of a Benzyl Group as a Traceless
Redox-active Hydrogen Donor**

A protocol for the *anti*-Markovnikov hydrofunctionalization of alkenes, that utilizes a benzyl group as a traceless redox-active hydrogen donor, has been developed. Under copper catalysis and in the presence of CF₃- or N₃-containing hypervalent iodine reagents, a series of homoallylic alcohol derivatives could be regioselectively hydrofunctionalized. A similar principle was also applied to the hydrofunctionalization of alkenols.