

# Backbone $^1\text{H}$ , $^{13}\text{C}$ , and $^{15}\text{N}$ resonance assignments of the N-terminal domain of FKBP38 (FKBP38NTD)

Kang, Cong Bao; Ye, Hong; Vivekanandan, Subramanian; Simon, Bernd; Sattler, Michael; Yoon, Ho Sup

2006

Kang, C. B., Ye, H., Vivekanandan, S., Simon, B., Sattler, M., & Yoon, H. S. (2006). Backbone  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{15}\text{N}$  resonance assignments of the N-terminal domain of FKBP38 (FKBP38NTD). *Journal of Biomolecular NMR*, 36(S1), 37.

<https://hdl.handle.net/10356/79968>

<https://doi.org/10.1007/s10858-006-9001-5>

---

© 2006 Springer. This is the author created version of a work that has been peer reviewed and accepted for publication by *Journal of Biomolecular NMR*, Springer. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: <http://dx.doi.org/10.1007/s10858-006-9001-5>.

*Downloaded on 23 Mar 2023 10:32:36 SGT*

Letter to the Editor

## Backbone $^1\text{H}$ , $^{13}\text{C}$ , and $^{15}\text{N}$ resonance assignments of the N-terminal domain of FKBP38 (FKBP38NTD)

*Cong Bao Kang<sup>a,†</sup>, Hong Ye<sup>a,†</sup>, Subramanian Vivekanandan<sup>a</sup>, Bernd Simon<sup>b</sup>, Michael Sattler<sup>b</sup>, Ho Sup Yoon<sup>a,\*</sup>*

*<sup>a</sup>School of Biological Science, Nanyang Technological University, 60 Nanyang Drive, Singapore 637665, Singapore; <sup>b</sup>Structural Biology Group, European Molecular Biology Laboratory, Heidelberg, Germany*

*\*To whom correspondence should be addressed. E-mail: hsyoon@ntu.edu.sg*

*<sup>†</sup>These authors equally contributed to this work.*

FK-506 binding protein 38 (FKBP38) interacts with Bcl-2/Bcl-Xl and helps them localize at the mitochondrial membrane (Shirane and Nakayama, 2003). The down-regulation of FKBP38 was shown to influence on the stability of Bcl-2 and consequently induce apoptotic cell death (Kang et al., 2005). FKBP38 is a unique protein among the FKBP family members. Unlike other members in the FKBP family, FKBP38 appears to have no FK-506 binding activity (Shirane and Nakayama, 2003). Thus, to understand the function of FKBP38 at molecular level, as the first step, we performed a NMR study on FKBP38, and here we report the NMR resonance assignments of the N-terminal domain of FKBP38, which includes the FK-506 binding domain and the flanking N-terminal residues. Nearly all of the backbone  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{15}\text{N}$  resonances were assigned (~99%). Assignments of the side chain atoms beyond  $\text{C}^\beta$  atoms are about 80% complete. The assignments have been deposited with BMRB accession number 6923.

**Supplementary material** is available in electronic format at <http://www.dx.doi.org/10.1007/s10858-006-9001-5>.