Dispersal from microbial biofilms

Barraud, Nicolas; Kjelleberg, Staffan; Rice, Scott A.

2015

Barraud, N., Kjelleberg, S., & Rice, S. A. (2015). Dispersal from microbial biofilms. *Microbiology Spectrum*, 3(6). doi:10.1128/microbiolspec.MB-0015-2014

<https://hdl.handle.net/10356/80430>

<https://doi.org/10.1128/microbiolspec.MB-0015-2014>

© 2015 American Society for Microbiology. This paper was published in *Microbiology Spectrum* with permission of American Society for Microbiology. The published version is available at: [<http://dx.doi.org/10.1128/microbiolspec.MB-0015-2014>]. One print or electronic copy may be made for personal use only. Systematic or multiple reproduction, distribution to multiple locations via electronic or other means, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper is prohibited and is subject to penalties under law.

Downloaded on 26 Apr 2023 19:29:08 SGT

Dispersal from Microbial Biofilms

NICOLAS BARRAUD,¹ STAFFAN KJELLEBERG,^{1,2} and SCOTT A. RICE^{1,2}

¹Centre for Marine Bio-Innovation and School of Biotechnology and

Biomolecular Sciences, The University of New South Wales, Sydney, NSW 2052, Australia;

²The Singapore Centre for Environmental Life Sciences Engineering, and the School of Biological Sciences, Nanyang Technological University, Singapore 639798

ABSTRACT One common feature of biofilm development is the active dispersal of cells from the mature biofilm, which completes the biofilm life cycle and allows for the subsequent colonization of new habitats. Dispersal is likely to be critical for species survival and appears to be a precisely regulated process that involves a complex network of genes and signal transduction systems. Sophisticated molecular mechanisms control the transition of sessile biofilm cells into dispersal cells and their coordinated detachment and release in the bulk liquid. Dispersal cells appear to be specialized and exhibit a unique phenotype different from biofilm or planktonic bacteria. Further, the dispersal population is characterized by a high level of heterogeneity, reminiscent of, but distinct from, that in the biofilm, which could potentially allow for improved colonization under various environmental conditions. Here we review recent advances in characterizing the molecular mechanisms that regulate biofilm dispersal events and the impact of dispersal in a broader ecological context. Several strategies that exploit the mechanisms controlling biofilm dispersal to develop as applications for biofilm control are also presented.

INTRODUCTION

For all organisms the ability to spread and colonize new habitats is crucial to ensure species continuity and prevent extinction (1). In sessile organisms this constraint has led to the evolution of a motile, dispersal phase in their life cycle, which in plants and corals involves the release of differentiated and often phenotypically diverse seeds or propagules. Similarly, sessile microbial biofilms have developed mechanisms to release differentiated, highly motile dispersal cells into the bulk liquid.

Dispersal has profound ecological consequences, allowing biofilm populations to spread and colonize new surfaces as well as to avoid overcrowding and regenerate bacteria in the biofilm core. These benefits represent strong evolutionary pressures for dispersal,

propagation, and rejuvenation of biofilms. The ecological drivers of dispersal are complex, because dispersal is both costly (2) and risky when dispersal cells face an uncertain and uncontrolled environment (3). However, the inability to disperse may ultimately lead to the collapse of any sessile biological community through the accumulation of social cheaters, such as those that disturb metabolic cooperation (4), or catastrophic events at one location resulting in either complete loss of the population leading to extinction or in genetic bottlenecks as a consequence of significant population die-off (5).

Sloughing is a passive process of cell loss from the biofilm, which may be a consequence of cell division at the biofilm–bulk liquid interface or shear forces that remove cells. In contrast, active or seeding dispersal is coordinated via regulatory systems and is energy dependent, and dispersal cells exhibit a distinct phenotype. Dispersal relies on a number of cues, both environmental and self-produced signals. Regulatory responses to changes in environmental conditions potentially enhance the chances of successful release of propagules and colonization of new surfaces. In addition, signals and cues from other biological organisms can also influence microbial dispersal. Cells that have

Received: 9 September 2014, **Accepted:** 30 September 2014,
Published: 20 November 2015

Editors: Mahmoud Ghannoum, Case Western Reserve University, Cleveland, OH; Matthew Parsek, University of Washington, Seattle, WA; Marvin Whiteley, University of Texas at Austin, Austin, TX; and Pranab Mukherjee, Case Western Reserve University, Cleveland, OH

Citation: Barraud N, Kjelleberg S, Rice SA. 2015. Dispersal from microbial biofilms. *Microbiol Spectrum* 3(6):MB-0015-2014. doi:10.1128/microbiolspec.MB-0015-2014.

Correspondence: Staffan Kjelleberg, s.kjelleberg@unsw.edu.au

© 2015 American Society for Microbiology. All rights reserved.

dispersed from biofilms exhibit a specialized phenotype that is distinct from that of biofilm and planktonic cells. Moreover, the dispersal subpopulation often comprises a high level of heterogeneity, as a consequence of cell differentiation and phase variation as well as genetic diversification through mutations. The production of such variation in dispersal cells has been hypothesized to increase population fitness, e.g., enhanced colonization and survival due to complementarity and selection effects associated with the formation of such variants.

This article aims to describe features that are common to the dispersal process across species and presents specific examples to define those features. The underlying regulatory and signaling systems are examined in terms of both the extracellular cues and the internal signal transduction cascades that control active dispersal. Further, the consequences of dispersal for the microbial population are addressed. Finally, because dispersal offers unique opportunities to control biofilms, several strategies currently being developed to exploit dispersal pathways and design novel biofilm control measures are described.

THE MOTILE PHASE OF THE BIOFILM LIFE CYCLE

It is now well established that biofilm formation occurs through a series of stages that resemble developmental processes in multicellular eukaryotes. Recent studies have characterized specific regulators involved at each stage of development (6). Initial adhesion to a surface is thus controlled by nutrient availability, surface sensing, and quorum sensing (QS) (7). While flagella-mediated motility may play a role for planktonic bacteria to reach a surface and attach, the initiation of biofilm formation often leads to the loss of flagella (8). Biofilm bacteria are essentially sessile, embedded in a matrix of extracellular polymeric substances (EPSs), with the exception of a subpopulation of cells using surface motility that migrate within the biofilm to form the caps of three-dimensional (3D) microcolonies (9). In the model organism *Pseudomonas aeruginosa*, maturation of the biofilm involves production of specific matrix components and sequentially activated regulatory systems (10). These mechanisms include positive and negative feedback regulations, which are pivotal components of developmental biology. For example, the polysaccharide synthesis locus (Psl) is necessary for *P. aeruginosa* cells to attach onto a surface and subsequently serves to recruit incoming bacteria into the biofilm, establishing a feed-forward loop amplifying biofilm attachment and

maturation (11). Over the course of the developmental process, bacteria adopt a succession of different phenotypes. While the biofilm phenotype appears adaptable to changes in environmental conditions, notably through stress responses, a core genetic program takes the biofilm through a series of checkpoints, eventually leading to maturation and dispersal of the biofilm.

During maturation, biofilms establish complex 3D structures comprised of differentiated bacteria, for instance exhibiting different expression of metabolic or EPS genes, and steep nutrient and oxygen (O₂) gradients, rendering the biofilm environment and the bacterial populations highly heterogeneous (12–14). The final stage of biofilm development involves the coordinated release of differentiated, motile, chemotactic cells known as dispersal cells (15). These specialized cells can colonize new surfaces and restart the biofilm life cycle. In several bacteria, biofilm dispersal correlates with the programmed death of a subpopulation of cells in mature microcolonies (16). Surviving cells are then able to escape the biofilm, leaving behind hollow structures in the biofilm. In many species, dispersal typically occurs from mature biofilm microcolonies and is preceded by localized death and lysis of cells in the center of these structures (16–22) (Fig. 1). Bacteria were observed to disperse from larger microcolonies, e.g., with a diameter greater than 40 µm and depth greater than 10 µm, but not from smaller ones (17, 23). This suggests that a specific maturation stage, as defined by the microcolony size and most likely linked to the establishment of specific gradients, e.g., nutrients or O₂, is required to trigger dispersal events. In *P. aeruginosa*, cell lysis has been linked to the activation of a superinfective prophage (16), while in the marine bacterium *Pseudoalteromonas tunicata*, the expression and activity of an autolytic protein caused cell death in mature biofilm microcolonies (24).

It has been shown that dispersal can be prevented by disrupting the cells' ability to generate energy (25), and thus, the surviving cells may benefit from nutrients released from lysed cells, and this energy is used to activate dispersal responses. This is supported by microscopic observations of biofilms showing that mature microcolonies undergo a brief "seething" stage, where cells become highly motile within microcolonies, suggesting that biofilm EPSs have been fully solubilized and that cells have regained motility (17, 26) (Figs. 1 and 2). Dispersal cells then escape by coordinated evacuation from biofilm structures (26), resulting in the hollowing of biofilm microcolony structures leaving behind voids that are typically observed during the dispersal stage for many biofilms. Dispersal cells are characterized by downregulation of

FIGURE 1 Microscopic images of biofilm microcolonies during seeding dispersal. (A-B) Motile cells appear in mature biofilm microcolonies. (A) Single frame of a mature microcolony. (B) Picture showing the average of 30 frames captured over a 1-second period. The highly motile cells “average” out and appear blurred in the center of the microcolony, demonstrating the extent of the motile region (white arrow in panels A and B). The sessile “wall” region is indicated by the black arrows in panel A (taken from reference 26, with permission from the publisher). (C) Live/dead staining of a 7-day-old biofilm reveals patterns of cell death inside biofilm structures that occur simultaneously with biofilm dispersal, as indicated by the formation of hollow biofilm structures. Live cells are green and dead cells are red (adapted from reference 56, copyright © American Society for Microbiology). (D) XZ cross-view of the biofilm in panel C (XY view) at the location indicated by the white line. [doi:10.1128/microbiolspec.MB-0015-2014.f1](https://doi.org/10.1128/microbiolspec.MB-0015-2014.f1)

genes that encode for the sessile biofilm phenotype, such as exopolysaccharides and fimbriae, and upregulation of genes encoding factors important for the motile lifestyle, including flagella and chemotaxis (27–30).

REGULATION AND COORDINATION IN SEEDING DISPERSAL

The release of dispersal cells from biofilms occurs in a precisely coordinated manner and in response to changes

in environmental conditions. Growth in a biofilm can lead to stress; for instance, as the biofilm grows in size, bacteria in the lower layers will experience reduced access to nutrients and electron acceptors available from the bulk liquid interface or will accumulate waste products and toxins. Stress cues including nutrient starvation, either carbon (31, 32), phosphate (33), or iron sources (34), can trigger dispersal responses. Depletion of O₂ (35, 36) and addition of the metabolism inhibitor carbonylcyanide *m*-chlorophenylhydrazone (CCCP) (37)

FIGURE 2 Effectors of biofilm dispersal. Bacteria within the center of microcolonies induce a number of mechanisms to degrade and solubilize the biofilm EPS matrix and extracellular appendages such as fimbriae that immobilize cells. When the interior of the microcolony becomes fluid, cells begin to show signs of motility, and a breach is made in the microcolony wall through which dispersal cells are released. [doi:10.1128/microbiolspec.MB-0015-2014.f2](https://doi.org/10.1128/microbiolspec.MB-0015-2014.f2)

were also found to induce dispersal. In mature biofilms, dispersal may also be triggered by specific responses to the presence of soluble signals derived from protozoan predators, which are produced whether in the presence or absence of bacteria (unpublished observation). An increase in nutrient availability, such as carbon (38) or iron sources (39), can also trigger dispersal and lead to mass detachment events. Possibly, dispersal in response to nutrient upshift could represent a strategy to enhance chances of successful colonization by dispersal cells by ensuring an adequate supply of nutrient sources. Interestingly, biofilms of the plant pathogen *Xylella fastidiosa* were found to use cell-cell signaling to coordinate dispersal events in response to sensing the presence of a transmission vector, e.g., leafhopper insects (40). In this case it is clear that precise regulation and rapid signal transduction are necessary for the bacteria to trigger dispersal processes upon sensing the presence of the insect before it leaves the plant surface. Transmission vectors for dispersal and spreading of bacterial pop-

ulations are common in nature. In a recent study, a nonmotile *Xanthomonas* sp. was found to associate with a motile *Paenibacillus vortex*, in a hitchhiking strategy, to spread and disperse on plant surfaces (41).

Cell-cell signaling is known to play a major role in regulating dispersal in various species (42). For example, QS via *N*-acyl homoserine lactones has been implicated in dispersal in *Vibrio* species (43, 44), *Serratia marcescens* (45), and *Rhodobacter sphaeroides* (46). In *Staphylococcus aureus*, the *agr*-mediated QS system was found to control biofilm dispersal (47). In *Vibrio cholerae*, QS regulates detachment (48), and add-back of cholera QS autoinducer-1 (CAI-1) and autoinducer-2 (AI-2) resuscitate dormant cells and induce dispersal of biofilm aggregates in environmental samples (49). In *Xanthomonas campestris*, cell-cell signaling via the diffusible signal *cis*-11-methyl-2-dodecenoic acid triggers dispersal responses (50). This class of signal molecule has more recently been shown to be conserved across a broader range of bacteria, where it induces dispersal

(51), including cis-2-decenoic acid, a potent dispersal trigger in various bacterial species (23). Further, the addition of QS inhibitors, such as furanones, can induce detachment of biofilm bacteria and prevent biofilm formation (52, 53). The role of cell-cell signaling in regulating dispersal may allow biofilm bacteria to induce the switch at an appropriate stage of the developmental process, e.g., once a threshold density is reached, or to transmit a dispersal signal from one location to another within the biofilm population.

Several research groups have also focused their efforts on identifying signal molecules produced in mature cultures entering stationary phase that were thought to trigger dispersal events. The diffusible fatty acid cis-2-decenoic acid mentioned above was first identified after fractionation of *P. aeruginosa* spent medium that was active at inducing dispersal (23). In *Bacillus subtilis*, the readdition of a conditioned medium from late biofilm cultures was also found to induce biofilm disassembly, and D-amino acids were tentatively identified as active compounds, suggesting a role in regulating dispersal (54). However, later studies from the same team revealed that the observed effects were due to a mutation in the experimental strain which caused addition of D-amino acids to reduce metabolism in these biofilms; thus, dispersal in these studies was found to rely entirely on reduced metabolism (55), which is similar to the starvation-induced dispersal for some species as described above.

In *P. aeruginosa*, studies of cell death and dispersal events led to the identification of nitric oxide (NO) as a major signal regulating dispersal (56). NO was found to be produced in mature biofilms and trigger detachment events. NO was also found to induce dispersal in a wide range of monospecies biofilms and mixed-species biofilms (reviewed in 57). Thus, biofilm dispersal appears to be controlled by a variety of cues and signals, some of which, such as N-acyl homoserine lactones, play dual roles in biofilm formation as well as dispersal (7).

One important system that is involved in integrating these signals is the regulatory network centered on the secondary messenger cyclic di-GMP, because it controls both attachment and dispersal (reviewed in 58, 59). Mainly found in Gram-negative bacteria, c-di-GMP is a central element of a signaling network that integrates one or multiple signals (input) sensed by a bacterium and activates cellular effectors (output), resulting in either biofilm attachment or dispersal. Intracellular levels of c-di-GMP are controlled through the opposing activities of diguanylate cyclases (DGCs) for the synthesis of c-di-GMP, and phosphodiesterases (PDEs) for its

degradation. These are encoded by conserved GGDEF-domain- and EAL- or HDGYP-domain-containing genes, respectively. Many bacterial genomes encode multiple DGCs and PDEs often associated with other putative signaling domains, suggesting that their enzymatic activities may be responsive to a range of environmental cues, including NO, O₂, and nutrients. Recently, cAMP, which was previously known to control the stringent response, was found to play a role in the regulation of biofilm formation and dispersal (25, 60). The role that nucleotide secondary messengers such as c-di-GMP and cAMP play in signal transduction in bacteria is being increasingly recognized (61, 62; cf. online census http://www.ncbi.nlm.nih.gov/Complete_Genomes/SignalCensus.html), which is perhaps not surprising given the abundance of nucleotides in the intracellular milieu and their relationship to metabolic activity, a key dispersal trigger.

A link between NO and c-di-GMP in the regulation of biofilm dispersal was first established in *P. aeruginosa* (29) and was then also described in *Shewanella woodyi* (63) and *Legionella pneumophila* (64). In *P. aeruginosa* several c-di-GMP-specific PDEs have been identified that appear to be involved in NO-mediated dispersal, including DipA, RbdA, and NbdA (65, 66). In some bacteria, the sensor for NO has been identified. For example, *S. woodyi* encodes a heme NO/O₂-binding (HNOX) protein, which when complexed with NO, binds to and activates a PDE enzyme, resulting in dispersal. HNOX domains are conserved hemoproteins that are highly sensitive to NO, producing responses at femtomolar levels in *Clostridium botulinum* (67). They are found in several Gram-negative and Gram-positive bacterial genomes and are often associated with a DGC or PDE (68). However, many bacterial strains known to disperse in response to NO, including *P. aeruginosa* and *Escherichia coli*, do not have HNOX-domain-encoding proteins, suggesting that other systems can sense NO signals. In *E. coli*, a redox sensor associated with a c-di-GMP PDE, EcDOS was shown to increase PDE activity upon binding of NO, O₂, and carbon monoxide (69). EcDOS was later shown to be part of a two-gene operon for the control c-di-GMP in response to O₂ levels where sensing of O₂ activates both DGC and PDE activity. This suggests a feedback control mechanism, similar to a thermostat, to regulate biofilm formation and dispersal precisely, depending on O₂ levels (70).

In *P. aeruginosa* the methyl-accepting protein BdlA has been found to be involved in the dispersal response to both nutrient upshift (28) and NO (29) and interacts with a PDE, possibly by stabilizing the enzyme, resulting in lower intracellular levels of c-di-GMP and enhanced

dispersal (65). Recently, an interesting mechanism of feedback regulation was uncovered when it was found that BdlA needed to be cleaved before activating dispersal. Further, cleavage of BdlA, involving the chaperone ClpD and protease ClpP, requires high levels of c-di-GMP (71). Thus, high levels of c-di-GMP, which are typical of biofilm cells, appear to be a prerequisite to induce dispersal events in mature biofilms in this strain.

EFFECTORS OF DISPERSAL

Upon sensing a dispersal cue, bacteria can activate a range of cellular effectors that allow them to break their bonds to the biofilm, including enzymes and surfactants that will degrade the biofilm EPS, resulting in dispersal (Fig. 2).

First, bacteria can produce and secrete enzymes that degrade the biofilm EPS matrix, the composition of which varies but is mainly comprised of polysaccharides, proteins, nucleic acids, and lipids. In the nonmotile species *Aggregatibacter actinomycetemcomitans*, a screen of transposon mutant biofilms deficient in the ability to release cells identified dispersinB, an endogenous β -N-acetylglucosaminidase capable of degrading the matrix polysaccharides, which was later found to induce biofilm dispersal in a range of bacterial species (72, 73). In *P. aeruginosa*, alginate is an essential component of the matrix, and early studies revealed that increased expression of alginate lyase leads to biofilm detachment (74). Furthermore, it was found that DNA is a major constituent of the EPS matrix, and biofilm dispersal could be induced upon treatment with DNase in *P. aeruginosa* and *S. aureus* (75, 76). DNase was also found to be endogenously produced and secreted in *Bacillus licheniformis* biofilms to induce dispersal (77). In *S. aureus*, extracellular protease activity mediated by the agr QS system was also found to trigger biofilm detachment (47). In *Pseudomonas fluorescens* (33), *Pseudomonas putida* (78), and *P. aeruginosa* (25), starvation-induced dispersal was linked to activation of a periplasmic cysteine protease, LapG. When activated, LapG has been shown to cleave the surface adhesion LapA to release cells from the surface (78). In *P. aeruginosa*, *in vitro* batch and continuous-flow biofilm assays revealed that a *lapG* knockout mutant strain was unable to disperse in response to nutrient starvation, O₂ depletion, or NO donor signals compared to a wild type strain (unpublished data). This suggests that dispersal events in response to these various cues all integrate via a c-di-GMP signaling cascade that relies on LapG activation to induce biofilm detachment.

Second, the connecting biopolymers that maintain bacteria in the biofilm may be modulated by the secretion of chemicals that specifically alter adhesion appendages. Studies of *E. coli* suggested that dispersal bacteria are coated with a protein, called dispersin, that counteracts fimbriae-mediated aggregative adherence (79). Further, dispersal of *E. coli* biofilms from human epithelial cells was associated with alterations of bundle-forming type IV pili structures from thin to much longer and thicker bundles that resulted in loss of adherence and aggregation (80). To reduce surface tension and allow cells to detach, biofilm bacteria also produce amphipathic molecules, such as rhamnolipids, to induce dispersal (81). In *P. putida*, endogenous biosurfactants called putisolvins produced upon entry into stationary growth phase were able to induce biofilm detachment (82). In this scenario, the dispersal signal is amplified within an entire subpopulation of cells, where neighboring cells may degrade the matrix surrounding cells that may not have yet switched to a dispersal phenotype.

Finally, the transition from a biofilm to a planktonic mode of growth involves the activation of flagella (swimming and swarming motility) or pili (twitching motility) in motile microorganisms. Thus, flagella synthesis genes were found to be expressed in dispersed cells of *E. coli* and *P. aeruginosa* biofilms (26, 27, 38, 83), while type IV pili were found to be activated in biofilm bacteria treated with the dispersal signal NO (29). Further, the analysis of microscopy movies of *P. aeruginosa* biofilms revealed that attached bacteria use type IV pili to adopt a vertical orientation and mediate surface detachment (84).

Overall, a variety of strategies have been observed for detachment of bacteria from biofilms between different species but also in the same species. It is likely that the mechanisms used will depend on the growth conditions, including the conditional composition of the matrix, as well as the age of the biofilm.

DISPERSAL CELLS PRESENT A SPECIALIZED PHENOTYPE

Biofilm populations often display a high level of phenotypic heterogeneity (13), and not surprisingly, the dispersal subpopulation released from mature biofilms also reveals a high level of phenotypic variation. The variable phenotypes of dispersal cells are the result of a combination of changes in gene expression (transient phenotypic differences) and altered genotypes (permanent phenotypic changes). This diversity in biofilms and dispersal cells has been linked to increased fitness and

improved ability to colonize a range of habitats. Dispersal bacteria appear to be affected in key traits including attachment, metabolism, and antimicrobial resistance, as well as virulence and motility (30, 85–87).

Two studies of *P. aeruginosa* and *Streptococcus mutans* have compared stationary-phase planktonic cultures, resuspended biofilm cells, and bacteria collected from the biofilm effluent, and found differences in attachment properties and growth rates (30, 87). The entire dispersal population appeared to grow at rates similar to biofilm cells, both of which grew more slowly than planktonic cultures. While dispersal cells may have an overall slower metabolism, in *P. tunicata*, *P. aeruginosa*, and *S. mutans*, the dispersal populations have been found to utilize a wider range of nutrient sources (85–87). When variants isolated from biofilm effluents were analyzed separately, individual variants showed either decreased or increased metabolic rates compared to the parental strain (85–87). Further, dispersal cells in *P. aeruginosa* and *S. mutans* attached better and formed biofilms with greater biomass compared to both planktonic and resuspended biofilm cells (30, 87). Another study showed enhanced attachment and biofilm formation in dispersal cells of *Marinobacter hydrocarbonoclasticus* compared to planktonic cultures (88). This ability was linked to reshaping of the cell envelope and mobilization of storage reserves of cellular fatty acids and alcohol esters. Finally, increased resistance to antimicrobials and stress has been observed in biofilm-dispersed cells compared to planktonic cultures. Bacteria released from *S. mutans* biofilms had greater tolerance to chlorhexidine and acid treatments compared to stationary-phase planktonic cells (87). However, when a range of antibiotic treatments were tested by using a disc diffusion method, no difference between the dispersal and planktonic subpopulations was found in *P. aeruginosa* (30). When effluents from dual cultures of *P. aeruginosa* and *Burkholderia cepacia* biofilms were analyzed, the dispersal cells showed enhanced resistance to chlorine compared to planktonic chemostat cultures, and this resistance was proportional to the size of the detached clusters (89).

The physiological plasticity of dispersal bacteria is in part due to a high level of heterogeneity, resulting in the formation of phenotypic variants. Individual variants have been isolated from biofilm effluents with a frequency ranging from 5 to 60% of the dispersal subpopulation, which increased as the biofilm matured, in various strains including *P. aeruginosa*, especially from undomesticated clinical isolate biofilms (86, 90) and *S. marcescens* (91). In *P. tunicata*, a mutant unable to

undergo cell lysis and dispersal released fewer variants in their effluent, with an overall reduced ability to form biofilms, reduced motility, and homogeneous growth rate. In contrast, wild type biofilms showed extensive variation and overall increased values in all three aspects (85). Interestingly, when all variants collected from the dispersal effluent were grown in mixed cultures and tested for resistance to predation, the assemblage of variants showed increased resistance compared to monocultures grown from individual variants or the parental strain (92). The effect of self-generated genetic variation on increasing resistance of the mono-species biofilm population is also observed when genetic diversity is increased at the species level in mixed-community biofilms (93).

Dispersal cells appear to have a specialized phenotype, different from biofilm bacteria and also clearly distinct from cells grown in planktonic cultures (Fig. 3). The dispersal phenotype appears to show various degrees of stability resulting from phase variation to more permanent genetic alterations. Variants that were isolated from biofilm effluents, e.g., in *P. aeruginosa* (86) or *S. marcescens* (91), were found to be stable and maintain a distinct phenotype after subculturing. The physiological characterizations of entire dispersal populations have been performed immediately after detachment events, within hours following induction or seeding dispersal, and it is not entirely clear whether the observed specialized traits, such as antibiotic resistance (e.g., persisters) or metabolic activity may be stable or transient and whether reculturing of these cells would lead to a phenotype closer to planktonic cells.

Genotypic diversity has been observed in dispersal cells; for example, in *S. marcescens* biofilms, variants isolated from the biofilm effluent were found to result from a single mutation in a regulatory gene (92), which likely represents a favorable genetic switch as opposed to *de novo* mutation. Recently, whole-genome deep sequencing of biofilm dispersal populations revealed single nucleotide mutations in several hot spots, indicative of parallel evolution, but no extensive genetic variation could be observed (94). The latter observation is surprising given the high frequency of phenotypic variation observed.

Thus, it appears that the dispersal population harbors a high level of heterogeneity. However, there has been no investigation of whether this diversity may be directly linked to dispersal or its regulation or whether it was already found within the biofilm, although we can detect such variants within the biofilm biomass prior to dispersal (S.A. Rice et al., unpublished). The generation

FIGURE 3 Physiological traits of planktonic, biofilm, and dispersal cells. Symbols are defined in [Figure 2](#). [doi:10.1128/microbiolspec.MB-0015-2014.f3](https://doi.org/10.1128/microbiolspec.MB-0015-2014.f3)

of phenotypic diversity in dispersal bacteria has been linked to a range of triggers including oxidative stress (22, 95), activation of a superinfective filamentous phage that is also linked to reactive oxygen and nitrogen species activity (96), as well as anaerobic conditions (97). In *P. aeruginosa*, the mismatch repair system was found to generate genetic mutants after repair of oxidative stress-induced DNA damage (95). The emergence of stable phenotypic variants was also associated with the activation of a superinfective prophage by nitrosative stress (56, 96). In *P. tunicata*, an autolysis protein, AlpP, was shown to generate production of hydrogen peroxide, which led to phenotypic diversification and increased fitness (85). Indeed, deletion of the filamentous phage, the AlpP protein, and RecA-mediated recombination resulted in the loss of variant formation and an increase in sensitivity of the biofilms formed by the mutants when exposed to stressors.

ECOLOGICAL AND EVOLUTIONARY ASPECTS OF BIOFILM DISPERSAL

From an evolutionary perspective, biofilm dispersal represents an important adaptive strategy with profound impact on the survival and fitness of bacteria. The nature of dispersal, whether it benefits the biofilm population as a whole or only individual dispersal cells, is still not entirely clear. Because biofilms display a high level of cell differentiation and show close similarities to multicellular organisms (98), biofilm development may share

some similarities with the evolution and transition to multicellularity of higher organisms. In this context, dispersal has been considered an important step for the transition to multicellularity (99) and has been explained within the framework of each of the two main theories of multilevel (group) (100) and kin (individual) (101) selection. Group selection implies dispersal is an altruistic event that conserves the genetic inheritance of the population. In this scenario, diversity in the dispersal subpopulation should be restricted to reversible phenotypic switches and not derive from genetic mutations. In contrast, dispersal as an event selected at the individual level, kin selection, correlates with a high level of genetic variation within dispersal cells and thus a gradual change in genetic content or function relative to the parental population.

These questions of genetic evolution and dispersal in biofilms have important implications, for instance, in the context of the emergence of antibiotic resistance, a significant concern of our modern society. Biofilms play a major role in the resistance of bacteria toward antimicrobials and antibiotics, because bacteria in biofilms often display a high level of tolerance, via the generation of persister cells (102) or by accumulating resistant mutants via spontaneous mutations (103) as well as horizontal transfer of resistance genes (104). A direct link between increased genetic diversity specifically associated with a dispersal program rather than as a result of diversity that occurred within the biofilm has not been firmly established yet. In contrast, specific diversity in

the dispersal population appears to result from stable cell differentiation or phase variation and thus derive from a program encoded within bacterial genomes to maximize chances of colonization and survival.

The lungs of cystic fibrosis (CF) patients chronically infected with bacterial pathogens, which are mostly clonal in nature due to the apparently infrequent transmission between patients, offer a rare opportunity to observe evolutionary adaptation of bacterial populations. Genomic analyses of *P. aeruginosa* infections in CF patients over 8 years showed strong positive selection signals (ratio of nonsynonymous to synonymous changes per site, dN/dS >1) (105). Later analyses of bacterial infections over 35 years, representing 200,000 generations, revealed that after a period of adaptation *P. aeruginosa* stabilizes its phenotype in CF lungs and then applies a negative selection for mutations (106, 107). Thus, it is clear that clonal populations in a restricted environment undergo genetic evolution, which is either positively or negatively selected. While bacterial infections in CF patients' lungs are often associated with the biofilm mode of growth, and thus chronic infections are likely to go through dispersal events, the role that biofilm dispersal plays in generating genetic mutations over time remains unclear. Another study specifically analyzed genetic changes in *B. cepacia* biofilms through successive dispersal events *in vitro*. Beneficial changes occurred in the population, leading to mixed communities of stable variants that were more productive (colony-forming units number) than any monoculture (108).

Further, the mixed-variant biofilms obtained in these experiments resembled that of CF infections, suggesting that parallel evolution occurs *in vitro* and in CF lungs (109). In particular, evolved *B. cepacia* populations harbored mutations in the *wsp* operon regulating c-di-GMP signaling, suggesting that the mutants were affected in their ability to form biofilms and disperse (109). This supports a close association between the evolution of strains in the CF lungs and biofilm dispersal events. Overall, it is likely that dispersal plays a critical role in evolutionary processes, because it allows spreading of any new phenotypic trait to colonize new habitats where competition with the parental population is limited. Interestingly, the importance of biofilm dispersal in evolutionary processes may be observed in the apparent evolutionary conservation of the regulatory pathways that mediate dispersal. For example, multiple components of the NO-mediated dispersal signaling pathways from biofilms, NO/c-di-GMP, are also conserved in higher multicellular organisms, where the NO/cyclic GMP system regulates diverse physiological functions in

mammals, regulating vasodilation, platelet aggregation, and sensory systems, as well as in plants, regulating development and pathogen defense responses (110). These similarities suggest an intriguing relationship between biofilm dispersal signaling and the evolution of eukaryotic regulatory pathways (29).

Intriguingly, bacteriophages have been associated with phenotypic diversity in the dispersal population. In particular, in *P. aeruginosa* a filamentous phage, Pf4, which is present as a prophage, is highly expressed during biofilm development compared to planktonic cells (8). The activity of the Pf4 phage was linked to killing and lysis of a subpopulation of cells within biofilms (16) and the emergence of phenotypic variants (96). Addition of the superinfective phage to planktonic cultures resulted in the formation of morphotypic variants, which were not observed in uninfected control cultures (111). Further, the Pf4 phage was found to increase fitness and virulence in a murine infection environment (111). However, it is not yet understood whether the Pf4 phage superinfective switch is coregulated with dispersal pathways. In *P. aeruginosa*, modulating c-di-GMP levels had no effect on phage superinfection, and both the wild type and a prophage mutant dispersed equally upon NO addition, suggesting that the NO dispersal pathway is not dependent on the Pf4 phage (unpublished data). A lytic RNA-containing bacteriophage, PP7, has also been implicated in regulating phenotypic variation in *P. aeruginosa*, leading to diversification and the evolution of a small-rough colony phenotype (112).

Dispersal is thought to be beneficial to biofilms for several reasons and, in particular, allows microbial populations to spread and survive. Arguably, dispersal may be considered the most important phase of biofilm development. Mathematical models have suggested that dispersal mostly benefits a population in an open space, while it can lead to its collapse in restricted environments (5). Dispersal is beneficial when a population faces catastrophic events that may eradicate a colony at a particular location, while leaving the location habitable for new organisms.

The lack of detailed studies of biofilm dispersal in natural systems leaves a number of questions unanswered. Can biofilm dispersal be detrimental as the result of utilizing crucial resources while increasing competition in a local environment and thus ultimately damaging the parental biofilm population? Do biofilms adopt a strategy of mass dispersal in harsh environments, risking extinction by misallocating energy resources to dispersal cells that lose the inherent protection of the biofilm? In the context of infection, a

number of reports revealed that dispersal is a critical strategy for pathogens to successfully colonize a host. A functioning dispersal response appears to be crucial to ensure transmission of disease in both animals (113, 114) and plants, for instance, transmission of the plant pathogen *X. campestris* on the Chinese radish (50). A recent study found that *P. aeruginosa* mutant strains impaired in their ability to disperse from biofilms had reduced virulence and reduced ability to cause persistent infections in both murine and plant *in vivo* infection models (115). These beneficial effects of dispersal in a context of infection may be due to the association of dispersal signaling pathways with activation of virulence in some pathogens, e.g., via a decrease in c-di-GMP levels leading to expression of an endo- β -1,4-mannanase in *X. campestris* (50) and activation of type III secretion in *P. aeruginosa* (116).

Further studies will be required to fully grasp the role of dispersal in complex ecological systems involving multispecies biofilms in highly heterogeneous habitats.

Metagenomics and molecular analysis-based studies appear highly suitable to reveal complex patterns of diversity in microbial communities but require precise temporally and geographically linked sampling points to track and analyze biofilm dispersal processes in natural environments (117). While not biofilm specific, some elegant model studies have been performed to determine the impact of dispersal events on microbial community composition and changes, which showed that environmental filtering, functional plasticity, and competition are all important mechanisms influencing the fate of dispersed communities (118).

OPPORTUNITIES FOR BIOFILM CONTROL

Manipulation of the biofilm development program by inducing dispersal signals has emerged as a strategy for developing novel biofilm control measures in recent years (Fig. 4). One of the first strategies was to treat biofilms with EPS-degrading agents to disperse attached

FIGURE 4 A range of strategies targeting dispersal have been developed to control biofilms and biofilm-related infections. (a) BdcA protein with enhanced c-di-GMP binding (124). (b) Oroidin and synthetic derivatives that were identified as potent dispersal inducers after screening chemical libraries (125). (c) Diffusible fatty acid signal cis-2-decenoic acid (23). (d) Furanone 30, a synthetic QS inhibitor derived from natural furanone compounds isolated from the red alga *Delisea pulchra* (131). (e) Fimbrilide-nitroester with dual action QS inhibition and NO release (134). (f) β -lactam-NO prodrugs for the targeted delivery of NO to infectious biofilms (132). (g) Controlled delivery of NO using nanoparticles (135). (h) Catalytic generation of NO from endogenous nitrite sources to disperse and prevent biofilm for long-term applications (138). doi:10.1128/microbiolspec.MB-0015-2014.f4

cells. These treatments were later combined with antibiotics to eradicate the dispersed cells, for instance dispersinB in combination with the broad-spectrum biocide triclosan (119), or alginate lyase together with the aminoglycoside tobramycin (120). Given the diversity in the components of biofilm EPS matrices (121), enzyme-based treatments are likely to be biofilm specific, which can be beneficial when designing narrow-spectrum treatments.

The recent discoveries of the central role of the secondary messenger c-di-GMP in regulating the transition between biofilm and free-swimming cells across many bacterial species attracted strong interest. Enzymes involved in c-di-GMP turnover appear very potent for controlling the switch, and the global signaling network has become a primary objective for developing novel antibiofilm measures. The first proof of concept studies of targeting c-di-GMP to treat infections showed that *in vivo* manipulation of c-di-GMP levels can effectively clear (by decreasing c-di-GMP) or prolong (by increasing c-di-GMP) *P. aeruginosa* infections in murine models (122, 123). Further, in an effort to develop a potential therapeutic drug, a BdcA protein was modified to enhance its c-di-GMP binding, thus reducing the intracellular c-di-GMP concentration. When added to biofilms *in vitro*, the modified BdcA protein caused nearly complete dispersal (124).

Improved high-throughput assays for biofilm dispersal allowed for the screening of large numbers of synthetic compounds and are particularly amenable to screening chemical libraries. Using this approach, 2-aminoimidazole derivatives targeting QS were designed and found to disperse established biofilms (125, 126). Studies of aerobic granules, a type of suspended biofilms used for wastewater treatment, revealed that the fatty acid 2-decenoic acid, which was previously identified to regulate dispersal in *P. aeruginosa* and a range of single-species biofilms, can also trigger dispersal in these mixed-species biofilms (127). In this context, strategies to inhibit 2-decenoic acid signaling may be useful to maintain granule integrity and sludge performance (127). QS signals and inhibitors have also been examined for their potential to enhance performance outputs in wastewater treatment biological reactors (128, 129), as well as for the treatment of biofilm-related infections and bacterial virulence (130, 131).

Of particular interest is NO, a simple and versatile dispersal signal that is highly conserved across biofilm species and activates c-di-GMP signaling to induce dispersal. Much progress has been made in recent years to design efficient NO delivery strategies, making it an

outstanding candidate for novel therapeutic strategies. A range of NO-based technologies have been developed that offer a versatile range of solutions to control biofilms (reviewed in 57), which include a new class of β -lactam-NO prodrugs for the targeted delivery of NO to infectious biofilms (132, 133), dual quorum-sensing antagonists–NO releasing compounds (134), and novel polymers for sustained delivery of NO to prevent biofilm formation in a nontoxic fashion (135). NO-based treatments have been applied to the treatment of fouling on water filtration membranes (136, 137). A promising approach for long-term prevention and dispersal of biofilms in industrial and clinical settings is the use of copper-based catalytic technology to reduce endogenous nitrite ions to continuously produce NO at the surface (138). Recently, the first clinical trial was conducted to evaluate the use of low-dose inhaled NO gas combined with standard antibiotic therapy for the disruption of *P. aeruginosa* biofilms in patients with CF (139). The results demonstrated that patients who received NO gas at 5–10 ppm (~200 nM NO) showed significant reductions in the number of *Pseudomonas* biofilm aggregates compared to patients who received a placebo. These data suggest that using NO as an adjunctive therapy may be highly beneficial for the treatment of CF-related biofilm infections.

CONCLUDING REMARKS

Advances in biofilm and molecular biology studies have led to the elucidation of key signaling molecules and genes that govern biofilm development and dispersal processes. The existence of a developmental program appears clearer, but the challenge remains to address how these molecules and genes work together as control systems to generate patterns in space and time in biofilms. The biofilm mode of growth represents the predominant lifestyle for bacteria in environmental, industrial, and medical settings. A precise understanding of biofilm development and dispersal is crucial to better control microbial communities, with profound implications for global health, and to overcome the threat of antibiotic resistance as well as to develop innovative solutions for improved management of environmental microbes.

ACKNOWLEDGMENTS

Conflicts of interest: We disclose no conflicts.

REFERENCES

1. Ronce O. 2007. How does it feel to be like a rolling stone? Ten questions about dispersal evolution. *Annu Rev Ecol Evol Syst* 38:231–253.

2. Nadell CD, Bassler BL. 2011. A fitness trade-off between local competition and dispersal in *Vibrio cholerae* biofilms. *Proc Natl Acad Sci USA* 108:14181–14185.
3. Kisdi E. 2002. Dispersal: risk spreading versus local adaptation. *Am Nat* 159:579–596.
4. Kerr B, Neuhauser C, Bohannan BJ, Dean AM. 2006. Local migration promotes competitive restraint in a host-pathogen ‘tragedy of the commons.’ *Nature* 442:75–78.
5. Gyllenberg M, Parvonen K, Dieckmann U. 2002. Evolutionary suicide and evolution of dispersal in structured metapopulations. *J Math Biol* 45:79–105.
6. Petrova OE, Sauer K. 2009. A novel signaling network essential for regulating *Pseudomonas aeruginosa* biofilm development. *PLoS Pathog* 5:e1000668. doi:10.1371/journal.ppat.1000668.
7. Petrova OE, Sauer K. 2012. Sticky situations: key components that control bacterial surface attachment. *J Bacteriol* 194:2413–2425.
8. Whiteley M, Bangera MG, Bumgarner RE, Parsek MR, Teitzel GM, Lory S, Greenberg EP. 2001. Gene expression in *Pseudomonas aeruginosa* biofilms. *Nature* 413:860–864.
9. Klausen M, Aaes-Jorgensen A, Molin S, Tolker-Nielsen T. 2003. Involvement of bacterial migration in the development of complex multicellular structures in *Pseudomonas aeruginosa* biofilms. *Mol Microbiol* 50:61–68.
10. Yang L, Hu Y, Liu Y, Zhang J, Ulstrup J, Molin S. 2011. Distinct roles of extracellular polymeric substances in *Pseudomonas aeruginosa* biofilm development. *Environ Microbiol* 13:1705–1717.
11. Irie Y, Borlee BR, O’Connor JR, Hill PJ, Harwood CS, Wozniak DJ, Parsek MR. 2012. Self-produced exopolysaccharide is a signal that stimulates biofilm formation in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 109:20632–20636.
12. Lenz AP, Williamson KS, Pitts B, Stewart PS, Franklin MJ. 2008. Localized gene expression in *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol* 74:4463–4471.
13. Stewart PS, Franklin MJ. 2008. Physiological heterogeneity in biofilms. *Nat Rev Microbiol* 6:199–210.
14. Williamson KS, Richards LA, Perez-Osorio AC, Pitts B, McInerney K, Stewart PS, Franklin MJ. 2012. Heterogeneity in *Pseudomonas aeruginosa* biofilms includes expression of ribosome hibernation factors in the antibiotic-tolerant subpopulation and hypoxia-induced stress response in the metabolically active population. *J Bacteriol* 194:2062–2073.
15. McDougald D, Rice SA, Barraud N, Steinberg PD, Kjelleberg S. 2012. Should we stay or should we go: mechanisms and ecological consequences for biofilm dispersal. *Nat Rev Microbiol* 10:39–50.
16. Webb JS, Thompson LS, James S, Charlton T, Tolker-Nielsen T, Koch B, Givskov M, Kjelleberg S. 2003. Cell death in *Pseudomonas aeruginosa* biofilm development. *J Bacteriol* 185:4585–4592.
17. Tolker-Nielsen T, Brinch UC, Ragas PC, Andersen JB, Jacobsen CS, Molin S. 2000. Development and dynamics of *Pseudomonas* sp. biofilms. *J Bacteriol* 182:6482–6489.
18. Entcheva-Dimitrov P, Spormann AM. 2004. Dynamics and control of biofilms of the oligotrophic bacterium *Caulobacter crescentus*. *J Bacteriol* 186:8254–8266.
19. Lawrence JR, Chenier MR, Roy R, Beaumier D, Fortin N, Swerhone GD, Neu TR, Greer CW. 2004. Microscale and molecular assessment of impacts of nickel, nutrients, and oxygen level on structure and function of river biofilm communities. *Appl Environ Microbiol* 70:4326–4339.
20. Manteca A, Fernandez M, Sanchez J. 2005. A death round affecting a young compartmentalized mycelium precedes aerial mycelium dismantling in confluent surface cultures of *Streptomyces antibioticus*. *Microbiology* 151:3689–3697.
21. Bayles KW. 2007. The biological role of death and lysis in biofilm development. *Nat Rev Microbiol* 5:721–726.
22. Mai-Prochnow A, Lucas-Elío P, Egan S, Thomas T, Webb JS, Sanchez-Amat A, Kjelleberg S. 2008. Hydrogen peroxide linked to lysine oxidase activity facilitates biofilm differentiation and dispersal in several Gram-negative bacteria. *J Bacteriol* 190:5493–5501.
23. Davies DG, Marques CN. 2009. A fatty acid messenger is responsible for inducing dispersion in microbial biofilms. *J Bacteriol* 191:1393–1403.
24. Mai-Prochnow A, Evans F, Dalisay-Saludes D, Stelzer S, Egan S, James S, Webb JS, Kjelleberg S. 2004. Biofilm development and cell death in the marine bacterium *Pseudoalteromonas tunicata*. *Appl Environ Microbiol* 70:3232–3238.
25. Huynh TT, McDougald D, Klebensberger J, Al Qarni B, Barraud N, Rice SA, Kjelleberg S, Schleheck D. 2012. Glucose starvation-induced dispersal of *Pseudomonas aeruginosa* biofilms is cAMP and energy dependent. *PLoS One* 7:e42874. doi:10.1371/journal.pone.0042874.
26. Purevdorj-Gage B, Costerton WJ, Stoodley P. 2005. Phenotypic differentiation and seeding dispersal in non-mucoid and mucoid *Pseudomonas aeruginosa* biofilms. *Microbiology* 151:1569–1576.
27. Sauer K, Camper AK, Ehrlich GD, Costerton JW, Davies DG. 2002. *Pseudomonas aeruginosa* displays multiple phenotypes during development as a biofilm. *J Bacteriol* 184:1140–1154.
28. Morgan R, Kohn S, Hwang SH, Hassett DJ, Sauer K. 2006. BdlA, a chemotaxis regulator essential for biofilm dispersal in *Pseudomonas aeruginosa*. *J Bacteriol* 188:7335–7343.
29. Barraud N, Schleheck D, Klebensberger J, Webb JS, Hassett DJ, Rice SA, Kjelleberg S. 2009. Nitric oxide signaling in *Pseudomonas aeruginosa* biofilms mediates phosphodiesterase activity, decreased cyclic di-GMP levels, and enhanced dispersal. *J Bacteriol* 191:7333–7342.
30. Rollet C, Gal L, Guzzo J. 2009. Biofilm-detached cells, a transition from a sessile to a planktonic phenotype: a comparative study of adhesion and physiological characteristics in *Pseudomonas aeruginosa*. *FEMS Microbiol Lett* 290:135–142.
31. Gjermansen M, Ragas P, Sternberg C, Molin S, Tolker-Nielsen T. 2005. Characterization of starvation-induced dispersion in *Pseudomonas putida* biofilms. *Environ Microbiol* 7:894–906.
32. Schleheck D, Barraud N, Klebensberger J, Webb JS, McDougald D, Rice SA, Kjelleberg S. 2009. *Pseudomonas aeruginosa* PAO1 preferentially grows as aggregates in liquid batch cultures and disperses upon starvation. *PLoS One* 4:e5513. doi:10.1371/journal.pone.0005513.
33. Newell PD, Boyd CD, Sondermann H, O’Toole GA. 2011. A c-di-GMP effector system controls cell adhesion by inside-out signaling and surface protein cleavage. *PLoS Biol* 9:e1000587. doi:10.1371/journal.pbio.1000587.
34. Banin E, Brady KM, Greenberg EP. 2006. Chelator-induced dispersal and killing of *Pseudomonas aeruginosa* cells in a biofilm. *Appl Environ Microbiol* 72:2064–2069.
35. Thormann KM, Saville RM, Shukla S, Spormann AM. 2005. Induction of rapid detachment in *Shewanella oneidensis* MR-1 biofilms. *J Bacteriol* 187:1014–1021.
36. An S, Wu J, Zhang LH. 2010. Modulation of *Pseudomonas aeruginosa* biofilm dispersal by a cyclic-di-GMP phosphodiesterase with a putative hypoxia-sensing domain. *Appl Environ Microbiol* 76:8160–8173.
37. Saville RM, Rakshe S, Haagensen JA, Shukla S, Spormann AM. 2011. Energy-dependent stability of *Shewanella oneidensis* MR-1 biofilms. *J Bacteriol* 193:3257–3264.
38. Sauer K, Cullen MC, Rickard AH, Zeef LA, Davies DG, Gilbert P. 2004. Characterization of nutrient-induced dispersion in *Pseudomonas aeruginosa* PAO1 biofilm. *J Bacteriol* 186:7312–7326.
39. Musk DJ, Banko DA, Hergenrother PJ. 2005. Iron salts perturb biofilm formation and disrupt existing biofilms of *Pseudomonas aeruginosa*. *Chem Biol* 12:789–796.

40. Chatterjee S, Wistrom C, Lindow SE. 2008. A cell-cell signaling sensor is required for virulence and insect transmission of *Xylella fastidiosa*. *Proc Natl Acad Sci USA* 105:2670–2675.
41. Hagai E, Dvora R, Havkin-Blank T, Zelinger E, Porat Z, Schulz S, Helman Y. 2014. Surface-motility induction, attraction and hitchhiking between bacterial species promote dispersal on solid surfaces. *ISME J* 8:1147–1151.
42. Solano C, Echeverez M, Lasa I. 2014. Biofilm dispersion and quorum sensing. *Curr Opin Microbiol* 18c:96–104.
43. Zhu J, Mekalanos JJ. 2003. Quorum sensing-dependent biofilms enhance colonization in *Vibrio cholerae*. *Dev Cell* 5:647–656.
44. Kim SM, Park JH, Lee HS, Kim WB, Ryu JM, Han HJ, Choi SH. 2013. LuxR homologue SmcR is essential for *Vibrio vulnificus* pathogenesis and biofilm detachment, and its expression is induced by host cells. *Infect Immun* 81:3721–3730.
45. Rice SA, Koh KS, Queck SY, Labbate M, Lam KW, Kjelleberg S. 2005. Biofilm formation and sloughing in *Serratia marcescens* are controlled by quorum sensing and nutrient cues. *J Bacteriol* 187:3477–3485.
46. Puskas A, Greenberg EP, Kaplan S, Schaefer AL. 1997. A quorum-sensing system in the free-living photosynthetic bacterium *Rhodobacter sphaeroides*. *J Bacteriol* 179:7530–7537.
47. Boles BR, Horswill AR. 2008. *agr*-Mediated dispersal of *Staphylococcus aureus* biofilms. *PLoS Pathog* 4:e1000052. doi:10.1371/journal.ppat.1000052.
48. Hammer BK, Bassler BL. 2003. Quorum sensing controls biofilm formation in *Vibrio cholerae*. *Mol Microbiol* 50:101–104.
49. Bari SM, Roky MK, Mohiuddin M, Kamruzzaman M, Mekalanos JJ, Faruque SM. 2013. Quorum-sensing autoinducers resuscitate dormant *Vibrio cholerae* in environmental water samples. *Proc Natl Acad Sci USA* 110:9926–9931.
50. Dow JM, Crossman L, Findlay K, He YQ, Feng JX, Tang JL. 2003. Biofilm dispersal in *Xanthomonas campestris* is controlled by cell-cell signaling and is required for full virulence to plants. *Proc Natl Acad Sci USA* 100:10995–11000.
51. Deng Y, Wu J, Tao F, Zhang LH. 2011. Listening to a new language: DSF-based quorum sensing in Gram-negative bacteria. *Chem Rev* 111:160–173.
52. Hentzer M, Riedel K, Rasmussen TB, Heydorn A, Andersen JB, Parsek MR, Rice SA, Eberl L, Molin S, Hoiby N, Kjelleberg S, Givskov M. 2002. Inhibition of quorum sensing in *Pseudomonas aeruginosa* biofilm bacteria by a halogenated furanone compound. *Microbiology* 148:87–102.
53. Lonn-Stensrud J, Landin MA, Benneche T, Petersen FC, Scheie AA. 2009. Furanones, potential agents for preventing *Staphylococcus epidermidis* biofilm infections? *J Antimicrob Chemother* 63:309–316.
54. Kolodkin-Gal I, Romero D, Cao S, Clardy J, Kolter R, Losick R. 2010. D-Amino acids trigger biofilm disassembly. *Science* 328:627–629.
55. Leiman SA, May JM, Lebar MD, Kahne D, Kolter R, Losick R. 2013. D-Amino acids indirectly inhibit biofilm formation in *Bacillus subtilis* by interfering with protein synthesis. *J Bacteriol* 195:5391–5395.
56. Barraud N, Hassett DJ, Hwang SH, Rice SA, Kjelleberg S, Webb JS. 2006. Involvement of nitric oxide in biofilm dispersal of *Pseudomonas aeruginosa*. *J Bacteriol* 188:7344–7353.
57. Barraud N, Kelso MJ, Rice SA, Kjelleberg S. 2014. Nitric oxide: a key mediator of biofilm dispersal with applications in infectious diseases. *Curr Pharm Design* 21:31–42.
58. Sondermann H, Shikuma NJ, Yildiz FH. 2012. You've come a long way: c-di-GMP signaling. *Curr Opin Microbiol* 15:140–146.
59. Römling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a universal bacterial second messenger. *Microbiol Mol Biol Rev* 77:1–52.
60. Kalivoda EJ, Brothers KM, Stella NA, Schmitt MJ, Shanks RM. 2013. Bacterial cyclic AMP-phosphodiesterase activity coordinates biofilm formation. *PLoS One* 8:e71267. doi:10.1371/journal.pone.0071267.
61. Galperin MY, Higdon R, Kolker E. 2010. Interplay of heritage and habitat in the distribution of bacterial signal transduction systems. *Mol Biosyst* 6:721–728.
62. Gomelsky M. 2011. cAMP, c-di-GMP, c-di-AMP and now cGMP: bacteria use them all! *Mol Microbiol* 79:562–565.
63. Liu N, Xu Y, Hossain S, Huang N, Coursolle D, Gralnick JA, Boon EM. 2012. Nitric oxide regulation of cyclic di-GMP synthesis and hydrolysis in *Shewanella woodyi*. *Biochemistry* 51:2087–2099.
64. Carlson HK, Vance RE, Marletta MA. 2010. H-NOX regulation of c-di-GMP metabolism and biofilm formation in *Legionella pneumophila*. *Mol Microbiol* 77:930–942.
65. Roy AB, Petrova OE, Sauer K. 2012. The phosphodiesterase DipA (PA5017) is essential for *Pseudomonas aeruginosa* biofilm dispersion. *J Bacteriol* 194:2904–2915.
66. Li Y, Heine S, Entian M, Sauer K, Frankenberg-Dinkel N. 2013. NO-induced biofilm dispersion in *Pseudomonas aeruginosa* is mediated by an MHYT domain-coupled phosphodiesterase. *J Bacteriol* 195:3531–3542.
67. Nioche P, Berka V, Vipond J, Minton N, Tsai AL, Raman CS. 2004. Femtomolar sensitivity of a NO sensor from *Clostridium botulinum*. *Science* 306:1550–1553.
68. Plate L, Marletta MA. 2013. Nitric oxide-sensing H-NOX proteins govern bacterial communal behavior. *Trends Biochem Sci* 38:566–575.
69. Tanaka A, Takahashi H, Shimizu T. 2007. Critical role of the heme axial ligand, Met⁹⁵, in locking catalysis of the phosphodiesterase from *Escherichia coli* (Ec DOS) toward cyclic diGMP. *J Biol Chem* 282:21301–21307.
70. Tuckerman JR, Gonzalez G, Sousa EH, Wan X, Saito JA, Alam M, Gilles-Gonzalez MA. 2009. An oxygen-sensing diguanylate cyclase and phosphodiesterase couple for c-di-GMP control. *Biochemistry* 48:9764–9774.
71. Petrova OE, Sauer K. 2012. Dispersion by *Pseudomonas aeruginosa* requires an unusual posttranslational modification of BdlA. *Proc Natl Acad Sci USA* 109:16690–16695.
72. Kaplan JB, Meyenhofer MF, Fine DH. 2003. Biofilm growth and detachment of *Actinobacillus actinomycetemcomitans*. *J Bacteriol* 185:1399–1404.
73. Kaplan JB, Raganath C, Velliagounder K, Fine DH, Ramasubbu N. 2004. Enzymatic detachment of *Staphylococcus epidermidis* biofilms. *Antimicrob Agents Chemother* 48:2633–2636.
74. Boyd A, Chakrabarty AM. 1994. Role of alginate lyase in cell detachment of *Pseudomonas aeruginosa*. *Appl Environ Microbiol* 60:2355–2359.
75. Whitchurch CB, Tolker-Nielsen T, Ragas PC, Mattick JS. 2002. Extracellular DNA required for bacterial biofilm formation. *Science* 295:1487.
76. Mann EE, Rice KC, Boles BR, Endres JL, Ranjit D, Chandramohan L, Tsang LH, Smeltzer MS, Horswill AR, Bayles KW. 2009. Modulation of eDNA release and degradation affects *Staphylococcus aureus* biofilm maturation. *PLoS One* 4:e5822. doi:10.1371/journal.pone.0005822.
77. Nijland R, Hall MJ, Burgess JG. 2010. Dispersal of biofilms by secreted, matrix degrading, bacterial DNase. *PLoS One* 5:e15668. doi:10.1371/journal.pone.0015668.
78. Gjermansen M, Nilsson M, Yang L, Tolker-Nielsen T. 2010. Characterization of starvation-induced dispersion in *Pseudomonas putida* biofilms: genetic elements and molecular mechanisms. *Mol Microbiol* 75:815–826.
79. Sheikh J, Czczulin JR, Harrington S, Hicks S, Henderson IR, Le Bouguenec C, Gounon P, Phillips A, Nataro JP. 2002. A novel dispersin protein in enteroaggregative *Escherichia coli*. *J Clin Invest* 110:1329–1337.
80. Knutton S, Shaw RK, Anantha RP, Donnenberg MS, Zorngani AA. 1999. The type IV bundle-forming pilus of enteropathogenic *Escherichia coli* undergoes dramatic alterations in structure associated with bacterial adherence, aggregation and dispersal. *Mol Microbiol* 33:499–509.

81. Boles BR, Thoendel M, Singh PK. 2005. Rhamnolipids mediate detachment of *Pseudomonas aeruginosa* from biofilms. *Mol Microbiol* 57:1210–1223.
82. Kuiper I, Lagendijk EL, Pickford R, Derrick JP, Lamers GEM, Thomas-Oates JE, Lugtenberg BJJ, Bloemberg GV. 2004. Characterization of two *Pseudomonas putida* lipopeptide biosurfactants, putisolvin I and II, which inhibit biofilm formation and break down existing biofilms. *Mol Microbiol* 51:97–113.
83. Jackson DW, Suzuki K, Oakford L, Simecka JW, Hart ME, Romeo T. 2002. Biofilm formation and dispersal under the influence of the global regulator CsrA of *Escherichia coli*. *J Bacteriol* 184:290–301.
84. Gibiansky ML, Conrad JC, Jin F, Gordon VD, Motto DA, Mathewson MA, Stopka WG, Zelasko DC, Shrout JD, Wong GC. 2010. Bacteria use type IV pili to walk upright and detach from surfaces. *Science* 330:197.
85. Mai-Prochnow A, Ferrari BC, Webb JS, Kjelleberg S. 2006. Ecological advantages of autolysis during the development and dispersal of *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol* 72:5414–5420.
86. Woo JK, Webb JS, Kirov SM, Kjelleberg S, Rice SA. 2012. Biofilm dispersal cells of a cystic fibrosis *Pseudomonas aeruginosa* isolate exhibit variability in functional traits likely to contribute to persistent infection. *FEMS Immunol Med Microbiol* 66:251–264.
87. Liu J, Ling JQ, Zhang K, Wu CD. 2013. Physiological properties of *Streptococcus mutans* UA159 biofilm-detached cells. *FEMS Microbiol Lett* 340:11–18.
88. Vaysse PJ, Sivadon P, Goulas P, Grimaud R. 2011. Cells dispersed from *Marinobacter hydrocarbonoclasticus* SP17 biofilm exhibit a specific protein profile associated with a higher ability to reinitiate biofilm development at the hexadecane-water interface. *Environ Microbiol* 13:737–746.
89. Behnke S, Parker AE, Woodall D, Camper AK. 2011. Comparing the chlorine disinfection of detached biofilm clusters with those of sessile biofilms and planktonic cells in single- and dual-species cultures. *Appl Environ Microbiol* 77:7176–7184.
90. Kirov SM, Webb JS, O'May CY, Reid DW, Woo JK, Rice SA, Kjelleberg S. 2007. Biofilm differentiation and dispersal in mucoid *Pseudomonas aeruginosa* isolates from patients with cystic fibrosis. *Microbiology* 153:3264–3274.
91. Koh KS, Lam KW, Alhede M, Queck SY, Labbate M, Kjelleberg S, Rice SA. 2007. Phenotypic diversification and adaptation of *Serratia marcescens* MG1 biofilm derived morphotypes. *J Bacteriol* 189:119–130.
92. Koh KS, Matz C, Tan CH, Le HL, Rice SA, Marshall DJ, Steinberg PD, Kjelleberg S. 2012. Minimal increase in genetic diversity enhances predation resistance. *Mol Ecol* 21:1741–1753.
93. Lee KW, Periasamy S, Mukherjee M, Xie C, Kjelleberg S, Rice SA. 2014. Biofilm development and enhanced stress resistance of a model, mixed-species community biofilm. *ISME J* 8:894–907.
94. McElroy KE, Hui JG, Woo JK, Luk AW, Webb JS, Kjelleberg S, Rice SA, Thomas T. 2014. Strain-specific parallel evolution drives short-term diversification during *Pseudomonas aeruginosa* biofilm formation. *Proc Natl Acad Sci USA* 111:E1419–E1427.
95. Boles BR, Singh PK. 2008. Endogenous oxidative stress produces diversity and adaptability in biofilm communities. *Proc Natl Acad Sci USA* 105:12503–12508.
96. Webb JS, Lau M, Kjelleberg S. 2004. Bacteriophage and phenotypic variation in *Pseudomonas aeruginosa* biofilm development. *J Bacteriol* 186:8066–8073.
97. Fang H, Toyofuku M, Kiyokawa T, Ichihashi A, Tateda K, Nomura N. 2013. The impact of anaerobiosis on strain-dependent phenotypic variations in *Pseudomonas aeruginosa*. *Biosci Biotechnol Biochem* 77:1747–1752.
98. Webb JS, Givskov M, Kjelleberg S. 2003. Bacterial biofilms: prokaryotic adventures in multicellularity. *Curr Opin Microbiol* 6:578–585.
99. Hochberg ME, Rankin DJ, Taborsky M. 2008. The coevolution of cooperation and dispersal in social groups and its implications for the emergence of multicellularity. *BMC Evol Biol* 8:238.
100. Boots M, Meador M. 2007. Local interactions select for lower pathogen infectivity. *Science* 315:1284–1286.
101. Wild G, Gardner A, West SA. 2009. Adaptation and the evolution of parasite virulence in a connected world. *Nature* 459:983–986.
102. Barraud N, Buson A, Jarolimek W, Rice SA. 2013. Mannitol enhances antibiotic sensitivity of persister bacteria in *Pseudomonas aeruginosa* biofilms. *PLoS One* 8:e84220. doi:10.1371/journal.pone.0084220.
103. Hoiby N, Bjarnsholt T, Givskov M, Molin S, Ciofu O. 2010. Antibiotic resistance of bacterial biofilms. *Int J Antimicrob Agents* 35:322–332.
104. Roberts AP, Mullany P. 2010. Oral biofilms: a reservoir of transferable, bacterial, antimicrobial resistance. *Expert Rev Anti-Infect Ther* 8:1441–1450.
105. Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller SI, Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV. 2006. Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. *Proc Natl Acad Sci USA* 103:8487–8492.
106. Yang L, Jelsbak L, Marvig RL, Damkiaer S, Workman CT, Rau MH, Hansen SK, Folkesson A, Johansen HK, Ciofu O, Hoiby N, Sommer MO, Molin S. 2011. Evolutionary dynamics of bacteria in a human host environment. *Proc Natl Acad Sci USA* 108:7481–7486.
107. Folkesson A, Jelsbak L, Yang L, Johansen HK, Ciofu O, Hoiby N, Molin S. 2012. Adaptation of *Pseudomonas aeruginosa* to the cystic fibrosis airway: an evolutionary perspective. *Nat Rev Microbiol* 10:841–851.
108. Poltak SR, Cooper VS. 2011. Ecological succession in long-term experimentally evolved biofilms produces synergistic communities. *ISME J* 5:369–378.
109. Traverse CC, Mayo-Smith LM, Poltak SR, Cooper VS. 2013. Tangled bank of experimentally evolved *Burkholderia* biofilms reflects selection during chronic infections. *Proc Natl Acad Sci USA* 110:E250–E259.
110. Wendehenne D, Pugin A, Klessig DF, Durner J. 2001. Nitric oxide: comparative synthesis and signaling in animal and plant cells. *Trends Plant Sci* 6:177–183.
111. Rice SA, Tan CH, Mikkelsen PJ, Kung V, Woo J, Tay M, Hauser A, McDougald D, Webb JS, Kjelleberg S. 2009. The biofilm life cycle and virulence of *Pseudomonas aeruginosa* are dependent on a filamentous prophage. *ISME J* 3:271–282.
112. Brockhurst MA, Buckling A, Rainey PB. 2005. The effect of a bacteriophage on diversification of the opportunistic bacterial pathogen, *Pseudomonas aeruginosa*. *Proc Biol Sci* 272:1385–1391.
113. Hall-Stoodley L, Stoodley P. 2005. Biofilm formation and dispersal and the transmission of human pathogens. *Trends Microbiol* 13:7–10.
114. Chamot-Rooke J, Mikaty G, Malosse C, Soyer M, Dumont A, Gault J, Imhaus AF, Martin P, Trellet M, Clary G, Chafey P, Camoin L, Nilges M, Nassif X, Dumenil G. 2011. Posttranslational modification of pili upon cell contact triggers *N. meningitidis* dissemination. *Science* 331:778–782.
115. Li Y, Petrova OE, Su S, Lau GW, Panmanee W, Na R, Hassett DJ, Davies DG, Sauer K. 2014. BdlA, DipA and induced dispersion contribute to acute virulence and chronic persistence of *Pseudomonas aeruginosa*. *PLoS Pathog* 10:e1004168. doi:10.1371/journal.ppat.1004168.
116. Moscose JA, Mikkelsen H, Heeb S, Williams P, Filloux A. 2011. The *Pseudomonas aeruginosa* sensor RetS switches type III and type VI secretion via c-di-GMP signalling. *Environ Microbiol* 13:3128–3138.
117. Besemer K, Singer G, Hodl I, Battin TJ. 2009. Bacterial community composition of stream biofilms in spatially variable-flow environments. *Appl Environ Microbiol* 75:7189–7195.

118. Székely AJ, Berga M, Langenheder S. 2013. Mechanisms determining the fate of dispersed bacterial communities in new environments. *ISME J* 7:61–71.
119. Darouiche RO, Mansouri MD, Gawande PV, Madhyastha S. 2009. Antimicrobial and antibiofilm efficacy of triclosan and dispersinB combination. *J Antimicrob Chemother* 64:88–93.
120. Lappia JW, Griswold KE. 2013. Alginate lyase exhibits catalysis-independent biofilm dispersion and antibiotic synergy. *Antimicrob Agents Chemother* 57:137–145.
121. Flemming HC, Wingender J. 2010. The biofilm matrix. *Nat Rev Microbiol* 8:623–633.
122. Byrd MS, Pang B, Hong W, Waligora EA, Juneau RA, Armbruster CE, Weimer KE, Murrah K, Mann EE, Lu H, Sprinkle A, Parsek MR, Kock ND, Wozniak DJ, Swords WE. 2011. Direct evaluation of *Pseudomonas aeruginosa* biofilm mediators in a chronic infection model. *Infect Immun* 79:3087–3095.
123. Christensen LD, van Gennip M, Rybtke MT, Wu H, Chiang WC, Alhede M, Hoiby N, Nielsen TE, Givskov M, Tolker-Nielsen T. 2013. Clearance of *Pseudomonas aeruginosa* foreign-body biofilm infections through reduction of the cyclic di-GMP level in the bacteria. *Infect Immun* 81:2705–2713.
124. Ma Q, Yang Z, Pu M, Peti W, Wood TK. 2011. Engineering a novel c-di-GMP-binding protein for biofilm dispersal. *Environ Microbiol* 13:631–642.
125. Rogers SA, Huigens RW, 3rd, Cavanagh J, Melander C. 2010. Synergistic effects between conventional antibiotics and 2-aminoimidazole-derived antibiofilm agents. *Antimicrob Agents Chemother* 54:2112–2118.
126. Frei R, Breitbach AS, Blackwell HE. 2012. 2-Aminobenzimidazole derivatives strongly inhibit and disperse *Pseudomonas aeruginosa* biofilms. *Angew Chem-Int Edit* 51:5226–5229.
127. Cai PJ, Xiao X, He YR, Li WW, Yu L, Yu HQ. 2013. Disintegration of aerobic granules induced by *trans*-2-decanoic acid. *Bioresour Technol* 128:823–826.
128. Yeon KM, Cheong WS, Oh HS, Lee WN, Hwang BK, Lee CH, Beyenal H, Lewandowski Z. 2009. Quorum sensing: a new biofouling control paradigm in a membrane bioreactor for advanced wastewater treatment. *Environ Sci Technol* 43:380–385.
129. Tan CH, Koh KS, Xie C, Tay M, Zhou Y, Williams R, Ng WJ, Rice SA, Kjelleberg S. 2014. The role of quorum sensing signalling in EPS production and the assembly of a sludge community into aerobic granules. *ISME J*. [Epub ahead of print.] doi:10.1038/ismej.2013.240.
130. Hentzer M, Givskov M. 2003. Pharmacological inhibition of quorum sensing for the treatment of chronic bacterial infections. *J Clin Invest* 112:1300–1307.
131. Hentzer M, Wu H, Andersen JB, Riedel K, Rasmussen TB, Bagge N, Kumar N, Schembri MA, Song Z, Kristoffersen P, Manefield M, Costerton JW, Molin S, Eberl L, Steinberg P, Kjelleberg S, Hoiby N, Givskov M. 2003. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J* 22:3803–3815.
132. Barraud N, Kardak BG, Yepuri NR, Howlin RP, Webb JS, Faust SN, Kjelleberg S, Rice SA, Kelso MJ. 2012. Cephalosporin-3'-diazoniumdiolates: targeted NO-donor prodrugs for dispersing bacterial biofilms. *Angew Chem-Int Edit* 51:9057–9060.
133. Yepuri NR, Barraud N, Shah Mohammadi N, Kardak BG, Kjelleberg S, Rice SA, Kelso MJ. 2013. Synthesis of cephalosporin-3'-diazoniumdiolates: biofilm dispersing NO-donor prodrugs activated by β -lactamase. *Chem Commun* 49:4791–4793.
134. Kutty SK, Barraud N, Pham A, Iskander G, Rice SA, Black DS, Kumar N. 2013. Design, synthesis, and evaluation of fimbrolide-nitric oxide donor hybrids as antimicrobial agents. *J Med Chem* 56:9517–9529.
135. Duong HT, Jung K, Kutty SK, Agustina S, Adnan NN, Basuki JS, Kumar N, Davis TP, Barraud N, Boyer C. 2014. Nanoparticle (star polymer) delivery of nitric oxide effectively negates *Pseudomonas aeruginosa* biofilm formation. *Biomacromolecules*. [Epub ahead of print.] doi:10.1021/bm500422v.
136. Barraud N, Storey MV, Moore ZP, Webb JS, Rice SA, Kjelleberg S. 2009. Nitric oxide-mediated dispersal in single- and multi-species biofilms of clinically and industrially relevant microorganisms. *Microb Biotechnol* 2:370–378.
137. Barnes RJ, Bandi RR, Wong WS, Barraud N, McDougald D, Fane A, Kjelleberg S, Rice SA. 2013. Optimal dosing regimen of nitric oxide donor compounds for the reduction of *Pseudomonas aeruginosa* biofilm and isolates from wastewater membranes. *Biofouling* 29:203–212.
138. Ren H, Wu J, Xi C, Lehnert N, Major T, Bartlett RH, Meyerhoff ME. 2014. Electrochemically modulated nitric oxide (NO) releasing biomedical devices via copper(II)-tri(2-pyridylmethyl)amine mediated reduction of nitrite. *ACS Appl Mater Interfaces* 6:3779–3783.
139. Cathie K, Howlin RP, Barraud N, Carroll MP, Clarke SC, Connott GJ, Cornelius V, Daniels TW, Duignan C, Feelisch M, Fernandez B, Hall-Stoodley L, Jefferies JMC, Kelso MJ, Kjelleberg S, Legg JP, Pink S, Rice SA, Rogers G, Salib RJ, Smith C, Stoodley P, Sukhtankar P, Webb JS, Faust SN. 2014. Low dose nitric oxide as adjunctive therapy to reduce antimicrobial tolerance of *Pseudomonas aeruginosa* biofilms in the treatment of patients with cystic fibrosis: report of a proof of concept clinical trial. *Am J Respir Crit Care Med* 189: A2843.