Simple QED- and QCD-like models at finite density

Pawlowski, Jan M.; Stamatescu, Ion-Olimpiu; Zielinski, Christian

2015

Pawlowski, J. M., Stamatescu, I.-O., & Zielinski, C. (2015). Simple QED- and QCD-like models at finite density. *Physical Review D*, 92, 014508-.

<https://hdl.handle.net/10356/81129>

<https://doi.org/10.1103/PhysRevD.92.014508>

© 2015 American Physical Society. This paper was published in *Physical Review D* and is made available as an electronic reprint (preprint) with permission of American Physical Society. The published version is available at: [<http://dx.doi.org/10.1103/PhysRevD.92.014508>]. One print or electronic copy may be made for personal use only. Systematic or multiple reproduction, distribution to multiple locations via electronic or other means, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper is prohibited and is subject to penalties under law.

Downloaded on 26 Apr 2023 20:33:04 SGT

Simple QED- and QCD-like models at finite densityJan M. Pawłowski,^{1,2} Ion-Olimpiu Stamatescu,^{1,3} and Christian Zielinski^{1,4*}¹*Institut für Theoretische Physik, Universität Heidelberg,
Philosophenweg 16, 69120 Heidelberg, Germany*²*ExtreMe Matter Institute EMMI, GSI, Planckstraße 1, D-64291 Darmstadt, Germany*³*FEST, Schmeilweg 5, 69118 Heidelberg, Germany*⁴*Division of Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore*

(Received 3 April 2014; published 30 July 2015)

In this paper we discuss one-dimensional models reproducing some features of quantum electrodynamics and quantum chromodynamics at nonzero density and temperature. Since a severe sign problem makes a numerical treatment of QED and QCD at high density difficult, such models help to explore various effects peculiar to the full theory. Studying them gives insights into the large density behavior of the Polyakov loop by taking both bosonic and fermionic degrees of freedom into account, although in one dimension only the implementation of a global gauge symmetry is possible. For these models we evaluate the respective partition functions and discuss several observables as well as the Silver Blaze phenomenon.

DOI: [10.1103/PhysRevD.92.014508](https://doi.org/10.1103/PhysRevD.92.014508)

PACS numbers: 11.15.Ha, 12.38.Gc, 12.38.-t

I. INTRODUCTION

One of the open challenges of lattice gauge theory is the *ab initio* treatment of full quantum chromodynamics (QCD) at finite density and low temperature. The fermion determinant is rendered complex and rapidly oscillating after the introduction of a finite chemical potential μ . The same holds for a wide spectrum of theories at nonzero density. The resulting near-cancellations make an evaluation of expectation values extremely challenging. Several methods for meeting the sign problem have been advanced, but are either limited in their applicability or are about to be tested for QCD, see e.g. Refs. [1–11].

In the past, studies of models of QCD have been proved insightful [12–19], including in the special case of heavy quarks [20–22]. The interest in the present and many other models discussed in the literature is that they provide a testing ground for new simulation algorithms to be applied to the full theory. Examples can be found in Refs. [18,19,21,23].

In this paper, we construct and study models that exhibit certain characteristic properties of quantum electrodynamics (QED) and quantum chromodynamics at nonzero μ and temperature T . These models are formulated on a one-dimensional lattice using staggered fermions [24–27]. In comparison to the full theories these models have several simplifications. In particular, as it is not possible to define a plaquette variable in one dimension, there is no Yang-Mills action and the models can only respect a global gauge symmetry. Nonetheless, the introduction of a bosonic field allows us to go beyond models with only fermionic degrees of freedom. Moreover, by the introduction of a suitable

bosonic action S_g we can mimic some features of Yang-Mills theory, which cannot be directly translated to one dimension.

We note that our models generalize the one-link models presented in Ref. [21]. With an explicit incorporation of bosonic degrees of freedom with a corresponding action, a particular form of the fermion matrix and an integration over conjugacy classes we are able to construct a novel low-dimensional QCD-like model, which extends and cross-checks previous work.

As a result these models allow us to investigate some universal phenomena also found in other models from a different perspective. In particular our findings presented in Sec. III shine new light on the behavior of the Polyakov loop at large densities.

The resulting partition function can be fully integrated for a U(1) gauge group. In the case of SU(3) we are left with an integral expression, whose sign problem is manageable and which can be numerically evaluated. We discuss some observables and investigate the Silver Blaze phenomenon [28] in these models. A theory exhibits Silver Blaze behavior, if in the zero-temperature limit $T \rightarrow 0$ observables become independent of the chemical potential for $\mu \leq \mu_{\text{crit}}$, where μ_{crit} is some critical onset value of the chemical potential. In realistic models $\mu_{\text{crit}} = m_q^{\text{phys}}$ corresponds to the physical fermion mass, or, more generally, the mass of the lowest excitation with nonvanishing quark number.

We organize the paper as follows: first we introduce a QED-like model in Sec. II. We derive a closed expression for the partition function and discuss the dependence of some observables on chemical potential and temperature. In Sec. III we deal with the case of QCD and derive an integral expression, which can be numerically evaluated. In Sec. IV we discuss and summarize our findings.

*Corresponding author.
zielinski@pmail.ntu.edu.sg

II. A SOLUBLE, QED-LIKE MODEL AT NONZERO DENSITY

For the construction of a QED-like model we formulate an Abelian U(1) lattice gauge theory on a finite one-dimensional lattice with staggered fermions and couple them to a chemical potential, following a similar ansatz as employed in previous works.

A. Partition function

The action we employ mimics the usual compact lattice QED in one dimension. The lattice is assumed to have a lattice spacing of a and an extension of N sites, where N is assumed to be even. We set $a = 1$, i.e., we measure all dimensionful quantities in appropriate powers of a . We consider a single staggered fermion field and couple it to a chemical potential μ . The temperature is identified with the inverse of the lattice extension $T = N^{-1}$. The action $S = S_f + S_g$ consists of the fermionic part

$$S_f = \sum_{t,\tau=1}^N \bar{\chi}(t) K(t, \tau) \chi(\tau), \quad (1)$$

and the pure bosonic part

$$S_g = \beta \sum_{t=1}^N \left[1 - \frac{1}{2} (U_t + U_t^\dagger) \right]. \quad (2)$$

Here $\bar{\chi}$ and χ denote the staggered fermion field, $K(t, \tau)$ the fermion matrix, $\beta = 1/e^2$ the inverse coupling constant and $U_t \in \text{U}(1)$ the link variables. The fermion matrix reads

$$K(t, \tau) = \frac{1}{2} (U_t e^{\mu} \delta_{t+1,\tau} - U_t^\dagger e^{-\mu} \delta_{t-1,\tau}) + m \delta_{t\tau}, \quad (3)$$

with m denoting the mass of the fermion. The introduction of the chemical potential μ follows the prescription by Hasenfratz and Karsch [29]. Furthermore we impose an antiperiodic boundary condition for the fermionic field. After integrating out the fermionic degrees of freedom, the partition function reads

$$Z = \int \prod_{t=1}^N dU_t \det K e^{-S_g}. \quad (4)$$

In this case the fermion determinant can be evaluated analytically using identity (1) derived in Ref. [30]. We find

$$2^N \det K = e^{N\mu} \prod_t U_t + e^{-N\mu} \prod_\tau U_\tau^\dagger + 2\rho_+, \quad (5)$$

where we have introduced

$$\rho_\pm = \lambda_+ \pm \lambda_-, \quad \lambda_\pm = \frac{1}{2} \left(m \pm \sqrt{1 + m^2} \right)^N. \quad (6)$$

Note that Eq. (5), like full QED, satisfies the identity

$$\det K(\mu) = [\det K(-\mu^*)]^*, \quad (7)$$

which shows that in general the fermion determinant is complex for $\mu > 0$. We parametrize the link variables as $U_t = \exp(i\phi_t)$ in terms of algebra-valued fields $\phi_t \in [0, 2\pi)$. The corresponding U(1)-Haar measure reads

$$\int dU_t = \int_0^{2\pi} \frac{d\phi_t}{2\pi}. \quad (8)$$

With this parametrization, the action in Eq. (2) takes the form

$$S_g = \beta \sum_{t=1}^N (1 - \cos \phi_t). \quad (9)$$

This allows us to integrate the partition function given in Eq. (4) by using the expression we derived for the fermion determinant in Eq. (5), to find

$$Z = \frac{e^{-\beta N}}{2^{N-1}} [\rho_+ I_0^N(\beta) + \cosh(N\mu) I_1^N(\beta)]. \quad (10)$$

Here I_n denotes the modified Bessel functions of the first kind. As a cross-check we verified that Z reduces to the previously derived partition function in Ref. [21] for $N = 1$ up to a normalization constant, which depends on m and β .

B. Observables

Given the final expression for the partition function in Eq. (10), we can easily calculate any observable of interest. The density follows from $\langle n \rangle = N^{-1} \partial_\mu \log Z$, the respective susceptibility is defined as $\langle \chi_n \rangle = \partial_\mu \langle n \rangle$ and the fermion condensate is given by $\langle \bar{\chi} \chi \rangle = N^{-1} \partial_m \log Z$. For the density we then find,

$$\langle n \rangle = \frac{\sinh(N\mu) I_1^N(\beta)}{\rho_+ I_0^N(\beta) + \cosh(N\mu) I_1^N(\beta)}. \quad (11)$$

Again this expression reduces to the known result in Ref. [21] for $N = 1$. The fermion condensate follows as,

$$\langle \bar{\chi} \chi \rangle = \frac{(1 + m^2)^{-1/2} \rho_- I_0^N(\beta)}{\rho_+ I_0^N(\beta) + \cosh(N\mu) I_1^N(\beta)}. \quad (12)$$

By directly evaluating the respective path integral expression, we find for the Polyakov loop $\mathcal{P} = \prod_t U_t$ the expectation value

$$\langle \mathcal{P} \rangle = \frac{e^{-\beta N}}{2^N Z} [2\rho_+ I_1^N(\beta) + e^{N\mu} I_2^N(\beta) + e^{-N\mu} I_0^N(\beta)]. \quad (13)$$

The conjugate Polyakov loop $\mathcal{P}^\dagger = \prod_t U_t^\dagger$ follows from a simple symmetry argument as $\langle \mathcal{P}^\dagger \rangle_\mu = \langle \mathcal{P} \rangle_{-\mu}$, cf. Ref. [18].

In Fig. 1 we show the density given by Eq. (11), the fermion condensate by Eq. (12) and the Polyakov loop by Eq. (13) as functions of the chemical potential μ . We see

FIG. 1 (color online). Observables in the U(1) model. (a) Density, condensate and normalized susceptibility; (b) Polyakov loop and conjugate Polyakov loop.

that already for $T = 1/N = 1/10$ the observables only show a weak dependence on the chemical potential below some critical value μ_{crit} . This shows how the Silver Blaze behavior [28] becomes apparent in this model, which holds strictly in the limit $T \rightarrow 0$.

Close to $\mu \approx \mu_{\text{crit}}$ we also observe a fast increase or decrease of the observables before reaching the saturation regime. Note that the limits $\mu \rightarrow \infty$ and $\beta \rightarrow 0$ do not commute. Density and condensate behave as one would expect. The expectation values of $\mathcal{P}$ and $\mathcal{P}^\dagger$ approach

$$\langle \mathcal{P} \rangle \rightarrow \left[\frac{I_2(\beta)}{I_1(\beta)} \right]^N, \quad \langle \mathcal{P}^\dagger \rangle \rightarrow \left[\frac{I_0(\beta)}{I_1(\beta)} \right]^N, \quad (14)$$

for $\mu \rightarrow \infty$. The Polyakov loop quickly drops to a typically small value with increasing μ while the conjugate Polyakov loop grows to a saturation value which diverges when $\beta \rightarrow 0$.

III. A QCD-LIKE MODEL AT NONZERO DENSITY

Now we extend the previous model to the non-Abelian gauge group SU(3). By restricting the integration over the full gauge group to the respective conjugacy classes of SU(3), we will be able to reduce the partition function to an integral expression with a manageable sign problem.

A. Partition function

Our starting point is again the path integral expression for the partition function in Eq. (4), where now the pure bosonic part of the action reads

$$S_g = \beta \sum_{t=1}^N \left[1 - \frac{1}{6} \text{Tr}_c (U_t + U_t^\dagger) \right]. \quad (15)$$

Here $\beta = 6/g^2$ denotes the inverse coupling, $U_t \in \text{SU}(3)$ the link variables and Tr_c a trace in color space. Furthermore we replace the fermion matrix by

$$K(t, \tau) = \frac{1}{2} (\sigma_+ U_t e^\mu \delta_{t+1, \tau} - \sigma_- U_t^\dagger e^{-\mu} \delta_{t-1, \tau}) + m \delta_{t\tau}, \quad (16)$$

with $\sigma_\pm = \frac{1}{2}(1 \pm \sigma_3)$ and the third Pauli matrix $\sigma_3 = \text{diag}(1, -1)$. In the loop expansion this suppresses back steps, thus simulating a special feature of Wilson fermions. This choice results in a factorization of the fermion determinant of the form

$$\det K = \det_{t,c} \mathcal{K}_f \cdot \det_{t,c} \mathcal{K}_b, \quad (17)$$

where we introduced

$$\begin{aligned} \det_{t,c} \mathcal{K}_f &= \det_{t,c} \left(m \delta_{t\tau} + \frac{1}{2} U_t e^\mu \delta_{t+1, \tau} \right), \\ \det_{t,c} \mathcal{K}_b &= \det_{t,c} \left(m \delta_{t\tau} - \frac{1}{2} U_t^\dagger e^{-\mu} \delta_{t-1, \tau} \right). \end{aligned} \quad (18)$$

Here $U_{N+1} = -U_1$ and $\det_{t,c}$ refers to a determinant in position and color space.

In the following we restrict ourselves to observables which only depend on the conjugacy class of the link variables. We then replace the integration over the full gauge group SU(3) with an integration over these conjugacy classes. This idea and the factorization given in Eq. (17) were also previously exploited in a one link model in Ref. [21]. We thus parametrize the links by

$$U_t = \text{diag}(e^{i\phi_t}, e^{i\vartheta_t}, e^{-i(\phi_t + \vartheta_t)}), \quad (19)$$

with $\phi_t, \vartheta_t \in (-\pi, \pi]$. Ignoring a normalization constant, the Haar measure is given by $dU_t \propto J(\phi_t, \vartheta_t) d\phi_t d\vartheta_t$ with

$$\begin{aligned} J(\phi_t, \vartheta_t) &= \sin^2 \left(\frac{\phi_t - \vartheta_t}{2} \right) \\ &\times \sin^2 \left(\frac{\phi_t + 2\vartheta_t}{2} \right) \sin^2 \left(\frac{2\phi_t + \vartheta_t}{2} \right), \end{aligned} \quad (20)$$

while the bosonic part of the action takes the form

$$S_g = \beta \sum_{t=1}^N \left[1 - \frac{1}{3} (\cos \phi_t + \cos \vartheta_t + \cos (\phi_t + \vartheta_t)) \right]. \quad (21)$$

The determinant in position space has a simple structure and can be analytically evaluated, e.g. directly or by resummation of the loop expansion for Eq. (16). For the remaining determinant in color space we use the identity

$$\det_c(\mathbb{1} + \alpha U_t) = 1 + \alpha \text{Tr}_c U_t + \alpha^2 \text{Tr}_c U_t^{-1} + \alpha^3, \quad (22)$$

valid for all $A \in \text{SL}_3(\mathbb{C})$, see Ref. [21]. We can express the result in terms of the (conjugate) Polyakov loop

$$\begin{aligned} \det_{t,c} \mathcal{K}_f &= m^{3N} \det_c \left(\mathbb{1} + \xi_f \prod_t U_t \right) \\ &= m^{3N} (1 + \xi_f \mathcal{P} + \xi_f^2 \mathcal{P}^\dagger + \xi_f^3), \end{aligned} \quad (23)$$

with $\xi_f = [\kappa \exp(\mu)]^N$ and hopping parameter $\kappa = 1/(2m)$. The Polyakov loop $\mathcal{P}$ and conjugate Polyakov loop $\mathcal{P}^\dagger$ are defined by

$$\mathcal{P} = \text{Tr}_c \prod_{t=1}^N U_t, \quad \mathcal{P}^\dagger = \text{Tr}_c \prod_{t=1}^N U_t^\dagger. \quad (24)$$

Analogously, we find

$$\det_{t,c} \mathcal{K}_b = m^{3N} (1 + \xi_b \mathcal{P}^\dagger + \xi_b^2 \mathcal{P} + \xi_b^3), \quad (25)$$

with $\xi_b = [\kappa \exp(-\mu)]^N$. We observe that, as in the QED-like model, the fermion determinant in Eq. (17) satisfies the relation in Eq. (7).

Putting all pieces together, we find that the partition function of the model reads

$$\begin{aligned} Z &= m^{6N} \int_{-\pi}^{\pi} \left[\prod_t d\phi_t d\vartheta_t J(\phi_t, \vartheta_t) \right] \\ &\times (1 + \xi_f \mathcal{P} + \xi_f^2 \mathcal{P}^\dagger + \xi_f^3) \\ &\times (1 + \xi_b \mathcal{P}^\dagger + \xi_b^2 \mathcal{P} + \xi_b^3) e^{-S_g}, \end{aligned} \quad (26)$$

where an irrelevant numerical normalization constant has been dropped. The measure term $J(\phi_t, \vartheta_t)$ was given in Eq. (20), S_g was introduced in Eq. (21).

B. Observables

Considering Eq. (26) as a partition function for a model of QCD, we can derive integral expressions for the density, the susceptibility and the fermion condensate by taking corresponding derivatives of $\log Z$. For the Polyakov loop and the conjugate Polyakov loop we insert a $\frac{1}{2}\mathcal{P}$ or a $\frac{1}{2}\mathcal{P}^\dagger$ term in Eq. (26).

The resulting integral expressions are numerically evaluated. Typical examples of these observables can be found in Fig. 2. The density and the condensate show similar qualitative behavior to the corresponding observables in the U(1) model, where we now find $\langle n \rangle \rightarrow 3$ for $\mu \rightarrow \infty$.

The Polyakov loop and conjugate Polyakov loop show some nontrivial behavior. Close to the critical onset μ_{crit} , we find peaks in $\langle \mathcal{P} \rangle$ and $\langle \mathcal{P}^\dagger \rangle$ with the peak in the conjugate Polyakov loop appearing at smaller μ . Similar behavior was previously observed in a simulation of a gauge theory with exceptional group G_2 [31], a strong coupling limit in HQCD [23], a three-dimensional effective theory of nuclear matter [22] and in recent studies of one-dimensional QCD [18,19]. The drop of the Polyakov loop at high density is easily understood as an effect of saturation, while the displacement of the peaks has a dynamical basis, see, e.g. Ref. [23].

Despite making use of a different approach, in general we find good qualitative agreement with the results reported in Refs. [18,19] after dropping S_g , i.e. for $\beta = 0$.

FIG. 2 (color online). Observables in the SU(3) model. (a) Density and condensate; (b) Polyakov loop and conjugate Polyakov loop.

IV. CONCLUSIONS

In this paper we have constructed one-dimensional lattice models resembling QED and QCD to investigate the finite density and finite temperature regime. Despite the drastic simplifications in these models, they capture some essential physical properties expected from the full theory and show an interesting behavior of the Polyakov loop. We found that they—like their four-dimensional continuous counterparts—exhibit the Silver Blaze property in the zero temperature limit $N \rightarrow \infty$. The μ -dependence of the SU(3) (conjugate) Polyakov loop $\mathcal{P}(\mathcal{P}^\dagger)$ shows the peculiar μ -dependence also found in other approximations of QCD.

The models presented here can also serve as a starting point for the construction of more elaborated models.

ACKNOWLEDGMENTS

This work is supported by the Helmholtz Alliance HA216/EMMI and by ERC-AdG-290623. J. M. P. thanks the Yukawa Institute for Theoretical Physics, Kyoto University, where this work was completed during the YITP-T-13-05 on 'New Frontiers in QCD'. I.-O. S. thanks the Deutsche Forschungsgemeinschaft by STA 283/16-1 and C. Z. thanks Nanyang Technological University for support.

-
- [1] P. de Forcrand, *Proc. Sci.*, LAT2009 (2009) 010 [[arXiv:1005.0539](#)].
 - [2] M. P. Lombardo, *Mod. Phys. Lett. A* **22**, 457 (2007).
 - [3] G. Aarts, *Proc. Sci.*, LAT2009 (2009) 024 [[arXiv:0910.3772](#)].
 - [4] G. Aarts, L. Bongiovanni, E. Seiler, D. Sexty, and I.-O. Stamatescu, *Eur. Phys. J. A* **49**, 89 (2013).
 - [5] G. Aarts, *Proc. Sci.*, LATTICE2012 (2012) 017 [[arXiv:1302.3028](#)].
 - [6] A. Schmidt, Y. D. Mercado, and C. Gattringer, *Proc. Sci.*, LATTICE2012 (2012) 098 [[arXiv:1211.1573](#)].
 - [7] Y. D. Mercado, C. Gattringer, and A. Schmidt, *Comput. Phys. Commun.* **184**, 1535 (2013).
 - [8] A. Alexandru, M. Faber, I. Horvath, and K.-F. Liu, *Phys. Rev. D* **72**, 114513 (2005).
 - [9] A. Alexandru and U. Wenger, *Phys. Rev. D* **83**, 034502 (2011).
 - [10] K. Langfeld, B. Lucini, and A. Rago, *Phys. Rev. Lett.* **109**, 111601 (2012).
 - [11] K. Langfeld and J. M. Pawłowski, *Phys. Rev. D* **88**, 071502 (2013).
 - [12] N. Bilic, H. Gausterer, and S. Sanielevici, *Phys. Rev. D* **37**, 3684 (1988).
 - [13] N. Bilic and K. Demeterfi, *Phys. Lett. B* **212**, 83 (1988).
 - [14] K. Splittorff and J. J. M. Verbaarschot, *Phys. Rev. Lett.* **98**, 031601 (2007).
 - [15] K. Splittorff and J. J. M. Verbaarschot, *Phys. Rev. D* **75**, 116003 (2007).
 - [16] K. Splittorff and J. J. M. Verbaarschot, *Phys. Rev. D* **77**, 014514 (2008).
 - [17] L. Ravagli and J. J. M. Verbaarschot, *Phys. Rev. D* **76**, 054506 (2007).
 - [18] J. Bloch, F. Bruckmann, and T. Wettig, *J. High Energy Phys.* **10** (2013) 140.
 - [19] J. Bloch, F. Bruckmann, and T. Wettig, *Proc. Sci.*, LATTICE 2013 (2013) 194 [[arXiv:1310.6645](#)].
 - [20] R. De Pietri, A. Feo, E. Seiler, and I.-O. Stamatescu, *Phys. Rev. D* **76**, 114501 (2007).
 - [21] G. Aarts and I.-O. Stamatescu, *J. High Energy Phys.* **09** (2008) 018.
 - [22] M. Fromm, J. Langelage, S. Lottini, M. Neuman, and O. Philipsen, *Phys. Rev. Lett.* **110**, 122001 (2013).
 - [23] E. Seiler, D. Sexty, and I.-O. Stamatescu, *Phys. Lett. B* **723**, 213 (2013).
 - [24] J. B. Kogut and L. Susskind, *Phys. Rev. D* **11**, 395 (1975).
 - [25] T. Banks, L. Susskind, and J. B. Kogut, *Phys. Rev. D* **13**, 1043 (1976).
 - [26] T. Banks, S. Raby, L. Susskind, J. Kogut, D. R. T. Jones, P. N. Scharbach, and D. K. Sinclair (Cornell-Oxford-Tel Aviv-Yeshiva Collaboration), *Phys. Rev. D* **15**, 1111 (1977).
 - [27] L. Susskind, *Phys. Rev. D* **16**, 3031 (1977).
 - [28] T. D. Cohen, *Phys. Rev. Lett.* **91**, 222001 (2003).
 - [29] P. Hasenfratz and F. Karsch, *Phys. Lett. B* **125**, 308 (1983).
 - [30] L. G. Molinari, *Linear Algebra Appl.* **429**, 2221 (2008).
 - [31] A. Maas, L. von Smekal, B. Wellegehausen, and A. Wipf, *Phys. Rev. D* **86**, 111901 (2012).