Impact of Using Different Promoters and Matrix Attachment Regions on Recombinant Protein Expression Level and Stability in Stably Transfected CHO Cells

Ho, Steven C. L.; Mariati; Yeo, Jessna H. M.; Fang, Shiyi Goh; Yang, Yuansheng

2014

Ho, S. C. L., Mariati, Yeo, J. H. M., Fang, S. G., & Yang, Y. (2015). Impact of Using Different Promoters and Matrix Attachment Regions on Recombinant Protein Expression Level and Stability in Stably Transfected CHO Cells. *Molecular Biotechnology*, 57(2), 138-144.

<https://hdl.handle.net/10356/81465>

<https://doi.org/10.1007/s12033-014-9809-2>

© 2014 Springer Science+Business Media New York. This is the author created version of a work that has been peer reviewed and accepted for publication by *Molecular Biotechnology*, Springer Science+Business Media New York. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [<http://dx.doi.org/10.1007/s12033-014-9809-2>].

Downloaded on 26 Apr 2023 23:26:38 SGT

Molecular Biotechnology

Impact of using different promoters and matrix attachment regions on recombinant protein expression level and stability in stably transfected CHO cells --Manuscript Draft--

Manuscript Number:	MOBI-D-14-00247R1
Full Title:	Impact of using different promoters and matrix attachment regions on recombinant protein expression level and stability in stably transfected CHO cells
Article Type:	Research Papers
Keywords:	matrix attachment region (MAR), CMV promoter, SV40 promoter, CHO elongation factor 1 α promoter, stable recombinant protein expression, CHO cells
Corresponding Author:	Yuansheng Yang, PhD Bioprocessing Technology Institute Singapore, SINGAPORE
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Bioprocessing Technology Institute
Corresponding Author's Secondary Institution:	
First Author:	Steven C.L. Ho
First Author Secondary Information:	
Order of Authors:	Steven C.L. Ho
	Mariati Mariati
	Jessna H.M. Yeo
	Shiyi Goh Fang
	Yuansheng Yang, PhD
Order of Authors Secondary Information:	
Abstract:	<p>High level and stable expression are required for therapeutic protein production in mammalian cells. Three commonly used promoters from the simian virus 40 (SV40), the CHO elongation factor 1α gene (EF1α) and the human cytomegalovirus major immediate early gene (CMV) and two matrix attachment regions from the chicken lysozyme gene (cMAR) and the human interferon β (iMAR) were evaluated for enhancing recombinant gene expression level and stability in stably transfected CHO cells. In the absence of MAR elements, the SV40 promoter gave lower expression level but higher stability than the EF1α promoter and the CMV promoter. The inclusion of MAR elements did not increase the integrated gene copies for all promoters but did enhance expression level for only the SV40 promoter. The enhanced gene expression was due to an increase in mRNA levels. Neither MAR elements enhance gene expression stability during long term culture. The combinations of SV40 promoter and MAR elements are the best for obtaining both high expression level and stability. The information presented here would be valuable to those developing vectors for generation of CHO cell lines with stable and high productivity.</p>
Response to Reviewers:	<p>Reviewer #1: The paper entitled "Impact of using different promoters and matrix attachment regions on recombinant protein expression level and stability in stably transfected CHO cells" is a useful publication that investigates the influence of different matrix attachment factors and promoter combinations in order to maximize the expression and stability of protein expression levels in CHO cells. The experiments are interesting in that they will aid further investigators by elucidating a strategy to maximize protein expression under a given set of parameters- notably showing that some combinations have little virtue. These results are interesting when taken in context with other publications that have perhaps not given the same overall impression to the over-arching utility of matrix attachment regions. The authors</p>

	<p>conclude that the less commonly used (w.r.t. to CMV) promoter, SV40 is the 'best', especially in terms of long term cell line protein production. It would be of interest to see these experiments repeated in other cell lines, such as the commonly used HEK line. In addition to this, the authors suggest that gene silencing is in part due to methylation of CG dinucleotides- improving on the SV40 promoter by mutating out these remaining dinucleotides would be an interesting experiment.</p> <p>Overall, the paper is interesting, useful and clearly written.</p> <p>Author response: We thank the reviewer for all the valuable comments. We tried to evaluate different vectors in HEK293 cells previously. We were not be able to generate stably transfected pools as all cells died during selection probably because the mutated neomycin was too weak for HEK293 cells to survive.</p> <p>Regarding removal of the remaining CG dinucleotides in the SV40 promoter, we also tried in our lab. The CG-free SV40 promoter had much lower expression than the wild type SV40 promoter in transient transfections. Stably transfected pools were not be able to be generated by using the mutated CG-free SV40 due to its weak strength. Further work to generate CG-free SV40 without compromised strength is in progress in our lab. We shared this information in the discussion in the revised manuscript.</p> <p>Reviewer #2: The authors report a study comparing the affect of two dfferent MAR elements on the level and stability of GFP expressed in CHO cells under the control of three different promoters (CMV, SV-40 and E1a). Although the use of MAR elements has been well documented, the observation that only expression of GFP from the SV-40 promoter is significantly improved by addition of MAR elements in the vector is novel. Given the widespread use of E1a and CMV promoter vectors the observation is of general interest. However it would be considerably strengthened if the result was exemplified with at least a second example, preferably a protein relevant to the production of biopharmaceuticals e.g. antibody or Fc-fusion protein.</p> <p>Author response: We agree with the reviewer it will be more convincing to test more relevant therapeutic proteins. As it will take more than 6 months to isolate clones and do stability testing for a second protein which may be beyond the time frame given by the journal to revise this manuscript, we would like to do it in future studies.</p>
--	---

For submission to Molecular Biotechnology

Impact of using different promoters and matrix attachment regions on recombinant protein expression level and stability in stably transfected CHO cells

Steven C.L. Ho¹, Mariati¹, Jessna H.M. Yeo¹, Shiyi Goh Fang¹, Yuansheng Yang^{1,2*}

¹ Bioprocessing Technology Institute, Agency for Science, Technology and Research (A*STAR), 20 Biopolis Way, #06-01 Centros, Singapore 138668

²School of Chemical and Biomedical Engineering, Nanyang Technological University, N1.2-B2-33, 62 Nanyang Avenue, Singapore 637459

* Correspondence to:
Yuansheng Yang
Phone: (65)-64070825, Fax: (65)-64789561
Email: yang_yuansheng@bti.a-star.edu.sg

Acknowledgements

This work was supported by the Biomedical Research Council/Science and Engineering Research Council of A*STAR (Agency for Science, Technology and Research), Singapore.

Title

Impact of using different promoters and matrix attachment regions on recombinant protein expression level and stability in stably transfected CHO cells

Abstract

High level and stable expression are required for therapeutic protein production in mammalian cells. Three commonly used promoters from the simian virus 40 (SV40), the CHO elongation factor 1 α gene (EF1 α) and the human cytomegalovirus major immediate early gene (CMV) and two matrix attachment regions from the chicken lysozyme gene (cMAR) and the human interferon β (iMAR) were evaluated for enhancing recombinant gene expression level and stability in stably transfected CHO cells. In the absence of MAR elements, the SV40 promoter gave lower expression level but higher stability than the EF1 α promoter and the CMV promoter. The inclusion of MAR elements did not increase the integrated gene copies for all promoters but did enhance expression level for only the SV40 promoter. The enhanced gene expression was due to an increase in mRNA levels. Neither MAR elements enhance gene expression stability during long term culture. The combinations of SV40 promoter and MAR elements are the best for obtaining both high expression level and stability. The information presented here would be valuable to those developing vectors for generation of CHO cell lines with stable and high productivity.

Keywords: matrix attachment region (MAR), CMV promoter, SV40 promoter, CHO elongation factor 1 α promoter, stable recombinant protein expression, CHO cells

Introduction

Mammalian cells, such as Chinese hamster ovary cells (CHO), are predominant for commercial production of therapeutic proteins because of their capacity to perform proper protein folding, assembly, and post-translational modifications (1, 2). Generating a therapeutic protein producing cell line starts with transfecting the mammalian host cells with a plasmid vector carrying the gene for the respective therapeutic protein. Subsequent selection is performed to isolate stably transfected clones with high productivity and long-term stable production. Expression stability also needs to be maintained without the aid of any selection pressure as these drugs add to production costs and can complicate downstream purification (3). Recombinant protein productivity and stability in a clone is influenced by both the composition of the plasmid vector and the site of vector integration on the chromosome (4-6). Apart from the gene of interest, components of the plasmid include promoters, polyadenylation signals and other expression augmenting elements like matrix attachment regions (MAR).

A commonly used promoter for high level recombinant protein production in mammalian cells is the human cytomegalovirus major immediate early gene promoter (CMV) (7).

While CMV is a strong promoter, there are reports that the promoter is intrinsically susceptible to silencing, resulting in declined productivity during long term culture (3, 8-16). Promoters derived from the simian virus 40 (SV40) and CHO elongation factor 1 α gene (EF1 α) are also strong for therapeutic protein production in mammalian cells (7, 17). EF1 α promoter used in conjunction with flanking regions of the CHO EF1 α gene was more active in CHO cells than using CMV and SV40 promoters alone (17). Protein

production instability in CHO cells when using the SV40 promoter has been reported (18, 19). Some studies indicated that promoters of endogenous mammalian genes like the EF1 α can be more resistant to silencing than viral promoters (20-25). Although there are many separate reports of promoter studies, it can be difficult to make comparisons of the promoters when each report is performed in differing culture conditions and cell lines. A comparison of transgene expression level and stability in stably transfected CHO cell clones under similar conditions without selection pressure would aid in choosing between the CMV, SV40 and EF1 α promoters for recombinant protein production in CHO cells.

Matrix attachment regions (MARs) are DNA elements which may be involved in anchoring DNA/chromatin to the nuclear matrix to define the boundaries of independent chromatin domains (26, 27). MARs were reported to shield transgenes from chromosomal position effects and increase transgene expression level in stably transfected cell lines (28-37). There are conflicting reports on whether MAR elements can prevent transgenes from silencing in stably transfected cell lines. Inclusion of MARs into viral vectors increased their resistance to gene silencing (38, 39). When plasmid vectors, the preferred vectors for safe production of therapeutic proteins in mammalian cells were used, enhanced resistance to silencing was observed for the SV40 promoter but not for the CMV promoter (19, 37), suggesting that the effect of MARs on gene silencing may be dependent on the promoters.

Among many MAR elements which have been identified to date, chicken lysozyme (cMAR) and human interferon β (iMAR) MAR elements have shown to be very effective

at enhancing transgene expression in stably transfected cells (29, 37). In this work, we evaluated the SV40, EF1 α and CMV promoters for recombinant protein expression level and stability in stably transfected CHO cells. The impact of including cMAR and iMAR with the above three promoters were studied as well. The information collected would benefit those choosing promoters and MAR elements during plasmid vector designs for generating cell lines with both high expression level and long-term expression stability.

Materials and Methods

Vector construction

The CMV containing bicistronic vector without MAR was constructed based on a previously described IRES-mediated tricistronic vector (40) with the LC-IRESwt-HC region replaced with EGFP cDNA. The remaining bicistronic vectors with or without MAR were then constructed by replacement of the CMV promoter with either EF1 α or SV40 promoter and insertion of cMAR or iMAR immediately upstream of the promoter and downstream of the SpA. The CMV and SV40 promoter were cloned from pcDNA3.1(+) (Life Technologies, Carlsbad, CA). The EF1 α promoter with sequence corresponding to the region from -463 to +1010 (relative to the transcription start site of +1) of the CHO EF1 α gene (NCBI: AY188393.1) was isolated from CHO K1 cells. The iMAR and cMAR were cloned from a pEPI-1 vector (41) and a pPAG1 vector (37), respectively. All restriction enzymes used in the vector construction were purchased from New England Biolabs (Ipswich, MA). The DH5 α TM competent cells were purchased from Life Technologies.

Cell culture and media

CHO K1 cells (American Type Culture Collection, Manassas, VA) were maintained in medium consisting of Dulbecco's modified Eagle's medium (DMEM) + GlutaMaxTM (Life Technologies) supplemented with 10% fetal bovine serum (FBS) (Sigma-Aldrich, St. Louis, MO). Regular passaging was carried out every 3 to 4 days by diluting cells to 2×10^5 cells/mL. Cell density and viability were measured using the trypan blue exclusion method on an automated Cedex counter (Innovatis, Bielefeld, Germany).

Generation of stably transfected cell lines

Triplicate transfections were performed for each vector using Nucleofector I system according to manufacturer's instructions (Lonza, Cologne, Germany). 5×10^6 cells were transfected with 5 μ g of linearized plasmids in each transfection. The transfected cells were then resuspended in 2 mL of pre-warmed maintenance medium in 6-well tissue culture plates. At 24 h post-transfection, maintenance medium was removed and 2 mL of maintenance medium containing 800 μ g/mL of G418 was added into each well for selection of stable transfectants. Stably transfected pools were obtained after 2 to 4 weeks depending on the vectors. Six clones were randomly isolated from each pool to obtain a total of 18 clones for each vector using limiting dilution method in 96-well tissue culture plates.

Stability testing

The clones isolated for each vector were passaged in 6-well tissue culture plates in the absence of G418 for 8 weeks. MFI for each clone before and after stability testing were measured with the FACS Calibur. Retention of EGFP expression for a clone was calculated as the ratio of MFI of the clone measured at the end of stability testing to the intensity at the start of stability testing.

Analysis of relative gene copies and mRNA levels

The relative EGFP gene copies and mRNA levels were determined using real-time quantitative PCR (qRT-PCR) as described previously(42). Genomic DNA and total RNA were isolated from 5×10^6 cells using the Gentra Puregene Cell Kit (Qiagen, Hilden, Germany) and the RNAqueous-4PCR kit (Ambion, Austin, TX), respectively. EF1 α and β -actin were used to normalize the variation in input amount and quality of RNA and DNA, respectively.

Results

Evaluating recombinant protein expression level using different promoters and impact of including MARs

We first evaluated SV40, EF1 α and CMV promoters and their use with cMAR and iMAR for expression level in stably transfected CHO cells. Bicistronic vectors expressing an enhanced green fluorescence reporter protein (EGFP) and a mutated neomycin phosphotransferase (mNTP) selection marker under the control of SV40, EF1 α or CMV promoter were first constructed (Fig. 1). Copies of cMAR and iMAR were then inserted at the 5' and 3' flanking regions of the expression cassette in each vector. This configuration was shown to be more effective in enhancing stable transgene expression than using only a single copy of MAR in previous studies (30, 35). Use of the mNTP with reduced enzyme activity is for enhancing the selection stringency for high producing cells as only clones with greater transcriptional activity or more copies of the integrated vector can survive the selection process (40, 43, 44).

CHO K1 cells were transfected with each vector followed by drug selection for stable transfectants. Only tens of clones survived the selection in transfected pools generated using the vectors without MAR and stably transfected pools were obtained in about four weeks. In contrast, hundreds of clones survived the selection in transfected pools generated using vectors containing MAR elements and stably transfected pools were obtained in about 2 weeks. EGFP expression in stably transfected pools were quantified by measuring the mean fluorescence intensity (MFI). The SV40 promoter only vector gave the lowest EGFP expression, 2- and 3-fold less than that from the EF1 α and CMV only vectors respectively (Fig. 2A). Application of cMAR and iMAR on the SV40 promoter (SV40+cMAR and SV40+iMAR) enhanced EGFP expression by about 3- and 4-fold, respectively. Inclusion of MARs failed to enhance EGFP expression for EF1 α and CMV promoters. Stably transfected pools generated using SV40+cMAR/iMAR vectors exhibited higher EGFP expression than those generated using any of the EF1 α and CMV vectors. qRT-PCR analysis indicated that application of MARs on different promoters did not change the integrated EGFP gene copies compared to the use of each promoter alone. However, MARs enhanced EGFP mRNA levels when they were applied on the SV40 promoter but had no effect on the EGFP mRNA levels when they were applied on the other two promoters.

Comparison of the different promoters and MAR elements on EGFP expression were subsequently repeated in stably transfected clones (Fig. 2B). MFI varied dramatically between clones generated using the same vectors for all pools. Consistent with the results in stably transfected pools, both the average MFI and maximal EGFP expression in the

SV40 promoter only clones were less than those generated using the EF1 α and CMV promoters. Similar to the stable pools, application of cMAR and iMAR on the SV40 promoter enhanced both average and maximal EGFP expression levels in clones but exhibited no effect when used with EF1 α and CMV promoters.

Evaluating long-term recombinant protein expression stability using different promoters and impact of including MARs

We next evaluated the long-term transgene expression stability of the three promoters and MAR elements by determining the retention of EGFP expression after eight weeks of culturing in the absence of selection pressure. A precise definition of stable production varies depending on application but typically clones which maintain above 70% of their starting productivity are considered to be stable (45). Using 70% retention as a cutoff, 11 out of 18 clones generated using a SV40 promoter were stable after eight weeks of passaging (Fig. 3). The 18 clones retained an average of 79% of their starting EGFP expression level. A reverse correlation between the expression level and stability was observed for these SV40 clones ($R^2=0.6574$, Supplementary Fig. 1). All stable clones had relatively low EGFP expressions. In contrast, none of clones generated using EF1 α and CMV promoters were stable, retaining an average of only 28% and 22% of their starting levels, respectively. Addition of cMAR and iMAR on any promoters did not result in enhanced expression stability but instead decreased expression stability in clones for the SV40 and EF1 α promoter. On average, clones generated using SV40+cMAR/iMAR vectors maintained about 50% their original expression level, 30% less than those generated using SV40 only but still greater than all the EF1 α and CMV clones. Among

18 clones generated using the SV40+cMAR and SV40+iMAR vectors, four and five clones respectively were still stable after the eight weeks. In contrast to those clones generated using SV40 only, no correlation between expression level and stability was observed for clones generated using other vectors ($R^2=0.0002$ to 0.2428 , Supplementary Fig. 1). Some stable clones generated using SV40+cMAR/iMAR also exhibited high EGFP expressions. For instance, the highest producing clone generated using the SV40+cMAR vector and the second highest producing clone generated using the SV40+iMAR vector had 73% and 107% retention of expression after eight weeks culturing.

To understand why the SV40 promoter gave more stable expression than other promoters, nine SV40 clones and nine CMV clones of varying EGFP retention were sorted based on their retention levels and analyzed for changes in EGFP mRNA levels and gene copies. The SV40 clones retained EGFP expression ranging from 23% to 115% (Fig. 4A), and the CMV clones had EGFP retention ranging from 1 to 51% (Fig. 4B). At the end of stability testing, only the SV40 clone A1 had a significant drop in EGFP gene copies to below 50%. Of the remaining eight clones, four clones maintained over 70% EGFP expression and mRNA levels. None of the CMV clones retained over 70% of EGFP expression or mRNA levels, despite six clones still retaining more than 70% of their gene copies. The relative changes in EGFP expression levels correlated to the changes in EGFP mRNA levels for all clones. These results suggest that the SV40 promoter is more resistant to transcriptional silencing than the CMV promoter.

Discussion

Previous studies indicated that endogenous mammalian promoters could provide more stable gene expression than the viral promoters. Interestingly, we found that the SV40 promoter gave the highest expression stability in stably transfected CHO cell clones, while none of eighteen clones generated using EF1 α and CMV were stable. Analysis of the changes in EGFP gene copies and mRNA levels in SV40 and CMV clones suggested that the SV40 promoter was more resistant to transcriptional silencing. DNA methylation is one of mechanisms causing transcriptional silencing. Both CMV and EF1 α are crowded with CG dinucleotides, containing 31 and 141 CGs respectively, while the SV40 promoter has only 6 CG dinucleotides (Supplementary Fig. 2). The lower number of CGs may be one reason why the SV40 promoter is more resistant to transcriptional silencing due to DNA methylation. Another possible reason is that SV40 contains two SP1 transcription factor binding sites (Supplementary Fig. 2), which was previously proposed to inhibit DNA methylation (46). [Mutation of CGs in the retroviral vector long terminal repeat had repressed promoter silencing in embryonic stem cells](#) (47). [We attempted to remove the CGs within the SV40 promoter to enhance its resistance to transcriptional silencing. The CG-free SV40 promoter had lower expression than the wild-type SV40 in transient transfections and failed to survive the selection process for stable expression \(unpublished data\). The mutant SV40 we generated was likely too weak for our application and further efforts are currently underway to develop CG-free SV40 promoters without compromising the gene expression levels.](#)

Addition of either cMAR or iMAR enhanced the stable gene expression level for the SV40 promoter but had no effect on EF1 α and CMV. Addition of the two MAR elements did not enhance long-term expression stability for any promoters. Among all the tested vectors, SV40+cMAR and SV40+iMAR were the best for obtaining both high expression level and stability in CHO cells. All previous studies based on SV40 promoter demonstrated that MARs were able to enhance stable gene expression (27, 32-37, 48). Fewer studies of MAR were performed on the CMV promoter and the results were conflicting (31, 35, 49). Our results were consistent with the study done by Lonza in which the inclusion of cMAR into their glutamine synthetase (GS) expression vector, which is commonly used for mammalian expression in the industry, did not enhance monoclonal antibody expression levels (49). A recent study demonstrated that the activity of MAR elements were most pronounced for the chromosomal positions with low expression potential but had negligible effects in the case of highly active chromosomal sites (5). We speculate that effect of MAR on transgene expression may be less effective when strong promoters and/or vectors with more stringent selection of high expression levels are used. Inclusion of MARs to a previously well optimized vector design would likely yield less benefit compared to adding to a simpler basic vector. Further studies are needed to investigate and identify the optimal plasmid vector design and context to maximize the effect of including MAR elements to improve expression level and stability.

References

1. Ho, S. C. L., Tong, Y. W. and Yang, Y. S. (2013) Generation of monoclonal antibody-producing mammalian cell lines. *Pharm. Bioprocess.* 1, 71-87.
2. Wurm, F. M. (2004) Production of recombinant protein therapeutics in cultivated mammalian cells. *Nat. Biotechnol.* 22, 1393-1398.
3. Barnes, L. M., Bentley, C. M. and Dickson, A. J. (2001) Characterization of the stability of recombinant protein production in the GS-NS0 expression system. *Biotechnol. Bioeng.* 73, 261-270.
4. Festenstein, R., Tolaini, M., Corbella, P., Mamalaki, C., Parrington, J., Fox, M., Miliou, A., Jones, M. and Kioussis, D. (1996) Locus control region function and heterochromatin-induced position effect variegation. *Science* 271, 1123-1125.
5. Goetze, S., Baer, A., Winkelmann, S., Nehlsen, K., Seibler, J., Maass, K. and Bode, J. (2005) Performance of genomic bordering elements at predefined genomic loci. *Mol. Cell Biol.* 25, 2260-2272.
6. Jordan, A., Defechereux, P. and Verdin, E. (2001) The site of HIV-1 integration in the human genome determines basal transcriptional activity and response to Tat transactivation. *Embo J.* 20, 1726-1738.
7. Rita Costa, A., Elisa Rodrigues, M., Henriques, M., Azeredo, J. and Oliveira, R. (2010) Guidelines to cell engineering for monoclonal antibody production. *Eur. J. Pharm. Biopharm.* 74, 127-138.
8. Dorai, H., Corisdeo, S., Ellis, D., Kinney, C., Chomo, M., Hawley-Nelson, P., Moore, G., Betenbaugh, M. J. and Ganguly, S. (2012) Early prediction of instability of chinese

- hamster ovary cell lines expressing recombinant antibodies and antibody-fusion proteins. *Biotechnol. Bioeng.* 109, 1016-1030.
9. Fann, C. H., Guirgis, F., Chen, G., Lao, M. S. and Piret, J. M. (2000) Limitations to the amplification and stability of human tissue-type plasminogen activator expression by Chinese hamster ovary cells. *Biotechnol. Bioeng.* 69, 204-212.
 10. He, L., Winterrowd, C., Kadura, I. and Frye, C. (2012) Transgene copy number distribution profiles in recombinant CHO cell lines revealed by single cell analyses. *Biotechnol. Bioeng.* 109, 1713-1722.
 11. Jun, S. C., Kim, M. S., Hong, H. J. and Lee, G. M. (2006) Limitations to the development of humanized antibody producing Chinese hamster ovary cells using glutamine synthetase-mediated gene amplification. *Biotechnol. Progr.* 22, 770-780.
 12. Kim, M., O'Callaghan, P. M., Droms, K. A. and James, D. C. (2011) A mechanistic understanding of production instability in CHO cell lines expressing recombinant monoclonal antibodies. *Biotechnol. Bioeng.* 108, 2434-2446.
 13. Kim, S. J., Kim, N. S., Ryu, C. J., Hong, H. J. and Lee, G. M. (1998) Characterization of chimeric antibody producing CHO cells in the course of dihydrofolate reductase-mediated gene amplification and their stability in the absence of selective pressure. *Biotechnol. Bioeng.* 58, 73-84.
 14. Osterlehner, A., Simmeth, S. and Goepfert, U. (2011) Promoter Methylation and Transgene Copy Numbers Predict Unstable Protein Production in Recombinant Chinese Hamster Ovary Cell Lines. *Biotechnol. Bioeng.* 108, 2670-2681.

15. Strutzenberger, K., Borth, N., Kunert, R., Steinfellner, W. and Katinger, H. (1999) Changes during subclone development and ageing of human antibody-producing recombinant CHO cells. *J. Biotechnol.* 69, 215-226.
16. Yang, Y., Mariati, Chusainow, J. and Yap, M. G. (2010) DNA methylation contributes to loss in productivity of monoclonal antibody-producing CHO cell lines. *J. Biotechnol.* 147, 180-185.
17. Deer, J. R. and Allison, D. S. (2004) High-level expression of proteins in mammalian cells using transcription regulatory sequences from the Chinese hamster EF-1 alpha gene. *Biotechnol. Bioeng. Prog.* 20, 880-889.
18. Paredes, V., Park, J. S., Jeong, Y., Yoon, J. and Baek, K. (2013) Unstable expression of recombinant antibody during long-term culture of CHO cells is accompanied by histone H3 hypoacetylation. *Biotechnol. Lett.* 35, 987-993.
19. Galbete, J. L., Bucetaz, M. and Mermoud, N. (2009) MAR elements regulate the probability of epigenetic switching between active and inactive gene expression. *Mol. Biosys.* 5, 143-150.
20. Byun, H. M., Suh, D. C., Jeong, Y. S., Wee, H. S., Kim, J. M., Kim, W. K., Ko, J. J., Kim, J. S., Lee, Y. B. and Oh, Y. K. (2005) Plasmid vectors harboring cellular promoters can induce prolonged gene expression in hematopoietic and mesenchymal progenitor cells. *Biochem. Biophys. Res. Commun.* 332, 518-523.
21. Damdindorj, L., Karnan, S., Ota, A., Takahashi, M., Konishi, Y., Hossain, E., Hosokawa, Y. and Konishi, H. (2012) Assessment of the long-term transcriptional activity of a 550-bp-long human beta-actin promoter region. *Plasmid* 68, 195-200.

22. Gill, D. R., Smyth, S. E., Goddard, C. A., Pringle, I. A., Higgins, C. F., Colledge, W. H. and Hyde, S. C. (2001) Increased persistence of lung gene expression using plasmids containing the ubiquitin C or elongation factor 1 alpha promoter. *Gene Ther.* **8**, 1539-1546.
23. Gopalkrishnan, R. V., Christiansen, K. A., Goldstein, N. I., DePinho, R. A. and Fisher, P. B. (1999) Use of the human EF-1 alpha promoter for expression can significantly increase success in establishing stable cell lines with consistent expression: a study using the tetracycline-inducible system in human cancer cells. *Nucleic Acids Res.* **27**, 4775-4782.
24. Teschendorf, C., Warrington, K. H., Siemann, D. W. and Muzyczka, N. (2002) Comparison of the EF-1 alpha and the CMV promoter for engineering stable tumor cell lines using recombinant adeno-associated virus. *Anticancer Res.* **22**, 3325-3330.
25. Chan, K. K. K., Wu, S. M., Nissom, P. M., Oh, S. K. W. and Choo, A. B. H. (2008) Generation of high-level stable transgene expressing human embryonic stem cell lines using Chinese hamster elongation factor-1 alpha promoter system. *Stem Cells Dev.* **17**, 825-836.
26. Bode, J., Benham, C., Knopp, A. and Mielke, C. (2000) Transcriptional augmentation: Modulation of gene expression by Scaffold/Matrix-Attached regions (S/MAR elements). *Crit. Rev. Eukar. Gene* **10**, 73-90.
27. Harraghy, N., Gaussin, A. and Mermoud, N. (2008) Sustained transgene expression using MAR elements. *Curr. Gene Ther.* **8**, 353-366.

28. Araki, Y., Hamafuji, T., Noguchi, C. and Shimizu, N. (2012) Efficient Recombinant Production in Mammalian Cells Using a Novel IR/MAR Gene Amplification Method. *PLoS ONE* 7, e41787.
29. Kim, J. D., Yoon, Y., Hwang, H. Y., Park, J. S., Yu, S., Lee, J., Baek, K. and Yoon, J. (2005) Efficient selection of stable Chinese hamster ovary (CHO) cell lines for expression of recombinant proteins by using human interferon beta SAR element. *Biotechnol. Progr.* 21, 933-937.
30. Kim, J. M., Kim, J. S., Park, D. H., Kang, H. S., Yoon, J., Baek, K. and Yoon, Y. (2004) Improved recombinant gene expression in CHO cells using matrix attachment regions. *J. Biotechnol.* 107, 95-105.
31. Otte, A. P., Kwaks, T. H. J., van Blokland, R. J. M., Sewalt, R. G. A. B., Verhees, J., Klaren, V. N. A., Siersma, T. K., Korse, H. W. M., Teunissen, N. C., Botschuijver, S., van Mer, C. and Man, S. Y. (2007) Various expression-augmenting DNA elements benefit from STAR-select, a novel high stringency selection system for protein expression. *Biotechnol. Progr.* 23, 801-807.
32. Wang, F., Wang, T. Y., Tang, Y. Y., Zhang, J. H. and Yang, X. J. (2012) Different matrix attachment regions flanking a transgene effectively enhance gene expression in stably transfected Chinese hamster ovary cells. *Gene* 500, 59-62.
33. Wang, T. Y., Yang, R., Qin, C. A., Wang, L. and Yang, X. J. (2008) Enhanced expression of transgene in CHO cells using matrix attachment region. *Cell Biol. Int.* 32, 1279-1283.

34. Wang, T. Y., Zhang, J. H., Jing, C. Q., Yang, X. J. and Lin, J. T. (2010) Positional effects of the matrix attachment region on transgene expression in stably transfected CHO cells. *Cell Biol. Int.* 34, 141-145.
35. Zahn-Zabal, M., Kobr, M., Girod, P. A., Imhof, M., Chatellard, P., de Jesus, M., Wurm, F. and Mermod, N. (2001) Development of stable cell lines for production or regulated expression using matrix attachment regions. *J. Biotechnol.* 87, 29-42.
36. Girod, P. A., Nguyen, D. Q., Calabrese, D., Puttini, S., Grandjean, M., Martinet, D., Regamey, A., Saugy, D., Beckmann, J. S., Bucher, P. and Mermod, N. (2007) Genome-wide prediction of matrix attachment regions that increase gene expression in mammalian cells. *Nat. Methods* 4, 747-753.
37. Girod, P. A., Zahn-Zabal, M. and Mermod, N. (2005) Use of the chicken lysozyme 5' matrix attachment region to generate high producer CHO cell lines. *Biotechnol. Bioeng.* 91, 1-11.
38. Dang, Q., Auten, J. and Plavec, I. (2000) Human beta interferon scaffold attachment region inhibits de novo methylation and confers long-term, copy number-dependent expression to a retroviral vector. *J. Virol.* 74, 2671-2678.
39. Moreno, R., Martinez, I., Petriz, J., Nadal, M., Tintore, X., Gonzalez, J. R., Gratacos, E. and Aran, J. M. (2011) The beta-Interferon Scaffold Attachment Region Confers High-Level Transgene Expression and Avoids Extinction by Epigenetic Modifications of Integrated Provirus in Adipose Tissue-Derived Human Mesenchymal Stem Cells. *Tissue Eng. Part C-Methods* 17, 275-287.

40. Ho, S. C. L., Bardor, M., Feng, H. T., Mariati, Tong, Y. W., Song, Z. W., Yap, M. G. S. and Yang, Y. S. (2012) IRES-mediated Tricistronic vectors for enhancing generation of high monoclonal antibody expressing CHO cell lines. *J. Biotechnol.* 157, 130-139.
41. Piechaczek, C., Fetzer, C., Baiker, A., Bode, J. and Lipps, H. J. (1999) A vector based on the SV40 origin of replication and chromosomal S/MARs replicates episomally in CHO cells. *Nucleic Acids Res.* 27, 426-428.
42. Chusainow, J., Yang, Y. S., Yeo, J. H., Toh, P. C., Asvadi, P., Wong, N. S. and Yap, M. G. (2009) A study of monoclonal antibody-producing CHO cell lines: what makes a stable high producer? *Biotechnol. Bioeng.* 102, 1182-1196.
43. Sautter, K. and Enenkel, B. (2005) Selection of high-producing CHO cells using NPT selection marker with reduced enzyme activity. *Biotechnol. Bioeng.* 89, 530-538.
44. Ng, S. K., Wang, D. I. C. and Yap, M. G. S. (2007) Application of destabilizing sequences on selection marker for improved recombinant protein productivity in CHO-DG44. *Metab. Eng.* 9, 304-316.
45. Bailey, L. A., Hatton, D., Field, R. and Dickson, A. J. (2012) Determination of Chinese hamster ovary cell line stability and recombinant antibody expression during long-term culture. *Biotechnol. Bioeng.* 109, 2093-2103.
46. Senigl, F., Plachy, J. and Hejnar, J. (2008) The core element of a CpG island protects avian sarcoma and leukosis virus-derived vectors from transcriptional silencing. *J. Virol.* 82, 7818-7827.
47. Swindle, C. S., Kim, H. G. and Klug, C. A. (2004) Mutation of CpGs in the murine stem cell virus retroviral vector long terminal repeat represses silencing in embryonic stem cells. *J. Biol. Chem.* 279, 34-41.

48. Harraghy, N., Regamey, A., Girod, P. A. and Mermoud, N. (2011) Identification of a potent MAR element from the mouse genome and assessment of its activity in stable and transient transfections. *J. Biotechnol.* 154, 11-20.
49. Kalwy, S. (2005) Towards stronger gene expression - a promoter's tale, in Presentation done at: 19th European Society for Animal Cell Technology (ESACT) meeting, Harrogate, England.

Figure captions

Fig. 1. Schematic representation of vectors for evaluating promoters and the impact of MAR on recombinant protein expression level and stability in CHO cells. SV40, simian virus 40 promoter; EF1 α , Chinese hamster elongation factor-1 α promoter; CMV, human cytomegalovirus IE gene promoter; iMAR, human interferon β matrix-attachment region (MAR) element; cMAR, chicken lysozyme MAR element; IRESatt, attenuated encephalomyocarditis virus (EMCV) internal ribosome entry site with reduced translation efficiency; SpA, simian virus 40 early polyadenylation signal; EGFP, enhanced green fluorescence protein cDNA; mNPT, mutated neomycin phosphotransferase cDNA with amino acid D at 261 changed to G (43).

Fig. 2. Use of different promoters and the impact of MAR on gene expression level in stably transfected pools (A) and clones (B). Three stably transfected pools were generated for each vector. To characterize each stably transfected pool, 2 mL of cultures at a density of 2×10^5 cells/mL were seeded into each well of 6-well plates. Cells were collected at day 3 and measured for the EGFP mean fluorescence intensity (MFI) with the FACS Calibur and EGFP gene copies and mRNA levels using quantitative real-time PCR (qRT-PCR). Results in Fig. 2A were presented as the EGFP MFI (black bar), mRNA levels (gray bar), and gene copies (white bar) normalized to those from the SV40 promoter. Each value represents the average and standard deviation of three independent stably transfected pools. Six clones each were isolated from three separately transfected pools for a total of eighteen clones for each vector. Each dot in Fig. 2B represents the MFI of each clone normalized to the average MFI of the eighteen clones generated using the SV40 promoter. Horizontal bars signify the average value of the 18 clones for each

vector. Mean values significantly different (two-tailed Student's t-test) between the vectors containing MAR and the vector without MAR for each promoter indicated by asterisks “*” ($p < 0.05$).

Fig. 3. Retention of EGFP expression in stably transfected clones generated using different vectors. Each dot represents the percentage of retention of GFP expression for one clone. The horizontal bars signify the average value of 18 clones for each vector. Mean values significantly different (two-tailed Student's t-test) between the vectors containing MAR and the vector without MAR for each promoter indicated by asterisks “*” ($p < 0.05$).

Fig. 4. Relative changes in EGFP expression (black bar), mRNA levels (gray bar) and gene copies (white bar) in SV40 and CMV clones during stability testing. The relative changes in EGFP expression, mRNA levels and gene copies were calculated as the ratio of EGFP MFI, mRNA levels and gene copies of a clone measured at week 8 to the starting level for the same clone measured at week 0. Each value represents the average of two measurements.

Fig. S1. Relationship between the normalized EGFP expression level before stability testing and retention of EGFP expression at the end of stability testing of clones generated using different promoters and combinations of promoter and MARs.

Fig. S2. DNA sequences of the SV40, EF1 α and CMV promoters. The CG dinucleotide is highlighted in red. The underlined sequences in the SV40 promoter are SP1 transcription factor binding sites.

Fig. 1.

Fig. 2.
(A)

(B)

Fig. 3.

Fig. 4
(A)

(B)

Fig. S1
(A)

(B)

(C)

(D)

(E)

(F)

(G)

(H)

(I)

Fig. S2.

(A)

GTGGAATGTGTGTCAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCAGAAGTA
TGCAAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTCCCCAGGCTCCCCA
GCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCCT
AACTCCGCCCCATCCCGCCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCT
GACTAATTTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCTGCCTCTGAGCTATTCCAG
AAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTTGCAAAAAGCT

(B)

GGATGGCGGGGCTGACGTGGGAGGTGGCCTCCACGGGAAGGGACACC CGGATCTCGAC
ACAGCCTTGGCAGTGGAGTCAGGAAGGGTAGGACAGATTCTGGA CGCCCTCTTGGCCAG
TCCTCACCGCCCCACCCC CGATGGAGCCGAGAGTAATTCATACAAAAGGAGGGATCGCC
TTCGCCCCCTGGGAATCCCAGGGACCGT CGCTAAATTCTGGCCGCGCTCCCAGCCCGGAA
CCGCTGTGCCCGCCCAGCGCGGCGGGAGGAGCCTGCGCCTAGGGCGGATCGCGGGTGG
CGGGAGAGCACAAAGCCACAGTCCCCGCGCGGTGGGGGAGGGGCGCGCTGAGCGGGGGCC
CGGGAGCCAGCGCGGGGCAAACCTGGGAAAGTGGTGTGTGTGTGCTGGCTCCGCCCTCTTC
CCGAGGGTGGGGGAGAA CGGTATAAAAGTGCGGTAGTCCGCGTTGGA CGTTCTTTTTCGC
AACCGGGTTTGC CGTCAGAACCGCAGGTGAGTGGCGGGTGTGGCCTCCGCGGGGCCCGGGCT
CCCTCCTTTGAGCGGGGTGGGACCGCCGTGCGGGTGTGTGTGTGCGCGGGCTTCTCTGCGA
GCGTTCCCGCCCTGGATGGCGGGCTGTGCGGGAGGGCGAGGGGGGAGGCCTGGCGGCG
GCCCCGGGAGCCTCGCCTCGTGTGGGCGTGAGGCCTAGCGTGGCTTCGCCCCCGCGCG
TGCCACCGCGGC CGCGCTTTGCTGTCTGCCCGGCTGCCCTCGATTGCCCTGCCCGCGGCC
CGGGCCAACAAAGGGAGGGCGTGAGCTGGCTGGTAGGGAGCCCCCGTAGTCCGCATGTC
GGGCAGGGAGAGCGGCAGCAGTGGGGGGGGGACCGGGCCCGCCCGTCCCGCAGCACAT
GTC CGACCGCCCGCTGGA CGGGTAGCGGCCTGTGTCTCTGATAAGGCGGC CGGGCGGTGGG
TTTTAGATGCCCGGGTT CAGGTGGCCCCCGGGTCCCGGCCCGGTCTGGCCAGTACCCCGTA
GTGGCTTAGCTCCGAGGAGGGCGAGCCCGCCCGCCCGGCACCAGTTGCGTGCGCGGAAA
GATGGCCGCTCCCGGGCCCTGTAGCAAGGAGCTCAAAATGGAGGACCGCGGCAGCCCGGC
GGAGCGGGGCGGGTGAGTACCCACACAAAGGAAGAGGGCCTTGCCCCCTCGCGCGCCCGC
TGCTTCCTGTGACCCCGTGGTGTACCGGCCGCACTTCAGTCACCCCGGGCGCTCTTTCG
GAGCACCGCTGGCCTCCGCTGGGGGAGGGGATCTGTCTAATGGCGTTGGAGTTTGCTCA
CATTTGGTGGGTGGAGACTGTAGCCAGGCCAGCCTGGCCATGGAAGTAATTCTTGGAAT
TTGCCCATTTTGGAGTTTGGAGCGAAGCTGATTGACAAAGCTGCTTAGCCGTTCAAAGGT
ATTCTT CGAACTTTTTTTTTTAAGGTGTTGTGAAAACCACCGCTAATTCAAATCCAAC

(C)

TTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTA CGGGGTCATTAGTTCATA
GCCCATATATGGAGTTC CGCGTTACATAACTTA CGGTAAATGGCCCGCCTGGCTGACCG
CCCAACCGACCCC CGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAA CGCCAAT
AGGGACTTTCCATTGA CGTCAATGGGTGGAGTATTTA CGGTAAACTGCCCCTTGGCAG
TACATCAAGTGTATCATATGCCAAGTA CGCCCCCTATTGA CGTCAATGA CGGTAAATGG
CCCGCCTGGCATTATGCCAGTACATGACCTTATGGGACTTTCTACTTGGCAGTACAT
CTA CGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCAATGGGC
GTGGATAGCGGTTTGACTCACGGGGATTTC AAGTCTCCACCCCATTGACGTCAATGGG

AGTTTGTTTTGGCACCAAAATCAACGGGACTTTCCAAAATGTGTAACAACCTCGCCCC
ATTGACGCAAATGGGCGGTAGGCGTGTAAGGTGGGAGGTCTATATAAGCAGAGCTC