Sulfoxidation of alkenes and alkynes with NFSI as a radical initiator and selective oxidant

Zhang, Yuexia; Wong, Zeng Rong; Wu, Xingxing; Lauw, Sherman Jun Liang; Huang, Xuan; Webster, Richard David; Chi, Yonggui Robin

2017

Zhang, Y., Wong, Z. R., Wu, X., Lauw, S. J. L., Huang, X., Webster, R. D., et al. (2017).
Sulfoxidation of alkenes and alkynes with NFSI as a radical initiator and selective oxidant.
Chemical Communications, 53(1), 184-187.

<https://hdl.handle.net/10356/82060>

<https://doi.org/10.1039/C6CC08631D>

© 2017 The Royal Society of Chemistry. This is the author created version of a work that has been peer reviewed and accepted for publication by Chemical Communications, The Royal Society of Chemistry. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [<http://dx.doi.org/10.1039/C6CC08631D>].

Downloaded on 26 Apr 2023 08:55:27 SGT

Journal Name

COMMUNICATION

Sulfoxidation of Alkenes and Alkynes with NFSI as a Radical Initiator and Selective Oxidant

Received 00th January 20xx,
Accepted 00th January 20xx

Yuexia Zhang,^{a†} Zeng Rong Wong,^{a†} Xingxing Wu,^a Sherman J. L. Lauw,^a Xuan Huang,^a Richard D. Webster^a and Yonggui Robin Chi^{*ab}

DOI: 10.1039/x0xx00000x

www.rsc.org/

Sulfoxides are important functional molecules. We develop a short-route (one-pot) synthesis of this class of molecules by reacting thiols with alkenes or alkynes under mild and metal-free conditions. N-fluorobenzenesulfonimide (NFSI) is used to play dual roles: as a radical initiator for a thiol-ene/-yne reaction to form sulfide adducts, and as efficient oxidant for conversion of the sulfides formed in situ to sulfoxides. Over-oxidation of the sulfoxides to sulfones is avoided in our approach.

The sulfoxide moiety is a functional group often found in pharmaceuticals and natural products, such as Esomeprazole,¹ Armodafinil² and Alliin.³ Sulfoxides are also used as ligands for metal catalysts with applications in a large set of reactions, including Pummerer⁴ and Mislow-Evans rearrangements.⁵ To date, the synthesis of sulfoxides typically involves oxidation of sulfides using peroxides⁶ and hypervalent iodine reagent⁷ as the oxidants with the assistance of transition metal catalysts.⁸ One challenge of this otherwise widely used two-step protocol lies with over-oxidation of sulfoxides to sulfones.⁶ Here we report a new method that selectively converts alkenes and alkynes to sulfoxides in a one-pot metal-free operation (Figure 1). N-fluorobenzenesulfonimide (NFSI), a shelf-stable reagent,⁹ is used as a radical initiator and a selective oxidant. No over-oxidation of sulfoxides to sulfones was observed (Figure 1).

Notably, Yadav¹⁰ and Lei¹¹ have recently reported reactions of alkenes and thiols under dioxygen to form β -keto sulfoxides and β -oxy sulfoxides. In their reactions, oxidation of the alkene carbon by oxygen occurred concurrently. The NFSI reagent used in our reactions has often been employed as electrophilic

fluorinating agent,¹² aminating reagent,¹³ or a combined amino and fluorine source.¹⁴ In radical reactions (SET reactions), it can generate aminyl radical for Cu-catalyzed radical amination of olefins, as reported by Kanai,¹⁵ Zhang¹⁶ and Studer.¹⁷ Zhang and co-workers also developed radical aminofluorination of styrenes using NFSI.¹⁸ In addition, Ritter reported Pd-catalyzed direct radical arene amination with NFSI,¹⁹ and Studer developed radical aminooxygenation of alkenes with NFSI in combination with TEMPONa.²⁰

Figure 1 Our strategy of NFSI-enabled selective one-pot access to sulfoxides from alkenes/alkynes and thiols

Our initial design was to combine NFSI and thiol to generate thiyl and aminyl radicals for aminosulfidation of alkenes. We first studied reactions between alkene **1a** and thiol **2a** with NFSI as a potential radical generator in the presence of K_2CO_3 or K_3PO_4 as the base (Table 1, entry 1). The expected alkene aminosulfidation adduct was not formed. Instead, we found alkene sulfoxidation adduct **3a** as the product in 11% yield (entry 1). Additional evaluation of the base effects (Table 1, entries 2-6) indicated that 1,8-diazabicycloundec-7-ene (DBU) performed the best, giving **3a** in 72% yield (Table 1, entry 6). Several other solvents studied here did not work as well as toluene (Table 1, entries 7-10). The yield of **3a** (based on **1a**) could be further improved by increasing the loadings of thiol **2a** and NFSI (Table 1, entry 11).

^a Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371 (Singapore). E-mail: robinchi@ntu.edu.sg

^b Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering, Key Laboratory of Green Pesticide and Agricultural Bioengineering, Ministry of Education, Guizhou University, Huaxi District, Guiyang 550025 (China).

[†] These authors contributed equally to this work.

Electronic Supplementary Information (ESI) available: Experimental procedures, mechanistic details, spectroscopic data and copies of ¹H and ¹³C NMR spectra. See DOI: 10.1039/x0xx00000x

COMMUNICATION

Journal Name

Table 1 Screening of reaction conditions for the reaction of **1a** with **2a** in the presence of NFSI.^a

Entry	Base	Solvent	Yield [%] ^b
1	K ₂ CO ₃ or K ₃ PO ₄	toluene	11
2	KOMe	toluene	23
3	NaOtBu	toluene	29
4	DIPEA	toluene	43
5	DABCO	toluene	66
6	DBU	toluene	72
7	DBU	THF	52
8	DBU	dioxane	31
9	DBU	CH ₃ CN	26
10	DBU	EA	48
11 ^c	DBU	toluene	87

^a Conditions: **1a** (0.2 mmol, 1.0 equiv.), **2a** (0.24 mmol, 1.2 equiv.), NFSI (130 mol%) and base (50 mol%) in 3.0 mL of solvent at rt for 2~5 h; ^b Isolated yield based on **1a**. ^c **2a** (0.4 mmol, 2.0 equiv.), NFSI (0.36 mmol, 1.8 equiv.).

With an acceptable condition (Table 1, entry 11) on hand, we proceeded to explore the scope of the reaction (Table 2). We first examined the reaction between various alkenes and thiol **2a** (**3a-3p**). Styrenes with different substituents or substitution patterns on the phenyl ring worked effectively (**3a-3l**). Internal alkenes, such as indene (**3m**), 1,2-dihydronaphthalene (**3n**), *trans*- β -methylstyrene (**3o**) were also effective substrates. Notably, no sulfoxidation products were obtained when aliphatic alkenes were used. Substituents (e.g., CH₃ and Br) could be placed on the phenyl unit of thiol **2a** (**3q**, **3r**), while 2-naphthyl thiol (**3s**) was also tolerated.

We next found that alkynes could be used as substrates (to replace alkenes) under the same condition, affording unsaturated sulfoxides as the products (Table 3). The reactions proceeded smoothly for aryl alkynes with either electron-withdrawing or electron-donating substituents present on the aromatic ring (**4a-4e**) to afford the desired products (**5a-5e**) in moderate to excellent yields. Alkyl alkynes (**4f**, **4i-j**) and alkynes bearing an ester (**4g**) or silyl (**4h**) moiety were also effective substrates.²¹ The structures of two vinylsulfoxide products (**E-5a**, **E-5d**) were unambiguously assigned via X-ray crystallography.

Table 2 Substrate scope for reactions of alkenes with thiols in the presence of NFSI.^a

^a Conditions as in Table 1, entry 11; yields (after SiO₂ chromatography purification) were based on olefin **1**. ^bDiastereoselective ratio (d.r.) was determined via ¹H NMR analysis of the product mixture.

The proposed reaction pathway is illustrated in Scheme 1. Initially, removal of one electron from thiol **2a** by oxidant NFSI^{12a,12b} generates radical cation **A** that can be subsequently transformed to thiyl radical **B** after a proton transfer (Scheme 1a). Thereafter, addition of radical **B** to alkene **1** forms radical intermediate **C**. A hydrogen atom transfer (HAT) from a molecule of **2a** converts **C** to sulfide **D** with the regeneration of the thiyl radical intermediate **B** which returns back into the propagation step (scheme 1b). Finally, the sulfide **D** is oxidized to sulfoxide **3** by NFSI. One possible pathway (see SI for another possibility) for the oxidation of sulfide **D** involves nucleophilic addition of the sulfide to the electrophilic fluorine atom of NFSI to afford intermediate **E** (Scheme 1c). Substitution of the fluorine atom of **E** by water forms **G** that then transforms to sulfoxide **3** after a proton transfer process.

Table 3 Substrate scope for reactions of alkynes with thiols in the presence of NFSI.^a

^a Alkyne (0.2 mmol), **2a** (0.4 mmol), NFSI (0.36 mmol) and DBU (50 mol%), in toluene (3 mL) at rt for 16h; yields (after SiO₂ chromatography purification) were based on alkyne. ^b The ratio of E/Z was determined by the weight of isolated E and Z products.

a) Radical initiation

b) Chain propagation

c) Oxidation

Scheme 1 Proposed reaction pathway.

To investigate the reaction mechanism, we performed multiple experiments (Scheme 2). The addition of TEMPO inhibited the formation of **3a** (Scheme 2a). In the presence of 150 mol% TEMPO, the formation of sulfoxide product **3a** was completely suppressed. The aminooxygenation adduct **6** previously reported by Studer²⁰ was observed in about 5% yield.

Adducts **7**, **8** and **9** (89% of combined yields based on thiol **2a**) were previously reported by Greci and co-workers in a study on the reaction between **2a** and TEMPO.²² The use of diphenyl disulfide **7** in place of thiophenol **2a** did not lead to desired product **3a** (Scheme 2b), suggesting that under the oxidative condition of NFSI the disulfide adduct did not convert to thiol. The oxygen atom in the sulfoxide product came from the trace amount of H₂O present in the solvent, as suggested by an ¹⁸O isotope labelling experiment (Scheme 2c). Additional experiments by adjusting the loadings of NFSI (Scheme 2d) suggest that both the thiol-ene reaction and oxidation reaction are enabled by NFSI. When 30 mol % of NFSI was added, only sulfide adduct **10** was observed. Sulfide **10** could then be transformed under standard conditions to sulfoxide **3a**.²³ These experiments (Scheme 2a, b) support the formation of radical intermediate as proposed in Scheme 1a and 1b. The results (Scheme 2c, d) showed that the formation of sulfide (**D**, scheme 1c) and its oxidation occurs in a stepwise process.

(a) TEMPO Trapping Experiment

(b) PhSSPh Used Instead of PhSH

(c) ¹⁸O Isotope Labelling Experiment

(d) Formation and Oxidation of Sulfide Enabled by NFSI

Scheme 2 Experiments to probe mechanism.

In summary, we have developed a facile synthesis of sulfoxides from alkenes and alkynes in the presence of NFSI. Sulfoxides are useful structural moieties for construction of natural products and other functional molecules. Our approach provides a metal-free and concise synthesis of this class of molecules. The substrates and reagents used in our approach are either inexpensive or readily available with reasonable cost. NFSI is used as a radical initiator (to initiate the sulfide formation) and a selective oxidant (to transform sulfides to sulfoxides). The oxygen atom in the sulfoxide products was determined to originate from trace water in the reaction mixture, providing a simple way for the incorporation of ^{18}O isotope to sulfoxide-containing molecules. Further development of related radical reactions, including catalytic versions mediated by carbene organic catalysts, is in progress.

Acknowledgements

We thank Dr. Yongxin Li (NTU) and Dr. Rakesh Ganguly for assistance with X-ray structure analysis; We acknowledge financial support by the Singapore National Research Foundation (NRF-NRFI2016-06), the Ministry of Education of Singapore (MOE2013-T2-2-003), Nanyang Technological University (NTU, Singapore), China's Ministry of Education, National Key program for Basic Research (No. 2010CB 126105), Thousand Talent Plan (700059143302), National Natural Science Foundation of China (No. 21132003; No. 21472028). Guizhou Province Returned Oversea Student Science and Technology Activity Program, Science and Technology of Guizhou Province, and Guizhou University.

Notes and references

- C. M. Spencer, D. Faulds, *Drugs*. 2000, **60**, 321-329.
- K. P. Garnock-Jones, S. Dhillon, L. J. Scott, *CNS Drugs*. 2009, **23**, 793-803.
- A. Stoll, E. Seebeck, *Advances in Enzymology and Related Areas of Molecular Biology, Volume 11*. 2006, 377-400.
- For selected reviews on Pummer rearrangements, see (a) S. K. Bur, A. Padwa, *Chem. Rev. (Washington, DC, U. S.)* 2004, **104**, 2401-2432; (b) K. S. Feldman, *Tetrahedron* 2006, **62**, 5003-5034; (c) L. H. Smith, S. C. Coote, H. F. Sneddon, D. J. Procter, *Angew. Chem. Int. Ed.* 2010, **49**, 5832-5844.
- C. M. Rojas, in *Molecular Rearrangements in Organic Synthesis*, John Wiley & Sons, Inc, 2015, pp. 569-626.
- See the review: K. Kaczorowska, Z. Kolarska, K. Mitka, P. Kowalski, *Tetrahedron* 2005, **61**, 8315-8327.
- (a) M. Xia, Z. C. Chen, *Synth. Commun.* 1997, **27**, 1315-1320; (b) B. Yu, C. X. Guo, C. L. Zhong, Z. F. Diao, L. N. He, *Tetrahedron Lett.* 2014, **55**, 1818-1821.
- (a) M. Matteucci, G. Bhalay, M. Bradley, *Org. Lett.* 2003, **5**, 235-237; (b) W. Dai, J. Li, B. Chen, G. Li, Y. Lv, L. Wang, S. Gao, *Org. Lett.* 2013, **15**, 5658-5661; (c) X. F. Wu, *Tetrahedron Lett.* 2012, **53**, 4328-4331; (d) M. Bagherzadeh, M. M. Haghdoust, A. Shahbazirad, *J. Coord. Chem.* 2012, **65**, 591-601.
- X. Yang, T. Wu, R. J. Phipps, F. D. Toste, *Chem. Rev. (Washington, DC, U. S.)*. 2015, **115**, 826-870.
- T. Keshari, V. K. Yadav, V. P. Srivastava, L. D. S. Yadav, *Green. Chem.* 2014, **16**, 3986-3992.
- H. Wang, Q. Lu, C. Qian, C. Liu, W. Liu, K. Chen, A. Lei, *Angew. Chem. Int. Ed.* 2016, **55**, 1094-1097.
- (a) X. Dong, W. Yang, W. Hu, J. Sun, *Angew. Chem. Int. Ed.* 2015, **54**, 660-663; (b) N. Kuhl, M. N. Hopkinson, L. Wencel-Delord, F. Glorius, *Angew. Chem. Int. Ed.* 2012, **51**, 10236-10254; (c) D. D. Steiner, N. Mase, C. F. Barbas, 3rd, *Angew. Chem. Int. Ed.* 2005, **44**, 3706-3710; (d) T. D. Beeson, D. W. C. MacMillan, *J. Am. Chem. Soc.* 2005, **127**, 8826-8828.
- (a) K. Sun, Y. Li, T. Xiong, J. Zhang, Q. Zhang, *J. Am. Chem. Soc.* 2011, **133**, 1694-1697; (b) J. Trenner, C. Depken, T. Weber, A. Breder, *Angew. Chem. Int. Ed.* 2013, **52**, 8952-8956; (c) T. Xiong, Y. Li, Y. Lv, Q. Zhang, *Chem. Commun.* 2010, **46**, 6831-6833.
- (a) Y. Li, N. Lou, L. Gan, *Org. Lett.* 2015, **17**, 524-527; (b) S. Qiu, T. Xu, J. Zhou, Y. Guo, G. Liu, *J. Am. Chem. Soc.* 2010, **132**, 2856-2857.
- K. Kaneko, T. Yoshino, S. Matsunaga, M. Kanai, *Org. Lett.* 2013, **15**, 2502-2505.
- J. Sun, G. Zheng, T. Xiong, Q. Zhang, J. Zhao, Y. Li, Q. Zhang, *ACS Catal.* 2016, **6**, 3674-3678.
- B. Zhang, A. Studer, *Org. Lett.* 2014, **16**, 1790-1793.
- H. Zhang, Y. Song, J. Zhao, J. Zhang, Q. Zhang, *Angew. Chem. Int. Ed.* 2014, **53**, 11079-11083.
- G. B. Boursalian, M. Y. Ngai, K. N. Hojczyk, T. Ritter, *J. Am. Chem. Soc.* 2013, **135**, 13278-13281.
- Y. Li, M. Hartmann, C. G. Daniliuc, A. Studer, *Chem. Commun.* 2015, **51**, 5706-5709.
- When internal alkyne **4k** was used as the substrate, product **5k** was isolated in low yield

- P. Carloni, E. Damiani, M. Iacussi, L. Greci, P. Stipa, D. Cauzi, C. Rizzoli, P. Sgarabotto, *Tetrahedron* 1995, **51**, 12445-12452.
- When the electron-deficient aryl thiol 2,3,5,6-tetrafluorobenzenethiol was used, no desired sulfoxide was isolated, instead sulfide **3u** was isolated in 98% yield. This might be due to the electron-deficient sulfide not being easy to oxidize.