Carbene-catalyzed desymmetrization of 1,3-diols: access to optically enriched tertiary alkyl chlorides

Li, Bao-Sheng; Wang, Yuhuang; Proctor, Rupert S. J.; Jin, Zhichao; Chi, Yonggui Robin

2016

Li, B.-S., Wang, Y., Proctor, R. S. J., Jin, Z., & Chi, Y. R. (2016). Carbene-catalyzed desymmetrization of 1,3-diols: access to optically enriched tertiary alkyl chlorides. *Chemical Communications*, 52(53), 8313-8316.

<https://hdl.handle.net/10356/82223>

<https://doi.org/10.1039/C6CC03345H>

© 2016 The Royal Society of Chemistry. This is the author created version of a work that has been peer reviewed and accepted for publication by *Chemical Communications*, The Royal Society of Chemistry. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [<http://dx.doi.org/10.1039/C6CC03345H>].

Downloaded on 26 Apr 2023 22:14:55 SGT

Carbene-catalyzed desymmetrization of 1,3-diols: access to optically enriched tertiary alkyl chlorides

Bao-Sheng Li,^{a,†} Yuhuang Wang,^{a,†} Rupert S. J. Proctor,^a Zhichao Jin^a and Yonggui Robin Chi^{a,b*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

The introduction of a chlorine atom to a carbon center in an enantioselective manner *via* conventional C-Cl bond formation is difficult. Here we report a new approach to this class of tertiary alkyl chlorides with high optical purities. Instead of forming a new C-Cl bond, our approach involves carbene-catalyzed desymmetrization of 2-chloro-1,3-diols as the key step to set up the chiral carbon center with excellent enantiomeric excess.

Introduction

The chlorine atom is widely found in natural products and synthetic molecules with interesting properties.¹ As such, its introduction to molecules, especially in an enantioselective fashion, has received considerable attention. In particular, it remains challenging to prepare optically enriched tertiary alkyl chlorides with the chlorine atom connected to a stereogenic carbon center. To date the dominant approach has been to form a new carbon-chlorine bond *via* asymmetric catalysis (Scheme 1a). For instance, asymmetric chlorination of carbonyl compounds (such as α -keto esters) *via* enol or enolate intermediates is the most well-established method.^{2,3} Chiral Lewis acid catalysts² and amine^{3a} or phosphoric acid^{3b} organic catalysts have been explored for this type of reaction, as disclosed by, Shibata,^{2a} Shibatomi,^{2b,c} Bartoli,^{3a} and Antilla.^{3c} Our laboratory is interested in exploring carbene organic catalysts for rapid access to functional molecules.⁴ Herein, we report a new approach for effective access to optically enriched tertiary alkyl chlorides *via* N-heterocyclic carbene (NHC)-catalyzed desymmetrization of 1,3-diols.⁵ Our method utilizes a readily available substrate with a chlorine atom pre-installed on a pre-stereogenic carbon center (Scheme 1b). The postulated reaction pathway involves selective formation of an ester at one of the two alcohol units of substrate **1** *via* the catalytic generation of key azolium intermediate **I**.⁶ The intramolecular hydrogen bonding between the two alcohol groups of the 1,3-diol substrate^{7,8} likely plays an important role in this process.⁹ The optically enriched alkyl chloride product **3**

bears an alcohol functional group from the substrate and can readily undergo further transformations.

a) Access to chiral tertiary alkyl chlorides *via* catalytic C-Cl bond formation (literature)

b) Access to optically enriched tertiary alkyl chlorides *via* catalytic desymmetrization (this work)

Scheme 1 The synthesis of optically enriched chiral tertiary alkyl chlorides

Notably, NHCs as versatile organic catalysts are most often used for new bond formations.¹⁰ The use of NHCs for kinetic resolution has also received considerable attention in recent years.¹¹ In contrast, desymmetrization through NHC organocatalysis¹² has thus far received much less study. One nice work in this direction was reported by Scheidt, in which addition of an azolium enolate intermediate to one of the ketone moieties of a symmetric 1,3-diketone compound led to enantiomerically enriched products.^{12a,b} Rovis^{12c} and You have reported desymmetrization through a Stetter reaction, respectively.^{12d} In the area of diol desymmetrization, conversion of non-chlorine-containing diols has mainly been achieved with chiral Lewis acid catalysts¹³ and more recently with chiral phosphoric acid,¹⁴ quinidine-derived^{15a} or sulfide^{15b} organic catalysts.

^a Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore, 637371 (Singapore, E-mail: robinchi@ntu.edu.sg)

^b State Key Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering Ministry of Education, Guizhou University, Guiyang, 550025 (China)

* These authors contributed equally to this work.

† Electronic Supplementary Information (ESI) available: Characterization data and experimental procedures. See DOI: 10.1039/x0xx00000x

Results and discussion

Table 1 Optimization of reaction conditions^a

A: Ar = Mes, R' = H, X = BF₄
 B: Ar = C₆H₅, R' = H, X = BF₄
 C: Ar = C₆H₅, R' = Br, X = BF₄
 D: Ar = C₆H₅, R' = NO₂, X = BF₄
 E: Ar = C₆F₅, R' = H, X = Cl
 F: Ar = 2,4,6-trichloro-phenyl, R' = H, X = BF₄

entry	cat.	R	3 yield (%) ^c	3 er ^d
1	A	C ₆ H ₅	95	84:16
2	A	4-Cl-C ₆ H ₅	90	80:20
3	A	4-CH ₃ O-C ₆ H ₅	92	76:24
4	A	C ₆ H ₅ CH=CH	85	72:28
5	A	2-CH ₃ -C ₆ H ₅	87	84:16
6	A	1-Naphthyl	93	89:11
7	B	1-Naphthyl	95	90:10
8	C	1-Naphthyl	90	77:23
9	D	1-Naphthyl	93	87:13
10	E	1-Naphthyl	94	87:13
11	F	1-Naphthyl	95	95:5
12	F (2 mol%) ^b	1-Naphthyl	90	95:5

^a All reactions of **1a** (0.10 mmol, 20.1 mg) with aldehyde **2** (0.2 mmol, 31.2 mg) were carried out with 5 mol% **A-F** and 10 mol% Cs₂CO₃ in THF (1.0 mL) for 16 h. ^b Catalyst **F** of 2 mol% was used (48 h). ^c Isolated yield. ^d Er of **3** was determined by chiral HPLC analysis. Mes, 1,3,5-trimethylbenzene.

Results of our condition optimization are briefed in Table 1, using diol **1a** as the model substrate. Triazolium NHC pre-catalyst **A**, developed by Bode,¹⁶ was first used to identify a suitable aldehyde to be the acylation agent (entries 1-12). We were very encouraged to find the formation of **3a** in 95% yield and 84:16 er when benzaldehyde was used (entry 1, R = Ph). Further studies of aldehyde substrates found that other aryl aldehydes and enals could effectively lead to the products **3a-3j** in good yield and a variety of enantiomeric ratios. The use of alkyl aldehydes, however, led to little formation of the desired product. Replacing the phenyl group of benzaldehyde with a naphthyl unit led to better results than with other aldehydes examined here (entry 6). We then proceeded to use naphthal as the acylation reagent to search for optimal NHC catalysts (entries 7-12), and found that the use of NHC pre-catalyst **F**, first developed by Rovis,¹⁷ afforded **3a** with 95% yield and 95:5 er (entry 11). Decreasing the catalyst loading of **F** to 2 mol% led to no apparent changes in er, albeit a longer reaction time

(entry 12). Increasing or decreasing the reaction temperature both led to decreased ee (see SI), suggesting that the hydrogen bonding might play important roles in our desymmetrization reactions.

Table 2 Examples of Ar substituents for the diol substrates^a

^a Conditions as in Table 1, entry 11 unless otherwise specified; isolated yield after column chromatography. ^b The corresponding fluorine atom-containing substrate was used.

The scope of the reaction was then examined. We first evaluated the diol substrates by replacing the phenyl group of **1a** with other (hetero) aryl substituents (Table 2). Installing various substituents on the phenyl ring **3b-3j** of the diols had no significant effect on the reaction outcomes, and the corresponding products were obtained with excellent yields and enantiomeric ratios. The phenyl unit of **1a** could also be replaced by a 1-naphthyl **3h** or 2-thiophenyl **3i** group. Furthermore, we found that when the Cl atom was replaced with an F atom, the reaction worked effectively to give **3j** with 95% yield and a somewhat lower 85:15 er under the same conditions without further optimization.

We next evaluated diol substrates bearing non-Ar substituents (olefins and alkynes), in order to provide rich opportunities for further functionalization. As indicated in Table 3, products **4a-4h** containing alkene units with a variety of substitution patterns could be obtained with excellent enantioselectivities and yields. Substrates with a cyclic substituent on the double bond terminus gave the corresponding products **4i-4l** with excellent yields and selectivities. For products **4i** and **4j**, containing a benzocycle, a higher enantioselectivity was achieved with the five-membered ring than with its six-membered analogue. When a benzopyran-like structure was incorporated, the product **4k** was obtained with good results as well. A cyclohexene-containing substrate was then examined and the expected product **4m** was afforded with good yield and er. Likewise, substrates with a Z-olefin substituent gave the desired product **4n** with similar yield and selectivity. Moreover, when the

alkene moiety of the diol substrate was replaced with an alkyne, the product **4p** was formed with yield and er comparable to those of the alkene-containing products such as **4a**. However, the substrates with the alkyl substituent gave the monoester products (**4q-4s**) with moderate er. Our attempt to prepare 1,3-diol substrate by replacing the allylic substituent with an aryl unit was unsuccessful (as below) due to the tertiary benzyl or allylic chloride is sensitive for reduction of 1,3-diester to 1,3-diol. The absolute configuration of the products was confirmed by optical rotation comparison of a derivative of **3j** with a known compound.¹⁸

Table 3 Examples of diol substrates containing alkene or alkyne substituents^a

^a Conditions as in Table 1 entry 11 unless otherwise specified; isolated yield after column chromatography.

Notably, further catalytic esterification of the product **3a** under the standard condition could occur slowly to form diester adduct **3a''** (in 20% yield). This additional esterification did not lead to change of ee for **3a**. This result suggested that the second acylation reaction (to form **3a''**) was not a kinetic resolution process. The enantiomeric excess of **3a** obtained in our reaction exclusively came from catalytic desymmetrization of **1a** (not from kinetic resolution of the mono-ester **3a**). Intramolecular transesterification of **3a** (to form the other enantiomer **3a'**) was not observed (eq. 1), indicating ester **3a** is stable (no ester-exchange) under our catalytic conditions.

The optically enriched products from our reaction can readily undergo further transformations (Scheme 2). For example, oxidation of the free alcohol unit of **3a** (to form **5**) followed by intramolecular Friedel-Crafts acylation effectively afforded indanone **6**, which is a common scaffold in functional molecules. Our phenol containing product **3e** can undergo oxidation followed by intramolecular addition/dearomatization to form spirocyclic ether adduct **10** (Scheme 2b).¹⁹

Scheme 2 Synthetic transformations

Conclusions

In summary, we have developed a new approach for effective access to tertiary alkyl chlorides with high enantiomeric excess. Instead of forming a new C-Cl bond, we obtain the sterically challenging chiral carbon center connected to a chlorine atom through carbene-catalyzed desymmetrization of 1,3-diols. The optically enriched products from our reaction can readily undergo further transformations. Since our method bypasses the difficulties in forming challenging C-Cl bonds enantioselectively, we expect a broad set of chiral alkyl chlorides to be easily obtainable by using our metal-free organic catalytic methods.

Acknowledgements

We acknowledge support from Singapore's National Research Foundation (NRF), Ministry of Education (MOE), Nanyang Technological University (NTU), and China's National Key Program for Basic Research (No. 2010CB 126105), Thousand Talent Plan, National Natural Science Foundation (No. 21132003; No. 21472028), Guizhou Province Returned Oversea Student Science and Technology Activity Program, and Guizhou University.

Notes and references

- (a) T. Dias, I. Brito, L. Moujir, N. Paiz, J. Darias and M. Cueto, *J. Nat. Prod.*, 2005, **68**, 1677; (b) A. R. Díaz-Marrero, I. Brito, J. M. de la Rosa, J. Darias and M. Cueto, *Tetrahedron*, 2008, **64**, 10821; (c) K. Motohashi, M. Sue, K. Furihata, S. Ito and Haruo. Seto, *J. Nat. Prod.*, 2008, **71**, 595.
- (a) N. Shibata, J. Kohno, K. Takai, T. Ishimaru, S. Nakamura, T. Toru and S. Kanemasa, *Angew. Chem., Int. Ed.*, 2005, **44**, 4204; (b) K. Shibatomi, A. Narayama, Y. Soga, T. Muto, S. Iwasa, *Org. Lett.*, 2011, **13**, 2944; (c) K. Shibatomi, Y. Soga, A. Narayama, I. Fujisawa and S. Iwasa, *J. Am. Chem. Soc.*, 2012, **134**, 9836.

- 3 (a) G. Bartoli, M. Bosco, A. Carlone, M. Locatelli, P. Melchiorre and L. Sambri, *Angew. Chem., Int. Ed.*, 2005, **44**, 6219; (b) W. Zheng, Z. Zhang, M. J. Kaplan and J. C. Antilla, *J. Am. Chem. Soc.*, 2011, **133**, 3339.
- 4 For activation of ester: (a) L. Hao, Y. Du, H. Lv, X. Chen, H. Jiang, Y. Shao and Y. R. Chi, *Org. Lett.*, 2012, **14**, 2154; (b) Z. Fu, J. Xu, T. Zhu, W. W. Y. Leong and Y. R. Chi, *Nature Chem.*, 2013, **5**, 835; For activation of C-C bond: (c) B.-S. Li, Y. Wang, Z. Jin, P. Zheng, R. Ganguly and Y. R. Chi, *Nat. Commun.*, 2015, **6**, 6207; (d) B.-S. Li, Y. Wang, Z. Jin and Y. R. Chi, *Chem. Sci.*, 2015, **6**, 6008; For other activation: (e) J. Mo, X. Chen and Y. R. Chi, *J. Am. Chem. Soc.*, 2012, **134**, 8810; (f) J. Zhang, C. Xing, B. Tiwari and Y. R. Chi, *J. Am. Chem. Soc.*, 2013, **135**, 811; (g) Y. Zhang, Y. Du, Z. Huang, J. Xu, X. Wu, Y. Wang, M. Wang, S. Yang, R. D. Webster and Y. R. Chi, *J. Am. Chem. Soc.*, 2015, **137**, 2416.
- 5 For representative metal-catalyzed desymmetrization of 1,3-diols: (a) C. Roux, M. Candy, J.-M. Pons, O. Chuzel and C. Bressy, *Angew. Chem., Int. Ed.*, 2014, **53**, 766; (b) M. S. Hong, T. W. Kim, B. Jung and S. H. Kang, *Chem. Eur. J.*, 2008, **14**, 3290; (c) B. M. Trost and T. Mino, *J. Am. Chem. Soc.*, 2003, **125**, 2410; (d) C. A. Lewis, B. R. Sculimbrene, Y. Xu and S. J. Miller, *Org. Lett.*, 2005, **7**, 3021; (e) T. Honjo, M. Nakao, S. Sano, M. Shiro, K. Yamaguchi, Y. Sei and Y. Nagao, *Org. Lett.*, 2007, **9**, 509; (f) S. Sano, T. Tsumura, N. Tanimoto, T. Honjo, M. Nakao and Y. Nagao, *Synlett*, 2010, 256; For representative Organo-catalyzed desymmetrization of 1,3-diols: (g) A. Sakakura, S. Umemura and K. Ishihara, *Adv. Synth. Catal.*, 2011, **353**, 1938; (h) J. Merad, P. Borkar, T. B. Yenda, C. Roux, J.-Marc Pons, J.-L. Parrain, O. Chuzel and C. Bressy, *Org. Lett.*, 2015, **17**, 2118; (i) S. Yamada, T. Misono, Y. Iwai, A. Masumizu and Y. Akiyama, *J. Org. Chem.*, 2006, **71**, 6872; (j) Z. Chen, and J. Sun, *Angew. Chem., Int. Ed.*, 2013, **52**, 13593.
- 6 (a) B. E. Maki, A. Chan, E. M. Phillips and K. A. Scheidt, *Org. Lett.*, 2007, **9**, 371; (b) J. Guin, S. De Sarkar, S. Grimme and A. Studer, *Angew. Chem., Int. Ed.*, 2008, **47**, 8727; (c) S. De Sarkar, S. Grimme and A. Studer, *J. Am. Chem. Soc.*, 2010, **132**, 1190; (d) P.-C. Chiang and J. W. Bode, *Org. Lett.*, 2011, **13**, 2422; (e) X. Zhao, K. Z. Ruhl and T. Rovis, *Angew. Chem., Int. Ed.*, 2012, **51**, 12330; (f) J. Mo, X. Chen and Y. R. Chi, *J. Am. Chem. Soc.*, 2012, **134**, 8810; (g) X. Chen, S. Yang, B.-A. Song and Y. R. Chi, *Angew. Chem., Int. Ed.*, 2013, **52**, 11134; (h) X.-Y. Chen, F. Xia, J.-T. Cheng and S. Ye, *Angew. Chem., Int. Ed.*, 2013, **52**, 10644; (i) M. Wang, Z. Huang, J. Xu and Y. R. Chi, *J. Am. Chem. Soc.*, 2014, **136**, 1214. For similar ammonium intermediate via organocatalysis: (j) G. Liu, M. E. Shirley, K. N. Van, R. L. McFarlin and D. Romo, *Nature Chem.*, 2013, **5**, 1049; (k) S. Vellalath, K. N. Van and D. Romo, *Angew. Chem., Int. Ed.*, 2013, **52**, 13688.
- 7 Recent selected reviews on desymmetrization of diols, see: (a) C. E. Müller and P. R. Schreiner, *Angew. Chem., Int. Ed.*, 2011, **50**, 6012; (b) Á. Enríquez-García and E. P. Kündig, *Chem. Soc. Rev.*, 2012, **41**, 7803; (c) M. D. Díaz-de-Villegas, J. A. Gálvez, R. Badorrey and M. P. López-Ram-de-Viú, *Chem.-Eur. J.*, 2012, **18**, 13920.
- 8 For recent reports on asymmetric desymmetrization of 2,2-disubstituted 1,3-diols, see: (a) B. Jung and S. H. Kang, *Proc. Natl. Acad. Sci., USA* 2007, **104**, 1472; (b) B. Jung, M. S. Hong and S. H. Kang, *Angew. Chem., Int. Ed.*, 2007, **46**, 2616; (c) J. Y. Lee, Y. S. You and S. H. Kang, *J. Am. Chem. Soc.*, 2011, **133**, 1772; (d) D. W. Tay, G. Y. C. Leung and Y.-Y. Yeung, *Angew. Chem., Int. Ed.*, 2014, **53**, 5161; (e) Z. Ke, C. K. Tan, F. Chen and Y.-Y. Yeung, *J. Am. Chem. Soc.*, 2014, **136**, 5627; (f) Z. Chen and J. Sun, *Angew. Chem., Int. Ed.*, 2013, **52**, 13593; (g) S.-S. Meng, Y. Liang, K.-S. Cao, L. Zou, X.-B. Lin, H. Yang, K. N. Houk and W.-H. Zheng, *J. Am. Chem. Soc.*, 2014, **136**, 12249; (h) W. Yang, Y. Liu, S. Zhang and Q. Cai, *Angew. Chem., Int. Ed.*, 2015, **54**, 8805.
- 9 (a) J. Chen and Y. Huang, *Nat. Commun.*, 2015, **5**, 3437; (b) J. Chen, S. Meng, L. Wang, H. Tang and Y. Huang, *Chem. Sci.*, 2015, **6**, 4184; (c) L. Wang, J. Chen and Y. Huang, *Angew. Chem., Int. Ed.*, 2015, **54**, 15414.
- 10 For selected reviews on NHC catalysis, see: (a) D. Enders and T. Balensiefer, *Acc. Chem. Res.*, 2004, **37**, 534; (b) K. Zeitler, *Angew. Chem., Int. Ed.*, 2005, **44**, 7506; (c) V. Nair, S. Vellalath and B. P. Babu, *Chem. Soc. Rev.*, 2008, **37**, 2691; (d) T. Rovis, *Chem. Lett.*, 2008, **37**, 2; (e) A. T. Biju, N. Kuhl and F. Glorius, *Acc. Chem. Res.*, 2011, **44**, 1182; (f) D. T. Cohen and K. A. Scheidt, *Chem. Sci.*, 2012, **3**, 5336; (g) X. Bugaut and F. Glorius, *Chem. Soc. Rev.*, 2012, **41**, 3511; (h) J. Izquierdo, G. E. Hutson, D. T. Cohen and K. A. Scheidt, *Angew. Chem., Int. Ed.*, 2012, **51**, 11686; (i) S. D. Sarkar, A. Biswas, R. C. Samanta and A. Studer, *Chem. Eur. J.*, 2013, **19**, 4664; (j) S. J. Ryan, L. Candish and D. W. Lupton, *Chem. Soc. Rev.*, 2013, **42**, 4906; (k) J. Douglas, G. Churchill and A. D. Smith, *Synthesis*, 2012, **44**, 2295; (l) J. Izquierdo, G. E. Hutson, D. T. Cohen and K. A. Scheidt, *Angew. Chem., Int. Ed.*, 2012, **51**, 11686; (m) M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485; (n) P. Chauhan and D. Enders, *Angew. Chem., Int. Ed.*, 2014, **53**, 1485; (o) D. M. Flanagan, F. Romanov-Michailidis, N. A. White and T. Rovis, *Chem. Rev.*, 2015, **15**, 9307.
- 11 For NHC catalyzed resolution of alcohol: (a) S. Kuwano, S. Harada, B. Kang, R. Oriez, Y. Yamaoka, K. Takasu and K.-I. Yamada, *J. Am. Chem. Soc.*, 2013, **135**, 11485; (b) S. Lu, S. B. Poh, W.-Y. Siau and Y. Zhao, *Angew. Chem., Int. Ed.*, 2013, **52**, 1731; (c) S. Lu, S. B. Poh and Y. Zhao, *Angew. Chem., Int. Ed.*, 2014, **53**, 11041; (d) S. Iwahana, H. Iida and E. Yashima, *Chem.-Eur. J.*, 2011, **17**, 8009.
- 12 (a) M. Wadamoto, E. M. Phillips, T. E. Reynolds and K. A. Scheidt, *J. Am. Chem. Soc.*, 2007, **129**, 10098; (b) B. E. Maki, A. Chan, E. M. Phillips and K. A. Scheidt, *Org. Lett.*, 2007, **9**, 371; (c) Q. Liu and T. Rovis, *J. Am. Chem. Soc.*, 2006, **128**, 2552; (d) M.-Q. Jia, C. Liu and S.-L. You, *J. Org. Chem.*, 2012, **77**, 10996.
- 13 (a) B. Jung and S. H. Kang, *Proc. Natl. Acad. Sci., USA* 2007, **104**, 1472; (b) B. Jung, M. S. Hong and S. H. Kang, *Angew. Chem., Int. Ed.*, 2007, **46**, 2616; (c) J. Y. Lee, Y. S. You and S. H. Kang, *J. Am. Chem. Soc.*, 2011, **133**, 1772.
- 14 (a) Z. Chen and J. Sun, *Angew. Chem., Int. Ed.*, 2013, **52**, 13593; (b) S.-S. Meng, Y. Liang, K.-S. Cao, L. Zou, X.-B. Lin, H. Yang, K. N. Houk and W.-H. Zheng, *J. Am. Chem. Soc.*, 2014, **136**, 12249; For catalytic asymmetric desymmetrization of 1,3-diol derivatives: (c) Z. Chen, B. Wang, Z. Wang, G. Zhu and J. Sun, *Angew. Chem., Int. Ed.*, 2013, **52**, 2027; (d) Z. Wang, Z. Chen and J. Sun, *Angew. Chem., Int. Ed.*, 2013, **52**, 6685; (e) Z. Wang, F. K. Sheong, H. Sung, H. Y. Ian, D. Williams, Z. Lin and J. Sun, *J. Am. Chem. Soc.*, 2015, **137**, 5895.
- 15 (a) D. W. Tay, G. Y. C. Leung and Y.-Y. Yeung, *Angew. Chem., Int. Ed.*, 2014, **53**, 5161; (b) Z. Ke, C. K. Tan, F. Chen and Y.-Y. Yeung, *J. Am. Chem. Soc.*, 2014, **136**, 5627.
- 16 M. He, J. R. Struble and J. W. Bode, *J. Am. Chem. Soc.*, 2006, **128**, 8418.
- 17 H. U. Vora, J. R. Moncecchi, O. Epstein and T. Rovis, *J. Org. Chem.*, 2008, **73**, 9727.
- 18 For details, please see the Supporting Information.
- 19 (a) S. Nozoe, M. Morisaki, K. Tsuda, Y. Iitaka, N. Takahashi, S. Tamura, K. Ishibashi and M. Shirasaka, *J. Am. Chem. Soc.*, 1965, **87**, 4968; (b) M. Entzeroth, A. J. Blackman, J. S. Mynderse and R. E. Moore, *J. Org. Chem.*, 1985, **50**, 1255; (c) S. Aoki, Y. Watanabe, M. Sanagawa, A. Setiawan, N. Kotoku and M. Kobayashi, *J. Am. Chem. Soc.*, 2006, **128**, 3148; (d) I. A. Katsoulis, G. Kythreoti, A. Papakyriakou, K. Koltsida, P. Anastasopoulou, C. I. Stathakis, I. Mavridis, T. Cottin, E. Saridakis and D. Vourloumis, *ChemBioChem*, 2011, **12**, 1188.