Spin-Orbit State-Selective C—I Dissociation Dynamics of the $\text{CH}_3\text{I}^+ \tilde{X}$ Electronic State Induced by Intense Few-Cycle Laser Fields

Zhengrong Wei,¹ Jialin Li,¹ Soo Teck See,¹ and Zhi-Heng Loh^{1,2,*}

¹ *Division of Chemistry and Biological Chemistry, and Division of Physics and Applied Physics, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore*

² *Centre for Optical Fibre Technology, The Photonics Institute, Nanyang Technological University, Singapore 639798, Singapore*

* Author to whom correspondence should be addressed. E-mail: zhiheng@ntu.edu.sg

Abstract

Studies of ultrafast molecular dynamics induced by intense laser fields can unravel new approaches to manipulating chemical reactions in the strong-field regime. Here, we show that intense few-cycle laser pulses can induce the spin-orbit state-selective C—I dissociation of the iodomethane cation (CH_3I^+) in the $\tilde{X}$ electronic state. Irradiation of CH_3I by 6-fs laser pulses with peak intensities of $1.9 \times 10^{14} \text{ W/cm}^2$ followed by femtosecond extreme ultraviolet (XUV) probing of the iodine $4d$ core-level transitions reveals dissociation of the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{1/2}$ state with a time constant of $0.76 \pm 0.16 \text{ ps}$. By contrast, the $\tilde{X}^2E_{3/2}$ spin-orbit ground state does not exhibit any appreciable dissociation on the picosecond timescale. The observed spin-orbit state-selective dissociation of the $\tilde{X}$ state is rationalized in terms of the laser-induced coupling to the $\tilde{A}$ state. Our results suggest that the intense-laser control of photodissociation channels can be potentially extended to spin-orbit-split states.

TOC graphic

Advances in tabletop femtosecond laser technologies enable routine access to laser pulses with peak intensities exceeding $\sim 1 \text{ PW/cm}^2$.¹ From the chemical perspective, the high electric-field strengths that are furnished by these intense laser pulses can be harnessed to control the outcomes of chemical reactions.²⁻⁹ For example, branching ratios of nonadiabatic photodissociation dynamics have been manipulated via the dynamic Stark effect³ and light-induced conical intersections.⁸ The realization of various strong-field control schemes requires an understanding of how molecules interact with laser fields in the nonperturbative, strong-field regime. Early investigations of strong-field laser-molecule interactions focused on the simplest molecule, H_2^+ , whose dissociation in intense laser fields has been extensively studied by both experiment and theory.¹⁰⁻¹⁵ These studies elucidate new phenomena such as bond softening, bond hardening and Coulomb explosion. Beyond H_2^+ , efforts have been undertaken to unravel the interaction of strong laser fields with multielectron molecules.¹⁶⁻²³ Compared to the case of H_2^+ , these studies reveal complex angle-dependent ionization yields,^{24,25} participation of multiple channels in strong-field ionization,^{26,27} multielectron nonadiabatic dynamics,^{22,23} and a variety of molecular dynamics.^{17,28} Recent investigations have also unraveled the timescales for strong-field dissociative ionization,²⁹⁻³² a key parameter in the understanding of ultrafast photochemical reactions.³³

Here, we employ femtosecond XUV absorption spectroscopy³⁴ to investigate the strong-field-induced dissociation dynamics of iodomethane (CH_3I). As a prototypical polyatomic molecule, the photodissociation dynamics of its neutral excited states in the weak-field regime has been the subject of numerous investigations.² In the presence of intense laser fields, dissociative ionization of CH_3I^+ and multiple Coulomb explosion pathways of CH_3I^{n+} ($n > 1$) are revealed by velocity map imaging of various fragments and measurements of the kinetic energy release for different dissociation channels.³⁵⁻³⁷ In the present study, we focus on elucidating the ultrafast CH_3I

dissociation dynamics in the presence of intense laser fields. We observe spin-orbit state-selective dissociation of the $\text{CH}_3\text{I}^+ \tilde{X}$ state – while the $^2E_{3/2}$ spin-orbit ground state is stable with respect to C—I dissociation, the $^2E_{1/2}$ spin-orbit excited state dissociates on a timescale of 0.76 ± 0.16 ps. In addition, dissociation of the $\text{CH}_3\text{I}^+ \tilde{A} \ ^2A_1$ state occurs with a time constant of 86 ± 11 fs. The observed dissociation dynamics are rationalized in terms of the coupling between the $\tilde{X}$ and $\tilde{A}$ potential energy surfaces induced by the intense laser field. Finally, our results suggest that both the $\text{CH}_3\text{I}^+ \tilde{X} \ ^2E_{3/2}$ and $^2E_{1/2}$ states undergo an ultrafast structural rearrangement on a time scale of 75 ± 8 fs immediately upon their formation.

Strong-field ionization is effected by the irradiation of neutral CH_3I molecules by intense laser pulses with a carrier wavelength of 786 nm (1.58-eV photon energy) and a pulse duration of 5.6 fs FWHM. The few-cycle pulse, with pulse energy of 0.17 mJ, is loosely focused by a 1-m focal length spherical mirror to a 100- μm beam waist to yield a peak intensity of 1.9×10^{14} W/cm². The sample target comprises 14 mbar of CH_3I in a 3-mm path length quasi-static gas cell that is heated to 353 K. Details of the experimental apparatus can be found in ref. 38. The XUV differential absorption spectra collected as a function of time delay is shown in Figure 1a. The negative differential absorption signal (ΔA) that is observed in the probe photon energy range of $\sim 50 - 52$ eV is due to bleaching of the CH_3I ground-state $^1A_1 \rightarrow 4d_j^{-1}\sigma_{\text{C-I}}^*$ transition³⁹ ($j = 5/2$ and $3/2$) that accompanies the depletion of the neutral CH_3I population by strong-field ionization. On the other hand, the positive ΔA signal in the 46 – 49-eV range can be assigned to the ground-state CH_3I^+ parent ion and the atomic I ($^2P_{3/2}$) and I^+

Figure 1. (a) Differential XUV absorption spectra collected as a function of pump-probe time delay. (b) Fit of the differential XUV absorption spectrum at 2 ps to multiple Gaussians. The red symbols are the data and the solid black line is the fit. Contributions from absorption of CH_3I^+ , neutral CH_3I , neutral I and singly charged I^+ are color-coded.

(3P_2) photoproducts produced by the intense laser-molecule interaction. Pronounced modulations of the transition energies about the neutral CH_3I depletion region and, to a lesser extent, about the CH_3I^+ parent ion peak at 47.0 eV (see below for assignment) originate from coherent vibrational motions launched by the intense laser field.³⁸

The ΔA spectrum collected at 2-ps time delay along with the fits to the XUV transitions of the various species is shown in Figure 1b. The peak positions and spectral assignments of the observed XUV transitions are summarized in Table 1. In assigning the differential XUV absorption spectrum, our strategy is to consider the species that are most likely to form under our experimental conditions. We start by identifying the transitions that arise from the depletion of the neutral CH_3I and the formation of the CH_3I^+ parent ion. Next, we identify the absorption peaks that are due to atomic I ($^2P_{3/2}$) and I^+ (3P_2), which are produced as a result of dissociation in the low-lying ($\tilde{X}$

Table 1. Observed XUV absorption peaks for CH_3I^+ ($\tilde{X}$), I, I^+ , and CH_3I ($\tilde{X}$), their assignments to the I $4d$ core-level transitions, (REV 2.3B) and the transition energies obtained from the literature.

Species	Peak position (eV)	Peak position, lit. (eV)	Transition
CH_3I^+ ($\tilde{X}$)	46.6	46.45 ^a	$^2E_{1/2} \rightarrow 4d_{5/2}^{-1}$
	47.0	47.08 ^a	$^2E_{3/2} \rightarrow 4d_{5/2}^{-1}$
	48.1	48.19 ^a	$^2E_{1/2} \rightarrow 4d_{3/2}^{-1}$
	48.7	48.82 ^a	$^2E_{3/2} \rightarrow 4d_{3/2}^{-1}$
	52.6	—	$^2E_{3/2} \rightarrow 4d_{5/2}^{-1}e_{3/2}^{-1}\sigma_{\text{C-I}}^*$ $^2E_{1/2} \rightarrow 4d_{5/2}^{-1}e_{1/2}^{-1}\sigma_{\text{C-I}}^*$
	54.2	—	$^2E_{3/2} \rightarrow 4d_{3/2}^{-1}e_{3/2}^{-1}\sigma_{\text{C-I}}^*$ $^2E_{1/2} \rightarrow 4d_{3/2}^{-1}e_{1/2}^{-1}\sigma_{\text{C-I}}^*$
CH_3I ($\tilde{X}$)	50.6	50.62 ^b	$^1A_1 \rightarrow 4d_{5/2}^{-1}\sigma_{\text{C-I}}^*$
	52.3	52.34 ^b	$^1A_1 \rightarrow 4d_{3/2}^{-1}\sigma_{\text{C-I}}^*$
I ($^2P_{3/2}$)	46.0	45.94 ^c	$^2P_{3/2} \rightarrow ^2D_{5/2}$
	47.6	47.64 ^c	$^2P_{3/2} \rightarrow ^2D_{3/2}$
I^+ (3P_2)	47.3	47.32 ^c	$^3P_2 \rightarrow ^3D_3$
	49.0	48.95 ^c	$^3P_2 \rightarrow ^3P_2$
	49.2	49.17 ^c	$^3P_2 \rightarrow ^3D_1$

^a Calculated based on literature values for the valence and I $4d$ core level binding energies of CH_3I (see the Supporting Information). ^b Ref. 39. ^c Ref. 40.

and $\tilde{A}$) electronic states of CH_3I^+ .⁴¹ With this approach, we are able to account for all the features that appear in the differential XUV absorption spectrum. Higher charged states of CH_3I^{n+} and I^{n+} ($n \geq 2$) do not seem to be present in detectable quantities. For example, the I^{2+} ($^4S_{3/2}$) ground state possesses a pronounced absorption peak at 48.93 eV,⁴⁰ which is not observed in the differential XUV absorption spectrum. Finally, we do not consider the formation of atomic I $^2P_{1/2}$ species, which originates from the dissociation of the $\tilde{B}$ state that is located $\sim 5 - 6$ eV above the

$\tilde{X}$ state of CH_3I^+ in the Franck-Condon region.⁴¹ Given our laser peak intensity, we do not expect any appreciable population of the $\tilde{B}$ state by strong-field ionization.

Assignment of the features observed in the positive ΔA spectral region requires knowledge of the transition energies of the CH_3I^+ ion. These transition energies can be estimated within the framework of the single-particle picture,⁴² which considers XUV transition to involve the promotion of an electron from the I $4d$ orbital to the single vacancy in the HOMO orbital of e symmetry. Using the original atomic I $4d_j^{-1}$ notation, these are the $^2E_{1/2} \rightarrow 4d_{5/2}^{-1}$, $^2E_{3/2} \rightarrow 4d_{5/2}^{-1}$, $^2E_{1/2} \rightarrow 4d_{3/2}^{-1}$, and $^2E_{3/2} \rightarrow 4d_{3/2}^{-1}$ transitions, whose transition energies are calculated to be 46.45, 47.08, 48.19, and 48.82 eV, respectively (see the Supporting Information). These predicted transition energies agree well with the peaks observed at 46.6, 47.0, 48.1, and 48.7 eV.

Aside from the four peaks that can be assigned to the CH_3I^+ parent ion, a peak at 46.0 eV is also present in the ΔA spectrum. Based on its absence in the ΔA spectrum at earlier time delays, is assigned it to the $^2P_{3/2} \rightarrow ^2D_{5/2}$ transition of the atomic I $^2P_{3/2}$ state, in good agreement with the literature value of 45.94 eV.⁴⁰ The accompanying $^2P_{3/2} \rightarrow ^2D_{3/2}$ transition, reported to be at 47.64 eV, is observed at 47.6 eV.^{40,43} It is noteworthy that the I $^2P_{1/2} \rightarrow ^2D_{3/2}$ transition at 46.70 eV is not observed in our experiment, which suggests that the strong-field dissociative ionization of CH_3I , under our experimental conditions, yields neutral atomic I only in the $^2P_{3/2}$ state. In addition to neutral atomic I ($^2P_{3/2}$), a shoulder at 47.3 eV that is evident in the ΔA spectrum collected at 2 ps time delay (Figure 1b) can be assigned to the $^3P_2 \rightarrow ^3D_3$ transition of I^+ , in good agreement with the transition energy of 47.32 eV that is reported in the literature.⁴⁰ Other transitions for the I^+ 3P_2 state with significant oscillator strengths, such as the $^3P_2 \rightarrow ^3P_2$ (48.95 eV) and $^3P_2 \rightarrow ^3D_1$ (49.17 eV),⁴⁰ are also observed at 49.0 eV and 49.2 eV, respectively. XUV

transitions with large oscillator strengths that originate from the 3P_1 and 3P_0 states of I^+ , e.g., the $^3P_1 \rightarrow ^3D_2$ and the $^3P_0 \rightarrow ^3P_1$ transitions at 47.80 and 47.15 eV, respectively, are notably absent. This observation suggests that the I^+ fragment is produced solely in the 3P_2 ground state during the dissociative ionization of CH_3I .

The negative peaks at 50.6 eV and 52.3 eV in the ΔA spectra, which arise from the $I\ 4d_j \rightarrow \sigma^*$ transitions ($j = 5/2$ and $3/2$) of depleted CH_3I , have relative amplitudes that deviate significantly from the 1.5:1 ratio that is expected based on the $(2j + 1)$ -degeneracy of the $I\ 4d$ core-hole states. This observation suggests that a product of strong-field ionization of CH_3I exhibits absorption features that partially overlap with the $I\ 4d_{3/2} \rightarrow \sigma^*$ transition of CH_3I . Following previous work on the XUV probing of strong-field-ionized Br_2 ,⁴⁴ we assign the features with positive ΔA to the $I\ 4d_j \rightarrow \sigma^*$ transitions ($j = 5/2$ and $3/2$) of CH_3I^+ ; curve fitting yields absorption peaks at 52.6 and 54.2 eV and a relative ratio of $1.7 \pm 0.1:1$ for the peak areas. The blueshift of the $I\ 4d_j \rightarrow \sigma^*$ transition energies by 2.0 eV upon ionization points to the preferential stabilization of the $I\ 4d$ core level over the valence σ^* level upon the removal of an electron from the outermost valence e orbital of CH_3I . Note that this curve fitting takes into account the negative ΔA signal that arises from the perturbed free-induction decay (PFID) of the $I\ 4d_{5/2}^{-1} \rightarrow 6pe$ transition at 54.3 eV; previous experiments on atomic species show that measuring the temporal evolution of the PFID feature yields autoionization lifetimes.^{45,46}

Figure 2 shows the temporal evolution of the ΔA signals collected for the $\tilde{X}^+ \ ^2E_{3/2}$ and $^2E_{1/2}$ states of the CH_3I^+ parent ion at the $^2E_{3/2} \rightarrow 4d_{5/2}^{-1}$ (47.0 eV, Figure 2a) and $^2E_{1/2} \rightarrow 4d_{3/2}^{-1}$ (48.1 eV, Figure 2b) transitions, respectively, along with the growth of the neutral I ($^2P_{3/2}$) and the singly charged I^+ (3P_2) species observed at their respective $^2P_{3/2} \rightarrow ^2D_{5/2}$ (46.0 eV,

Figure 2. Temporal traces of the ΔA signal for the (a) $\text{CH}_3\text{I}^+ \tilde{X}^2E_{3/2}$, (b) $\text{CH}_3\text{I}^+ \tilde{X}^2E_{1/2}$, (c) neutral I ($^2P_{3/2}$), (d) singly charged I^+ (3P_2) species, and the (e) spectral first-moments calculated about the transitions for the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{3/2}$ and $\text{CH}_3\text{I}^+ \tilde{X}^2E_{1/2}$ states at 47.0 and 48.7 eV, respectively. The dashed red lines are the data and the solid black lines are the fits.

Figure 2c) and $^3P_2 \rightarrow ^3D_3$ (47.3 eV, Figure 2d) transitions. The positive ΔA signals for both the $\tilde{X}^2E_{3/2}$ and $^2E_{1/2}$ states of CH_3I^+ appear within the <10-fs instrument response of the experimental apparatus. Global fitting of the time traces for the $\tilde{X}^+ ^2E_{3/2}$ and $^2E_{1/2}$ states of CH_3I^+ as well as the I $^2P_{3/2}$ state shows that both parent ion species exhibit a rapid decay with a time constant of 80 ± 20 fs. Moreover, the slow decay observed for the $^2E_{1/2}$ spin-orbit-excited state of CH_3I^+ is correlated with the growth of the I $^2P_{3/2}$ population with a time constant of 0.76

± 0.16 ps; note that the growth of the I $^2P_{3/2}$ population does not exhibit any fast component on the sub-100-fs timescale. In addition, the time- dependent spectral first- moments computed about the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{3/2} \rightarrow 4d_{5/2}^{-1}$ and $^2E_{1/2} \rightarrow 4d_{3/2}^{-1}$ transitions both reveal a blueshift that is characterized by a time constant of 75 ± 8 fs (Figure 2e); the oscillatory features that appear for the $\tilde{X}^2E_{3/2} \rightarrow 4d_{5/2}^{-1}$ transition are due to vibrational wave packet dynamics.³⁸ Finally, in contrast to the slow appearance of the I $^2P_{3/2}$ state, the I $^+ ^3P_2$ signal rises with a time constant of 86 ± 11 fs.

From previous spectroscopic measurements on the CH_3I^+ parent ion,⁴¹ the dissociation channels and their dissociation energies^{47,48} that are relevant to the present work are:

The prompt appearance of the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{3/2}$ and $^2E_{1/2}$ states within the <10-fs instrument response time suggests that they originate directly from the strong-field ionization of CH_3I instead of internal conversion from the electronically excited states of CH_3I^+ .⁴⁹ To gain insight into the ensuing ultrafast C—I dissociation dynamics, we first consider the correlation between the appearance time of the I $^2P_{3/2}$ state and the decay of the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{1/2}$ state, both of which can be fit to the same time constant of 0.76 ± 0.16 ps. Despite the 80 ± 20 -fs decay observed for both $\tilde{X}^2E_{3/2}$ and $^2E_{1/2}$ states of CH_3I^+ , note that such a timescale is not reflected in the increase of the I $^2P_{3/2}$ population (see below). These observations suggest that the I $^2P_{3/2}$ species is produced exclusively from the dissociation of the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{1/2}$ spin-orbit excited state. The

$\tilde{X}^2E_{3/2}$ spin-orbit ground state, on the other hand, does not exhibit any appreciable dissociation on the picosecond timescale of the present experiment.

Previous studies by mass-analyzed threshold ionization (MATI) spectroscopy have revealed selective dissociation of larger iodoalkane ions, $C_3H_7I^+$ and $C_4H_9I^+$, in their spin-orbit ground state.^{50,51} In those cases, following selective photoexcitation to the spin-orbit excited state, strong coupling between the spin-orbit-split states leads to efficient internal conversion to the spin-orbit ground state, from which statistical dissociation occurs. In contrast, the sub-picosecond C—I dissociation dynamics observed in the present work occurs selectively from the $CH_3I^+ \tilde{X}^2E_{1/2}$ spin-orbit excited state.

One possible way to rationalize this counterintuitive result is to consider the intense laser-induced coupling between the ion $\tilde{X}$ and $\tilde{A}$ states. Figure 3 shows a schematic illustration of the potential energy curves for the $CH_3I^+ \tilde{A}$ state and for both $\tilde{X}^2E_{3/2}$ and $\tilde{X}^2E_{1/2}$ states along the C—I dissociation coordinate. Ionization of CH_3I near the peak of the intense few-cycle laser pulse envelope (at time t_0 in the upper panel of Figure 3) results in CH_3I^+ ions formed in a mixture of $\tilde{X}$ and $\tilde{A}$ states. The former is bound and therefore does not lead to C—I dissociation. On the other hand, the $\tilde{A}$ state is dissociative. As it undergoes dissociation in the presence of a laser pulse, the wave packet will arrive in a region for which the $\tilde{A} - \tilde{X}$ gap is resonant with the photon energy of the laser pulse. In fulfilling the resonance condition, stimulated emission from the $\tilde{A}$ state to the $\tilde{X}$ state occurs. The mean photon energy of the laser pulse used in our experiments is 1.58 eV, computed based on the carrier wavelength. Figure 3 shows that this resonance condition occurs at shorter C—I bond distances for the $\tilde{A} - \tilde{X}^2E_{1/2}$ gap than the $\tilde{A} - \tilde{X}^2E_{3/2}$ gap; a dressed states picture yields the same result (see the Supporting Information). When the wave packet arrives at

Figure 3. Schematic illustration of the dissociative wave packet dynamics along the C—I coordinate for the $\tilde{A}$ and $\tilde{X}$ electronic states (bottom panel), mapped to different points in time within the intensity envelope of the strong-field ionization pump pulse (top panel). t_0 occurs near the peak of the intensity envelope, whereas the intensity is vanishingly small at t_2 . At t_1 , laser-induced coupling leads to the stimulated emission of the $\tilde{A}$ state dissociative wave packet to the $\tilde{X}^2E_{1/2}$ state, from which dissociation occurs to give $I(^2P_{3/2})$.

the $\tilde{A} - \tilde{X}^2E_{1/2}$ crossing (at time t_1), the same laser pulse that earlier induced strong-field ionization now drives stimulated emission, resulting in the transfer of population from the $\tilde{A}$ state to the $\tilde{X}^2E_{1/2}$ state. Since the nuclear kinetic energy is conserved in this process, the wave packet continues to propagate along the $\tilde{X}^2E_{1/2}$ state potential to give the atomic $I(^2P_{3/2})$ dissociation product. Because the laser pulses used in the experiments are of few-cycle duration, it is conceivable that the laser field strength is insufficient to induce the same stimulated emission process by the time the wave packet arrives at the $\tilde{A} - \tilde{X}^2E_{3/2}$ crossing (at time t_2). As such, selective dissociation of the $\tilde{X}^2E_{1/2}$ spin-orbit excited state is observed. Note that even though the nuclear wave packet gains considerable kinetic energy from dissociating along the $\tilde{A}$ state potential prior to stimulated emission to the $\tilde{X}$ state, this kinetic energy must subsequently be used

to overcome the barrier to dissociation on the $\tilde{X}$ state potential. As a result, the dissociation time of 0.76 ± 0.16 ps is longer than that for a purely repulsive dissociation.

According to the mechanism proposed above, the spin-orbit selectivity would vanish when intense laser pulses of longer durations are employed. Indeed, previous strong-field experiments performed with ~ 100 -fs pulses revealed dissociation of both $^2E_{3/2}$ and $^2E_{1/2}$ spin-orbit states of CH_3I^+ .³⁶ Future experiments that scan the XUV probe pulse in time within the envelope of the strong-field ionization pulse can be used to track the laser-induced coupling in real time. Further insight into the origin of the spin-orbit state-selective dissociation can also be obtained from theoretical wave packet propagation on field-dressed potential energy surfaces. We note, however, that such studies are nontrivial as they require the inclusion of the time-varying electric field of the few-cycle pulse and the spin-orbit splitting between the $\text{CH}_3\text{I}^+ \tilde{X} \ ^2E_{3/2}$ and $^2E_{1/2}$ states, as well as accurate values for the $\tilde{X} \rightarrow \tilde{A}$ transition dipole moments as a function of C—I distance.

The above explanation for the dissociation of CH_3I^+ to give atomic I ($^2P_{3/2}$) naturally accounts for the appearance of the $\text{I}^+ \ ^3P_2$ signal on the sub-100-fs timescale: the portion of the wave packet on the $\text{CH}_3\text{I}^+ \tilde{A} \ (^2A_1)$ state that does not cross over to the $\tilde{X} \ (^2E_{1/2})$ state yields $\text{I}^+ \ (^3P_2)$ as the dissociation product. The observed sub-100-fs timescale is consistent with the featureless $\tilde{A}$ band in the photoelectron spectrum of CH_3I ,⁵² which suggests prompt dissociation along the shallow $\text{CH}_3\text{I}^+ \tilde{A}$ state potential. The XUV differential absorption spectrum at early time delays, however, does not reveal any signatures that are associated with the $\tilde{A}$ state. According to our estimates, the XUV transitions of the $\tilde{A}$ state to the $4d_{5/2}^{-1}$ and $4d_{3/2}^{-1}$ states should be located at 44.7 and 46.4 eV, respectively. The indiscernible $\tilde{A}$ state XUV absorption suggests a vanishing

Franck-Condon overlap between the weakly bound $\tilde{A}$ state and the repulsive $\text{I } 4d_j^{-1}$ ($j = 5/2$ or $3/2$) core-excited states.

Aside from the sub-picosecond dissociation dynamics of the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{1/2}$ state, an additional 80 ± 20 -fs decay component is observed for both the CH_3I^+ parent ion $^2E_{3/2} \rightarrow 4d_{5/2}^{-1}$ (47.0 eV, Figure 2a) and $^2E_{1/2} \rightarrow 4d_{3/2}^{-1}$ (48.1 eV, Figure 2b) transitions. This decay cannot originate from the dissociation of the $\text{CH}_3\text{I}^+ \tilde{X}$ state because such a process should be accompanied by the corresponding growth of the atomic I ($^2P_{3/2}$) population on a similar timescale, which is not observed. Instead, we attribute the decay of the ΔA signals to the spectral shift that is observed about the two transitions (Figure 2e) due to the identical timescales for both processes (80 ± 20 fs for the decay vs. 75 ± 8 fs for the spectral shift). The spectral shift is indicative of ultrafast structural rearrangement of CH_3I^+ in the $\tilde{X}$ state. Possible candidates include vibrational relaxation along the ν_1 C—H symmetric stretch⁵³ at 2938 cm^{-1} and the $E \times e$ Jahn-Teller distortion that results from the coupling of the $\tilde{X}$ electronic state to the doubly degenerate vibrational modes.^{53,54} Further theoretical investigations involving ab initio wave packet propagation can be used to identify the origin of the sub-100-fs structural rearrangement.

In conclusion, femtosecond XUV absorption spectroscopy reveals sub-picosecond dissociation of the C—I bond for the $\text{CH}_3\text{I}^+ \tilde{X}^2E_{1/2}$ state whereas the $\tilde{X}^2E_{3/2}$ state does not undergo any discernible dissociation on the picosecond timescale. The spin-orbit state-selective dissociation is rationalized in terms of the laser-induced coupling between the $\tilde{X}$ and $\tilde{A}$ electronic states of CH_3I^+ . The one-photon crossing between the two electronic states is located nearer to the Franck-Condon region for the spin-orbit excited state than the ground state. Since the few-cycle laser pulse acts as a temporal gate that confines the laser-induced coupling of ion electronic states

to within its pulse envelope, selective dissociation is observed from the spin-orbit-excited state because of the shorter time needed for the $\tilde{A}$ state wave packet to travel from the Franck-Condon region to the one-photon crossing with the $\tilde{X}^2E_{1/2}$ state. While previous studies have demonstrated the ability of intense laser fields to manipulate the branching ratios of photodissociation reactions involving different electronic states, our results suggest that such control can be potentially extended to spin-orbit-split states. Employing a second few-cycle laser pulse that is appropriately delayed in time with respect to the first strong-field ionization pulse allows the dissociating $\tilde{A}$ state wave packet to couple selectively to either the $\tilde{X}^2E_{1/2}$ state or the $\tilde{X}^2E_{3/2}$ state, hence steering the dissociation dynamics on a predetermined spin-orbit state.

Acknowledgments

This work is supported by a NTU start-up grant, the Ministry of Education Academic Research Fund (MOE2014-T2-2-052 and RG105/17), and the award of a Nanyang Assistant Professorship to Z.-H.L.

Supporting Information Available: Calculation of the iodine $4d$ XUV transition energies of CH_3I^+ , evaluation of accuracy of the observed XUV transitions, and dressed state potentials relevant to the spin-orbit state-selective dissociation of the $\text{CH}_3\text{I}^+ \tilde{X}$ state.

References

- (1) Brabec, T.; Krausz, F. Intense Few-Cycle Laser Fields: Frontiers of Nonlinear Optics. *Rev. Mod. Phys.* **2000**, *72*, 545-591.
- (2) Zhong, D.; Zewail, A. H. Femtosecond Real-Time Probing of Reactions. 23. Studies of Temporal, Velocity, Angular, and State Dynamics from Transition States to Final Products by Femtosecond-Resolved Mass Spectrometry. *J. Phys. Chem. A* **1998**, *102*, 4031-4058.
- (3) Sussman, B. J.; Townsend, D.; Ivanov, M. Y.; Stolow, A. Dynamic Stark Control of Photochemical Processes. *Science* **2006**, *314*, 278-281.
- (4) Solá, I. R.; González-Vázquez, J.; de Nalda, R.; Bañares, L. Strong Field Laser Control of Photochemistry. *Phys. Chem. Chem. Phys.* **2015**, *17*, 13183-13200.
- (5) Sheehy, B.; DiMauro, L. F. Atomic and Molecular Dynamics in Intense Optical Fields. *Annu. Rev. Phys. Chem.* **1996**, *47*, 463-494.
- (6) Posthumus, J. H. The Dynamics of Small Molecules in Intense Laser Fields. *Rep. Prog. Phys.* **2004**, *67*, 623-665.
- (7) Levis, R. J.; Rabitz, H. A. Closing the Loop on Bond Selective Chemistry Using Tailored Strong Field Laser Pulses. *J. Phys. Chem. A* **2002**, *106*, 6427-6444.
- (8) Corrales, M. E.; González-Vázquez, J.; Balerdi, G.; Solá, I. R.; de Nalda, R.; Bañares, L. Control of Ultrafast Molecular Photodissociation by Laser-Field-Induced Potentials. *Nat. Chem.* **2014**, *6*, 785-790.
- (9) Codling, K.; Frasinski, L. J. Dissociative Ionization of Small Molecules in Intense Laser Fields. *J. Phys. B: At. Mol. Opt. Phys.* **1993**, *26*, 783-809.
- (10) McKenna, J.; Sayler, A. M.; Anis, F.; Gaire, B.; Johnson, N. G.; Parke, E.; Hua, J. J.; Mashiko, H.; Nakamura, C. M.; Moon, E.; Chang, Z.; Carnes, K. D.; Esry, B. D.; Ben-

- Itzhak, I. Enhancing High-Order Above-Threshold Dissociation of H_2^+ Beams with Few-Cycle Laser Pulses. *Phys. Rev. Lett.* **2008**, *100*, 133001.
- (11) Jiang, S.; Yu, C.; Yuan, G.; Wu, T.; Lu, R. Dissociation and Ionization of Quasi-Periodically Vibrating H_2^+ in Intense Few-Cycle Mid-Infrared Laser Fields. *Sci. Rep.* **2017**, *7*, 42086.
- (12) Giusti-Suzor, A.; Mies, F. H.; DiMauro, L. F.; Charron, E.; Yang, B. Dynamics of H_2^+ in Intense Laser Fields. *J. Phys. B: At. Mol. Opt. Phys.* **1995**, *28*, 309-339.
- (13) Frasinski, L. J.; Posthumus, J. H.; Plumridge, J.; Codling, K.; Taday, P. F.; Langley, A. J. Manipulation of Bond Hardening in H_2^+ by Chirping of Intense Femtosecond Laser Pulses. *Phys. Rev. Lett.* **1999**, *83*, 3625-3628.
- (14) Bucksbaum, P. H.; Zavriyev, A.; Muller, H. G.; Schumacher, D. W. Softening of the H_2^+ Molecular Bond in Intense Laser Fields. *Phys. Rev. Lett.* **1990**, *64*, 1883-1886.
- (15) Brannon, P. J.; Church, C. H.; Peters, C. W. Electric Field Induced Spectra of Molecular Hydrogen, Deuterium and Deuterium Hydride. *J. Mol. Spectrosc.* **1968**, *27*, 44-54.
- (16) Zhang, W.; Li, Z.; Lu, P.; Gong, X.; Song, Q.; Ji, Q.; Lin, K.; Ma, J.; He, F.; Zeng, H.; Wu, J. Photon Energy Deposition in Strong-Field Single Ionization of Multielectron Molecules. *Phys. Rev. Lett.* **2016**, *117*, 103002.
- (17) Xu, H.; Marceau, C.; Nakai, K.; Okino, T.; Chin, S.-L.; Yamanouchi, K. Communication: Two Stages of Ultrafast Hydrogen Migration in Methanol Driven by Intense Laser Fields. *J. Chem. Phys.* **2010**, *133*, 071103.
- (18) Wu, G.; Boguslavskiy, A. E.; Schalk, O.; Schuurman, M. S.; Stolow, A. Ultrafast Non-Adiabatic Dynamics of Methyl Substituted Ethylenes: The π^*3s Rydberg State. *J. Chem. Phys.* **2011**, *135*, 164309.

- (19) Smits, M.; de Lange, C. A.; Stolow, A.; Rayner, D. M. Dynamic Polarization in the Strong-Field Ionization of Small Metal Clusters. *Phys. Rev. Lett.* **2004**, *93*, 203402.
- (20) Mikosch, J.; Boguslavskiy, A. E.; Wilkinson, I.; Spanner, M.; Patchkovskii, S.; Stolow, A. Channel- and Angle-Resolved Above Threshold Ionization in the Molecular Frame. *Phys. Rev. Lett.* **2013**, *110*, 023004.
- (21) Mathur, D.; Rajgara, F. A.; Dharmadhikari, A. K.; Dharmadhikari, J. A. Strong-Field Ionization of Water by Intense Few-Cycle Laser Pulses. *Phys. Rev. A* **2008**, *78*, 023414.
- (22) Lezius, M.; Blanchet, V.; Rayner, D. M.; Villeneuve, D. M.; Stolow, A.; Ivanov, M. Y. Nonadiabatic Multielectron Dynamics in Strong Field Molecular Ionization. *Phys. Rev. Lett.* **2001**, *86*, 51-54.
- (23) Boguslavskiy, A. E.; Mikosch, J.; Gijsbertsen, A.; Spanner, M.; Patchkovskii, S.; Gador, N.; Vrakking, M. J. J.; Stolow, A. The Multielectron Ionization Dynamics Underlying Attosecond Strong-Field Spectroscopies. *Science* **2012**, *335*, 1336-1340.
- (24) Pavičić, D.; Lee, K. F.; Rayner, D. M.; Corkum, P. B.; Villeneuve, D. M. Direct Measurement of the Angular Dependence of Ionization for N₂, O₂, and CO₂ in Intense Laser Fields. *Phys. Rev. Lett.* **2007**, *98*, 243001.
- (25) Endo, T.; Matsuda, A.; Fushitani, M.; Yasuike, T.; Tolstikhin, O. I.; Morishita, T.; Hishikawa, A. Imaging Electronic Excitation of NO by Ultrafast Laser Tunneling Ionization. *Phys. Rev. Lett.* **2016**, *116*, 163002.
- (26) Smirnova, O.; Mairesse, Y.; Patchkovskii, S.; Dudovich, N.; Villeneuve, D.; Corkum, P.; Ivanov, M. Y. High Harmonic Interferometry of Multi-Electron Dynamics in Molecules. *Nature* **2009**, *460*, 972-977.

- (27) McFarland, B. K.; Farrell, J. P.; Bucksbaum, P. H.; Gühr, M. High Harmonic Generation from Multiple Orbitals in N₂. *Science* **2008**, *322*, 1232-1235.
- (28) Rajgara, F. A.; Krishnamurthy, M.; Mathur, D. Electron Rescattering and the Dissociative Ionization of Alcohols in Intense Laser Light. *J. Chem. Phys.* **2003**, *119*, 12224-12230.
- (29) Loh, Z.-H.; Leone, S. R. Ultrafast Strong-Field Dissociative Ionization Dynamics of CH₂Br₂ Probed by Femtosecond Soft X-Ray Transient Absorption Spectroscopy. *J. Chem. Phys.* **2008**, *128*, 204302.
- (30) Lin, M.-F.; Neumark, D. M.; Gessner, O.; Leone, S. R. Ionization and Dissociation Dynamics of Vinyl Bromide Probed by Femtosecond Extreme Ultraviolet Transient Absorption Spectroscopy. *J. Chem. Phys.* **2014**, *140*, 064311.
- (31) Lackner, F.; Chatterley, A. S.; Pemmaraju, C. D.; Closser, K. D.; Prendergast, D.; Neumark, D. M.; Leone, S. R.; Gessner, O. Direct Observation of Ring-Opening Dynamics in Strong-Field Ionized Selenophene Using Femtosecond Inner-Shell Absorption Spectroscopy. *J. Chem. Phys.* **2016**, *145*, 234313.
- (32) Chatterley, A. S.; Lackner, F.; Neumark, D. M.; Leone, S. R.; Gessner, O. Tracking Dissociation Dynamics of Strong-Field Ionized 1,2-Dibromoethane with Femtosecond XUV Transient Absorption Spectroscopy. *Phys. Chem. Chem. Phys.* **2016**, *18*, 14644-14653.
- (33) Zewail, A. H. Femtochemistry: Atomic-Scale Dynamics of the Chemical Bond. *J. Phys. Chem. A* **2000**, *104*, 5660-5694.
- (34) Loh, Z.-H.; Leone, S. R. Capturing Ultrafast Quantum Dynamics with Femtosecond and Attosecond X-ray Core-Level Absorption Spectroscopy. *J. Phys. Chem. Lett.* **2013**, *4*, 292-302.

- (35) Wang, Y.; Zhang, S.; Wei, Z.; Zhang, B. Velocity Map Imaging of Dissociative Ionization and Coulomb Explosion of CH₃I Induced by a Femtosecond Laser. *J. Phys. Chem. A* **2008**, *112*, 3846-3851.
- (36) Liu, H.; Yang, Z.; Gao, Z.; Tang, Z. Ionization and Dissociation of CH₃I in Intense Laser Field. *J. Chem. Phys.* **2007**, *126*, 044316.
- (37) Corrales, M. E.; Gitzinger, G.; González-Vázquez, J.; Lorient, V.; de Nalda, R.; Bañares, L. Velocity Map Imaging and Theoretical Study of the Coulomb Explosion of CH₃I under Intense Femtosecond IR Pulses. *J. Phys. Chem. A* **2012**, *116*, 2669-2677.
- (38) Wei, Z.; Li, J.; Wang, L.; See, S.-T.; Jhon, M. H.; Zhang, Y.; Shi, F.; Yang, M.; Loh, Z.-H. Elucidating the Origins of Multimode Vibrational Coherences of Polyatomic Molecules Induced by Intense Laser Fields. *Nat. Commun.* **2017**, *8*, 735.
- (39) Olney, T. N.; Cooper, G.; Brion, C. E. Quantitative Studies of the Photoabsorption (4.5–488 eV) and Photoionization (9–59.5 eV) of Methyl Iodide Using Dipole Electron Impact Techniques. *Chem. Phys.* **1998**, *232*, 211-237.
- (40) O’Sullivan, G.; McGuinness, C.; Costello, J. T.; Kennedy, E. T.; Weinmann, B. Trends in 4d-Subshell Photoabsorption along the Iodine Isonuclear Sequence: I, I⁺ and I²⁺. *Phys. Rev. A* **1996**, *53*, 3211-3226.
- (41) Marggi Poullain, S.; Chicharro, D. V.; González-Vázquez, J.; Rubio-Lago, L.; Bañares, L. A Velocity Map Imaging Study of the Photodissociation of the Methyl Iodide Cation. *Phys. Chem. Chem. Phys.* **2017**, *19*, 7886-7896.
- (42) Schirmer, J.; Braunstein, M.; Lee, M.-T.; McKoy, V. *VUV and Soft X-Ray Photoionization*; Plenum Press: New York; 1996.

- (43) Pettini, M.; Mazzoni, M.; Tozzi, G. P. Excitation of the Inner 4d Shell of Neutral Iodine. *Phys. Lett. A* **1981**, 82, 168-170.
- (44) Hosler, E. R.; Leone, S. R. Characterization of Vibrational Wave Packets by Core-Level High-Harmonic Transient Absorption Spectroscopy. *Phys. Rev. A* **2013**, 88, 023420.
- (45) Wang, H.; Chini, M.; Chen, S.; Zhang, C.-H.; He, F.; Cheng, Y.; Wu, Y.; Thumm, U.; Chang, Z. Attosecond Time-Resolved Autoionization of Argon. *Phys. Rev. Lett.* **2010**, 105, 143002.
- (46) Kaldun, A.; Blättermann, A.; Stooß, V.; Donsa, S.; Wei, H.; Pazourek, R.; Nagele, S.; Ott, C.; Lin, C. D.; Burgdörfer, J.; Pfeifer, T. Observing the Ultrafast Buildup of a Fano Resonance in the Time Domain. *Science* **2016**, 354, 738-741.
- (47) Walter, K.; Weinkauff, R.; Boesl, U.; Schlag, E. W. Molecular Ion Spectroscopy: Mass Selected, Resonant Two-Photon Dissociation Spectra of CH_3I^+ and CD_3I^+ . *J. Chem. Phys.* **1988**, 89, 1914-1922.
- (48) Song, Y.; Qian, X.-M.; Lau, K.-C.; Ng, C. Y.; Liu, J.; Chen, W. High-Resolution Energy-Selected Study of the Reaction $\text{CH}_3\text{X}^+ \rightarrow \text{CH}_3^+ + \text{X}$: Accurate Thermochemistry for the $\text{CH}_3\text{X}/\text{CH}_3\text{X}^+$ (X=Br, I) System. *J. Chem. Phys.* **2001**, 115, 4095-4104.
- (49) Eland, J. H. D.; Frey, R.; Kuestler, A.; Schulte, H.; Brehm, B. Unimolecular Dissociations and Internal Conversions of Methyl Halide Ions. *Int. J. Mass Spectrom.* **1976**, 22, 155-170.
- (50) Park, S. T.; Kim, S. K.; Kim, M. S. Vacuum-Ultraviolet Mass-Analyzed Threshold Ionization Spectra of Iodobutane Isomers: Conformer-Specific Ionization and Ion-Core Dissociation Followed by Ionization. *J. Chem. Phys.* **2001**, 115, 2492-2498.

- (51) Park, S. T.; Kim, S. K.; Kim, M. S. One-Photon Mass-Analyzed Threshold Ionization Spectroscopy of 1- and 2-Iodopropanes in Vacuum Ultraviolet. *J. Chem. Phys.* **2001**, *114*, 5568-5576.
- (52) Karlsson, L.; Jadrny, R.; Mattsson, L.; Chau, F. T.; Siegbahn, K. Vibrational and Vibronic Structure in the Valence Electron Spectra of CH₃X Molecules (X=F, Cl, Br, I, OH). *Phys. Scripta* **1977**, *16*, 225-234.
- (53) Grütter, M.; Michaud, J. M.; Merkt, F. Photoelectron Spectroscopic Study of the E⊗e Jahn–Teller Effect in the Presence of a Tunable Spin–Orbit Interaction. I. Photoionization Dynamics of Methyl Iodide and Rotational Fine Structure of CH₃I⁺ and CD₃I⁺. *J. Chem. Phys.* **2011**, *134*, 054308.
- (54) Lee, M.; Bae, Y. J.; Kim, M. S. K Selection in One-Photon Mass-Analyzed Threshold Ionization of CH₃I and CD₃I to the $\tilde{X}^2E_{3/2}$ State Cations. *J. Chem. Phys.* **2008**, *128*, 044310.