Maternal bonding in childhood moderates autonomic responses to distress stimuli in adult males

Dalsant, Arianna; Truzzi, Anna; Setoh, Peipei; Esposito, Gianluca

2015

Dalsant, A., Truzzi, A., Setoh, P., & Esposito, G. (2015). Maternal bonding in childhood moderates autonomic responses to distress stimuli in adult males. *Behavioural Brain Research*, 292, 428-431.

<https://hdl.handle.net/10356/84974>

<https://doi.org/10.1016/j.bbr.2015.06.026>

© 2015 Elsevier. This is the author created version of a work that has been peer reviewed and accepted for publication by Behavioural Brain Research, Elsevier. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [<http://dx.doi.org/10.1016/j.bbr.2015.06.026>].

Downloaded on 26 Apr 2023 21:33:28 SGT

Author Manuscript

Published as:

Dalsant A., Truzzi A., Setoh P., Esposito G.

Maternal bonding in childhood moderates autonomic responses to distress stimuli in adult males.

Behavioral Brain Research, 292, 428-431.

Maternal bonding in childhood moderates autonomic responses to distress stimuli in adult males

Arianna Dalsant^{a,1}, Anna Truzzi^{a,1}, Peipei Setoh^b, Gianluca Esposito^{a,b,*}

^a Affiliative Behaviour and Physiology Lab, Department of Psychology and Cognitive Science, University of Trento, Rovereto, Trento, Italy

^b Early Cognition Lab, Division of Psychology, School of Humanities and Social Sciences, Nanyang Technological University, Singapore

Highlights

- Autonomic nervous system responses to vocal distress are moderated by quality of participant maternal bonding during childhood.
- Participants who reported optimal maternal bonding showed greater calming response to distressful stimuli.
- Participants who reported non-optimal maternal bonding showed a heightened distress response to distressful stimuli.

Abstract

Mother-child bonding influences the development of cognitive and social skills. In this study we investigate how maternal attachment, developed in early childhood, modulates physiological responses to social stimuli later in life. Our results suggest that the autonomic nervous system's responses to vocal distress are moderated by the quality of participants' maternal bonding. In particular, participants with optimal maternal bonding showed a greater calming response to distressful stimuli whereas participants with non-optimal maternal bonding showed a heightened distress response.

Keywords:

Attachment, Parental Care, Physiological responses

Introduction

Attachment theory, initially described by Bowlby [1], states that humans, similar to other animal species, have an innate, evolutionarily driven biological predisposition towards social interaction. Bowlby proposed that infants actively and instinctively seek and maintain caregivers' (usually the mother) proximity and care (see for example [2]). Caregivers too, are instinctively drawn to respond and interact with their offspring [3]. Therefore, the bonds formed through attachment function to maintain security [1,4] by attenuating psychological and physiological stress reactivity [5,6] and by potentially serving as psychobiological regulators of felt security [4]. The quality of attachment developed with caregivers is crucial

because it influences both cognitive and socio-emotional development and later on, the development of adult personality. Studies have shown that high quality bonding with caregivers provides infants with an optimal relational experience that will help the child to better interact with his or her external social environment [1,4]. Early caregiving relationships influence physiological processes as well [7,8], e.g. by modulating brain sensitivity to stress and the body's capacity for managing stress-related metabolic demands [9]. These attachment-based physiological patterns of emotion regulation are thought to remain relatively stable over individuals' lifetimes, suggesting important links between adult attachment style and health related physiological processes [10]. However, there has been relatively little research looking at the normative psychobiological functions of the attachment system [11]. In this study we investigate how early attachment relationships modulate physiological responses to social stimuli. Our aim is to assess how participants, who had different interactional patterns with their parents during childhood, respond physiologically to vocal distress signals (i.e. human and animal distress vocalisations, namely cries) as adults. Specifically, we want to explore the subcortical responses mediated by the autonomic nervous system (ANS). The autonomic nervous system consists of two subsystems; (i) the parasympathetic system, which is active under resting-state or calm conditions and whose activation is manifested by a decrease in heart rate, (ii) the sympathetic nervous system, which is active in fight-or-flight arousal responses that prepare the organism to action and whose activation is manifested by an increase in heart rate and blood flow to skeletal muscles. Considering the special evolutionary salience of infant cries for women [12], we decided to test only men who were not parents. Our study is driven by two predictions: (P1) In response to distress stimuli there will be a greater calming response in participants who report better levels (higher quality) of parental bonding. (P2) Participants who report lower quality parental bonding will show an increased sympathetic response to distressful stimuli.

Methods

Participants

Forty-four non-parent male adults ($M = 24.7$, $SD = 5.06$) were recruited as participants through a University database of volunteers and by announcements posted on the University of Trento web site. All participants

were Caucasian, with Italian citizenship and had similar socio-economic status (middle–high level). All participants had heterosexual preferences and had previously shared the same household with two biological parents, at least until the age of 18. No participants had a history of psychiatric disorders or drug abuse. Informed consent was obtained from all participants. The study was conducted in agreement with the ethical principles stated in the Helsinki declaration.

Procedure

The study was conducted in two parts: the first part involved the completion of an online survey about attachment style (the parental bonding instrument (PBI)) and the second part was an experimental procedure that consisted of listening to audio files of distress cues while physiological responses were recorded at the Affiliative Behaviour and Physiology Lab in the Department of Psychology and Cognitive Science, University of Trento. After participants were briefed about the study, they gave their informed consent and were given instructions about the experimental procedure.

The PBI [13] is a 50-item self-report questionnaire developed to measure the principal parental dimensions of care and control, and is used in this study as a measure of the quality of parental bonding. In this questionnaire participants rate the parenting behavior they received from their parents during childhood. Demographic information such as age, gender, and education level was collected in addition to the PBI. The survey was created and disseminated using a web-based survey platform (maybe put which one? Survey monkey? Qualtrics?). Participants were required to answer each question on the survey, so there were no missing data for any participants. The PBI investigates individuals' attachment style toward their parents (we specifically used the mother form) by measuring two dimensions, (i) care (affection) and (ii) overprotection (level of parental control). While the concept of care seems universally valid, some studies have shown that there are cultural differences in the interpretation of the concept of overprotection. Indeed in some cultural groups, overprotection may be considered a positive value. For example, Raudino et al. [14] found that Italian mothers were more overprotective towards their offspring than English mothers. Therefore, it is likely that overprotection is normative in Italy, which is where our sample is taken from. Scores in these two dimensions are distributed as a continuum, which enabled us to identify four main groups in the population: high care and high overprotection, high care and low overprotection, low care and high overprotection, low care and low overprotection (see Fig. 1A). Participants were categorized in the four groups using the median splits.

Stimuli

The stimuli were thirty 15-s long audio clips of distressful vocalizations. There were ten audio clips of infant cries (average fundamental frequency f_0 = 410.27 Hz; f_0 max = 741.01 f_0 range = 667.84 Hz), adult woman cries (f_0 =

429.41 Hz; f_0 max = 742.10 Hz; f_0 range = 662.67 Hz) and animal cries, respectively

(average f_0 = 406.07 Hz; f_0 max = 603.82 Hz; f_0 = 662.67 Hz). Each audio clip was presented following a 10-s block of silence. All audio clips were organised into three different randomized sequences, and presentation order of the three sequences was counterbalanced across participants. The stimuli sequences were created using the open source software Audacity. Although, cries were interspaced with silence, the entire experimental experience was designed to simulate a continuously distressing situation. The level of distressfulness was tested in a pilot study, where 10 male participants (not included in the final sample) were asked to listen to and provide a behaviour judgement on the distress level felt (on a 7-point Likert scale) while listening to five different experimental sequences where the ratio between each block of crying and silence was manipulated. Participants judged the ratio of 15 s of crying to

10 s of silence as the most distressful (other options included; 5:5, 10:5, 10:10, 15:15; crying/silence in seconds).

Physiological recording and measures. Heart rate variability (HRV)

As an index of parasympathetic activity, we measured participants' heart rate variability (HRV) using a pulse oximeter (CONTEC CMS60D) placed on participants' left forefinger. Pulse oximeter as a measure of HRV has been shown to be sufficiently accurate when subjects are at rest [15], as well as when they are in non-stationary conditions (i.e. during the tilt test [16]). The standard deviation of absolute differences between successive intervals (SDSD) was used as an index of heart rate variability in time [17,18].

Nose temperature (T)

As index of sympathetic activity we measured participants' peripheral surface body temperature on the tip of the nose using a resistance thermometer (Applent at 4524) placed on participants' nose skin [19,20].

Results

Prior to data analysis, univariate and multivariate distributions of heart rate values and temperature values were examined for normality (using the Shapiro–Wilk test), homogeneity of variance (using the Levene's test), outliers, and influential cases [21]. All variables were found to be normally distributed. An analysis of variance (ANOVA) was used to determine if there was any effect of stimulus type (infant cry, woman cry and animal cry) on HRV or nose temperature. No main effect of stimulus type emerged on HRV, or on nose temperature (respectively: $F(1,77) = .001$, ns; $F(1,77) = .001$, ns).

Fig. 1(C–D) shows mean and SEM for both measures. Two way ANOVAs were performed to assess the effect of PBI-parent score and type of stimuli on each physiological response throughout the entire experimental session. A significant effect of PBI-parent score on participants' heart rate variability

($F(3,77) = 5.3$, $p = .002$, $\eta^2 = .08$) and nose temperature variability ($F(3,77) = 5.5$, $p = .002$, $\eta^2 = .06$) was found. In response to the distressful stimuli presented, participants with high scores in both care and overprotection showed greater heart rate variability compared with other participants, whereas participants with low scores in both care and overprotection showed a greater increase in surface temperature on the nose compared to other participants. Tukey HSD post hoc tests revealed significant differences in heart rate variability values for participants with high scores in both care and overprotection compared to participants in other categories. Significant differences were also found for nose temperature values in participants with low scores in both care and overprotection compared to participants in the other categories.

A. Dalsant et al. / Behavioural Brain Research 292 (2015) 428–431

Fig. 1. Physiological responses to distress stimuli in participants with different maternal bonding styles. (A) Four possible outcomes of the interaction between the two maternal bonding dimensions, care and overprotection. We divided participants depending on the levels, high (+) or low (-), they showed in these two dimensions (care/overprotection): (+/+) high care and high overprotection, (+/-) high care and low overprotection, (-/+) low care and high overprotection, (-/-) low care and low overprotection. (B) Mean of the heart rate variability index (SDSD) for each group in response to distress stimuli. (C) Nose mean temperature variation for each group in response to distress stimuli. * $p < .05$. All numerical data are shown as mean $\pm$ SEM.

Discussion

Previous studies [5,22] have highlighted that the attachment formed with caregivers plays a critical role in the development of children's emotion regulation – the process by which individuals modulate their positive and negative emotional responses to internal and external stimuli [23] – throughout their lives [9,10]. This study has shown how early parental bonding is related to adults' emotion regulation.

It is not easy to objectively evaluate parenting quality because parental bonding is a complex process (or phenomenon); it is informed by the common characteristics between parent and child, and by a history of interactions and mutual transactions that take place over time [24]. Despite this complexity, it is generally agreed that parenting is a process that involves biological, social

and relational components related to care and protecting one's offspring. The results of this study suggest that adults' perception of their own maternal bonding affects their emotional reactions to distressful situations. In line with our predictions, there are differential ANS activation patterns between individuals with different parental bonding styles. Specifically, participants with better early parental bonding — those high on both care and overprotection, were able to react to the distressful situation with enhanced calming responses compared to participants who were low on either care or overprotection. Thus, experiencing high levels of both care and overprotection during childhood appear to be necessary in order to display the most optimal response towards a stressful situation in adulthood. Participants with the lowest quality parental

bonding — those low on both care and overprotection, reacted to the distressful situation with elevated distress responses compared to participants who were high on either care or protection. This finding suggests that being high on just one of the dimensions was a protective factor against a heightened distress response during adversity. These findings underline the presence of a continuum in parental bonding effects, such that when an individual is exposed to a potentially aversive situation, only those individuals with the highest quality parental bonding style will display calming responses, whereas only those with the lowest quality parental bonding style will show heightened distress responses. Displaying a heightened distress response when facing negative stimuli is an indication of a sub-optimal emotion regulation system that is less able to cope with adversity. Higher quality early parental bonding leads to a greater control of bodily arousal in the face of distressful stimuli. Thus, as our results suggest, early parental bonding affects emotional regulation in adult males. This relationship between parental bonding style experienced during childhood and the emotional regulation abilities present in adulthood could also

be related to the development of psychopathology. An excessive readiness in the activation of a distress response has been found to be the physiological correlate of several psychopathologies (i.e. ADHD). On the contrary, optimal parental bonding in childhood could be a protective factor against psychopathology, because individuals may develop the ability to deal with aversive stimuli with a calming response.

Before concluding, we would like to acknowledge some limitations of our study as well as possible future directions. First, we tested only male participants. Future research should examine (i) how parental bonding affects emotion regulation in adult women, and (ii) how participants who are parents might differ on their physiological arousal and perception of maternal bonding compared to non-parents. Furthermore, in addition to self-reports, other objective measures should be used to evaluate levels of parental bonding. Finally, future research should examine how different cultures view maternal overprotection, and how such norms affect individuals' perception of their own parental bonding which, in turn, may affect physiological pathways and coping mechanisms.

References

- [1] J. Bowlby, *Attachment. Attachment and Loss*, Vol. I, Hogarth, London, 1969.
- [2] G. Esposito, S. Yoshida, R. Ohnishi, Y. Tsuneoka, M.D.C. Rostagno, S. Yokota, et al., Infant calming responses during maternal carrying in humans and mice, *Curr. Biol.* 23 (2013) 739–745.
- [3] G. Esposito, S. Valenzi, T. Islam, C. Mash, M.H. Bornstein, Immediate and selective maternal brain responses to own infant faces, *Behav. Brain Res.* 278 (2015) 40–43.
- [4] D.A. Sbarra, C. Hazan, Coregulation, dysregulation, self-regulation: an integrative analysis and empirical agenda for understanding adult attachment, separation, loss, and recovery, *Pers. Soc. Psychol. Rev.* 12 (2) (2008) 141–167.
- [5] L.M. Diamond, A.M. Hicks, Psychobiological perspectives on attachment: implications for health over the lifespan, in: J.A. Simpson, W.S. Rholes (Eds.), *Adult Attachment: New Directions and Emerging Issues*, Guilford Press, New York, 2004, pp. 240–263.
- [6] C. Schoenmaker, R. Huffmeijer, M.H. van IJzendoorn, M.J. Bakermans-Kranenburg, L. van den Dries, M. Linting, et al., Attachment and physiological reactivity to infant crying in young adulthood: dissociation between experiential and physiological arousal in insecure adoptees, *Phys. Behav.* 139 (2015) 549–556.
- [7] D. Glaser, Child abuse and neglect and the brain-a review, *J. Child Psychol. Psychiatry* 41 (1) (2000) 97–116.
- [8] M.J. Meaney, Maternal care, gene expression, and the transmission of individual differences in stress reactivity across generations, *Ann. Rev. Neurosci.* 24 (1) (2001) 1161–1192.
- [9] S.W. Porges, J.A. Doussard-Roosevelt, A.K. Maiti, Vagal tone and the physiological regulation of emotion, in: N. Fox (Ed.), *The Development of Emotion Regulation: Biological and Behavioral Considerations*, Monographs of the Society for Research in Child Development, 1994, pp. 167–186.
- [10] L.M. Diamond, A.M. Hicks, K. Otter-Henderson, Physiological evidence for repressive coping among avoidantly attached adults, *J. Soc. Pers. Relat.* 23 (2) (2006) 205–229.
- [11] T.F. Robles, H.S. Kane, The attachment system and physiology in adulthood: normative processes, individual differences, and implications for health, *J. Pers.* 82 (6) (2014) 515–527.

- [12] N. De Pisapia, M.H. Bornstein, P. Rigo, G. Esposito, S. De Falco, P. Venuti, Sex differences in directional brain responses to infant hunger cries, *Neuroreport* 24 (3) (2013) 142–146.
- [13] G. Parker, H. Tupling, L.B. Brown, Parental bonding instrument (PBI), *Br. J. Med. Psychol.* 52 (1979) 1–10.
- [14] A. Raudino, L. Murray, C. Turner, E. Tsampala, A. Lis, L. De Pascalis, et al., Child anxiety and parenting in England and Italy: the moderating role of maternal warmth, *J. Child Psychol. Psychiatry* 54 (12) (2013) 1318–1326.
- [15] A. Schäfer, J. Vagedes, How accurate is pulse rate variability as an estimate of heart rate variability? A review on studies comparing photoplethysmographic technology with an electrocardiogram, *Int. J. Cardiol.* 166 (1) (2013) 15–29.
- [16] E. Gil, M. Orini, R. Bailón, J.M. Vergara, L. Mainardi, P. Laguna, Photoplethysmography pulse rate variability as a surrogate measurement of heart rate variability during non-stationary conditions, *Physiol. Meas.* 31 (9) (2010) 1271.
- [17] J.M. Dekker, R.S. Crow, A.R. Folsom, P.J. Hannan, D. Liao, C.A. Swenne, et al., Low heart rate variability in a 2-minute rhythm strip predicts risk of coronary heart disease and mortality from several causes, *Circ* 102 (2000) 1239–1244.
- [18] M.G. Tsipouras, D. Fotiadis, Automatic arrhythmia detection based on time and time-frequency analysis of heart rate variability, *Comput. Methods Program Biomed.* 74 (2) (2004) 95–108.
- [19] G. Esposito, J. Nakazawa, S. Ogawa, R. Stival, A. Kawashima, D.L. Putnick, et al., Baby, you light-up my face: culture-general physiological responses to infants and culture-specific cognitive judgements of adults, *PLoS One* 9 (10) (2014) e106705.
- [20] G. Esposito, J. Nakazawa, S. Ogawa, R. Stival, D.L. Putnick, M.H. Bornstein, Using infrared thermography to assess emotional responses to infants, *Early Child Dev. Care* 185 (3) (2014) 438–447.
- [21] J. Fox, *Applied Regression Analysis, Linear Models, and Related Methods*, Sage Publications, Thousand Oaks, CA, US, 1997.
- [22] R.L. Repetti, S.E. Taylor, T.E. Seeman, Risky families: family social environments and the mental and physical health of offspring, *Psychol. Bull.* 128 (2) (2002) 330.

[23] R. Thompson, Emotion regulation: a theme in search of definition, in: N. Fox (Ed.), *The Development of Emotion Regulation: Biological and Behavioral Considerations*, 59, Monographs of the Society for Research in Child Development, 1994, pp. 225–252.

[24] M.H. Bornstein, C. Hahn, J.T. Suwalsky, O.M. Haynes, Socioeconomic Status, Parenting and Child Development: The Hollingshead Four-Factor Index of Social Status and the Socioeconomic Index of Occupations, in: M.H. Bornstein, R. Bradley (Eds.), *Socioeconomic Status, Parenting and Child Development*, 2003, pp. 29–82.