Rapid access to bicyclic δ -lactones via carbene-catalyzed activation and cascade reaction of unsaturated carboxylic esters

Fu, Zhenqian; Wu, Xingxing; Chi, Robin Yonggui

2016

Fu, Z., Wu, X., & Chi, R. Y. (2016). Rapid access to bicyclic δ -lactones via carbene-catalyzed activation and cascade reaction of unsaturated carboxylic esters. *Organic Chemistry Frontiers*, 3(2), 145-149.

<https://hdl.handle.net/10356/87048>

<https://doi.org/10.1039/C5QO00348B>

© 2016 The Partner Organisations. This is the author created version of a work that has been peer reviewed and accepted for publication by *Organic Chemistry Frontiers*, The Partner Organisations. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [<http://dx.doi.org/10.1039/C5QO00348B>].

Downloaded on 26 Apr 2023 17:50:09 SGT

Rapid Access to Iridoids via Carbene-Catalyzed Activation and Cascade Reaction of Unsaturated Carboxylic Esters

Zhenqian Fu, Xingxing Wu and Yonggui Robin Chi*

Abstract: Carboxylic esters are excellent choice of substrates in organic synthesis. Here we demonstrate that carbene-catalyzed activation of α,β -unsaturated ester can be developed to synthesis sophisticated multi-cyclic molecules in a single step cascade process. The iridoid-type lactone products are obtained with highly stereoselectivities, and can readily undergo further transformations.

Iridoids are a class of secondary metabolites found in a wide variety of plants and in some animals.^[1] This class of molecules and their natural or synthetic analogs usually consist of a cyclopentane ring fused to a six-membered oxygen heterocycle (Figure 1a). Significantly, these molecules have been found to exhibit useful bioactivity, such as neuroprotective effect,^[2a] anti-cancer,^[2b] anti-tumour,^[2c] anti-inflammatory,^[2d] anti-oxidant,^[2e] and anti-bacterial^[2f] activity. Considerable effort has been spent in the synthesis of this class of molecules via both biosynthetic approach^[3] and organic synthesis.^[4,5] However, the majority of methods involve relatively long synthetic steps, sometimes harsh conditions and often result in racemic products.^[5] Our laboratory is interested in catalytically activating readily available substrates for rapid (ideally single-step) asymmetric synthesis of functional molecules.^[6] We have demonstrated that stable esters^[7] and α,β -unsaturated esters^[8] can be activated by carbene organic catalysts for efficient and selective reactions.^[9] Here we report a single step access to optically enriched iridoids by using our catalytic ester activation approach (Figure 1b). Mechanistically, the reaction starts via the addition of a carbene catalyst to an α,β -unsaturated ester followed by a cascade process^[10,11] involving two Michael additions and a lactonization step. The bicyclic products were obtained in excellent yields,

diastereoselectivities and enantioselectivities.

Figure 1. Iridoids derivatives in natural products (a) and our synthetic approach (b). (Glc = β -glucopyranosyl; EWG = electron-withdrawing group)

[*] Dr. Z. Fu,^[+] X. Wu,^[+] Prof. Dr. Y. R. Chi, Division of Chemistry & Biological Chemistry, School of Physical & Mathematical Sciences, Nanyang Technological University, Singapore 637371 (Singapore)

E-mail: robinchi@ntu.edu.sg

Prof. Dr. Y. R. Chi, Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering, Key Laboratory of Green Pesticide and Agricultural Bioengineering, Ministry of Education, Guizhou University, Huaxi District, Guiyang 550025 (China)

Dr. Z. Fu, current address: Key Lab for Flexible Electronics and Institute of Advanced Materials, Jiangsu National Synergistic Innovation Center for Advanced Materials (SICAM), Nanjing Tech University, Nanjing 211816 (China)

[+] These authors contributed equally to this work.

[**] We acknowledge support from Singapore's National Research Foundation (NRF), Ministry of Education (MOE), Nanyang Technological University (NTU), and China's National Key Program for Basic Research (No. 2010CB126105), Thousand Talent Plan, National Natural Science Foundation (No. 21132003; No. 21472028), Guizhou Province Returned Oversea Student Science and Technology Activity Program, and Guizhou University.

We started by using unsaturated ester **1a** and enone malonate **2a** as model substrates to search for suitable conditions, as shown in Table 1. Achiral triazolium pre-catalyst **A**^[12] in the presence of DBU as a base and LiCl^[13,14] as an additive could mediate this reaction to afford proposed cascade product **3a** in 90% (entry 1). The use of chiral triazolium pre-catalyst (e.g. **B**^[15]) based on the scaffold of **A** led to **3a** with poor er (entry 2). We also found that catalyst **C**^[16] could promote this reaction with much improved er (entry 3). Replacing of the two hydrogen substituents in **C** with methyl units (catalyst **D**^[17]) could enhance the product er from 77:23 to 90:10, albeit with a lower yield (entry 4). The low yield in the reactions (entries 2-4) was due to the competing ester hydrolysis. We reasoned that increasing the nucleophilicity of the carbene catalyst could likely promote the desired reaction between the ester and the carbene. Thus we next replaced the methyl substituents on the N-mesityl group of catalyst **D** with methoxyl units to give catalyst **E**^[18]. By using catalyst **E** with DABCO (0.3 equivalent) and DBU (1.0 equivalent) as the base, the reaction proceeded

Table 1. Screening of reaction conditions for the reaction of **1a** with **2a**.^[a]

Entry	NHC	Yield [%] ^[b]	dr ^[b]	er ^[c]
1	A	90	20:1	-
2	B	55	20:1	57:43
3	C	43	10:1	77:23
4	D	25	12:1	90:10
5 ^d	E	88	11:1	91:9
6 ^d	F	74	12:1	96:4
7 ^{d,e}	F	92(85) ^f	12:1	96:4

[a] Standard condition: NHC precursor (20 mol%), **1a** (0.1 mmol), **2a** (0.1 mmol), DBU (1.0 equiv), LiCl (1.0 equiv), THF (1.0 mL), 4 Å MS, rt, 3 h. [b] Yields and diastereomeric ratio of **3a**, determined via ¹H NMR analysis of unpurified reaction mixtures. 1,3,5-trimethoxybenzene was used as an internal standard. Absolute configuration of product was determined via X-ray of **4o** (Table 3). [c] Determined via chiral phase HPLC analysis. [d] 0.3 eq DABCO was added, then 1.0 eq DBU was added after 20 min. [e] 0.2 eq HOBt was added and 1.2 eq **2a** and 1.4 eq DBU was used instead. [f] Isolated yield in parenthesis. DBU=1,8-diazabicyclo [5.4.0] undec-7-ene; Mes=mesityl (2, 4, 6-trimethylphenyl); DABCO=1,4-Diazabicyclo[2.2.2]octane; HOBt=Hydroxybenzotriazole.

Table 2. Scope of ester **1**.^[a]

R=H; 3a ; 85% yield; 12:1 d.r.; 96:4 e.r. R=OMe; 3b ; 40% yield; 20:1 d.r.; 95:5 e.r. R=F; 3c ; 62% yield; 17:1 d.r.; 95:5 e.r. R=Cl; 3d ; 84% yield; 11:1 d.r.; 96:4 e.r. R=Br; 3e ; 85% yield; 13:1 d.r.; 97:3 e.r. R=NO ₂ ; 3f ; 65% yield; 14:1 d.r.; 97:3 e.r.				
3g ; 85% yield, 7:1 d.r.; 98:2 e.r. 3h ; 75% yield, 18:1 d.r.; 96:4 e.r. 3i ; 88% yield, 13:1 d.r.; 96:4 e.r. 3j ; 88% yield, 13:1 d.r.; 96:4 e.r. 3k ; 74% yield, 20:1 d.r.; 95:5 e.r. 3l ; 49% yield, 15:1 d.r.; 94:6 e.r. 3m ; 50% yield, 10:1 d.r.; 94:6 e.r. 3n ; 53% yield, 20:1 d.r.; 75:25 e.r. 3o ; 58% yield, 7:1 d.r.; 53:47 e.r.				

[a] Conditions as in Table 1, entry 7; yields (after SiO₂ chromatography purification) were based on ester **1**.

smoothly to give **3a** in 88% yield, 11:1 dr and 91:9 er (entry 5). Replacing the methyl units in **E** with ethyl substituents (to give catalyst **F**) could further improve the reaction stereoselectivities (entry 6). The reaction yield (with catalyst **F**) could be further improved by using a catalytical amount of HOBt as an additive without affecting the product er (85% yield, 12:1 dr, 96:4 er; entry 7).

With an acceptable condition in hand (Table 1, entry 7), we next studied the generality of the ester substrate in our reaction by using β -enone malonate **2a** as a model substrate (Table 2). Installation of various substituents (**1b-h**) at the para-carbon of the β -phenyl group of **1** was all tolerated, except strong electron-releasing group (CH₃O, **1b**) where a lower yield was observed under standard conditions. Placing the substituent at different positions of the β -phenyl group of **1** has no apparent effect on the reaction outcome (**3e** and **3h**). The β -phenyl unit of **1** could be replaced by naphthyl (**1i**) or heteroaryl (**1j-k**) units. The β -aryl unit of unsaturated ester could also be replaced by an alkenyl (**1l**) or ester (**1m**) substituent. Esters with β -alkyl substituent (**1n**) and acetic ester (**1o**) could also undergo desired reactions, albeit with a dropped yield and er.

Table 3. Scope of β -enone malonate **2**.^[a]

R=OMe; 4a ; 72% yield; 20:1 d.r.; 94:6 e.r. R=Me; 4b ; 82% yield; 20:1 d.r.; 96:4 e.r. R=F; 4c ; 75% yield; 20:1 d.r.; 96:4 e.r. R=Cl; 4d ; 58% yield; 20:1 d.r.; 96:4 e.r. R=Br; 4e ; 56% yield; 19:1 d.r.; 96:4 e.r.				
4f ; 78% yield, 20:1 d.r.; 96:4 e.r. 4g ^b ; 49% yield, 20:1 d.r.; 94:6 e.r. 4h ^b ; 52% yield, 20:1 d.r.; 95:5 e.r. 4i ; 99% yield, 20:1 d.r.; 98:2 e.r. 4j ; 76% yield, 10:1 d.r.; 97:3 e.r. 4k ; 57% yield, 20:1 d.r.; 99:1 e.r. 4l ; 86% yield, 12:1 dr, 95:5 e.r. 4m ; 83% yield, 13:1 d.r.; 96:4 e.r. 4n ; 62% yield, 12:1 d.r.; 96:4 e.r. 4o ; 78% yield, 13:1 d.r.; 97:3 e.r. (X-ray)				

[a] Conditions as in Table 1, entry 7; yields (after SiO₂ chromatography purification) were based on ester **1**. [b] Reactions run with 30 mmol% NHC and 1.6 equivalents of enone.

The scope of the β -enone malonate was then evaluated (Table 3). Different (hetero)aryl substituents on the enone part were all

tolerated (**4a-h**). Remarkably, substituting on the α -carbon of the enone part did not affect the reaction outcome (**4i-j**). The tolerance of substituent on the enone α -carbon allows us to prepare the multi-cyclic Steroid derivative^[19] **4j** in 76% yield, 10:1 dr and 97:3 er. Not surprisingly, the two methyl alcohol ester groups of the malonate substrates could be changed to ethyl (**4m**) or benzyl (**4n**) alcohol ester units.

Table 4. Scope of ester **1** and enone **4**.^[a]

[a] Conditions as in Table 1, entry 7; yields (after SiO₂ chromatography purification) were based on ester **1**. [b] Reactions run at -78 °C. [c] After one recrystallization. [d] Reactions run with 30 mmol% NHC and 1.5 equivalents of ester. Yield (after SiO₂ chromatography purification) was based on enone **5**.

Our strategy of forming bicyclic product (**3** and **4**) with fused five and six-membered rings could be directly extended to prepare fused six and six-ring products. When γ -enone malonate **5** was used to react with the unsaturated ester substrate, fused bicyclic product **6** could be obtained with excellent yield, and er under the above conditions without further optimizations (Table 4). Interestingly, γ -nitro enone was also an effective substrate, leading to product **6e-g** bearing synthetically useful nitro group. The reaction of γ -nitro enone was performed at -78 °C to give **6e** in 87:13 er (99:1 er after a single recrystallization).

Scheme 1. Synthetic transformations of cascade product.

The optically enriched products from our catalytic reactions could undergo further transformations under simple conditions. For example, the enol ester carbon-carbon double bond of **3a** could be reduced to give **7** or oxidized to give **8**, bearing a significant core structure in many biologically active natural products^[20]. The nitro group of **6g** could be removed to give **9** (with a ring opening ester exchange of the lactone).

In summary, we have developed a highly efficient single-step approach for the preparation of multi-cyclic lactone products with excellent optical purity. Our reactions proceed through carbene-catalyzed activation of unsaturated esters as the key step, followed by a highly selective cascade processes. The multi-cyclic products from our catalytic reaction can readily undergo further transformations. Our study demonstrates the synthetic power in turning readily available and stable ester substrates to sophisticated products via an extremely short route, and shall encourage further development and application of carbene catalysis and ester activation.

Received: ((will be filled in by the editorial staff))

Published online on ((will be filled in by the editorial staff))

Keywords: iridoid • bicyclic ring • organocatalysis • N-heterocyclic carbene • cascade reaction

- [1] a) L. J. Elnaggar, J. L. Beal, *J. Nat. Prod.* **1980**, *43*, 649-707; b) E. L. Ghisalberti, *Phytomedicine* **1998**, *5*, 147-163; c) B. Dinda, S. Debnath, Y. Harigaya, *Chem. Pharm. Bull.* **2007**, *55*, 159-222; d) B. Dinda, S. Debnath, R. Banik, *Chem. Pharm. Bull.* **2011**, *59*, 803-833.
- [2] a) J. Xu, P. Zhao, Y. Q. Guo, C. F. Xie, D. Q. Jin, Y. G. Ma, W. B. Hou, T. J. Zhang, *Fitoterapia* **2011**, *82*, 1133-1136; b) S. Pandeti, K. Sharma, S. R. Bathula, N. Tadigoppula, *Phytomedicine* **2014**, *21*, 333-339; c) H. K. Hamdi, R. Castellon, *Biochem. Bioph. Res. Co.* **2005**, *334*, 769-778; d) W. J. Zhu, M. Q. Pang, L. Y. Dong, X. Y. Huang, S. M. Wang, L. L. Zhou, *Life Sci.* **2012**, *91*, 369-376; e) E. J. Jeong, K. Y. Lee, S. H. Kim, S. H. Sung, Y. C. Kim, *Eur. J. Pharmacol.* **2008**, *588*, 78-84; f) M. Modaresi, A. Delazar, H. Nazemiyeh, F. Fathi-Azad, E. Smith, M. M. Rahman, S. Gibbons, L. Nahar, S. D. Sarker, *Phytother. Res.* **2009**, *23*, 99-103.
- [3] F. Geu-Flores, N. H. Sherden, V. Courdavault, V. Burlat, W. S. Glenn, C. Wu, E. Nims, Y. H. Cui, S. E. O'Connor, *Nature*, **2012**, *492*, 138-142.
- [4] a) E. M. Phillips, M. Wadamoto, A. Chan, K. A. Scheidt, *Angew. Chem.* **2007**, *119*, 3167-3170; *Angew. Chem. Int. Ed.* **2007**, *46*, 3107-3110; b) Y. Li, X. Q. Wang, C. Zheng, S. L. You, *Chem. Commun.* **2009**, 5823-5825; c) L. Candish, D. W. Lupton, *Org. Lett.* **2010**, *12*, 4836-4839; d) D. Belmessieri, L. C. Morrill, C. Simal, A. M. Z. Slawin, A. D. Smith, *J. Am. Chem. Soc.* **2011**, *133*, 2714-2720; e) A. Biswas, S. De Sarkar, R. Frohlich, A. Studer, *Org. Lett.* **2011**, *13*, 4966-4969.
- [5] a) S. H. Lim, M. D. Curtis, P. Beak, *Org. Lett.* **2001**, *3*, 711-714; b) A. Srikrishna, M. S. Rao, *Tetrahedron Lett.* **2002**, *43*, 151-154; c) T. I. Richardson, J. A. Dodge, G. L. Durst, L. A. Pfeifer, J. Shah, Y. Wang, J. D. Durbin, V. Krishnan, B. H. Norman, *Bioorg. Med. Chem. Lett.* **2007**, *17*, 4824-4828; d) J. H. Joo, S. W. Youn, *Adv. Synth. Catal.* **2013**, *355*, 559-568; e) B. Zhou, Z. Luo, Y. C. Li, *Chem.-Eur. J.* **2013**, *19*, 4428-4431; f) X. W. Fan, Y. Cheng, *Org. Biomol. Chem.* **2014**, *12*, 123-131.
- [6] a) J. M. Mo, L. Shen, Y. R. Chi, *Angew. Chem.* **2013**, *125*, 8750-8753; *Angew. Chem. Int. Ed.* **2013**, *52*, 8588-8591; b) J. M. Mo, R. J. Yang, X. K. Chen, B. Tiwari, Y. R. Chi, *Org. Lett.* **2013**, *15*, 50-53; c) M. Wang, Z. J. Huang, J. F. Xu, Y. R. Chi, *J. Am. Chem. Soc.* **2014**, *136*, 1214-1217; d) T. S. Zhu, P. C. Zheng, C. L. Mou, S. Yang, B. A. Song, Y. R. Chi, *Nat. Commun.* **2014**, *5*, 5027; e) Y. X. Zhang, Y. Du, Z. J. Huang, J. F. Xu, X. X. Wu, Y. H. Wang, M. Wang, S. Yang, R. D. Webster, Y. R. Chi, *J. Am. Chem. Soc.* **2015**, *137*, 2416-2419.
- [7] a) L. Hao, Y. Du, H. Lv, X. Chen, H. Jiang, Y. Shao, Y. R. Chi, *Org. Lett.* **2012**, *14*, 2154-2157; b) L. Hao, S. Chen, J. Xu, B. Tiwari, Z. Fu,

- T. Li, J. Lim, Y. R. Chi, *Org. Lett.* **2013**, *15*, 4956-4959; c) L. Hao, C. W. Chuen, R. Ganguly, Y. R. Chi, *Synlett* **2013**, 1197-1200; d) Z. Fu, J. Xu, T. Zhu, W. W. Y. Leong, Y. R. Chi, *Nat. Chem.* **2013**, *5*, 835-839; e) S. Chen, L. Hao, Y. Zhang, B. Tiwari, Y. R. Chi, *Org. Lett.* **2013**, *15*, 5822-5825; f) Z. Fu, K. Jiang, T. S. Zhu, J. Torres, Y. R. Chi, *Angew. Chem.* **2014**, *126*, 6624-6628; *Angew. Chem. Int. Ed.* **2014**, *53*, 6506-6510.
- [8] a) J. Cheng, Z. Huang, Y. R. Chi, *Angew. Chem.* **2013**, *125*, 8754-8758; *Angew. Chem. Int. Ed.* **2013**, *52*, 8592-8596; b) J. Xu, Z. Jin, Y. R. Chi, *Org. Lett.* **2013**, *15*, 5028-5031; Also see: c) S. J. Ryan, L. Candish, D. W. Lupton, *J. Am. Chem. Soc.* **2009**, *131*, 14176-14177;
- [9] For selected examples: a) B. E. Maki, A. Chan, E. M. Phillips, K. A. Scheidt, *Org. Lett.* **2007**, *9*, 371-374; b) J. Guin, S. De Sarkar, S. Grimme, A. Studer, *Angew. Chem.* **2008**, *120*, 8855-8858; *Angew. Chem. Int. Ed.* **2008**, *47*, 8727-8730; c) B. E. Maki, K. A. Scheidt, *Org. Lett.* **2008**, *10*, 4331-4334; d) S. De Sarkar, S. Grimme, A. Studer, *J. Am. Chem. Soc.* **2010**, *132*, 1190-1191; e) S. De Sarkar, A. Studer, *Angew. Chem.* **2010**, *122*, 9452-9455; *Angew. Chem. Int. Ed.* **2010**, *49*, 9266-9269; f) Z. Q. Rong, M. Q. Jia, S. L. You, *Org. Lett.* **2011**, *13*, 4080-4083; g) F. G. Sun, L. H. Sun, S. Ye, *Adv. Synth. Catal.* **2011**, *353*, 3134-3138; h) B. Wanner, J. Mahatthananchai, J. W. Bode, *Org. Lett.* **2011**, *13*, 5378-5381; i) A. G. Kravina, J. Mahatthananchai, J. W. Bode, *Angew. Chem.* **2012**, *124*, 9568-9572; *Angew. Chem. Int. Ed.* **2012**, *51*, 9433-9436; j) J. Mahatthananchai, J. Kaeobamrung, J. W. Bode, *Acs Catal.* **2012**, *2*, 494-503; k) R. C. Samanta, B. Maji, S. De Sarkar, K. Bergander, R. Frohlich, C. Muck-Lichtenfeld, H. Mayr, A. Studer, *Angew. Chem.* **2012**, *124*, 5325-5329; *Angew. Chem. Int. Ed.* **2012**, *51*, 5234-5238; l) S. De Sarkar, A. Biswas, R. C. Samanta, A. Studer, *Chem.-Eur. J.* **2013**, *19*, 4664-4678; m) J. Mahatthananchai, J. W. Bode, *Acc. Chem. Res.* **2014**, *47*, 696-707; n) Z. J. Wu, F. Y. Li, J. Wang, *Angew. Chem.* **2015**, *127*, 1249-1253; *Angew. Chem. Int. Ed.* **2015**, *54*, 1629-1633.
- [10] For reviews, see: C. Grondal, M. Jeanty, D. Enders, *Nat. Chem.* **2010**, *2*, 167-178. For selected examples of NHC-catalyzed cascade reactions, see: a) V. Nair, S. Vellalath, M. Poonoth, E. Suresh, *J. Am. Chem. Soc.* **2006**, *128*, 8736-8737; b) P. C. Chiang, J. Kaeobamrung, J. W. Bode, *J. Am. Chem. Soc.* **2007**, *129*, 3520-3521; c) M. He, J. W. Bode, *J. Am. Chem. Soc.* **2008**, *130*, 418-419; d) B. Cardinal-David, D. E. A. Raup, K. A. Scheidt, *J. Am. Chem. Soc.* **2010**, *132*, 5345-5347; e) D. T. Cohen, B. Cardinal-David, K. A. Scheidt, *Angew. Chem.* **2011**, *123*, 1716-1720; *Angew. Chem. Int. Ed.* **2011**, *50*, 1678-1682; f) E. Sanchez-Larios, J. M. Holmes, C. L. Daschner, M. Gravel, *Synthesis* **2011**, 1896-1904; g) L. Candish, D. W. Lupton, *J. Am. Chem. Soc.* **2013**, *135*, 58-61; h) L. Candish, C. M. Forsyth, D. W. Lupton, *Angew. Chem.* **2013**, *125*, 9319-9322; *Angew. Chem. Int. Ed.* **2013**, *52*, 9149-9152; i) Z. J. Wu, X. Wang, F. Y. Li, J. C. Wu, J. Wang, *Org. Lett.* **2015**, *17*, 3588-3591.
- [11] Ye group recently reported a similar cascade reaction of enals via oxidative NHC catalysis, see: Z. Q. Liang, D. L. Wang, H. M. Zhang, S. Ye, *Org. Lett.* **2015**, ASAP, DOI: 10.1021/acs.orglett.5b02695.
- [12] K. B. Ling, A. D. Smith, *Chem. Commun.* **2011**, *47*, 373-375.
- [13] In the absence of LiCl, the product was formed with 49% NMR yield.
- [14] For selective studies using Lewis acid as additives in NHC catalysis, see: a) B. Cardinal-David, D. E. A. Raup, K. A. Scheidt, *J. Am. Chem. Soc.* **2010**, *132*, 5345-5347; b) D. E. A. Raup, B. Cardinal-David, D. Holte, K. A. Scheidt, *Nat. Chem.* **2010**, *2*, 766-771; c) A. M. ElSohly, D. A. Wespe, T. J. Poore, S. A. Snyder, *Angew. Chem.* **2013**, *125*, 5901-5906; *Angew. Chem. Int. Ed.* **2013**, *52*, 5789-5794; d) J. Qi, X. G. Xie, R. F. Han, D. H. Ma, J. Yang, X. G. She, *Chem.-Eur. J.* **2013**, *19*, 4146-4150; e) S. Bera, R. C. Samanta, C. G. Daniliuc, A. Studer, *Angew. Chem.* **2014**, *126*, 9776-9780; *Angew. Chem. Int. Ed.* **2014**, *53*, 9622-9626; f) H. Y. Dang, Z. T. Wang, Y. Cheng, *Org. Lett.* **2014**, *16*, 5520-5523.
- [15] P.-C. Chiang, M. Rommel, J. W. Bode, *J. Am. Chem. Soc.* **2009**, *131*, 8714-8718.
- [16] I. Piel, M. Steinmetz, K. Hirano, R. Frohlich, S. Grimme, F. Glorius, *Angew. Chem.* **2011**, *123*, 5087-5091; *Angew. Chem. Int. Ed.* **2011**, *50*, 4983-4987.
- [17] C. G. Goodman, M. M. Walker, J. S. Johnson, *J. Am. Chem. Soc.* **2015**, *137*, 122-125.
- [18] F. Liu, X. Bugaut, M. Schedler, R. Frohlich, F. Glorius, *Angew. Chem.* **2011**, *123*, 12834-12838; *Angew. Chem. Int. Ed.* **2011**, *50*, 12626-12630.
- [19] a) J. S. R. Hanson, *Nat. Prod. Rep.* **2010**, *27*, 887-899; b) J. P. Munafo Jr, T. J. Gianfagna, *Nat. Prod. Rep.* **2015**, *32*, 454-477; c) A. Ali, M. Asif, H. Khanam, A. Mashrai, M. A. Sherwani, M. Owais, S. Shamsuzzaman, *RSC Adv.* **2015**, *5*, 75964-75984.
- [20] a) R. Bandichhor, B. Nosse, O. Reiser, *Top. Curr. Chem.* **2005**, *243*, 43-72; b) R. R. A. Kitson, A. Millemaggi, R. J. K. Taylor, *Angew. Chem.* **2009**, *121*, 9590-9615; *Angew. Chem. Int. Ed.* **2009**, *48*, 9426-9451; c) Y. Li, G. Pattenden, *Nat. Prod. Rep.* **2011**, *28*, 429-440; d) S. Lee, J. H. Jang, G.-S. Hwang, D. H. Ryu, *J. Org. Chem.* **2013**, *78*, 770-775.