Extended Mellin integral representations for the absolute value of the gamma function

Privault, Nicolas

2018

Privault, N. (2018). Extended Mellin integral representations for the absolute value of the gamma function. *Analysis*, 38(1), 11-20.

<https://hdl.handle.net/10356/89152>

<https://doi.org/10.1515/anly-2017-0046>

© 2018 Walter de Gruyter GmbH, Berlin/Boston. This paper was published in *Analysis* and is made available as an electronic reprint (preprint) with permission of Walter de Gruyter GmbH, Berlin/Boston. The published version is available at: [<http://dx.doi.org/10.1515/anly-2017-0046>]. One print or electronic copy may be made for personal use only. Systematic or multiple reproduction, distribution to multiple locations via electronic or other means, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper is prohibited and is subject to penalties under law.

Downloaded on 26 Apr 2023 19:38:55 SGT

Nicolas Privault*

Extended Mellin integral representations for the absolute value of the gamma function

<https://doi.org/10.1515/anly-2017-0046>

Received September 10, 2017; accepted December 29, 2017

Abstract: We derive Mellin integral representations in terms of Macdonald functions for the squared absolute value $s \mapsto |\Gamma(a + is)|^2$ of the gamma function and its Fourier transform when $a < 0$ is non-integer, generalizing known results in the case $a > 0$. This representation is based on a renormalization argument using modified Bessel functions of the second kind, and it applies to the representation of the solutions of a Fokker–Planck equation.

Keywords: Gamma function, Mellin transform, Harman–Watson distribution

MSC 2010: 32A26, 33C10, 33B15

1 Introduction

The classical Mellin integral representation or Kontorovich–Lebedev transform, given by

$$\Gamma(z + is)\Gamma(z - is) = 4 \int_0^\infty \left(\frac{x}{2}\right)^{2z} K_{2is}(x) \frac{dx}{x}, \quad (1.1)$$

holds for $s \in \mathbb{R}$ and $z \in \mathbb{C}$ such that $\Re(z) > 0$, see, e.g., [6, p. 331, relation (26)]. Here

$$K_w(y) = \int_0^\infty e^{-y \cosh x} \cosh(wx) dx, \quad y > 0, w \in \mathbb{C}, \quad (1.2)$$

is the modified Bessel function of the second kind, or Macdonald function.

In this paper we extend the integral representation (1.1) to all $z \in \mathbb{C}$ such that $\Re(z)$ takes non-integer negative values, and we deduce related extensions for the Fourier transform of $s \mapsto \Gamma(a + is)$. In particular, when $\Re(z) \in (-1, 0)$ and $s \in \mathbb{R}$, we show that

$$\Gamma(z + is)\Gamma(z - is) = 4 \int_0^\infty \left(\frac{x}{2}\right)^{2z} K_{2is}(x) \left(1 + \left(\frac{2}{x}\right)^{2z} \frac{4z}{\Gamma(1 - 2z)} K_{2z}(x)\right) \frac{dx}{x}. \quad (1.3)$$

More generally, when $z \in \mathbb{C}$, with $\Re(z) \in (-n - 1, -n)$ and $s \in \mathbb{R}$, we have

$$\Gamma(z + is)\Gamma(z - is) = 4 \int_0^\infty \left(\frac{x}{2}\right)^{2z} K_{2is}(x) \left(1 + \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{4(k+z)}{k!\Gamma(1-k-2z)!} K_{2k+2z}(x)\right) \frac{dx}{x} \quad (1.4)$$

for any $n \in \mathbb{N} = \{0, 1, 2, \dots\}$, see Proposition 3.2 below. In particular, the introduction of additional terms in (1.3)–(1.4) ensures the integrability lacking in (1.1) when $\Re(z) < 0$.

*Corresponding author: Nicolas Privault, Division of Mathematical Sciences, School of Physical and Mathematical Sciences, Nanyang Technological University, SPMS-MAS-05-43, 21 Nanyang, 637371 Singapore, Singapore, e-mail: nprivault@ntu.edu.sg

On the other hand, Ramanujan showed in [9] that for $a > 0$, the Fourier transform of $s \mapsto |\Gamma(a + is)|^2$ satisfies the relation

$$\int_{-\infty}^{\infty} e^{-i\xi s} |\Gamma(a + is)|^2 ds = \sqrt{\pi} \Gamma(a) \Gamma(a + 1/2) (\cosh(\xi/2))^{-2a}. \quad (1.5)$$

This relation has been extended for all $a \in (-1, 0)$ as an integral expression in [3, Theorem 1.2].

As an application of (1.3)–(1.4), we deduce other integral expressions for (1.5) for all non-integer negative values of a . Namely, when $a \in (-1, 0)$, we derive the integral representation

$$\int_{-\infty}^{\infty} e^{-i\xi s} |\Gamma(a + is)|^2 ds = \frac{2\pi}{2^{2a}} \int_0^{\infty} x^{2a-1} \left(1 + \left(\frac{2}{x} \right)^{2a} \frac{4a}{\Gamma(1-2a)} K_{2a}(x) \right) e^{-x \cosh(\xi/2)} dx, \quad \xi \in \mathbb{R}, \quad (1.6)$$

for the Fourier transform of $s \mapsto |\Gamma(a + is)|^2$. More generally, when $a \in (-n-1, -n)$, we have

$$\int_{-\infty}^{\infty} e^{-i\xi s} |\Gamma(a + is)|^2 ds = \frac{2\pi}{2^{2a}} \int_0^{\infty} x^{2a-1} \left(1 + \left(\frac{2}{x} \right)^{2a} \sum_{k=0}^n \frac{4(a+k)}{k! \Gamma(1-k-2a)} K_{2a+2k}(x) \right) e^{-x \cosh(\xi/2)} dx \quad (1.7)$$

for any $\xi \in \mathbb{R}$ and $n \in \mathbb{N} = \{0, 1, 2, \dots\}$, see Proposition 4.1 below.

The integral representations (1.3)–(1.4) have applications to the solution of the Fokker–Planck type equation

$$\begin{cases} \frac{\partial U_p}{\partial t}(t, y) = y^2 \frac{\partial^2 U_p}{\partial y^2}(t, y) + y \frac{\partial U_p}{\partial y}(t, y) - (p^2 + y^2) U_p(t, y), & y, t > 0, \\ U_p(0, y) = y^p, \end{cases} \quad (1.8)$$

which originates from statistical physics, see [10]. The solution of (1.8) admits the integral expression

$$U_p(t, y) = \frac{2}{\pi^2} \int_0^{\infty} u \sinh(\pi u) K_{iu}(y) e^{-(p^2+u^2)t} \int_0^{\infty} x^{p-1} K_{iu}(x) dx du, \quad (1.9)$$

using the heat kernel of the operator $y^2 - y^2 \frac{\partial^2}{\partial y^2} - y \frac{\partial}{\partial y}$, see, e.g., [12]. From the representations (1.1), (1.4) and the Fubini theorem, (1.9) can be rewritten as the single integral representation

$$\begin{aligned} U_p(s, y) &= \frac{2}{\pi^2} \int_0^{\infty} u \sinh(\pi u) K_{iu}(y) e^{-(p^2+u^2)s} \int_0^{\infty} x^{p-1} K_{iu}(x) \left(1 + \sum_{k=0}^n \frac{2^{p+1}(p+2k)}{k! \Gamma(1-p-k)} K_{-p-2k}(x) \right) dx du \\ &\quad - e^{-p^2 s} \sum_{k=0}^n \frac{2^{p+1}(p+2k)}{k! \Gamma(1-p-k)!} K_{-p-2k}(y) \\ &= \frac{2^p}{2\pi^2} \int_0^{\infty} u \sinh(\pi u) e^{-(p^2+u^2)s} \left| \Gamma\left(\frac{p}{2} + i\frac{u}{2}\right) \right|^2 K_{iu}(y) du - \sum_{k=0}^n e^{4((p+2k)^2-p^2)s} \frac{2^{p+1}(p+2k)}{k! \Gamma(1-p-k)!} K_{-p-2k}(y), \end{aligned}$$

where $p \in (-2n-2, 0]$, $n \in \mathbb{Z}$. The function $U_p(s, y)$ is the Laplace transform of the Harman–Watson distribution, which appears in the analysis of exponential functionals of Brownian motion [4] and in option pricing, cf., e.g., [5, 8].

The above expression of $U_p(s, y)$ has been originally obtained in [10] using spectral expansions, however the derivation presented here is much simpler, since the argument used in [10] involves severe analytical difficulties in the computation of normalization constants via the use of Meijer functions, cf. [10, p. 1641].

This paper is organized as follows. In Section 2 we start by proving some asymptotic expansion and integrability results that are needed for the proof of both (1.3) and (1.6). The proof of the integral representation (1.4) and its extension to $\Re(z) \in (-n-1, -n)$ for all $n \in \mathbb{N}$ are given in Section 3. Finally, in Section 4 we derive the extension of the Fourier transform identity (1.6) to $\Re(z) \in (-n-1, -n)$ for all $n \in \mathbb{N}$, as a consequence of Proposition 3.2.

2 Asymptotic expansion and integrability

In this section we derive the asymptotic results needed for the proofs of (1.3)–(1.7) in Propositions 3.2 and 4.1 below. We will use the modified Bessel function of the first kind

$$I_z(x) = \sum_{k=0}^{\infty} \frac{1}{\Gamma(k+1)\Gamma(k+z+1)} \left(\frac{x}{2}\right)^{z+2k}, \quad x \in \mathbb{R}, z \in \mathbb{C}. \quad (2.1)$$

Lemma 2.1. For all $n \in \mathbb{N}$, $x \in \mathbb{R}$ and $z \in \mathbb{C}$, we have

$$\left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{k+z}{k!\Gamma(1-k-2z)} I_{2k+2z}(x) = \frac{\sin(2\pi z)}{2\pi} + \sum_{l=n+1}^{\infty} \frac{1}{l!} \left(\frac{x}{2}\right)^{2l} \sum_{k=0}^n \binom{l}{k} \frac{k+z}{\Gamma(1-2z-k)!\Gamma(k+l+2z+1)}. \quad (2.2)$$

Proof. From (2.1), we have

$$\begin{aligned} \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{k+z}{k!\Gamma(1-k-2z)} I_{2k+2z}(x) &= \sum_{k=0}^n \frac{k+z}{k!\Gamma(1-k-2z)} \sum_{l=0}^{\infty} \frac{1}{l!\Gamma(2k+l+2z+1)} \left(\frac{x}{2}\right)^{2k+2l} \\ &= \sum_{k=0}^n \frac{k+z}{k!\Gamma(1-k-2z)} \sum_{l=k}^{\infty} \frac{1}{(l-k)!\Gamma(k+l+2z+1)} \left(\frac{x}{2}\right)^{2l} \\ &= \sum_{l=0}^{\infty} \left(\frac{x}{2}\right)^{2l} \sum_{k=0}^{\min(n,l)} \frac{k+z}{k!\Gamma(1-k-2z)(l-k)!\Gamma(k+l+2z+1)}. \end{aligned} \quad (2.3)$$

Next, using Euler's reflection formula

$$\frac{1}{\Gamma(k+l+2z+1)} = -(-1)^{k+l} \frac{\sin(2\pi z)}{\pi} \Gamma(-2z-k-l), \quad k, l \in \mathbb{N},$$

cf., e.g., [1, p. 256, relation (6.1.17)], we get

$$\begin{aligned} &\sum_{l=0}^n \left(\frac{x}{2}\right)^{2l} \sum_{k=0}^l \frac{k+z}{k!(l-k)!\Gamma(1-k-2z)\Gamma(k+l+2z+1)} \\ &= \frac{\sin(2\pi z)}{\pi} \sum_{l=0}^n \left(\frac{x}{2}\right)^{2l} \frac{(-1)^l}{l!} \sum_{k=0}^l (-1)^k \binom{l}{k} (k+z) \frac{\Gamma(-2z-k-l)}{\Gamma(1-2z-k)} \\ &= \frac{\sin(2\pi z)}{\pi} \sum_{l=0}^n \left(\frac{x}{2}\right)^{2l} \frac{(-1)^l}{l!} \frac{\Gamma(-2z-2l)}{\Gamma(1-2z-k)} \sum_{k=0}^l \binom{l}{k} (k+z)(-2z-2l)_{l-k} (2z)_k, \end{aligned} \quad (2.4)$$

where $(p)_k = p(p+1)\cdots(p+k-1)$ is the shifted factorial, $p \in \mathbb{C}$ and $k \geq 1$, with $(p)_0 = 1$. Next, for all $l \geq 1$ and $z \in \mathbb{C}$, we check that

$$\begin{aligned} \sum_{k=0}^l \binom{l}{k} (k+z)(-2z-2l)_{l-k} (2z)_k &= z \sum_{k=0}^l \binom{l}{k} (-2z-2l)_{l-k} (2z)_k - k \sum_{k=0}^l \binom{l}{k} (-2z-2l)_{l-k} (2z)_k \\ &= z \sum_{k=0}^l \binom{l}{k} (-2z-2l)_{l-k} (2z)_k - \sum_{k=1}^l \frac{l!}{(l-k)!(k-1)!} (-2z-2l)_{l-k} (2z)_k \\ &= z \sum_{k=0}^l \binom{l}{k} (-2z-2l)_{l-k} (2z)_k + 2zl \sum_{k=0}^{l-1} \binom{l-1}{k} (-2z-2l)_{l-1-k} (2z+1)_k \\ &= z(-2l)_l + 2zl(-2l+1)_{l-1} \\ &= 0, \end{aligned} \quad (2.5)$$

where we used the Pfaff–Saalschütz binomial identity

$$(p+q)_l = \sum_{k=0}^l \binom{l}{k} (p)_k (q)_{l-k}, \quad p, q \in \mathbb{C}, l \in \mathbb{N},$$

see, e.g., [2, Theorem 2.2.6 and Remark 2.2.1]. As a consequence of (2.4) and (2.5), we get

$$\sum_{l=0}^n \left(\frac{x}{2}\right)^{2l} \sum_{k=0}^l \frac{k+z}{k!(l-k)!\Gamma(1-k-2z)\Gamma(k+l+2z+1)} = \frac{\sin(2\pi z)}{2\pi},$$

which allows us to rewrite (2.3) as

$$\left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{k+z}{k!\Gamma(1-k-2z)} I_{2k+2z}(x) = \frac{\sin(2\pi z)}{\pi} + \sum_{l=n+1}^{\infty} \left(\frac{x}{2}\right)^{2l} \sum_{k=0}^n \frac{k+z}{k!(l-k)!\Gamma(1-k-2z)\Gamma(k+2z+l+1)},$$

which in turn shows (2.2). $\square$

As a consequence of Lemma 2.1, we have the following estimates, which justify the existence of the integrals in (1.4)–(1.7). For example, (2.8) entails the estimate

$$x^{2a} + \frac{2^{2a+2}a}{\Gamma(1-2a)} K_{2a}(x) = o(x^\varepsilon), \quad x \rightarrow 0, \quad (2.6)$$

for any $\varepsilon \in (0, 2+2a) \cap (0, -2a)$, when $a \in (-1, 0)$, which ensures the integrability, as $x \rightarrow 0$, in (1.3) and (1.6).

Corollary 2.2. *Let $n \in \mathbb{N}$.*

(i) *For all $z \in \mathbb{C}$ such that $\Re(z) > -n-1$, we have*

$$\sum_{k=0}^n \frac{k+z}{k!\Gamma(1-k-2z)} I_{2k+2z}(x) = \frac{\sin(2\pi z)}{2\pi} \left(\frac{x}{2}\right)^{2z} + o(x^\varepsilon), \quad x \rightarrow 0, \quad (2.7)$$

for all $\varepsilon \in (0, 2n+2+2\Re(z))$.

(ii) *For all $z \in \mathbb{C}$ such that $-n-1 < \Re(z) < -n$, we have*

$$1 + \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{4(z+k)}{k!\Gamma(1-k-2z)} K_{2k+2z}(x) = o(x^{\varepsilon-2\Re(z)}), \quad x \rightarrow 0, \quad (2.8)$$

for all $\varepsilon \in (0, 2n+2+2\Re(z)) \cap (0, -2n-2\Re(z))$.

Proof. (i) Relation (2.7) follows from Lemma 2.1.

(ii) On the other hand, relation (2.1), with $-2\Re(z) - 2k \geq -2\Re(z) - 2n > \varepsilon > 0$, shows that

$$I_{-2z-2k}(x) = \sum_{l=0}^{\infty} \frac{1}{l!\Gamma(l-2z-2k+1)} \left(\frac{x}{2}\right)^{-2z-2k+2l} = o(x^\varepsilon), \quad x \rightarrow 0.$$

Hence, (2.7) and the identity

$$K_{2k+2z}(x) = \frac{\pi}{2\sin(2\pi z)} (I_{-2z-2k}(x) - I_{2k+2z}(x)), \quad x \in \mathbb{R}, \quad (2.9)$$

allow us to conclude (2.8). $\square$

The next integrability result is a consequence of Lemma 2.1, and it will be useful for the proofs of Propositions 3.1, 3.2 and 4.1 below.

Lemma 2.3. *Let $n \in \mathbb{N}$.*

(i) *For all $z \in \mathbb{C}$ such that $-n-1 < \Re(z)$, we have*

$$\sup_{s \in \mathbb{R}} \int_0^\infty |K_{is}(x)| \left| \frac{\sin(2\pi z)}{\pi} - 2\left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{k+z}{k!\Gamma(1-2z-k)} I_{2k+2z}(x) \right| \frac{dx}{x^{1-2z}} < \infty. \quad (2.10)$$

(ii) *For all $z \in \mathbb{C}$ such that $-n-1 < \Re(z) < -n$, we have*

$$\sup_{s \in \mathbb{R}} \int_0^\infty x^{2z-1} |K_{is}(x)| \left| 1 + \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{4(k+z)}{k!\Gamma(1-k-2z)} K_{2k+2z}(x) \right| dx < \infty. \quad (2.11)$$

Proof. (i) By relation (1.2), for all $\alpha > 0$, there exists a constant $c_\alpha > 0$ such that $\cosh x > c_\alpha x^\alpha$ for all $x > 0$, which shows that

$$|K_{is}(y)| \leq \int_0^\infty e^{-y \cosh x} dx \leq \int_0^\infty e^{-y c_\alpha x^\alpha} dx = \frac{\Gamma(1/\alpha)}{\alpha(c_\alpha)^{1/\alpha}} y^{-1/\alpha}, \quad y > 0, s \in \mathbb{R}.$$

Hence, using (2.7), for all $s \in \mathbb{R}$ and $\alpha > 1/\varepsilon$, we have

$$\int_0^1 |K_{is}(x)| \left| \frac{\sin(2\pi z)}{\pi} - \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{2z+2k}{k! \Gamma(1-k-2z)} I_{2k+2z}(x) \right| \frac{dx}{x^{1-2z}} \leq c \frac{\Gamma(1/\alpha)}{\alpha(c_\alpha)^{1/\alpha}} \int_0^1 \frac{dx}{x^{1-\varepsilon+1/\alpha}} < \infty, \quad s \in \mathbb{R},$$

for some constant $c > 0$. Next, the bound

$$|K_{is}(x)| \leq K_0(x), \quad x > 0, s \in \mathbb{R}, \quad (2.12)$$

that follows from relation (3.5) below and the equivalences

$$K_{is}(x) \approx e^{-x} \sqrt{\frac{\pi}{2x}}, \quad x \rightarrow \infty, s \in \mathbb{R}, \quad (2.13)$$

and

$$I_p(x) \approx \frac{e^x}{\sqrt{2\pi x}}, \quad x \rightarrow \infty, p \in \mathbb{C}, \quad (2.14)$$

show that

$$\sup_{s \in \mathbb{R}} \int_1^\infty |K_{is}(x)| \left| \frac{\sin(2\pi z)}{\pi} - 2\left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{k+z}{k! \Gamma(1-2z-k)} I_{2k+2z}(x) \right| \frac{dx}{x^{1-2z}} < \infty,$$

which yields (2.10) for all $z \in \mathbb{C}$ such that $\Re(z) > -n-1$.

(ii) Due to the equivalences (2.9) and (2.14), there exists also $c > 0$ such that

$$\left| 1 + \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{4(k+z)}{k! \Gamma(1-k-2z)} K_{2k+2z}(x) \right| \leq c x^{-2z-1/2} e^x \quad (2.15)$$

for sufficiently large $x > 0$ and all $z \in \mathbb{C}$, and this yields (2.11) by replacing the use of (2.7) with that of (2.8) in the proof of part (i) above. $\square$

3 Analytic continuation and integral representation

In the next proposition, using analytic continuation, we prove an integral representation formula that will be applied to the proof of (3.6) in Proposition 3.2 below.

Proposition 3.1. For all $n \in \mathbb{N}$, $s \in \mathbb{R}$ and $z \in \mathbb{C}$, with $\Re(z) > -n-1$, $\Re(z) \notin -\mathbb{N}$, we have

$$\begin{aligned} \Gamma(z+is)(z-is) &= \frac{2\pi}{\sin(2\pi z)} \sum_{k=0}^n \frac{k+z}{k!((z+k)^2+s^2)\Gamma(1-k-2z)} \\ &\quad + 2^{2-2z} \int_0^\infty K_{2is}(x) \left(1 - \frac{\pi}{\sin(2\pi z)} \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{2z+2k}{k! \Gamma(1-k-2z)} I_{2k+2z}(x) \right) \frac{dx}{x^{1-2z}}. \end{aligned} \quad (3.1)$$

Proof. Let $s \in \mathbb{R} \setminus \{0\}$. We will prove the equality

$$\sin(2\pi z) \Gamma(z+is) \overline{\Gamma(\bar{z}+is)} = \sin(2\pi z) \Gamma(z+is)(z-is) \quad (3.2)$$

$$\begin{aligned} &= \pi \sum_{k=0}^n \frac{2z+2k}{k!((z+k)^2+s^2)\Gamma(1-k-2z)} \\ &\quad + \frac{4}{2^{2z}} \int_0^\infty K_{2is}(x) \left(\sin(2\pi z) - \pi \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{2z+2k}{k! \Gamma(1-k-2z)} I_{2k+2z}(x) \right) \frac{dx}{x^{1-2z}}, \end{aligned} \quad (3.3)$$

for all $z = a+ib \in \mathbb{C}$ with $a > -n-1$, in the following three steps.

Step 1: Analyticity. In (3.2), the function

$$\sin(2\pi z)\Gamma(z + is)(z - is)$$

is analytic in $\{z : z + is \notin (-\mathbb{N}), z - is \notin (-\mathbb{N})\}$, and for each $k = 0, 1, \dots, n$, the function $((z + k)^2 + s^2)^{-1}$ is analytic in $z \in \mathbb{C} \setminus (-\mathbb{N})$. On the other hand, by Lemma 2.1, we can write the integrand in (3.2) as

$$\begin{aligned} x &\mapsto \frac{K_{is}(x)}{x^{1-2z}} \left(\frac{\sin(2\pi z)}{\pi} - \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{2z+2k}{k!\Gamma(1-k-2z)} I_{2k+2z}(x) \right) \\ &= -2K_{is}(x) \sum_{k=0}^n \frac{x^{2z-1}}{k!\Gamma(1-2z-k)} \sum_{l=n+1}^{\infty} \left(\frac{x}{2}\right)^{2l} \frac{k+z}{(l-k)!\Gamma(k+2z+l+1)} \\ &= -2K_{is}(x) \sum_{k=0}^n \frac{x^{2z-1}}{k!\Gamma(1-2z-k)\Gamma(k+2z)} \sum_{l=n+1}^{\infty} \left(\frac{x}{2}\right)^{2l} \frac{k+z}{(l-k)!(k+2z+l)\cdots(k+2z)}, \end{aligned}$$

where for each $k = 0, 1, \dots, n$ the partial derivatives of

$$z = a + ib \mapsto \sum_{l=n+1}^{\infty} \left(\frac{x}{2}\right)^{2l} \frac{k+z}{(l-k)!(k+l+2z)\cdots(k+2z)},$$

with respect to a and b , are locally uniformly bounded by integrable functions of $x \in \mathbb{R}_+$ from the bounds (2.12) and (2.13), by the same arguments used in the proof of Lemma 2.3.

Hence, we can exchange partial differentiation with respect to a and b with the integration in (3.3), showing that the Cauchy–Riemann conditions are satisfied by the integral (3.3), since all functions in the integrand are analytic in $z \in \mathbb{C}$. Consequently, all terms in (3.2) are analytic in

$$\{z \in \mathbb{C} : \Re(z) > -n-1, z + is \notin (-\mathbb{N}), z - is \notin (-\mathbb{N})\}.$$

Step 2: Equality (3.2) holds for all $s \in \mathbb{R} \setminus \{0\}$ and $z = a + ib \in (0, \infty) + i\mathbb{R}$. This follows from the integral representation (1.1), which reads

$$\Gamma(z + is)(z - is) = 4 \int_0^{\infty} \left(\frac{x}{2}\right)^{2z} K_{2is}(x) \frac{dx}{x},$$

provided that $a > 0$, and from the Mellin transform

$$4 \int_0^{\infty} K_{2is}(x) I_{2k+2z}(x) \frac{dx}{x} = \frac{1}{(z+k)^2 + s^2}, \quad (3.4)$$

which is valid whenever $a + k > 0$, cf., e.g., [6, p. 334, relation (44)].

Step 3. By analytic continuation, the relation (3.2) extends to

$$\{z \in \mathbb{C} : \Re(z) > -n-1, z + is \notin (-\mathbb{N}), z - is \notin (-\mathbb{N})\},$$

and we conclude by dividing (3.3) by $\sin(2\pi z)$ when $z \in \mathbb{C}$, with $\Re(z) > -n-1$ and $\Re(z) \notin -\mathbb{N}$. $\square$

Note that in the above proof we could also have used the unique continuation principle for real analytic functions of a , see, e.g., [7, Corollary 1.2.3]. However, real analyticity requires to check the growth rate of partial derivatives, which would have been more delicate.

Relation (1.1) can be recovered from the integral representation

$$K_{is}(y) = \frac{1}{2} \left(\frac{y}{2}\right)^{is} \int_0^{\infty} x^{-is-1} e^{-x-y^2/(4x)} dx, \quad s, y \in \mathbb{R}, \quad (3.5)$$

see [11, p. 183], using the Fubini theorem, as follows:

$$\begin{aligned}\Gamma(z+is)(z-is) &= \int_0^\infty x^{-2is-1} e^{-x} x^{z+is} \int_0^\infty y^{-1+z+2is} e^{-y} dy dx \\ &= \int_0^\infty \left(\frac{t}{2}\right)^{2z-1+2is} \int_0^\infty x^{-2is-1} e^{-x-t^2/(4x)} dx dt \\ &= 4 \int_0^\infty \left(\frac{t}{2}\right)^{2z} K_{2is}(t) \frac{dt}{t}, \quad s \in \mathbb{R},\end{aligned}$$

where we applied the change of variable $y = t^2/4x$. However, this argument is valid only for $\Re(z) > 0$ due to integrability restrictions in the exchange of integrals.

We are now able to extend the above argument to all $\Re(z) \in (-n-1, -n)$, $n \in \mathbb{N}$, in order to prove the integral representation (3.6), which also implies (1.3).

Proposition 3.2. *For all $n \in \mathbb{N}$, $s \in \mathbb{R}$ and $z \in \mathbb{C}$ such that $-n-1 < \Re(z) < -n$, we have*

$$\Gamma(z+is)(z-is) = 2 \int_0^\infty K_{2is}(x) \left(1 + \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{4(k+z)}{k!\Gamma(1-k-2z)!} K_{2k+2z}(x)\right) \left(\frac{2}{x}\right)^{1-2z} dx. \quad (3.6)$$

Proof. First we note that the integrability in (3.6) follows from the bound (2.10). Next, from relations (2.9), (3.1) and (3.4), with $-2\Re(z) - 2k \geq -2\Re(z) - 2n > 0$, we have

$$\begin{aligned}2^{2z-2}\Gamma(z+is)(z-is) &= \frac{\pi}{\sin(2\pi z)} \int_0^\infty K_{2is}(x) \left(\frac{\sin(2\pi z)}{\pi} - \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{2z+2k}{k!\Gamma(1-k-2z)} I_{2k+2z}(x)\right) \frac{dx}{x^{1-2z}} \\ &\quad + \frac{2^{2z}\pi}{2\sin(2\pi z)} \sum_{k=0}^n \frac{k+z}{k!(k+z)^2+s^2}\Gamma(1-k-2z) \\ &= \frac{\pi}{\sin(2\pi z)} \int_0^\infty K_{2is}(x) \left(\frac{\sin(2\pi z)}{\pi} - \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{2z+2k}{k!\Gamma(1-k-2z)} I_{2k+2z}(x)\right) \frac{dx}{x^{1-2z}} \\ &\quad + \frac{2^z\pi}{\sin(2\pi z)} \sum_{k=0}^n \frac{2z+2k}{k!\Gamma(1-k-2z)} \int_0^\infty K_{2is}(x) I_{-2z-2k}(x) \frac{dx}{x} \\ &= \int_0^\infty K_{2is}(x) \left(1 + \left(\frac{2}{x}\right)^{2z} \sum_{k=0}^n \frac{4(k+z)}{k!\Gamma(1-k-2z)} K_{2k+2z}(x)\right) \frac{dx}{x^{1-2z}}. \quad \square\end{aligned}$$

4 Fourier transform of $|\Gamma(a+is)|^2$

We begin by proving an integral representation for the Fourier transform of

$$s \mapsto |\Gamma(a+is)|^2, \quad a \in (-n-1, -n), \quad n \in \mathbb{N},$$

as a consequence of the integral representation (1.4) of Proposition 3.2.

Proposition 4.1. *Let $n \in \mathbb{N}$ and $a \in (-n-1, -n)$. For all $\xi \in \mathbb{R}$, we have*

$$\int_{-\infty}^\infty e^{-i\xi s} |\Gamma(a+is)|^2 ds = \frac{2\pi}{2^{2a}} \int_0^\infty x^{2a-1} \left(1 + \left(\frac{2}{x}\right)^{2a} \sum_{k=0}^n \frac{4(a+k)}{k!\Gamma(1-k-2a)} K_{2a+2k}(x)\right) e^{-x \cosh(\xi/2)} dx. \quad (4.1)$$

Proof. This result can be informally deduced from (3.6) in Proposition 3.2 and the Fourier–Gelfand formula

$$\int_{-\infty}^\infty \cos(2sy) e^{-i\xi s} ds = \pi(\delta(\xi/2-y) + \delta(\xi/2+y))$$

in distribution theory, where δ is the Dirac distribution at 0. However, with a view towards completeness, we provide a proof by approximation, following the approach used in the proof of [3, Theorem 1.1]. With the abbreviation

$$\Psi_n(x) := 1 + \left(\frac{2}{x}\right)^{2a} \sum_{k=0}^n \frac{4(a+k)}{k!\Gamma(1-k-2a)} K_{2a+2k}(x), \quad x \in \mathbb{R}, \quad (4.2)$$

we rewrite (3.6) as

$$|\Gamma(a+is)|^2 = \frac{4}{2^{2a}} \int_0^\infty x^{2a-1} K_{2is}(x) \Psi_n(x) dx, \quad s \in \mathbb{R},$$

for $a \in (-n-1, -n)$. Then, for any $\varepsilon > 0$, we have

$$\begin{aligned} \int_{-\infty}^\infty e^{-i\xi s} |\Gamma(a+is)|^2 ds &= \int_{-\infty}^\infty \lim_{\varepsilon \rightarrow 0} (e^{-2\varepsilon s^2 - i\xi s} |\Gamma(a+is)|^2) ds \\ &= \lim_{\varepsilon \rightarrow 0} \int_{-\infty}^\infty e^{-2\varepsilon s^2 - i\xi s} |\Gamma(a+is)|^2 ds \\ &= 2^{2-2a} \lim_{\varepsilon \rightarrow 0} \int_{-\infty}^\infty e^{-2\varepsilon s^2 - i\xi s} \int_0^\infty x^{2a-1} K_{2is}(x) \Psi_n(x) dx ds \\ &= 2^{2-2a} \lim_{\varepsilon \rightarrow 0} \int_0^\infty x^{2a-1} \Psi_n(x) \int_{-\infty}^\infty e^{-2\varepsilon s^2 - i\xi s} K_{2is}(x) ds dx, \end{aligned} \quad (4.3)$$

where the exchange of limit follows from the fact that $s \mapsto |\Gamma(a+is)|^2$ is a rapidly decreasing function in the Schwartz class, and the last equality comes from (2.11), which ensures the integrability required for the exchange of integrals. Next, from relation (1.2), written as

$$K_{2is}(x) = \int_0^\infty e^{-x \cosh y} \cos(2sy) dy, \quad x > 0, s \in \mathbb{R},$$

we find

$$\begin{aligned} \int_{-\infty}^\infty e^{-2\varepsilon s^2 - i\xi s} K_{2is}(x) ds &= \int_{-\infty}^\infty e^{-2\varepsilon s^2 - i\xi s} \int_0^\infty e^{-x \cosh y} \cos(2sy) dy ds \\ &= \int_{-\infty}^\infty e^{-x \cosh y} \int_0^\infty \cos(2sy) e^{-2\varepsilon s^2 - i\xi s} ds dy \\ &= \frac{1}{4} \sqrt{\frac{\pi}{2\varepsilon}} \int_{-\infty}^\infty e^{-x \cosh y} \left(e^{-\frac{1}{2\varepsilon}(y-\xi/2)^2} + e^{-\frac{1}{2\varepsilon}(y+\xi/2)^2} \right) dy, \quad x > 0. \end{aligned}$$

Hence, by (4.3), we obtain

$$\begin{aligned} \int_{-\infty}^\infty e^{-i\xi s} |\Gamma(a+is)|^2 ds &= 2^{-2a} \lim_{\varepsilon \rightarrow 0} \sqrt{\frac{\pi}{2\varepsilon}} \int_0^\infty x^{2a-1} \Psi_n(x) \int_{-\infty}^\infty e^{-x \cosh y} \left(e^{-\frac{1}{2\varepsilon}(y-\xi/2)^2} + e^{-\frac{1}{2\varepsilon}(y+\xi/2)^2} \right) dy dx \\ &= 2^{-2a} \lim_{\varepsilon \rightarrow 0} \sqrt{\frac{\pi}{2\varepsilon}} \int_0^\infty x^{2a-1} \Psi_n(x) \int_{-\infty}^\infty e^{-x \cosh(y+\xi/2) - \frac{1}{2\varepsilon} y^2} dy dx \\ &\quad + 2^{-2a} \lim_{\varepsilon \rightarrow 0} \sqrt{\frac{\pi}{2\varepsilon}} \int_0^\infty x^{2a-1} \Psi_n(x) \int_{-\infty}^\infty e^{-x \cosh(y-\xi/2) - \frac{1}{2\varepsilon} y^2} dy dx \end{aligned}$$

$$\begin{aligned}
&= 2^{-2a} \sqrt{\frac{\pi}{2}} \lim_{\varepsilon \rightarrow 0} \int_0^\infty x^{2a-1} \Psi_n(x) \int_{-\infty}^\infty e^{-x \cosh(z\sqrt{\varepsilon} + \xi/2) - z^2/2} dz dx \\
&\quad + 2^{-2a} \sqrt{\frac{\pi}{2}} \lim_{\varepsilon \rightarrow 0} \int_0^\infty x^{2a-1} \Psi_n(x) \int_{-\infty}^\infty e^{-x \cosh(z\sqrt{\varepsilon} - \xi/2) - z^2/2} dz dx \\
&= 2^{1-2a} \pi \int_0^\infty x^{2a-1} \Psi_n(x) e^{-x \cosh(\xi/2)} dx,
\end{aligned}$$

where the required integrability follows from the bounds (2.8) and (2.15). $\square$

In case $a > 0$, $\Psi_{-1}(x)$ in (4.2) is identically equal to 1 and the proof of Proposition 4.1 also yields the Mellin transform

$$\int_{-\infty}^\infty e^{-i\xi s} |\Gamma(a + is)|^2 ds = \frac{2\pi}{2^{2a}} \int_0^\infty x^{2a-1} e^{-x \cosh(\xi/2)} dx = \frac{2\pi}{2^{2a}} (\cosh(\xi/2))^{-2a} \Gamma(2a),$$

which recovers (1.5), cf. also [3, Theorem 1.1].

On the other hand, when $a = -1/2$, the Fourier transform of $s \mapsto |\Gamma(-1/2 + is)|^2$ can be explicitly computed as

$$\begin{aligned}
\int_{-\infty}^\infty e^{-i\xi s} |\Gamma(-1/2 + is)|^2 ds &= 4\pi \int_{-\infty}^\infty \frac{e^{-i\xi s}}{(1 + 4s^2) \cosh(\pi s)} ds \\
&= 4\pi \log(1 + e^{-\xi}) \cosh(\xi/2) + 2\pi \xi e^{-\xi/2},
\end{aligned}$$

see, e.g., [6, p. 32, relation (22)], whereas (4.1) yields

$$\int_{-\infty}^\infty e^{-i\xi s} |\Gamma(-1/2 + is)|^2 ds = 4\pi \int_0^\infty \left(\frac{1}{x} - K_{-1}(x) \right) e^{-x \cosh(\xi/2)} \frac{dx}{x},$$

where the integrability in 0 in the above integral follows from

$$\frac{1}{x} - K_{-1}(x) = o(x^\varepsilon), \quad x \rightarrow 0,$$

for any $\varepsilon \in (0, 1)$, cf. (2.6).

Funding: This research was supported by the Singapore MOE Tier 2 Grant MOE2016-T2-1-036

References

- [1] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Appl. Math. Ser. 55, U. S. Department of Commerce, Washington, 1964.
- [2] G. E. Andrews, R. Askey and R. Roy, *Special Functions*, Encyclopedia Math. Appl. 71, Cambridge University Press, Cambridge, 1999.
- [3] D. Chakrabarti and G. K. Srinivasan, On a remarkable formula of Ramanujan, *Arch. Math. (Basel)* **99** (2012), no. 2, 125–135.
- [4] A. Comtet, C. Monthus and M. Yor, Exponential functionals of Brownian motion and disordered systems, *J. Appl. Probab.* **35** (1998), no. 2, 255–271.
- [5] M. Craddock, On an integral arising in mathematical finance, in: *Nonlinear Economic Dynamics and Financial Modelling*, Springer, Cham (2014), 355–370.
- [6] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Tables of Integral Transforms. Vol. I*, McGraw-Hill, New York, 1954.
- [7] S. G. Krantz and H. R. Parks, *A Primer of Real Analytic Functions*, 2nd ed., Birkhäuser Adv. Texts Basler Lehrbücher, Birkhäuser, Boston, 2002.

- [8] C. Pintoux and N. Privault, The Dothan pricing model revisited, *Math. Finance* **21** (2011), no. 2, 355–363.
- [9] S. Ramanujan, Some definite integrals, *Messenger* **44** (1915), 10–18.
- [10] A. Schenzle and H. Brand, Multiplicative stochastic processes in statistical physics, *Phys. Rev. A* **20** (1979), no. 4, 1628–1647.
- [11] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Cambridge Math. Lib., Cambridge University Press, Cambridge, 1995.
- [12] S. B. Yakubovich, A class of polynomials and discrete transformations associated with the Kontorovich–Lebedev operators, *Integral Transforms Spec. Funct.* **20** (2009), no. 7–8, 551–567.