Stalagmites from western Thailand : preliminary investigations and challenges for palaeoenvironmental research

Chawchai, Sakonvan; Liu, Guangxin; Bissen, Raphael; Jankham, Kampanart;
Paisonjumlongsri, Warisa; Kanjanapayont, Pitsanupong; Chutakositkanon, Vichai;
Choowong, Montri; Pailoplee, Santi; Wang, Xianfeng

2018

Chawchai, S., Liu, G., Bissen, R., Jankham, K., Paisonjumlongsri, W., Kanjanapayont, P., . . .
Wang, X. (2018). Stalagmites from western Thailand : preliminary investigations and
challenges for palaeoenvironmental research. *Boreas*, 47(1), 367-376.
doi:10.1111/bor.12299

<https://hdl.handle.net/10356/92355>

<https://doi.org/10.1111/bor.12299>

© 2017 Collegium Boreas. All rights reserved. This paper was published by John Wiley &
Sons Ltd in *Boreas* and is made available with permission of Collegium Boreas.

Downloaded on 26 Apr 2023 17:03:17 SGT

Stalagmites from western Thailand: preliminary investigations and challenges for palaeoenvironmental research

1 SAKONVAN CHAWCHAI , GUANGXIN LIU, RAPHAEL BISSEN, KAMPANART JANKHAM, WARISA PAISONJUMLONGSRI, PITSANUPONG KANJANAPAYONT, VICHAI CHUTAKOSITKANON, MONTRI CHOOWONG, **2** SANTI PAILOPLEE AND XIANFENG WANG

BOREAS

Chawchai, S., Liu, G., Bissen, R., Jankham, K., Paisonjumlongsri, W., Kanjanapayont, P., Chutakositkanon, V., Choowong, M., Pailoplee, S. & Wang, X.: Stalagmites from western Thailand: preliminary investigations and challenges for palaeoenvironmental research. *Boreas*. <https://doi.org/10.1111/bor.12299>. ISSN 0300-9483.

Locating suitable caves and stalagmites for palaeoenvironmental and palaeoclimatic studies can be challenging. Isotopic geochemical analyses, albeit commonly performed for palaeoclimatic reconstruction, are also time consuming and costly. Therefore, petrographic and non-destructive morphological studies on speleothems are desirable to facilitate sample selection for further analysis. In this study, 20 caves were surveyed in Ban Rai district, Uthai Thani province in western Thailand. After external physical observations in the field, three stalagmite samples were collected from Tham Nam Cave to test their potential for palaeoclimatic research. Firstly, the stalagmites were scanned by X-ray computed tomography (CT scanning) and subsequently the CT images were compared with petrographic inspections. Columnar fabrics show the highest density, whereas closed and open dendritic fabrics have medium and the lowest densities, respectively. Layers near the top and bottom of the three stalagmites were dated by U-Th mass spectrometric techniques. All three samples were deposited between *c.* 87 and *c.* 105 ka ago; therefore, they are probably the oldest stalagmites that have been reported so far from mainland Southeast Asia. However, their physical features indicate that all the samples have suffered from ~~postdepositional~~ dissolution, and are unlikely to be suitable for palaeoclimatic research. The internal dissolution feature of stalagmites, however, cannot be identified by visual inspection of uncut samples. We hereby argue that CT images are useful to characterize stalagmite petrography, in particular fabric, porosity and density. Such features can be used to select the ideal plane of a stalagmite for sectioning, ~~and then~~ to maximize the chances of robust climatic reconstruction.

Sakonvan Chawchai (sakonvan.c@chula.ac.th), Kampanart Jankham, Warisa Paisonjumlongsri, Pitsanupong Kanjanapayont, Vichai Chutakositkanon, Montri Choowong and Santi Pailoplee, MESA Research Unit, Department of Geology, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand; Guangxin Liu and Xianfeng Wang, Earth Observatory of Singapore and Asian School of the Environment Nanyang Technological University, Singapore 639798, Singapore; Raphael Bissen, Department of Mining and Petroleum Engineering, Faculty of Engineering, Chulalongkorn University, Bangkok 10330, Thailand; received 24th August 2017, accepted 15th November 2017.

Speleothems are secondary carbonate deposits formed inside caves. Their fabrics, stable isotope ratios, trace element compositions and organic chemistry can record events that occur at the cave surface during their growth history (McDermott 2004; Fairchild *et al.* 2006). Several types of speleothems (e.g. flowstones, stalactites and stalagmites) have been used as palaeoenvironmental and palaeoclimatic archives to study the changes in local or regional temperature, vegetation and precipitation, thanks to their continuous or semi-continuous deposition, absolute and precise datability, occurrence over a wide range of latitudes and possible conservation of palaeoclimate signals on millennial, decadal and even annual time scales (e.g. Hendy 1971; ~~Banner *et al.* 2004~~; McDermott 2004; Fairchild *et al.* 2006; Fairchild & Treble 2009; Lachniet 2009; Mickler *et al.* 2004a, b; Wong & Breecker 2015). Stalagmites have been favourably used in past hydroclimate reconstruction because of their predominantly ~~column~~ shapes and relatively simple sequential depositions (e.g. Zhang *et al.* 2008; Cai *et al.* 2010; Chen *et al.* 2015; Sinha *et al.* 2015). However, it remains a challenge to find suitable caves and stalagmite samples for palaeoclimatic research because only pristine stalagmites, deposited under chemical equilibrium conditions, can faithfully record climate signals.

Mainland Southeast Asia (e.g. Thailand, Myanmar, Malaysia peninsula and Laos) is ~~strategically~~ located on the route of moisture transport of the Indian summer monsoon (ISM, Fig. 1A). The agrarian societies dominant in the region heavily rely on the monsoon rainfall. Changes in rainfall intensity and variability therefore have a great impact on people's livelihoods in mainland Southeast Asia (Loo *et al.* 2015). The projection of future hydroclimatic change in the region demands a good understanding of monsoon rainfall history (Cook *et al.* 2010; Raghavan *et al.* 2017; Thirumalai *et al.* 2017). Instrumental records of precipitation and temperature are however short (<50 years) and sparse in mainland Southeast Asia. Hence, palaeoclimate proxy data have to be applied to delineate the natural variability as well as anthropogenic contribution to climate change. Yet, such studies are still scarce in mainland Southeast Asia.

The beauty of the caves and speleothems in mainland Southeast Asia has attracted millions of tourists and visitors from around the world. Stalagmites from this region however have rarely been investigated for palaeoenvironmental research. In Thailand for instance, only two stalagmite records have been reported so far. In these studies, stalagmites from the Namjang Cave, Mae Hong Son province in northwestern Thailand (Fig. 1A), were

Fig. 1. A. Locations of Ban Rai district (red square) and Tham Nam Cave (white star) in Uthai Thani, western Thailand. B. Locations of the 20 surveyed caves in Ban Rai, Uthai Thani province.

used to reconstruct monsoon variability covering the last 1700 and 400 years, respectively (Cai *et al.* 2010; Muangsong *et al.* 2014). Stalagmite growth rates during the last century from Namjang have been correlated with meteorological data, showcasing the potential palaeoclimatic proxy of Thailand monsoon rainfall (Muangsong *et al.* 2014).

In recent years, there have been studies on diagenetic alteration (e.g. dissolution, recrystallization and neomorphism) in speleothems. This process results in the chemical properties of samples being altered, meaning that they no longer represent the original depositional conditions (Perrin *et al.* 2014; Scholz *et al.* 2014; Zhang *et al.* 2014; Bajo *et al.* 2016). The diagenetic alteration of speleothems can be observed through analysing petrography, internal microstratigraphy and trace elements in samples (Frisia *et al.* 2002; Railsback *et al.* 2002). These petrographic and non-destructive morphological studies are important to evaluate speleothems before further chemical analysis.

X-ray computed tomography (CT scanning) is a non-destructive technique for visualizing interior features of opaque solid objects by measuring the intensity of X-rays when they pass through the objects. The principle of computed tomography is to acquire multiple sets of views of an object over a range of angular orientations. These data are used to create two-dimensional images that are called slices (along the scan plane) and then provide

a three-dimensional image of a volume by obtaining a series of contiguous slices (Ketcham & Carlson 2001). CT scanning has previously been used in studying speleothem features, including porosity (Mickler *et al.* 2004a, b), macro-holes and postdepositional off axis holes (Zisu *et al.* 2012), and calcite density (Vanghi *et al.* 2015; Walczak *et al.* 2015), as well as in exploring evidence of diagenesis in speleothems (Bajo *et al.* 2016). It has been proposed that CT images of uncut stalagmites can provide essential information on their petrography to evaluate the potential impact on U-Th ages and isotopic geochemical analyses, which cannot be visually identified instead.

The aims of this study were to find stalagmite samples in Ban Rai district, Uthai Thani province, western Thailand (Fig. 1A, B), based on their external physical properties (e.g. shape, colour and texture), and analyse them with respect to their potential for palaeoclimatic research via CT scanning, petrographic studies and U-Th analysis.

Material and methods

Geological setting

The study areas are located in national parks and national conservation zones of Ban Rai district, Uthai Thani province, western Thailand (Fig. 1A, B). All caves in this area are overlain by host rock of argillaceous

limestone, which is of Ordovician age as indicated by fossil nautilus and crinoids (Department of Mineral Resources (DMR) 2009).

In this study, 20 caves were explored (Fig. 1A, B). Fifteen of them do not contain any stalagmites; three caves contain broken stalagmites that show visible macro-holes and irregular external surfaces; and the other two caves contain both broken and standing stalagmites. Based on their external features (nearly symmetrical, column shapes, no obvious corruptions), three standing stalagmite samples (BR1, BR2 and BR3) were collected from Tham Nam Cave, a 300-m-long cave (latitude 15.05°N, longitude 99.48°E; altitude 200 m a.s.l; Fig. 1A, B). The stalagmites were retrieved from a large chamber ~200 m from the cave entrance.

Climate in western Thailand is mainly influenced by the ISM, which brings in torrential rains from the Bay of Bengal. Additional precipitation can be contributed by tropical cyclones from the South China Sea in the

east during the summer. Mean annual rainfall in Uthai Thani province is about 1770 mm with 105 rain days on average. Mean annual temperature is around 28° C (data from Thai Meteorological Department, AD 1962–2015).

X-ray computed tomography (CT scanning) and petrography

Firstly, the stalagmites BR1, BR2 and BR3 (Fig. 2) were scanned with X-ray computed tomography (CT scanner) at King Chulalongkorn Memorial Hospital, Bangkok, Thailand. The X-ray computed tomography machine (GE Discovery 750 HD) was operated at a tube voltage of 140 kV and a current intensity of 80 mA. The CT scanner was set to a 0.625-mm slice thickness, with a 0.4-mm overlay between slice spacing. Reconstructed images were exported as Digital Imaging and Communications in Medicine (DICOM) files.

Fig. 2. A. Stalagmite samples BR1, BR2 and BR3. B. 2D images of the stalagmites from CT scanning (colour scale: blue colour represents the lowest density; green colour indicates medium density and red/orange colours represent highest density).

When X-rays scan an object, a portion of the radiation can be absorbed or reflected. Such interaction between the X-rays and the object is dependent upon the X-ray energy and the atomic structures of the object. The X-rays that pass through the sample are collected by a detector, which converts them into light radiations. The light radiations are then transformed by a digital camera and later processed by a computer into images (Ketcham & Carlson 2001).

CT numbers in Hounsfield units (HU) define the intensity of the transmitted X-ray beam. The HU is calculated from the relative value of the linear voxel attenuation coefficient (μ_{voxel}) normalized to the reference material water (μ_{water}), which is then multiplied by a magnifying integer constant (Mickler *et al.* 2004a, b): CT numbers in Hounsfield units (HU) = $K [(\mu_{\text{voxel}} - \mu_{\text{water}}) / \mu_{\text{water}}]$.

CT image processing and data analysis were performed using MeVisLab 2.6.2 software (<http://www.mevislab.de/>). CT images are shown both in colour (Fig. 2B), and as greyscale images of axial sections (Fig. 3), which indicate the stalagmite's density (Mickler *et al.* 2004a, b; Vanghi *et al.* 2015; Walczak *et al.* 2015); the denser the scanned stalagmites, the brighter the images in CT slices (Fig. 3).

After CT scanning, the plane along the growth axis with less variability in density was chosen for sectioning, and the samples were cut to characterize and describe the different fabrics of the stalagmites. Based on different CT

numbers (HU), 19 polished thin sections were made (Fig. 3).

U-Th dating

Eight subsamples of BR1, BR2 and BR3 were drilled along the growth layers near their top and bottom for U-Th series dating (Table 1; Fig. 3). Replicate tests were performed on each subsample. U and Th were then separated and purified following the chemical procedure described in Edwards *et al.* (1986). U-Th isotopic measurement was performed on a Thermo Finnigan Neptune Plus multi-collector inductively coupled plasma mass spectrometer (MC-ICP-MS) in the isotope geochemistry laboratory at the Earth Observatory of Singapore, Nanyang Technological University, Singapore. The U and Th isotopes were measured on a secondary electron multiplier using peak jumping mode (Shen *et al.* 2012; Cheng *et al.* 2013).

Results and discussion

Physical properties and CT images

Sample BR1 has a nearly symmetrical candle shape with a length of 31 cm and a diameter between 6 and 8 cm. The colour is yellowish grey. The rough outer surface shows some visible macro-holes (Fig. 2A). Along the

Fig. 3. Halved sections of stalagmites BR1, BR2 and BR3 compared to 2D images of uncut samples from CT scanning in greyscale. Higher densities are indicated by lighter greyscale colour. Red circles show subsample locations for U-Th dating and black rectangles show areas for thin sections.

Table 1. U and Th isotopic compositions (^{238}U , ^{234}U , ^{232}Th and ^{230}Th) and U-Th ages for subsamples of stalagmites BR1, BR2 and BR3. U decay constants: $\lambda_{238} = 1.55125 \times 10^{-10}$ (Jaffey *et al.* 1971) and $\lambda_{234} = 2.82206 \times 10^{-6}$ (Cheng *et al.* 2013). Th decay constant: $\lambda_{230} = 9.1705 \times 10^{-6}$ (Cheng *et al.* 2013). $\delta^{234}\text{U} = ((^{234}\text{U}/^{238}\text{U})_{\text{sample}} / (^{234}\text{U}/^{238}\text{U})_{\text{standard}} - 1) \times 1000$. $\delta^{234}\text{U}_{\text{initial}}$ was calculated based on ^{230}Th age (T), i.e. $\delta^{234}\text{U}_{\text{initial}} = \delta^{234}\text{U}_{\text{measured}} \times e^{\lambda_{234} \times T}$. Corrected ^{230}Th ages assume the initial $^{230}\text{Th}/^{232}\text{Th}$ atomic ratio of $4.4 \pm 2.2 \times 10^{-6}$. Those are the values for a material at secular equilibrium with the bulk earth $^{230}\text{Th}/^{232}\text{Th}$ value of 3.8. The errors are arbitrarily assumed to be 50%.

Sample	Number	^{238}U (ppb)	^{232}Th (ppt)	$^{230}\text{Th}/^{232}\text{Th}$ (atomic $\times 10^{-6}$)	$\delta^{234}\text{U}^*$ (measured)	$^{230}\text{Th}/^{238}\text{U}$ (activity)	^{230}Th age (years, uncorrected)	^{230}Th age (years, corrected)	$\delta^{234}\text{U}_{\text{initial}}^{**}$ (corrected)	^{230}Th age (years, corrected)
BR1 top	7A	78	30 562 $\pm$ 615	30 $\pm$ 1	132.7 $\pm$ 2.1	0.7102 $\pm$ 0.0023	104 684 $\pm$ 653	94 528 $\pm$ 7248	173 $\pm$ 4.4	94 464 $\pm$ 7248
BR1 top	7B	75	31 410 $\pm$ 631	27 $\pm$ 1	138.5 $\pm$ 2.1	0.6862 $\pm$ 0.0032	98 152 $\pm$ 798	87 343 $\pm$ 7728	177 $\pm$ 4.7	87 278 $\pm$ 7728
BR1 bottom	8A	44	2806 $\pm$ 57	172 $\pm$ 4	143.7 $\pm$ 2.0	0.6575 $\pm$ 0.0032	91 115 $\pm$ 729	89 553 $\pm$ 1317	185 $\pm$ 2.7	89 488 $\pm$ 1317
BR1 bottom	8B	40	1575 $\pm$ 32	273 $\pm$ 6	146.7 $\pm$ 1.9	0.6566 $\pm$ 0.0030	90 511 $\pm$ 690	89 534 $\pm$ 971	189 $\pm$ 2.6	89 470 $\pm$ 971
BR2 top	5A	84	26 950 $\pm$ 541	35 $\pm$ 1	144.4 $\pm$ 2.1	0.6777 $\pm$ 0.0029	95 378 $\pm$ 711	87 227 $\pm$ 5826	185 $\pm$ 4.0	84 162 $\pm$ 5826
BR2 top	5B	75	9403 $\pm$ 189	87 $\pm$ 2	149.7 $\pm$ 2.1	0.6617 $\pm$ 0.0026	91 172 $\pm$ 618	88 052 $\pm$ 2289	192 $\pm$ 2.9	87 978 $\pm$ 2289
BR2 bottom	6A	89	7334 $\pm$ 148	131 $\pm$ 3	145.3 $\pm$ 2.1	0.6576 $\pm$ 0.0023	90 917 $\pm$ 567	88 868 $\pm$ 1552	187 $\pm$ 2.8	88 803 $\pm$ 1552
BR2 bottom	6B	89	6498 $\pm$ 130	149 $\pm$ 3	146.5 $\pm$ 1.6	0.6576 $\pm$ 0.0021	90 748 $\pm$ 500	88 951 $\pm$ 1363	188 $\pm$ 2.2	88 887 $\pm$ 1363
BR3 Top big	1A	181	21 239 $\pm$ 428	106 $\pm$ 2	177.6 $\pm$ 2.1	0.7560 $\pm$ 0.0027	108 039 $\pm$ 732	105 253 $\pm$ 2096	239 $\pm$ 3.0	105 118 $\pm$ 2096
BR3 Top big	1B	180	10 042 $\pm$ 202	200 $\pm$ 4	179.9 $\pm$ 2.1	0.6791 $\pm$ 0.0023	90 822 $\pm$ 548	89 494 $\pm$ 1084	232 $\pm$ 3.0	89 429 $\pm$ 1084
BR3 Top small	2A	119	39 624 $\pm$ 797	36 $\pm$ 1	144.7 $\pm$ 2.1	0.7263 $\pm$ 0.0032	106 550 $\pm$ 859	98 117 $\pm$ 6042	191 $\pm$ 4.0	98 052 $\pm$ 6042
BR3 Top small	2B	116	29 348 $\pm$ 590	47 $\pm$ 1	157.3 $\pm$ 2.0	0.7146 $\pm$ 0.0028	101 698 $\pm$ 716	95 434 $\pm$ 4493	206 $\pm$ 4.0	95 369 $\pm$ 4493
BR3 bottom	3A	85	32 394 $\pm$ 650	31 $\pm$ 1	157.8 $\pm$ 2.0	0.7228 $\pm$ 0.0028	103 518 $\pm$ 720	93 990 $\pm$ 6809	206 $\pm$ 4.7	93 925 $\pm$ 6809
BR3 bottom small	3B	82	12 161 $\pm$ 244	78 $\pm$ 2	155.4 $\pm$ 2.0	0.7017 $\pm$ 0.0023	99 080 $\pm$ 596	95 443 $\pm$ 2639	203 $\pm$ 3.1	95 378 $\pm$ 2639
BR3 bottom big	4A	48	10 654 $\pm$ 214	53 $\pm$ 1	164.3 $\pm$ 1.9	0.7216 $\pm$ 0.0028	102 216 $\pm$ 716	96 731 $\pm$ 3948	216 $\pm$ 3.5	96 666 $\pm$ 3948
BR3 bottom big	4B	47	9754 $\pm$ 196	56 $\pm$ 1	161.2 $\pm$ 2.1	0.7148 $\pm$ 0.0028	101 154 $\pm$ 727	96 025 $\pm$ 3700	211 $\pm$ 3.6	95 960 $\pm$ 3700

axis, the halved samples display small cavities and discontinuities. The growth axis clearly shifted with time as the drip point moved laterally, giving rise to an asymmetrical stalagmite so that only one flank of the underlying edifice was partially coated (Fig. 3).

Sample BR2 is approximately 28 cm in height and ranges from 8 to 10 cm in diameter. The shape is a nearly symmetrical candle with a larger diameter in the middle part. Compared to BR1, the yellowish-grey BR2 exhibits darker colours on its weathered outer surface. The surface texture is irregular, rough and has some visible macro-holes at the bottom (Fig. 2A). The cut sample shows growth layers with darker bands interlayered with lighter bands. There are several obvious discontinuities (hiatus; Fig. 3).

Sample BR3 is significantly different from BR1 and BR2 in shape. It incorporates a smaller stalagmite in its lower portion. The length of the outer stalagmite is 36 cm and the diameter 8–11 cm. The inner stalagmite has a length of 26 cm and a diameter of 3–6 cm. Both stalagmites have an asymmetrical shape with the smallest diameter in their middle. The growth layers and growth axis of the inner stalagmite are unlike the outer one (Fig. 3). The boundary between the stalagmites is characterized by a discontinuity. The porosity is high on the flanks of the two stalagmites.

In Fig. 3, CT images are displayed in greyscale. Lighter greyscale values indicate higher densities. Lower densities are observed in areas with high porosity or a hiatus (Fig. 3). The cut samples of BR1 and BR3 show axis holes (diameter >1 mm) extending parallel to the growth axis (Fig. 4). The axis holes were also previously observed in CT scan images of uncut samples (Fig. 3). This feature can

be caused by an elongated dissolution process that affects the primary porosity of columnar calcite and is initiated particularly as elongated cavities along the boundaries between neighbouring crystals (Perrin *et al.* 2014). The dissolution process gradually enlarges secondary voids. Their width varies from a few micrometres to a few hundreds of micrometres. In the last stage, these dissolution cavities become connected to each other and form a framework of micro-holes in the central part of the stalagmite (Perrin *et al.* 2014). The cut sample of BR2 shows its central voids with depression downward across the layers (Fig. 4), where the CT image indicates lower densities. This feature may be caused by under-saturated or corrosive water cutting through the growth layers along micro-fractures inside ~~as off-axis holes~~ (Shtober-Zisu *et al.* 2014).

The internal dissolution feature and shift of growth axis suggest that the samples BR1, BR2 and BR3 have suffered significant depositional changes during or after their growths. These changes however cannot be readily predicted from the uncut stalagmites. The CT images conversely reveal that the samples have significant density variations and contain small and large voids (Fig. 3). Obviously, the CT images provide unique information about the internal structure of stalagmites that cannot be identified by visual inspection of the uncut specimens (Figs 2B, 3).

Petrography and density

Petrographic analysis of polished thin sections can facilitate the identification of different types of calcite fabrics in the stalagmites (Frisia 2015). We prepared 19

Fig. 4. Samples BR1 and BR2 showing axis holes elongated parallel to the growth axis with diameter ranging from 1 to 5 mm. In BR2, the marked feature shows downward bending of growth layers towards the holes.

Fig. 5. Fabric types in samples BR1, BR2 and BR3 under plane polarized light (PPL) and cross-polarized light (XPL).

Table 2. Characteristics of CT number (HU values), petrography, and palaeoenvironment of stalagmites.

This study		Vanghi <i>et al.</i> (2015)		Walczak <i>et al.</i> (2015) and Frisia (2015)			
HU values	Fabrics	HU values	Fabrics	HU values	Fabrics	Growth rate	Drip rate
<1200	Open dendritic	<226	Porous	<2000	Columnar microcrystalline, dendritic	Fast	Not stable and seasonally variable
		227–776	Dendritic				
1200–1400	Closed dendritic	77–1176	Closed dendritic				
1401–1550	Columnar	1177–1576	Columnar				
		1577–3071	Micrite	2000–2400	Open columnar	Medium	Constant with high drip rate
				>2400	Compacted columnar	Slow	Constant with low drip rate

polished thin sections from samples BR1, BR2 and BR3. Four fabric types can be observed (Fig. 5A–H, see positions of polished thin sections in Fig. 3). (i) Open dendritic fabric formed interlocking and elongated crystals with spatial distribution. This texture is characterized by high porosity (Frisia *et al.* 2000, 2002). (ii) Closed dendritic fabric has more equal dimensions and forms less porous lamina. (iii) Columnar fabric is characterized by large crystals growing parallel to the central axis. Their crystal boundaries are acute. In general, the columnar fabric is less porous than the dendritic fabric (Frisia 2015). (iv) Micritic layers are observed mainly on the edges of the stalagmite BR3.

In this study, variations in the crystalline fabrics as well as the porosities are represented in the CT numbers (HU) in Table 2. The open dendritic fabric shows a high porosity with the lowest density (<1200 HU). Closed dendritic fabric has a medium density (1200–1400 HU). Columnar fabric has the highest density (1401–1550 HU). The calcite fabrics and CT numbers (HU) of Tham Nam stalagmites were then compared to those previously reported in Vanghi *et al.* (2015), Walczak *et al.* (2015) and Frisia (2015) (Table 2). Stalagmites BR1, BR2 and BR3 display CT numbers (HU) <1550, and fabrics indicate variable drip rates and fast growth rates (Frisia 2015). Density in a speleothem is related to primary crystalline texture and porosity. Secondary porosity can be the result of dissolution and/or fractures that lead to lower density. Diagenesis can cause loss of primary texture, mineralogy and geochemical signatures (Perrin *et al.* 2014; Scholz *et al.* 2014; Zhang *et al.* 2014; Bajo *et al.* 2016). Thus, the present work, in agreement with previous studies (Vanghi *et al.* 2015; Walczak *et al.* 2015), suggests that the areas within stalagmites regarded suitable for palaeoclimatic study should exhibit primary crystalline texture, low porosity and CT numbers (HU) of 1550 or higher.

U-Th dating

All U-Th ages of the subsamples from stalagmites BR1, BR2 and BR3 are replicable within uncertainties, except

for 1A and 1B (Table 1; Fig. 3). In fact, in each pair of replicates, the difference between the mean ages is much smaller than the uncertainty (2σ), which suggests that the U-Th chemical separations and machine measurements are robust. The measured ^{238}U concentrations and $\delta^{234}\text{U}$ values can also be well replicated. However, the ^{232}Th content and values of the $^{230}\text{Th}/^{232}\text{Th}$ atomic ratio show considerable variation. This Th heterogeneity is probably due to an uneven distribution of detritus, which contains high Th contents. The discrepancy in U-Th ages between subsamples 1A and 1B near the top layer of stalagmite BR3 is probably caused by a diagenetic process (e.g. dissolution, recrystallization and neomorphism) because they were drilled in the low-density part of the sample (Fig. 3).

^{238}U concentrations in all the samples are lower than 200 ppb and ^{232}Th concentrations range from thousands to tens of thousands of ppt. The isotopic ratios of ^{230}Th to ^{232}Th are also low, typically <200 ppm. This unfortunately limits the age precisions, which largely depend on the ^{230}Th initial correction. A low ^{238}U concentration and particularly low $^{230}\text{Th}/^{232}\text{Th}$ values indicate that a significant portion of ^{230}Th does not originate from *in situ* decay of ^{238}U . The large uncertainty of the detrital $^{230}\text{Th}/^{232}\text{Th}$ in bulk earth values thus dominates the age uncertainties. Nevertheless, if the ^{232}Th concentration in the subsample is low and the measured $^{230}\text{Th}/^{232}\text{Th}$ value is high, the relative uncertainty of the corrected age can easily be reduced to ~1% or better (e.g. BR1 bottom (8B), Table 1). After the initial Th correction, all the U-Th ages of BR1, BR2 and BR3 are concentrated in a small interval, ranging between 87 and 105 ka. This confirms that they were deposited during the same time interval.

CT images, petrography and U-Th isotope data indicate the dissolution features in the internal structure of stalagmites BR1, BR2 and BR3. These can be caused by several scenarios, for example, (i) undersaturated or acidic dripping water with high content of pCO_2 (Perrin *et al.* 2014); (ii) an abruptly increasing dripping water rate, which prevents sufficient degassing to bring feeding water to equilibrium with CaCO_3 (Railsback *et al.* 2011);

or (iii) mixing of dissolution solution with several drip water sources (Scholz *et al.* 2014). Recent studies suggest that microbial activity can affect calcification through trapping and binding detrital particles and inducing calcite precipitation. This leads to breakdown dissolution, boring and residue micrite production (Jones 2001; Shtober-Zisu *et al.* 2014). Such processes can change the primary isotopic and elemental composition of a speleothem. These changes alter the geochemical signals, which may have an impact on the interpretation of the results obtained in palaeoenvironmental studies (Martín-García *et al.* 2009). Typically, stalagmite samples containing these features are no longer suitable for palaeoclimate studies. For speleothem age determination, carbonate samples must have remained completely closed with respect to loss or gain of U and Th isotopes. For example, Bajo *et al.* (2016) reported that U-Th ages could be overestimated in a stalagmite that has anomalously high $^{230}\text{Th}/^{238}\text{U}$ isotopic ratios resulting from preferential U loss in micro-void dissolution. Based on the CT images, petrography analysis and large uncertainty of $^{230}\text{Th}/^{232}\text{Th}$, we hereby decided not to perform any additional dating and geochemical analysis on the stalagmites from Tham Nam cave.

Conclusions

In the present work, stalagmites from Tham Nam cave, western Thailand, were evaluated for their palaeoclimatic research potential, through X-ray computed tomography (CT scanning), petrography, and U-Th dating analysis. The CT scanning data reveal that stalagmite density is associated with fabrics and distribution of porosity. The porosity/axis-holes/voids in the stalagmites are related to postdepositional dissolution processes and may have an influence on U-Th dating accuracy. The ages of stalagmites from the Tham Nam Cave range between 87 and 105 ka. Currently, they are the oldest stalagmites that have been reported from mainland Southeast Asia. However, the stalagmites show dissolution features, several discontinuities and large age uncertainty, and are thus probably not suitable for palaeoclimatic research. Consistent with previous work (Mickler *et al.* 2004a, b; Zisu *et al.* 2012; Vanghi *et al.* 2015; Walczak *et al.* 2015 and Bajo *et al.* 2016), this study suggests that CT images have a large yet underestimated potential in speleothem study. They can help to identify internal structures in samples that may compromise geochemical elemental and isotopic analyses. For stalagmites potentially suitable for palaeoenvironmental research, CT scanning can also facilitate selection of the right cutting location. This study hence sets an example for future speleothem exploration using CT scanning as a tool to examine petrological textures prior to any further analysis.

Acknowledgements. – S. Chawchai wishes to express appreciation for the financial support provided by the Ratchadaphiseksomphot Endowment Fund, part of the ‘Research Grant for New Scholar CU Researcher’s

Project’ and the Thailand Research Fund (MRG5980080). The authors would also like to thank Mr Siripong Tonongto, his family and PANDA CAMP for accommodation and assistance during the field survey. In addition, the authors want to thank Peerapong Sritangsirikul and officials at Ban Rai district, Uthai Thani province, for their help and hospitality as well as Dr Thanisa Tongbai (Diagnostic Radiologist) and Walaiporn Suksancharoen (nurse) at King Chulalongkorn Memorial Hospital for their advice and help. We also thank the editor and anonymous reviewers for their careful reading of our manuscript and their insightful comments and suggestions.

References

- Bajo, P., Hellstrom, J., Frisia, S., Drysdale, R., Black, J., Woodhead, J., Borsato, A., Zanchetta, G., Wallace, M. W., Regattieri, E. & Haese, F. 2016: ‘Cryptic’ diagenesis and its implications for speleothem geochronologies. *Quaternary Science Reviews* 148, 17–28.
- Cai, B., Pumijumnong, N., Tan, M., Muangsong, C., Kong, X., Jiang, X. & Nan, S. 2010: Effects of intraseasonal variation of summer monsoon rainfall on stable isotope and growth rate of a stalagmite from northwestern Thailand. *Journal of Geophysical Research: Atmospheres* 115, D21104.
- Cai, Y., Fung, I. Y., Edwards, R. L., An, Z., Cheng, H., Lee, J. E., Tan, L., Shen, C. C., Wang, X., Day, J. A., Zhou, W., Kelly, W. & Chiang, J. 2015: Variability of stalagmite-inferred Indian monsoon precipitation over the past 252,000 Y. *Proceedings of the National Academy of Sciences of the United States of America* 112, 2954–2959.
- Chen, F., Xu, Q., Chen, J., Birks, H. J. B., Liu, J., Zhang, S., Jin, L., An, C., Telford, R. J., Cao, X., Wang, Z., Zhang, X., Selvaraj, K., Lu, H., Li, Y., Zheng, Z., Wang, H., Zhou, A., Dong, G., Zhang, J., Huang, X., Bloemendal, J. & Rao, Z. 2015: East Asian summer monsoon precipitation variability since the last deglaciation. *Scientific Reports* 5, 11186.
- Cheng, H., Edwards, R. L., Shen, C. C., Polyak, V. J., Asmerom, Y., Woodhead, J., Hellstrom, J., Wang, Y., Kong, X., Spötl, C., Wang, X. & Calvin Alexander, E. Jr. 2013: Improvements in ^{230}Th dating, ^{230}Th and ^{234}U half-life values, and U-Th isotopic measurements by multi-collector inductively coupled plasma mass spectrometry. *Earth and Planetary Science Letters* 371, 82–91.
- Cook, E. R., Anchukaitis, K. J., Buckley, B. M., D’Arrigo, R. D., Jacoby, G. C. & Wright, W. E. 2010: Asian monsoon failure and mega drought during the last millennium. *Science* 328, 486–489.
- Department of Mineral Resources (DMR) 2009: *Geological Map of Thailand (Uthai Thani) Scale 1: 250000*. Ministry of Natural Resource and Environment, Bangkok, Thailand.
- Edwards, R. L., Chen, J. H. & Wasserburg, G. J. 1986: ^{238}U – ^{234}U – ^{230}Th – ^{232}Th systematics and the precise measurement of time over the past 500,000 years. *Earth and Planetary Science Letters* 81, 175–192.
- Fairchild, I. J. & Treble, P. C. 2009: Trace elements in speleothems as recorders of environmental change. *Quaternary Science Reviews* 28, 449–468.
- Fairchild, I. J., Frisia, S., Borsato, A. & Tooth, A. F. 2007: Speleothems. In Nash, D. J. & McLaren, S. J. (eds.): *Geochemical Sediments and Landscapes*, 200–245. Blackwell Publishing Ltd, Oxford.
- Fairchild, I. J., Smith, C. L., Baker, A., Fuller, L., Spötl, C., Matthey, D. & McDermott, F. 2006: Modification and preservation of environmental signals in speleothems. *Earth-Science Reviews* 75, 105–153.
- Frisia, S. 2015: Microstratigraphic logging of calcite fabrics in speleothems as tool for palaeoclimate studies. *International Journal of Speleology* 44, 1–16.
- Frisia, S., Borsato, A., Fairchild, I. J. & McDermott, F. 2000: Calcite fabrics, growth mechanisms, and environments of formation in speleothems from the Italian Alps and Southwestern Ireland. *Journal of Sedimentary Research* 70, 1183–1196.
- Frisia, S., Borsato, A., Fairchild, I. J., McDermott, F. & Selmo, E. M. 2002: Aragonite-calcite relationships in speleothems (Grotte de Clamouse, France): environment, fabrics, and carbonate geochemistry. *Journal of Sedimentary Research* 72, 687–699.

- Hendy, C. H. 1971: The isotopic geochemistry of speleothems—I. The calculation of the effects of different modes of formation on the isotopic composition of speleothems and their applicability as palaeoclimatic indicators. *Geochimica et Cosmochimica Acta* 35, 801–824.
- Jaffey, A. H., Flynn, K. F., Glendenin, L. E., Bentley, W. T. & Essling, A. M. 1971: Precision measurement of half-lives and specific activities of ^{235}U and ^{238}U . *Physical Review C* 4, 1889.
- Jones, B. 2001: Microbial activity in caves—a geological perspective. *Geomicrobiology Journal* 18, 345–357.
- Ketcham, R. A. & Carlson, W. D. 2001: Acquisition, optimization and interpretation of X-ray computed tomographic imagery: applications to the geosciences. *Computers and Geosciences* 27, 381–400.
- Lachniet, M. S. 2009: Climatic and environmental controls on speleothem oxygen-isotope values. *Quaternary Science Reviews* 28, 412–432.
- Loo, Y. Y., Billa, L. & Singh, A. 2015: Effect of climate change on seasonal monsoon in Asia and its impact on the variability of monsoon rainfall in Southeast Asia. *Geoscience Frontiers* 6, 817–823.
- Martín-García, R., Alonso-Zarza, A. M. & Martín-Pérez, A. 2009: Loss of primary texture and geochemical signatures in speleothems due to diagenesis: evidences from Castañar Cave, Spain. *Sedimentary Geology* 221, 141–149.
- McDermott, F. 2004: Palaeo-climate reconstruction from stable isotope variations in speleothems: a review. *Quaternary Science Reviews* 23, 901–918.
- Mickler, P. J., Banner, J. L., Stern, L., Asmerom, Y., Edwards, R. L. & Ito, E. 2004b: Stable isotope variations in modern tropical speleothems: evaluating equilibrium vs. kinetic isotope effects. *Geochimica et Cosmochimica Acta* 68, 4381–4393.
- Mickler, P. J., Ketcham, R. A., Colbert, M. W. & Banner, J. L. 2004a: Application of high-resolution X-ray computed tomography in determining the suitability of speleothems for use in palaeoclimatic, palaeohydrologic reconstructions. *Journal of Cave and Karst Studies* 66, 3–8.
- Muangsong, C., Cai, B., Pumijumnong, N., Hu, C. & Cheng, H. 2014: An annually laminated stalagmite record of the changes in Thailand monsoon rainfall over the past 387 years and its relationship to IOD and ENSO. *Quaternary International* 349, 90–97.
- ~~Muangsong, C., Cai, B., Pumijumnong, N., Hu, C. & Lei, G. 2016: Intra-seasonal variability of teak tree-ring cellulose $\delta^{18}\text{O}$ from northwestern Thailand: a potential proxy of Thailand summer monsoon rainfall. *The Holocene* 26, 1397–1405.~~
- Perrin, C., Prestimonaco, L., Servelle, G., Tilhac, R., Maury, M. & Cabrol, P. 2014: Aragonite–calcite speleothems: identifying original and diagenetic features. *Journal of Sedimentary Research* 84, 245–269.
- Raghavan, S. V., Liu, J., Nguyen, N. S., Vu, M. T. & Liong, S. Y. 2017: Assessment of CMIP5 historical simulations of rainfall over Southeast Asia. *Theoretical and Applied Climatology* 222, 1–14.
- Railsback, L. B., Dabous, A. A., Osmond, J. K. & Fleisher, C. J. 2002: Petrographic and geochemical screening of speleothems for U-series dating: an example from recrystallized speleothems from Wadi Sannur Cavern, Egypt. *Journal of Cave and Karst Studies* 64, 108–116.
- Railsback, L. B., Liang, F., Romaní, J. R. V., Grandal-d'Anglade, A., Rodríguez, M. V., Fidalgo, L. S., Mosquera, D. F., Cheng, H. & Edwards, R. L. 2011: Petrographic and isotopic evidence for Holocene long-term climate change and shorter-term environmental shifts from a stalagmite from the Serra do Courel of northwestern Spain, and implications for climatic history across Europe and the Mediterranean. *Palaeogeography, Palaeoclimatology, Palaeoecology* 305, 172–184.
- Scholz, D., Tolzmann, J., Hoffmann, D. L., Jochum, K. P., Spötl, C. & Riechelmann, D. F. 2014: Diagenesis of speleothems and its effect on the accuracy of $^{230}\text{Th}/\text{U}$ -ages. *Chemical Geology* 387, 74–86.
- Shen, C. C., Wu, C. C., Cheng, H., Edwards, R. L., Hsieh, Y. T., Gallet, S., Chang, C.-C., Li, T.-Y., Lam, D. D., Kano, A., Hori, M. & Spötl, C. 2012: High-precision and high-resolution carbonate ^{230}Th dating by MC-ICP-MS with SEM protocols. *Geochimica et Cosmochimica Acta* 99, 71–86.
- Shtober-Zisu, N., Schwarcz, H. P., Chow, T., Omelon, C. R. & Southam, G. 2014: Caves in caves: evolution of post-depositional macroholes in stalagmites. *International Journal of Speleology* 43, 323–334.
- Sinha, A., Kathayat, G., Cheng, H., Breitenbach, S. F., Berkelhammer, M., Mudelsee, M., Biswas, J. & Edwards, R. L. 2015: Trends and oscillations in the Indian summer monsoon rainfall over the last two millennia. *Nature Communications* 6, ncomms7309.
- ~~Sinha, A., Stott, L., Berkelhammer, M., Cheng, H., Edwards, R. L., Buckley, B., Aldenderfer, M. & Mudelsee, M. 2011: A global context for megadroughts in monsoon Asia during the past millennium. *Quaternary Science Reviews* 30, 47–62.~~
- Thirumalai, K., DiNezio, P. N., Okumura, Y. & Deser, C. 2017: Extreme temperatures in Southeast Asia caused by El Niño and worsened by global warming. *Nature Communications* 8, ncomms15531.
- Vanghi, V., Iriarte, E. & Aranburu, A. 2015: High resolution X-ray computed tomography for petrological characterization of speleothems. *Journal of Cave and Karst Studies* 77, 75–82.
- Walczak, I. W., Baldini, J. U., Baldini, L. M., McDermott, F., Marsden, S., Standish, C. D., Richards, D. A., Andreo, B. & Slater, J. 2015: Reconstructing high-resolution climate using CT scanning of unsectioned stalagmites: a case study identifying the mid-Holocene onset of the Mediterranean climate in southern Iberia. *Quaternary Science Reviews* 127, 117–128.
- Wong, C. I. & Breecker, D. O. 2015: Advancements in the use of speleothems as climate archives. *Quaternary Science Reviews* 127, 1–18.
- Zhang, H., Cai, Y., Tan, L., Qin, S. & An, Z. 2014: Stable isotope composition alteration produced by the aragonite-to-calcite transformation in speleothems and implications for palaeoclimate reconstructions. *Sedimentary Geology* 309, 1–14.
- Zhang, P., Cheng, H., Edwards, R. L., Chen, F., Wang, Y., Yang, X., Liu, J., An, C., Dai, Z., Zhou, J., Zhang, D., Jia, J., Jin, L. & Johnson, R. K. 2008: A test of climate, sun, and culture relationships from an 1810-year Chinese cave record. *Science* 322, 940–942.
- Zisu, N. S., Schwarcz, H. P., Konyer, N., Chow, T. & Noseworthy, M. D. 2012: Macroholes in stalagmites and the search for lost water. *Journal of Geophysical Research: Earth Surface* 117, F03020.