Abelian Manna model in three dimensions and below

Huynh, Hoai Nguyen.; Pruessner, Gunnar.

2012

Huynh, H. N., & Pruessner, G. (2012). Abelian Manna model in three dimensions and below. *Physical Review E*, 85(6), 061133-.

<https://hdl.handle.net/10356/95175>

<https://doi.org/10.1103/PhysRevE.85.061133>

© 2012 American Physical Society. This paper was published in *Physical Review E* and is made available as an electronic reprint (preprint) with permission of American Physical Society. The paper can be found at the following official DOI: [<http://dx.doi.org/10.1103/PhysRevE.85.061133>]. One print or electronic copy may be made for personal use only. Systematic or multiple reproduction, distribution to multiple locations via electronic or other means, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper is prohibited and is subject to penalties under law.

Downloaded on 26 Apr 2023 18:47:41 SGT

Abelian Manna model in three dimensions and below

Hoai Nguyen Huynh^{1,*} and Gunnar Pruessner^{2,†}

¹*Division of Physics and Applied Physics, School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang Link, Singapore 637371, Singapore*

²*Department of Mathematics, Imperial College London, 180 Queen's Gate, London SW7 2BZ, England, United Kingdom*

(Received 11 January 2012; revised manuscript received 22 May 2012; published 29 June 2012)

The Abelian Manna model of self-organized criticality is studied on various three-dimensional and fractal lattices. The exponents for avalanche size, duration, and area distribution of the model are obtained by using a high-accuracy moment analysis. Together with earlier results on lower-dimensional lattices, the present results reinforce the notion of universality below the upper critical dimension and allow us to determine the coefficients of an ϵ expansion. By rescaling the critical exponents by the lattice dimension and incorporating the random walker dimension, a remarkable relation is observed, satisfied by both regular and fractal lattices.

DOI: [10.1103/PhysRevE.85.061133](https://doi.org/10.1103/PhysRevE.85.061133)

PACS number(s): 05.70.Jk, 64.60.al, 05.65.+b

I. INTRODUCTION

Critical phenomena play an important role in our understanding of complex systems in nature. One of the key features of critical systems is the notion of universality [1]. It suggests common underlying mechanisms of seemingly different phenomena and also allows us to study complicated natural systems through analysis of simple (numerical) models. A huge number of numerical models has been proposed to study different features of critical systems. Traditionally, those systems require external fine tuning of a control parameter to the critical point. However, many models exhibit the features of the critical state without the need of tuning a control parameter, which is known as self-organized criticality [2]. While models of traditional critical phenomena have been very well studied both analytically and numerically, and important results including exact ones have been obtained, e.g., [3], little success has been achieved for self-organized critical phenomena. One example of such models is the Manna model [4] which has so far defied any attempt for an analytical approach but has been studied extensively numerically. Yet the question of universality seems to have been overlooked in the literature on a very fundamental point: Does the Manna model display the same critical behavior on different lattices of the same dimension? Recent extensive numerical studies [5] of the Abelian version of the Manna model [4,6,7] on various lattices in integer dimensions 1 and 2 provide very strong support for the model's universal behavior. The situation, however, is more complicated in noninteger dimensions [8]. Thus, it would be of great interest to see if the results in fractal dimensions can be reconciled with those in integer dimensions in a systematic manner.

Our motivation for this study is to provide a complete numerical picture of the Manna model. Three main results are reported: First, we confirm universality in three dimensions, i.e., the independence of critical exponents and moment ratios from the detailed structure of the underlying lattice. This allows us, second, to firmly estimate the coefficients of an ϵ expansion for the exponents. Third, we identify a general

scaling relation unifying critical behavior on regular and fractal lattices.

II. MODEL AND OBSERVABLES

The Abelian Manna model [4–7] is defined on a lattice of dimension d with N sites and linear extent L , where $N \propto L^d$ asymptotically. Each site i has a local degree of freedom $z_i \geq 0$ which can be thought of as the number of particles residing at that site. If $z_i > 1$, the site is said to be active or unstable, otherwise it is stable. The system evolves by driving and subsequently fully relaxing it. “Driving” means the system is “charged” by picking a site j randomly and uniformly and increasing z_j by 1. “Relaxation” means an unstable site i is picked randomly and uniformly; its particle number is reduced by 2 and the particle number of two of its q_i neighbors, which are chosen independently and at random (possibly the same one twice), is increased by 1, thereby possibly rendering them unstable. Dissipation takes place only when relaxing boundary sites transfer particles to $q^{(v)}$ virtual neighbors “outside” the lattice. These virtual neighbors cannot topple themselves and are chosen so that the topology of the finite lattice corresponds to that of an “offcut” from an infinite lattice. Relaxation of unstable sites continues until there are no unstable sites left. Only then the system is driven again, known as a separation of time scales. The number of topplings between two driving steps is the size s of the “avalanche,” and the number of distinct sites receiving at least one particle is the area a . The duration t of the avalanche is measured on the microscopic time scale, which advances in steps of $1/N_a$, where N_a is the instantaneous number of active sites, mimicking a Poissonian decay of active sites. The moments of the observables mentioned above are measured in the stationary state, which is reached after a generously estimated transient. A number of (asymptotic) key characteristics of the lattices are listed in Table I, such as the average number of neighbors $\bar{q}$, the average number of virtual neighbors $\bar{q}^{(v)}$ among sites with at least one virtual neighbor, and the particle density ζ in the stationary state.

The probability densities $\mathcal{P}^{(x)}(x, L)$ of the observables, $x \in \{s, a, t\}$, are expected to display finite size scaling,

$$\mathcal{P}^{(x)}(x, L) = a_x x^{-\tau_x} \mathcal{G}_x \left(\frac{x}{b_x L^{D_x}} \right), \quad (1)$$

*n.huynh10@imperial.ac.uk; <http://www3.ntu.edu.sg/home2008/hu0004en/>.

†g.pruessner@imperial.ac.uk; <http://www.ma.ic.ac.uk/pruess/>.

TABLE I. Key features of the three-dimensional ($d = 3$) lattices considered in this work. The random walker dimension d_w is 2 for every regular lattice [20]. The asymptotic site averaged number of nearest neighbors is $\bar{q}$, with boundary sites having on average $\bar{q}^{(v)}$ neighbors missing (virtual neighbors [5]). The stationary particle density, equal to the average density of (singly) occupied sites, is given by ζ .

Lattice	d	d_w	$\bar{q}$	$\bar{q}^{(v)}$	ζ
SC	3	2	6	1	[0.622325(1)]
BCC	3	2	8	4	[0.600620(2)]
BCCN	3	2	14	5	[0.581502(1)]
FCC	3	2	12	4	[0.589187(3)]
FCCN	3	2	18	5	[0.566307(3)]

provided that $L \gg 1$ and $x \gg x_0$ for some lower cutoff x_0 . The metric factors a_x and b_x are not expected to be universal [9], whereas the exponent D_x should only depend on the dimension and τ_x should only depend on the dimension and the boundary conditions [10]. The universal scaling function $\mathcal{G}_x$ is characterized below by moment ratios. For $n > \tau_x - 1$, the moments scale asymptotically like $\langle x^n \rangle \propto L^{\mu_n^{(x)}}$ with $\mu_n^{(x)} = D_x(1 + n - \tau_x)$. Historically, the exponents are denoted as D (for D_s), τ (for τ_s), z (for D_t), and α (for τ_t).

Five three-dimensional lattices and five fractal lattices are employed in this study. The three-dimensional lattices [11] are built upon the standard simple cubic (SC) lattice. The body centered cubic (BCC) and face centered cubic (FCC) lattices are also studied with next-nearest-neighbor interactions (BCCN and FCCN, respectively). The total numbers of sites N of all five lattices are chosen to be as close as possible to one another. Typically, six system sizes ranging from $N = 181^3$ to 1024^3 are used. A total of approximately 50 000 CPU hours has been spent on three-dimensional systems.

The key features of the fractal lattices are listed in Table II. Of those the lesser known semi-inverse square triadic Koch (SSTK) lattice [12,13] has Hausdorff dimension $d = \ln 5 / \ln 3$ and is shown in Fig. 1. Of particular interest is the Sierpinski tetrahedron (SITE) lattice, which is the three-dimensional version of the well-known Sierpinski gasket, based on tetrahedra instead of triangles. Its fractal dimension is $d = 2$ and thus allows a direct comparison to regular two-dimensional lattices. The strongly anisotropic extended Sierpinski gasket (EXGA, $d = \ln 6 / \ln 2$) is obtained by stacking L copies of a Sierpinski gasket on top of one another and applying periodic boundary conditions. Finally,

TABLE II. Key features of the fractal lattices studied, as listed in Table I for the three-dimensional ones. The random walker dimensions are exactly known or derived (or, in the case of SSTK, estimated) using the methods described in [8].

Lattice	d	d_w	$\bar{q}$	$\bar{q}^{(v)}$	ζ
SSTK	1.464...	2.552..	3	1	[0.8435(2)]
ARRO	1.584...	2.322..	7/3	1	[0.862(2)]
CRAB	1.584...	2.578..	7/3	1	[0.8794(6)]
SITE	2	2.584..	6	3	[0.7427(3)]
EXGA	2.584...	2.321..	6	2	[0.65640(8)]

FIG. 1. Semi-inverse square triadic Koch lattice at iteration 3.

the arrowhead (ARRO, $d = \ln 3 / \ln 2$) and the crab (CRAB, $d = \ln 3 / \ln 2$) lattices are the same as the ones used in [8]. Typically, four system sizes corresponding to iterations from 6 to 9 (5 to 8 for SSTK) are used for all fractal lattices. A total of approximately 75 000 CPU hours has been spent on fractal lattices.

III. RESULTS

Details of the Monte Carlo simulation, the fitting procedures, and the derivation of the error bars can be found in [5]. In short, individual moments $\langle x^n \rangle(L)$ are fitted against the system size $N \propto L^d$ to obtain the scaling exponent $\mu_n^{(x)}/d$ using a power law with corrections. For regular three-dimensional lattices, the form of the fitting function used for avalanche size and duration is

$$\langle x^n \rangle(N) = A_n^{(x)} N^{\mu_n^{(x)}/d} + B_n^{(x)} N^{\mu_n^{(x)}/d-0.25} \quad (2)$$

and for avalanche area is

$$\langle a^n \rangle(N) = A_n^{(a)} N^{\mu_n^{(a)}/d} + B_n^{(a)} N^{\mu_n^{(a)}/d-0.25} + C_n^{(a)} N^{\mu_n^{(a)}/d-0.5}. \quad (3)$$

For fractal lattices we use Eq. (2) in [8]:

$$\langle x^n \rangle(N) = A_n^{(x)} N^{\mu_n^{(x)}/d} + B_n^{(x)} N^{\mu_n^{(x)}/d-1} + C_n^{(x)} N^{\mu_n^{(x)}/d-2}. \quad (4)$$

The estimated scaling exponents $\mu_n^{(x)}$ are then linearly fitted against the moment orders n ($n = 2, 3, 4, 5$ for all except for $\mu_n^{(s)}$ in three dimensions with $n = 2, 3, 4$). The slope gives the exponent D_x , and the interception with the abscissa gives the exponent τ_x , except for three-dimensional lattices, whose estimate is obtained by employing the exact relation $D_s(2 - \tau_s) = 2$. Although the relative errors are as small as 3×10^{-3} , we were unable to adjust the fitting scheme to recover $\mu_1^{(s)} = 2$ within less than three standard deviations.

The quality of all data fitting reported in this work is assessed by the goodness of fit q [14], which is considered good if $q > 0.1$; otherwise the data are marked by $[-]$. Tables III and V summarize the estimated critical exponents, all obtained with $q > 0.5$. For regular lattices, our results compare well with the literature [15–17] (also [18,19] for absorbing state phase transitions), although some variability and discrepancy are observed, in particular for z , which may be explained by the use of slightly different model definitions (and dynamics) by other authors. A number of scaling relations (see below) are

TABLE III. Avalanche exponents of five three-dimensional lattices. The estimates for τ and $D(\tau - 1)$ are obtained from D via the exact scaling relation $D(2 - \tau) = 2$. Identities $D_a = d$ and $\mu_1^{(s)} = 2$ are used to validate the fitting scheme.

Lattice	d	d_w	D	τ	z	α	D_a	τ_a	$\mu_1^{(s)}$	$-\Sigma_s$	$-\Sigma_t$	$-\Sigma_a$
SC	3	2	3.38(2)	1.408(3)	1.779(7)	1.784(9)	3.04(5)	1.45(4)	2.0057(5)	1.38(2)	1.395(16)	1.36(13)
BCC	3	2	3.36(2)	1.404(4)	1.777(8)	1.78(1)	2.99(2)	1.444(18)	2.0030(5)	1.36(2)	1.390(19)	1.33(6)
BCCN	3	2	3.38(3)	1.408(4)	1.776(9)	1.783(11)	3.01(3)	1.44(3)	2.0041(6)	1.38(3)	1.39(2)	1.32(7)
FCC	3	2	3.35(4)	1.402(8)	1.765(16)	1.78(2)	3.1(2)	1.48(14)	2.0035(11)	1.35(4)	1.37(4)	1.5(5)
FCCN	3	2	3.38(4)	1.408(7)	1.781(14)	1.787(18)	3.00(4)	1.44(3)	2.0051(8)	1.38(4)	1.40(3)	1.32(9)
Overall	3	2	3.370(11)	1.407(2)	1.777(4)	1.783(5)	3.003(14)	1.442(12)	2.0042(3)		1.380(13)	
[15]	3	2	3.33	1.43	1.8							
[16]	3	2	3.302(10)		1.713(10)							
[17]	3	2	3.36(1)	1.41(1)	1.76(1)	1.78(2)						
[22]	3	2		1.41(2)	1.823(23)	1.77(4)						

confirmed, such as $-\Sigma_x = D_x(\tau_x - 1) = D_y(\tau_y - 1) = -\Sigma_y$ [16]. Overall estimates are included in Table III; the correlation of Σ_s , Σ_t , and Σ_a is taken into account by multiplying their respective errors by $\sqrt{3}$.

The moment ratios [5,21] $g_n^{(x)}$ (to leading order) are independent of the system size and characterize the scaling function $\mathcal{G}_x$. The fitting of moment ratios follows a similar procedure as the avalanche exponents, using

$$g_n^{(x)} = \mathcal{A}_n^{(x)} + \mathcal{B}_n^{(x)} N^{-0.25} + \mathcal{C}_n^{(x)} N^{-0.5}. \quad (5)$$

Together with the avalanche exponents, they provide very strong support for universality in regular lattices. Table IV lists the overall moment ratios based on five three-dimensional lattices.

Surprisingly (see [5] for the same phenomenon in one and two dimensions), the *amplitudes* of the leading order of the moments of the avalanche area seem to be universal. We found

$$\begin{aligned} \langle a^1 \rangle &= [0.202(4)] N^{\tilde{\mu}_1^{(a)}}, \\ \langle a^2 \rangle &= 0.0151(15) N^{\tilde{\mu}_2^{(a)}}, \\ \langle a^3 \rangle &= 0.0027(6) N^{\tilde{\mu}_3^{(a)}}, \\ \langle a^4 \rangle &= 0.00055(19) N^{\tilde{\mu}_4^{(a)}}, \\ \langle a^5 \rangle &= 0.00012(6) N^{\tilde{\mu}_5^{(a)}}, \end{aligned} \quad (6)$$

with universal $\tilde{\mu}_n^{(a)} = n + 1 - 1.4396(8)$ across the three-dimensional lattices introduced above. It is obviously crucial to consider $\langle a^n \rangle$ as a function of N , as fitting against $L = \lambda N^{1/d}$ leads to different amplitudes, because λ varies from lattice to lattice.

A. Regular lattices

All critical exponents including previous results [5] are summarized in Table VI. First, on regular lattices, a relation between D_x , τ_x , and the dimension d can be obtained by fitting

TABLE IV. Overall estimates of moment ratios for three-dimensional lattices.

Observable	x	$g_3^{(x)}$	$g_4^{(x)}$	$g_5^{(x)}$	$g_6^{(x)}$
Size	s	2.373(16)	7.76(17)	30.0(14)	121(8)
Duration	t	[4.164(6)]	[25.99(9)]	[201.4(12)]	1811(18)
Area	a	2.331(4)	7.30(5)	27.1(3)	113(2)

exponents against a proposed function $D_x = f_x(d)$ and $\tau_x = h_x(d)$. With six exponents six functions are to be determined, which, however, are related by scaling laws. They are

$$D(2 - \tau) = 2 \quad (7)$$

on regular lattices (exact [10]), $D_a = d$ (assumed to hold on regular lattices by [15,24], and in the present case confirmed for fractal lattices), and $D_x(\tau_x - 1) = -\Sigma_x$ with $\Sigma_a = \Sigma_s = \Sigma_t$ (narrow distribution assumption [25]). Using $\tau = 2 - 2/D$, $D_a = d$, $\tau_a = 1 + (D - 2)/d$, and $\alpha = 1 + (D - 2)/z$, there are thus only two functions to determine, which are best expressed in terms of $\epsilon = 4 - d$ since $d_c = 4$ is the upper critical dimension [19], where the exponents are known exactly. Writing $D = 4 - c_1^{(s)}\epsilon + c_2^{(s)}\epsilon^2 + \dots$, at most two amplitudes $c_i^{(s)}$ can reasonably be determined on the basis of the three data points available. A fit of D with only a linear term produces a very poor goodness of fit, which does not improve satisfactorily by including a term quadratic in ϵ . Omitting the quadratic gives

$$D = 4 - 0.654(6)\epsilon + 0.0079(10)\epsilon^3, \quad (8)$$

FIG. 2. Data of Table VI plotted in the form D/d vs d_w/d as suggested by Eq. (10). The dashed straight line is based on the estimates $a = [0.6061(5)]$ and $b = [0.7878(10)]$; the dotted line is the mean-field theory, $a = 0$, $b = 2$.

FIG. 3. Fit of the exponents in all dimensions (on regular and fractal lattices) against Eq. (11). The symbols represent the data in Table VI, the dashed lines are the fits as described in the text, Eq. (11) and Eq. (13), respectively.

with $q \approx 0.095$ [$c_1^{(s)} = 0.60(4)$, $c_2^{(s)} = -0.05(3)$, $c_3^{(s)} = -0.019(7)$ with three terms]. Similarly, $z = 2 - 0.239(4)\epsilon + 0.0056(6)\epsilon^3$, however with nearly vanishing goodness of fit.

B. Fractal lattices

Attempting to unify the above ϵ expansion obtained for regular lattices with the results for fractals with Hausdorff dimension d is bound to fail, which is immediately clear when comparing the exponents found for the fractal SITE lattice ($d = 2$, Table V) with those for the regular two-dimensional lattices or the ARRO with the CRAB lattice, Table V, which have the same Hausdorff dimension. As is well understood, the basic scaling relation Eq. (7) is valid only for regular lattices and has to be generalized to

$$D(2 - \tau) = d_w, \quad (9)$$

with random walker dimension $d_w \geq 2$ [8,20].

It turns out, however, that D is essentially a linear function of d and d_w :

$$D = ad + bd_w, \quad (10)$$

with the same coefficients a and b for both regular and fractal lattices, which can be extracted from the ϵ expansion obtained above with $d_c = 4 = 2d_w$, because $d_w = 2$ on regular lattices, $a = c_1^{(s)}$ and $b = 2(1 - c_1^{(s)})$, so that on the basis of Eq. (8) $a =$

$0.654(6)$ and $b = 0.692(12)$ or $a = 0.60(4)$ and $b = 0.80(8)$ depending on $c_1^{(s)}$. Fitting the data in Table VI against Eq. (10) gives $a = [0.550(4)]$, $b = [0.822(3)]$, and a fitting with the constraint $b = 2(1 - a)$ from exactly known values of exponents at $d = d_c = 4$ gives $a = [0.6061(5)]$, $b = [0.7878(10)]$. Figure 2 shows D/d as a function of d_w/d for all lattices listed in Table VI. As expected from Eq. (10), fractal and regular lattices display essentially the same linear dependence $D/d = a + bd_w/d$. Above the upper critical dimension $d_c = 4$ regular lattices attain their mean-field values, $D = 4$, $\tau = 3/2$ [23] and $d_w = 2$, therefore following Eq. (10) with $a = 0$ and $b = 2$. One may wonder whether fractal lattices with Hausdorff dimension greater than $d_c = 4$ have corresponding exponents $D = 2d_w$.

The choice of rescaling exponent D by dimension d of the lattice is not random, but rather a natural choice, given that we performed all fitting of $\mu_n^{(x)}$ against the number of sites N rather than the lattices' linear length L and multiplied the results by the Hausdorff dimension d . The gap exponents for the scaling in N is D/d , which, as it turns out, displays a very systematic dependence on d_w/d .

Further investigation shows that D/d fits very well to

$$\left(\frac{D}{d}\right)^2 (\tau - \tilde{a}) = \tilde{b}, \quad (11)$$

TABLE V. Avalanche exponents of five fractal lattices.

Lattice	d	d_w	D	τ	z	α	D_a	τ_a	$\mu_1^{(s)}$	$-\Sigma_s$	$-\Sigma_t$	$-\Sigma_a$
SSTK	1.464..	2.552..	2.94(3)	1.13(2)	1.817(17)	1.21(2)	1.466(5)	1.273(11)	2.551(6)	0.37(6)	0.38(4)	0.399(17)
ARRO	1.584..	2.322..	2.793(2)	1.173(2)	1.673(1)	1.280(2)	1.5847(3)	1.2985(6)	2.3103(4)	0.484(5)	0.468(3)	0.473(1)
CRAB	1.584..	2.578..	3.020(5)	1.151(4)	1.837(3)	1.237(4)	1.5847(8)	1.279(2)	2.5655(12)	0.456(11)	0.435(7)	0.443(3)
SITE	2	2.584..	3.232(6)	1.211(4)	1.870(4)	1.357(4)	1.9975(9)	1.339(2)	2.5533(6)	0.682(14)	0.667(8)	0.677(3)
EXGA	2.584..	2.321..	3.352(4)	1.312(3)	1.835(3)	1.581(3)	2.5895(6)	1.3915(8)	2.3000(2)	1.046(10)	1.066(6)	1.014(2)

TABLE VI. Summary of exponents in all dimensions.

Lattice	d	d_w	D	τ	z	α	D_a	τ_a	$\mu_1^{(s)}$	$-\Sigma$	Ref.
Regular	1	2	2.253(14)	1.112(6)	1.445(10)	1.18(2)	0.998(3)	1.259(11)	1.996(3)	0.26(2)	[5]
SSTK	1.464.. ^a	2.552..	2.94(3)	1.13(2)	1.817(17)	1.21(2)	1.466(5)	1.273(11)	2.551(6)	0.40(3)	Here
ARRO	1.584.. ^b	2.322..	2.7938(19)	1.1731(16)	1.6732(12)	1.2797(17)	1.5847(3)	1.2985(6)	2.3103(4)	0.4730(16)	Here+ [8]
CRAB	1.584.. ^b	2.578..	3.020(5)	1.151(4)	1.837(3)	1.237(4)	1.5847(8)	1.2793(17)	2.5655(12)	0.442(4)	Here+ [8]
Regular	2	2	2.750(6)	1.273(2)	1.532(8)	1.4896(96)	1.995(3)	1.382(3)	1.9993(5)	0.761(13)	[5]
SITE	2 ^c	2.584..	3.232(6)	1.211(4)	1.870(4)	1.357(4)	1.9975(9)	1.3388(14)	2.5533(6)	0.676(5)	
EXGA	2.584.. ^d	2.321..	3.352(4)	1.312(3)	1.835(3)	1.581(3)	2.5895(6)	1.3915(8)	2.3000(2)	1.020(3)	
Regular	3	2	3.370(11)	1.407(2)	1.777(4)	1.783(5)	3.003(14)	1.442(12)	2.0043(3)	1.380(13)	Here
Regular	4 ^e	2	4	1.5	2	2	4	1.5	2	2	[23]

^a $\ln 5 / \ln 3$.^b $\ln 3 / \ln 2$.^cFractal lattice.^d $\ln 6 / \ln 2$, strongly anisotropic.^eAt upper critical dimension $d_c = 4$ [19], exponents take the mean-field value [23].

with $\tilde{a} = 1.020(2)$ and $\tilde{b} = 0.481(3)$ for *all* lattices, which results in

$$D = 4 - 0.658(5)\epsilon + 0.00962(13)\epsilon^2 + 0.00161(3)\epsilon^3 + \dots, \quad (12)$$

using $D(2 - \tau) = d_w = 2$ for the regular ones.

The form of Eq. (11) was obtained by first fitting τ against $(D/d)^\kappa$, which gives a κ deviating from -2 by less than 2%. The coefficients $\tilde{a}$ and $\tilde{b}$ are then fitted according to Eq. (11). Figure 3 compares that relation to results for lattices in *all* dimensions. In the same manner, a similar relation can be obtained for z and α :

$$\left(\frac{z}{d}\right)^{\frac{3}{2}}(\alpha - \tilde{a}) = \tilde{b}, \quad (13)$$

with $[\tilde{a} = 0.936(2)]$ and $[\tilde{b} = 0.3768(12)]$.

The above results suggest that the scaling in N is more suitable for fractals than the scaling in L . We suspect this is related to L not capturing the chemical distance, which is the distance particles need to travel on the lattice, whereas L is measured as a Euclidean distance. By using d_w , which is sensitive to the chemical distance, and considering the scaling

against N , which is a well-defined measure of the size for any lattice, we are able to determine the relations above.

IV. CONCLUSION

In conclusion, we studied the Abelian Manna model on various three-dimensional and fractal lattices with the aim to provide a complete picture about the model below the upper critical dimension. The results confirm the consistent and robust universal behavior of the Manna model across different, regular lattices, which allowed us to produce an ϵ expansion of avalanche exponents below the upper critical dimension. A relation between critical exponents and lattice dimension is observed which systematically reconciles integer dimensional with fractal lattices.

ACKNOWLEDGMENTS

The authors are indebted to Andy Thomas, Dan Moore, and Niall Adams for running the Supercomputer at night (SCAN) computing facility at the Department of Mathematics of Imperial College London. HNH thanks Lock Yue Chew for his support.

- [1] H. E. Stanley, *Rev. Mod. Phys.* **71**, S358 (1999).
- [2] P. Bak, C. Tang, and K. Wiesenfeld, *Phys. Rev. Lett.* **59**, 381 (1987).
- [3] I. Syözi, *Prog. Theor. Phys.* **6**, 306 (1951).
- [4] S. S. Manna, *J. Phys. A: Math. Gen.* **24**, L363 (1991).
- [5] H. N. Huynh, G. Pruessner, and L. Y. Chew, *J. Stat. Mech.* (2011) P09024.
- [6] D. Dhar, *Physica A* **263**, 4 (1999).
- [7] D. Dhar, arXiv:cond-mat/9909009.
- [8] H. N. Huynh, L. Y. Chew, and G. Pruessner, *Phys. Rev. E* **82**, 042103 (2010).
- [9] V. Privman, P. C. Hohenberg, and A. Aharony, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. L.

- Lebowitz (Academic Press, New York, 1991), Vol. 14, Chap. 1, pp. 1–134.
- [10] H. Nakanishi and K. Sneppen, *Phys. Rev. E* **55**, 4012 (1997).
- [11] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Harcourt College Publishers, New York, 1976).
- [12] H. N. Huynh and L. Y. Chew, *Fractals* **19**, 141 (2011).
- [13] P. S. Addison, *Fractals and Chaos: An Illustrated Course* (Institute of Physics, London, 1997).
- [14] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes: The Art of Scientific Computing*, 3rd ed. (Cambridge University Press, Cambridge, 2007).
- [15] A. Ben-Hur and O. Biham, *Phys. Rev. E* **53**, R1317 (1996).
- [16] S. Lübeck, *Phys. Rev. E* **61**, 204 (2000).

- [17] R. Pastor-Satorras and A. Vespignani, [Eur. Phys. J. B](#) **19**, 583 (2001).
- [18] M. Alava and M. A. Muñoz, [Phys. Rev. E](#) **65**, 026145 (2002).
- [19] S. Lübeck and P. C. Heger, [Phys. Rev. Lett.](#) **90**, 230601 (2003).
- [20] D. ben-Avraham and S. Havlin, *Diffusion and Reactions in Fractals and Disordered System* (Cambridge University Press, Cambridge, 2000).
- [21] K. Binder, [Phys. Rev. Lett.](#) **47**, 693 (1981).
- [22] S. Lübeck and P. C. Heger, [Phys. Rev. E](#) **68**, 056102 (2003).
- [23] S. Lübeck, [Int. J. Mod. Phys. B](#) **18**, 3977 (2004).
- [24] A. Chessa, H. E. Stanley, A. Vespignani, and S. Zapperi, [Phys. Rev. E](#) **59**, R12 (1999).
- [25] H. J. Jensen, K. Christensen, and H. C. Fogedby, [Phys. Rev. B](#) **40**, 7425 (1989).