Electronic structure and optical properties of conjugated molecules : SAC-CI study

Ehara, Masahiro; Lu, Yunpeng; Promkatkaew, Malinee; Hannongbua, Supa

2012

Ehara, M., Lu, Y. P., Promkatkaew, M., & Hannongbua, S. (2012). Electronic structure and optical properties of conjugated molecules: SAC-CI study. International Conference Of Computational Methods In Sciences And Engineering 2009: (ICCMSE 2009), 1504(7), pp.848-851.

<https://hdl.handle.net/10356/96006>

<https://doi.org/10.1063/1.4771827>

© 2012 American Institute of Physics. his paper was published in International Conference Of Computational Methods In Sciences And Engineering 2009: (ICCMSE 2009) and is made available as an electronic reprint (preprint) with permission of American Institute of Physics. The paper can be found at the following official DOI:
[<http://dx.doi.org/10.1063/1.4771827>]. One print or electronic copy may be made for personal use only. Systematic or multiple reproduction, distribution to multiple locations via electronic or other means, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper is prohibited and is subject to penalties under law.

Electronic structure and optical properties of conjugated molecules: SAC-CI study

Masahiro Ehara, Lu Yun-peng, Malinee Promkatkaew, and Supa Hannongbua

Citation: [AIP Conf. Proc.](#) **1504**, 848 (2012); doi: 10.1063/1.4771827

View online: <http://dx.doi.org/10.1063/1.4771827>

View Table of Contents: <http://proceedings.aip.org/dbt/dbt.jsp?KEY=APCPCS&Volume=1504&Issue=1>

Published by the [American Institute of Physics](#).

Additional information on AIP Conf. Proc.

Journal Homepage: <http://proceedings.aip.org/>

Journal Information: http://proceedings.aip.org/about/about_the_proceedings

Top downloads: http://proceedings.aip.org/dbt/most_downloaded.jsp?KEY=APCPCS

Information for Authors: http://proceedings.aip.org/authors/information_for_authors

ADVERTISEMENT

Submit Now

Explore AIP's new open-access journal

- **Article-level metrics
now available**
- **Join the conversation!
Rate & comment on articles**

Electronic Structure and Optical Properties of Conjugated Molecules: SAC-CI Study

Masahiro Ehara,^{†§} Lu Yun-peng,[¶] Malinee Promkatkaew,^{‡#} and Supa Hannongbua^{‡#}

[†]*Institute for Molecular Science, 38 Myodaiji, Okazaki, 444-8585, Japan*

[§]*JST-CREST, Sanboncho-5, Chiyoda-ku, Tokyo 102-0075, Japan*

[¶]*Division of Chemistry and Biochemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang Link, Singapore 637371*

[‡]*Department of Chemistry, Faculty of Science, Kasetsart University, Bangkok 10900, Thailand*

[#]*Center of Nanotechnology, Kasetsart University, Bangkok 10900, Thailand*

Abstract. Electronic structure and optical properties of some organic conjugated molecules, chelating hetero-atomic conjugated ligands and some derivatives of the UVB blocking cinnamates, have been investigated by the SAC-CI method. The absorption and emission spectra of these molecules have been reproduced in high accuracy. The excited-state geometry relaxation was interpreted based on the electrostatic force theory. The present work provides a useful basis for the theoretical design predicting the optical properties of the molecules.

Keywords: SAC-CI, optical properties, hetero-atomic conjugated molecule, UVB blocking

PACS: 31.15.ag, 31.15.bw, 31.15.vq

INTRODUCTION

Chelating heteroatomic conjugated molecules, such as pyridine, benzoxazole, and benzothiazole derivatives, are technologically important compounds. An increasing research focus for heteroatomic conjugate molecules is their application in the area of organic optoelectronic materials, such as photoconducting materials, liquid crystals, and fluorophores. Heteroatomic conjugate molecules have also been widely used as cyclometalated ligands in heavy metal complexes because of their strong chelating capability and their characteristic optical properties based on π -conjugation. Heavy metal complexes, such as Ir(III) and Pt(II) complexes, have been proven to be excellent phosphorescent dyes for the organic light-emitting diodes.

Cinnamates also have received much attention, as they are the most widely used UVB blocking compounds among the various cosmetic sunscreen agents. Cinnamates achieve UVB blocking from a $\pi\pi^*$ absorption followed by a *cis-trans* isomerization at the propenyl double bond in the S_1 state and a relaxation to the ground state involving nonradiative decay. As a UVB blocking compound, the optical properties, in particular the photoabsorption efficiency in the UVB energy region (290–320 nm) is an important factor. In order to achieve the favorable optical properties, molecular design using the variation of the substituents has been extensively performed.

The SAC-CI (symmetry-adapted cluster-configuration interaction) method¹ has been shown to be reliable and useful for investigating a wide variety of molecular spectroscopic properties in many successful applications.² In this work, we have systematically investigated the electronic structure and optical properties of some conjugated molecules. The first group belongs to the chelating hetero-atomic conjugate molecules that are widely used for developing iridium (III) complexes, including 2-phenyl pyridine (PPY), 2-phenyl quinoline (PQ), and bonzo[h]quinoline (BZQ).³ The second group is the methoxy substituted cinnamates, which show excellent UVB blocking performance.⁴ The target molecules are ortho-(1), meta-(2), and para-(3) monomethoxy substituted cinnamates, and 2,4,5-(4) and 2,4,6-(5) trimethoxy cinnamates in both *cis*- and *trans*- isomers. The difference in electronic transitions among these molecules and the effect of methoxy substitution at the ortho-, meta-, and para-positions were analyzed.

RESULTS AND DISCUSSION

chelating hetero-atomic conjugated molecules

We studied the electronic structure and optical properties of the 13 chelating hetero-atomic conjugated molecules,³ categorized into three groups: I) PPY, PQ, and BZQ; II) DPO, BO, BON, and OP; and III) BT BTP, α BSN, β BSN THP, and BTTH, based on the heteroatoms in their molecular structures. In this article, we present the results of the molecules in group I.

Each compound in this group contains one N atom in the conjugated ring. Figure 1 illustrates the calculated geometries of PPY, PQ and BZQ in the ground state (S_0) and the lowest $\pi\pi^*$ excited state (S_1) with the CC bond lengths. In the S_0 state of PPY, a non-planar structure is most stable with the torsion angle of 20.4° ; the energy difference between planar and non-planar structure is about 0.2 kcal/mol. The PQ also has non-planar geometry at the S_0 state with $\approx 21.0^\circ$. These molecules become planar from a non-planar structure by the S_0 - S_1 excitation. For PPY, the inter-ring CC bond distance is 1.488 Å in S_0 and the bond shrinks significantly to 1.407 Å in S_1 . The bonds in the pyridyl and phenyl rings also become elongated or shortened alternatively by the S_0 - S_1 excitation. The same trend was also found in PQ.

FIGURE 1. Geometries of PPY, PQ, and BZQ in the S_0 and S_1 states. Values are bond lengths of the S_0 (S_1) states.

The geometry relaxation in the excited state can be interpreted using the electrostatic force (ESF) theory. As an example, we discuss the geometric changes in the S_1 state of PPY. In ESF theory, geometry relaxation in the excited state is due to the force acting on nuclei caused by changes in electron distribution. The differences in the SAC-CI electron density between the S_0 and S_1 states in PPY and BZQ are shown in Figure 2. In PPY, the electron density in the C-C (C_2 - C_3 , C_7 - C_{12} , C_7 - C_8) and C_2 = N_1 bonds in the phenyl ring and pyridyl ring decrease, while the density of the inter-ring C_2 - C_7 bond significantly increases. The EC force along bonds including the C_2 = N_1 and C_2 - C_3 bonds decreases because of the decrement of electron density, whereas the EC force is enhanced along the C_2 - C_7 , C_6 - N_1 , and C_3 - C_4 bonds. Consequently, the former bond lengths increase, and the latter bond lengths decrease. This geometry change enhances the conjugation between two rings and the S_1 structure of PPY becomes planar.

FIGURE 2. The SAC-CI electron density difference between the S_0 and S_1 states of PPY and BZQ. The region in the blue range has a decrement of electron density and that in the red range has an increment of electron density when a molecule is excited.

The experimental and theoretical absorption spectra of PPY and PQ are compared in Figure 3. The calculated spectra satisfactorily reproduced the position of the observed absorption peaks for these molecules. Based on the SAC-CI calculations, we assign the broad peak around 4.49 eV (276 nm) as the second and third transitions calculated as 4.49 eV and 4.69 eV. The peak around 5.14 eV (242 nm) is assigned as the transition with a vertical excitation energy of 5.15 eV. These transitions are mainly due to $\pi\pi^*$ transitions. The SAC-CI calculation for PQ

shows that the strongest transition occurs at an energy of 5.22eV. The experimental spectrum gives a strong indication that the absorption is very strong above 4.76eV (260nm). The absorption energies of 3.87eV and 4.49eV in our calculations are consistent with the structures at 3.87eV (320nm) and 4.35eV (285nm) in the experimental absorption spectra.

FIGURE 3. Theoretical absorption spectra of (a) PPY and (b) PQ compared with the experimental spectra.

UVB blocking cinnamates

The electronic structure and optical properties of the *cis*- and *trans*-isomers of methoxy substituted cinnamates have been investigated using the SAC-CI method.⁴ The molecules are ortho-(1), meta-(2), and para-(3) monomethoxy substituted cinnamates, and 2,4,5-(4) and 2,4,6-(5) trimethoxy cinnamates as shown in Figure 4. In this article, we focus on the results of the *trans* isomers of the compounds 1, 2, and 3. The trend of *cis*-isomers are similar to those of *trans*-isomers and the results of tri methoxy derivatives are referred to ref. [4].

FIGURE 4. Chemical structures of 2-ethylhexyl-cinnamate derivatives and the calculated model compounds of *trans*-isomers.

The S_0 and S_1 state geometries of the *trans*-isomers of ortho-(1), meta-(2), and para-(3) methoxy substituted compounds were shown in Figure 5. All the compounds, except for the *cis*-5 compound, had local minima in the coplanar structure of the *cis*- and *trans*-isomers in both the S_0 and S_1 states due to π -conjugation. In the S_0 state, the rotational energy barrier of the methoxy group is very low, although the planar structure is the most stable structure. The global minimum of the S_1 state is in the form of a nonplanar structure, but local minimum exist in the planar structure. The emission was observed from this local minimum in the planar structure, and in particular, a strong emission was observed for compound 2.⁵

FIGURE 5. CC and CO bond length of the S_0 and S_1 states (in parentheses) of *trans*-1, *trans*-2, and *trans*-3.

It was found that the C_1-C_2 , C_3-C_8 , $C_{10}-C_{12}$, $C_{12}-O_{14}$, and $O_{14}-C_{23}$ bond lengths decrease and the length of the other bonds increases. In the S_0 state, the carbon-carbon bond alternation exists for both the single and double bonds, but this bond alternation relaxes in the S_1 state. Since the excitation is relatively localized in the central unit, the prominent changes occur in the vinylenic unit. For example, in the *trans*-1 compound, the change in bond length is

$\Delta r = +0.048, -0.060$, and $+0.057$ Å for C₂–C₃, C₃–C₈, and C₈–C₁₀, respectively, while the other changes in bond length are within 0.022 Å.

The absorption spectra of the methoxy substituted cinnamates were calculated. The SAC-CI excitation spectra for the monomethoxy substituted molecules are compared with the experimental absorption spectra in Figure 6. For the monomethoxy substituted compounds, the excited states that contribute to the absorption have a different character that depends on the methoxy substituted position. In the case of the ortho-(*trans*-1) and meta-(*trans*-2) compounds, the S₃ state has the highest transition probability, and is characterized as being the transition from the next HOMO to LUMO (nH–L) transition, whereas, the highest transition possibility of the para-(*trans*-3) compound was calculated for the S₀→S₁ transition, and the excitation character is a HOMO–LUMO (H–L) transition. The agreement with the experimental values was satisfactory; the deviations were within about 10 nm.

FIGURE 6. SAC-CI absorption spectra of *trans*-1, 2 and 3 compounds compared with the experimental spectra.⁵

The emission energies of these molecules have also been calculated using the SAC-CI method. The stable geometries of the S₁ state were located using CIS followed by the SAC-CI calculations of the vertical emission energies. The calculated emission energies are in good agreement with the experimental data as shown in Table 1.

TABLE 1. Emission energy (ΔE), emission wavelength ($\lambda_{\max}$), oscillator strength (f) calculated using the SAC-CI method. Experimental values are cited from Ref. 5.

Mol.	SAC-CI			Expt.
	ΔE (eV)	$\lambda_{\max}$ (nm)	f	$\lambda_{\max}$ (nm) in hexane
<i>trans</i> -1	3.37	367	0.44	359
<i>trans</i> -2	3.62	343	0.29	350
<i>trans</i> -3	3.52	352	0.61	351

SUMMARY

Electronic structure and optical properties of chelating hetero-atomic conjugated ligands and some derivatives of the UVB blocking cinnamates have been investigated by the SAC-CI method. The absorption and emission spectra of these molecules have been reproduced in high accuracy. The present work provides a useful basis for the theoretical design predicting the optical properties of the molecules.

ACKNOWLEDGMENTS

This was supported by the grant from the JST-CREST, Scientific Research in Priority Areas “Molecular Theory for Real Systems” from the MEXT of Japan, the Next Generation Supercomputing Project, and the Molecular-Based New Computational Science Program, NINS.

REFERENCES

1. H. Nakatsuji, *Chem. Phys. Lett.* **59**, 362 (1978); H. Nakatsuji, *Chem. Phys. Lett.* **67**, 329 (1979).
2. M. Ehara, J. Hasegawa and H. Nakatsuji, *In Theory and Applications of Computational Chemistry* edited by C.E. Dykstra, G. Frenking, K.S. Kim, G.E. Scuseria, 1099 (Elsevier, Oxford, 2005).
3. L. Yun-peng and M. Ehara, *Theor. Chem. Acc.* in press.
4. T. M. Karpkird, S. Pattanaargson, and B. Albinsson, *Photochem. Photobiol. Sci.* in press.
5. M. Promkatkaew, S. Suramit, T. M. Karpkird, S. Namuangruk, M. Ehara, S. Hannongbua, submitted.