Tf 2 NH-catalyzed amide synthesis from vinyl azides and alcohols

Zhang, Feng-Lian; Zhu, Xu; Chiba, Shunsuke

2015

Zhang, F.-L., Zhu, X., & Chiba, S. (2015). Tf 2 NH-catalyzed amide synthesis from vinyl azides and alcohols. *Organic Letters*, 17(12), 3138-3141.

<https://hdl.handle.net/10356/98598>

<https://doi.org/10.1021/acs.orglett.5b01458>

© 2015 American Chemical Society. This is the author created version of a work that has been peer reviewed and accepted for publication by *Organic Letters*, American Chemical Society. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [<http://dx.doi.org/10.1021/acs.orglett.5b01458>].

Downloaded on 26 Apr 2023 20:36:23 SGT

Tf₂NH-Catalyzed Amide Synthesis from Vinyl Azides and Alcohols

Feng-Lian Zhang,[†] Xu Zhu,[†] and Shunsuke Chiba*

Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore

Supporting Information Placeholder

ABSTRACT: Triflimide (Tf₂NH) specifically catalyzed reactions of alcohols and vinyl azides, enabling efficient construction of amides with C-C bond formation through nucleophilic attack of vinyl azides onto the putative carbocation intermediates derived from alcohols.

The amide functionality is ubiquitous in various functional molecules of biological, pharmaceutical/medicinal, and materials importance.¹ It also serves as a versatile synthon for a variety of molecular transformations, especially for synthesis of nitrogen-containing molecules. Therefore, installation of the amide moiety onto organic frameworks is one of the most valuable processes in synthetic chemistry.² Nucleophilic attack of amide enolates onto suitable carbon electrophiles could be one of the most step-economical ways to introduce an acetamide unit through C-C bond formation (Scheme 1). Recent advancement in this field involves Kobayashi's direct aldol/Mannich reactions of (*N*-Boc)acylanisidides with barium phenoxide as a catalyst (Scheme 1-a)³ as well as direct Mannich reactions of unactivated tertiary amides catalyzed by trialkylsilyl triflate (Scheme 1-b).⁴ Scheidt recently reported base-mediated amide enolate addition to acylsilanes, which subsequently induces the Brook rearrangement to form the

Scheme 1. Reactions of Amide Enolates

(a) Direct aldol reactions of (*N*-Boc)acylanisidides by Ba(OAr)₂

(b) Direct Mannich reactions of tertiary amides by R₃SiOTf

(c) LDA-mediated addition of amide enolates to acylsilanes

corresponding carbanions, enabling efficient construction of tertiary β-hydroxy amides with alkyl halides (Scheme 1-c).⁵

As an alternative amide enolate equivalent, we have recently utilized vinyl azides. Nucleophilic attack of vinyl azides onto several carbon electrophiles such as *N*-Ts imines, aldehydes, and carbocations derived from alcohols in the presence of BF₃•OEt₂ forms iminodiazonium ions **A** with C-C bond formation (Scheme 2).^{6,7} Subsequent substituent migration generates nitrilium ions **B**, which are finally hydrolyzed to give amides. However, the process needs to use excess amounts (2 equivalents) of BF₃•OEt₂, that causes another procedure requirement of slow addition of vinyl azides to the solution to prevent decomposition of vinyl azides.⁸ Therefore, we have strived to develop a catalytic process of the acetamide installation using vinyl azides, that could be implemented under

Scheme 2. Amide Synthesis by Nucleophilic Attack of Vinyl Azides

• with *N*-Ts imines (Mannich type)

• with alcohols as a carbocation precursor

milder reaction conditions with easy and simple operation. Herein, we report realization of the catalytic acetamide incorporation with vinyl azides using triflimide ($\text{ Tf}_2\text{NH}$) as the catalyst for the reactions with a series of alcohols as a carbocation precursor.⁹

We became interested in construction of 3-indoloxindole skeletons via the reaction of 3-hydroxy-3-indoloxindoles with vinyl azides, as these scaffolds are prevalent in biologically active alkaloids.¹⁰ The recent seminal reports by Gong, Guo/Peng, and Ma demonstrated that 3-hydroxy-3-indoloxindoles could be activated by Brønsted acid catalysts to generate the corresponding carbocation equivalents, which are trapped by ketone enolate-type carbon nucleophiles.¹¹ Inspired by these findings, we surmised if direct installation of acetamide unit using vinyl azides is enabled by Brønsted acid catalysts. We commenced our study to screen Brønsted acids in the reactions of 3-hydroxy-3-indoloxindole **1a** and vinyl azide **2a** to target amide **3aa** (Table 1). The reactions with $\text{TsOH}\cdot\text{H}_2\text{O}$, BINOL-linked phosphoric acid, and $(\text{PhSO}_2)_2\text{NH}$ as the catalyst (10 mol %) resulted in the C-C bond formation but afforded ketone **4aa** in moderate yields as the major product (runs 1-3), in which almost no formation of desired amide **3aa** was observed.¹² The results indicated that these acids are incapable of enabling migration of the phenyl group from the iminodiazonium intermediate. On the other hand, use of $\text{CF}_3\text{SO}_3\text{H}$ rendered the process very rapid, consuming **1a** within 5 min to give 1:1 mixture of amide **3aa** and ketone **4aa** (run 4). We envisaged that strength of the acidity¹³ might be the key to enable selective formation of amide **3aa**. Further screening revealed that the reaction with triflimide ($\text{ Tf}_2\text{NH}$)¹⁴ (10 mol %) provided much better selectivity to afford amide **3aa** in 75% yield and ketone **4aa** in 15% yield (run 5). Slightly higher reaction temperature enhanced the selectivity further (runs 6 and 7), giving amide **3aa** as a sole product (86% yield) when the reaction was conducted in 1,2-dichloroethane (DCE) at 40 °C (run 7).

Table 1. Optimization of the reaction conditions^a

run	acid catalysts	conditions	time	3aa [%] ^b	4aa [%] ^b
1	$\text{TsOH}\cdot\text{H}_2\text{O}$	CH_2Cl_2 , rt	0.5 h	0	51
2	BINOL- $\text{P}(\text{O})\text{OH}^d$	CH_2Cl_2 , rt	2 h	0	43
3	$(\text{PhSO}_2)_2\text{NH}$	CH_2Cl_2 , rt	14 h	4	28
4	$\text{CF}_3\text{SO}_3\text{H}$	CH_2Cl_2 , rt	5 min	44	41
5	$(\text{CF}_3\text{SO}_2)_2\text{NH}$	CH_2Cl_2 , rt	20 min	75	15
6	$(\text{CF}_3\text{SO}_2)_2\text{NH}$	CH_2Cl_2 , reflux	10 min	80 (78) ^c	trace
7	$(\text{CF}_3\text{SO}_2)_2\text{NH}$	DCE, 40 °C	5 min	88 (86) ^c	0

^a All the reactions were carried out using 0.3 mmol of **1a** in the presence of 10 mol % of acid catalysts in solvent (0.1 M). ^b ¹H NMR yields. ^c Isolated yields. ^d ($\pm$)-1,1'-binaphthyl-2,2'-diyl hydrogenphosphate.

With the optimized reaction conditions in hand, we next investigated the substrate scope (Scheme 3). By varying the substituent R^1 of vinyl azides **2**, a series of aryl groups could be incorporated in the amide formation generally in good yields (for **3ab-3af**), while the reactions with methoxyphenyl-substituted vinyl azides **2b** and **2c** needed 30 mol % of $\text{ Tf}_2\text{NH}$ to complete the processes. The reaction with α -alkyl-substituted vinyl azide **2g** also required 30 mol % of $\text{ Tf}_2\text{NH}$ to give amide product **3ag** in 54% yield. In this case, the putative iminodiazonium ion possesses two different secondary alkyl substituents, while only formation of **3ag** via migration of the phenethyl group was observed probably because of bulkiness of the 3-indoloxindole moiety in another secondary alkyl group prevents its migration. The reaction with β -methyl substituted vinyl azide **1h** ($\text{R}^1 = \text{Ph}$, $\text{R}^2 = \text{Me}$) proceeded in diastereoselective manner, giving **3ah** in 78% yield with 3.3:1 dr ratio.¹⁵ As the substituents R^3 and R^4 on the nitrogens of 3-hydroxy-3-indoloxindoles **1**, not only benzyl but also phenyl (for **3ba**), hydrogen (for **3ca**), and methyl groups (for **3db**) could be installed. Amide **3db** is the key intermediate in synthesis of folicanthine by Gong,^{11c} in which the 4-methoxy-

Scheme 3. Substrate Scope^{a,b}

^a Unless otherwise noted, the reactions were conducted using alcohols **1** (0.20–0.34 mmol) with vinyl azides **2** (1.2 equiv) in the presence of Tf_2NH (10 mol %) in DCE (3 mL) under a N_2 atmosphere. See the Supporting Information for more details. ^b Isolated yields were recorded above. ^c 30 mol % of Tf_2NH was used. ^d 20 mol % of Tf_2NH was used. ^e Structure of the major isomer was described, which was determined by X-ray crystallographic analysis. See the Supporting Information.

phenylamide moiety was constructed via a stepwise sequence including: 1) synthesis of the corresponding ketone by BINOL-linked phosphoric acid-catalyzed reaction of 3-hydroxy-3-indoloxindole **1d** with enamide; 2) conversion of the resulting ketone into oxime; 3) the Beckmann rearrangement. The present strategy using vinyl azide **2b** provides an operationally straightforward installation of acetamide unit onto **1d**. A variety of substituents could be installed as R^5 and R^6 on the framework of 3-hydroxy-3-indoloxindoles **1** (for **3ea–3ja**).

Conversion of amides **3** into pyrroloindolinones **5** was demonstrated by careful treatment of **3** with LiAlH_4 in THF (Scheme 4).¹⁶ For synthesis of **5ac** having an allyl group on the nitrogen, exchange from 2-methoxyphenyl to allyl amide was first conducted (see the Supporting Information), that was followed by the reductive ring-closure.

Scheme 4. Construction of Pyrroloindolinone Structures^{a,b}

^a All the reactions were conducted using 0.08–0.1 mmol of amides **3** in THF (3 mL) under a N_2 atmosphere. See the Supporting Information for more details. ^b Isolated yields were recorded above. ^c **5ac** was synthesized from **3ac** via amide exchange from 2-methoxyphenyl to allyl via stepwise sequence followed by LiAlH_4 reduction. See the Supporting Information. PMP = 4-methoxyphenyl.

The present catalytic amide synthesis with vinyl azides could be successfully applied for other types of benzylic (for **6a–6c**), allylic (for **6d**), and propargylic alcohols (for **6e**), providing the corresponding amide products **7** of synthetic and medicinal relevance (Scheme 5). For example, 2-(9H-xanthenyl)acetamide **7c** functions as an acyl-CoA cholesterol acyltransferase (ACAT) inhibitor for treatment of atherosclerosis.¹⁷ β -Alkynyl-carboxamide **7e** is severed as a useful synthon for construction of pyrrolidone scaffolds.¹⁸

Scheme 5. Amide Synthesis from Alcohols **6** with Vinyl Azide **2a**.^{a,b}

^a All the reactions were conducted using alcohols **6** (0.30–0.32 mmol) with vinyl azide **2a** (1.2 equiv) and Tf₂NH (10 mol %) in DCE (3 mL) under a N₂ atmosphere. See the Supporting Information for more details. ^b Isolated yields were recorded above. ^c Alcohol **6d** was recovered in 20% yield.

In summary, we have developed Tf₂NH-catalyzed reactions of alcohols with vinyl azides, enabling a straightforward access of synthetically and medicinally useful amide derivatives. Further upgrading of this catalytic method to the asymmetric variant as well as application of Tf₂NH to other types of catalytic transformations is now under way in our laboratory.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures, characterization of new compounds, and CIF data of **3ah**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail: shunsuke@ntu.edu.sg

Notes

The authors declare no competing financial interest.

† F.-L. Z. and X. Z. contributed equally to this work.

ACKNOWLEDGMENT

This work was supported by funding from Nanyang Technological University and Singapore Ministry of Education and Singapore Ministry of Education (Academic Research Fund Tier 1: RG4/13).

REFERENCES

- (1) For a review, see: Arthur, G.; Breneman, C. M.; Liebman, J. F.; *The Amide Linkage: Selected Structural Aspects in Chemistry, Biochemistry, and Materials Science*, 12; John Wiley & Sons Inc.: New York, 2000.
- (2) For reviews, see: (a) Pattabiraman, V. J.; Bode, J. W. *Nature* **2011**, *480*, 471. (b) Allen, C. L. A.; Williams, J. M. J. *Chem. Soc. Rev.* **2011**, *40*, 3405. (c) Chen, C.; Hong, S. H. *Org. Biomol. Chem.* **2011**, *9*, 20. (d) Valeur, E.; Bradley, M. *Chem. Soc. Rev.* **2009**, *38*, 606. (e) Bode, J. W. *Curr. Opin. Drug Discovery Dev.* **2006**, *9*, 765. (f) Montalbetti, C. A. G. N.; Flaque, V. *Tetrahedron* **2005**, *61*, 10827. (g) Han, S.-Y.; Kim, Y.-A. *Tetrahedron* **2004**, *60*, 2447. (h) Humphrey, J. M.; Chamberlin, A. R. *Chem. Rev.* **1997**, *97*, 2243.
- (3) (a) Saito, S.; Tsubogo, T.; Kobayashi, S. *Chem. Commun.* **2007**, 1236. (b) Saito, S.; Kobayashi, S. *J. Am. Chem. Soc.* **2006**, *128*, 8704.
- (4) Kobayashi, S.; Kiyohara, H.; Yamaguchi, M. *J. Am. Chem. Soc.* **2011**, *133*, 708.
- (5) (a) Lettan II, R. B.; Galliford, C. V.; Woodward, C. C.; Scheidt, K. A. *J. Am. Chem. Soc.* **2009**, *131*, 8805. (b) Lettan II, R. B.; Reynolds, T. E.; Galliford, C. V.; Scheidt, K. A. *J. Am. Chem. Soc.* **2006**, *128*, 15566.
- (6) Zhang, F.-L.; Wang, Y.-F.; Lonca, G. H.; Zhu, X.; Chiba, S. *Angew. Chem. Int. Ed.* **2014**, *53*, 4390.
- (7) For azaheterocycle synthesis triggered by nucleophilic attack of vinyl azides onto *N*-unsaturated aldimines, see: Zhu, X.; Wang, Y.-F.; Zhang, F.-L.; Chiba, S. *Chem.-Asian J.* **2014**, *9*, 2458.
- (8) For reports on the reactions of vinyl azides under acidic conditions, see: (a) Hassner, A.; Ferdinandi, E. S.; Isbister, R. J. *J. Am. Chem. Soc.* **1970**, *92*, 1672. (b) Moore, H. W.; Shelden, H. R.; Weyler Jr., W. *Tetrahedron Lett.* **1969**, *10*, 1243.
- (9) For a review on functionalization of carbocations, see: Naredla, R. R.; Klumpp, D. A. *Chem. Rev.* **2013**, *113*, 6905.
- (10) For reviews, see: (a) Repka, L. M.; Reisman, S. E. *J. Org. Chem.* **2013**, *78*, 12314. (b) Ruiz-Sanchis, P.; Savina, S. A.; Albericio, F.; Alvarez, M. *Chem.-Eur. J.* **2011**, *17*, 1388. (c) Schmidt, M. A.; Movassaghi, M. *Synlett*, **2008**, 313. (d) Steven, A.; Overman, L. E. *Angew. Chem. Int. Ed.* **2007**, *46*, 5488.
- (11) (a) Tang, X.-D.; Li, S.; Guo, R.; Nie, J.; Ma, J.-A. *Org. Lett.* **2015**, *17*, 1389. (b) Song, J.; Guo, C.; Adele, A.; Yin, H.; Gong, L.-Z. *Chem.-Eur. J.* **2013**, *19*, 3319. (c) Zhang, Y.; Wang, S.-Y.; Xu, X.-P.; Jiang, R.; Ji, S.-J. *Org. Biomol. Chem.* **2013**, *11*, 1933. (d) Song, L.; Guo, Q.-X.; Li, X.-C.; Tian, J.; Peng, Y.-G. *Angew. Chem. Int. Ed.* **2012**, *51*, 1899. (e) Guo, C.; Song, J.; Huang, J.-Z.; Chen, P.-H.; Luo, S.-W.; Gong, L.-Z. *Angew. Chem. Int. Ed.* **2012**, *51*, 1046.
- (12) The reactions in entries 1–4 formed acetophenone and acetamide by the reactions of vinyl azide **2a** with the acid catalysts. See Table S1 in the Supporting Information for more details.
- (13) Koppel, I. A.; Taft, R. W.; Anvia, F.; Zhu, S.-Z.; Hu, L.-Q.; Sung, K.-S.; DesMarteau, D. D.; Yagupolskii, L. M.; Yagupolskii, Y. L.; Ignat'ev, N. V.; Kondratenko, N. V.; Volkonskii, A. Y.; Vlazov, V. M.; Notario, R.; Maria, P.-C. *J. Am. Chem. Soc.* **1994**, *116*, 3047.
- (14) For reviews on synthetic application of Tf₂NH, see: (a) Sun, J. *Triflimide in e-EROS: Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (DOI: 10.1002/047084289X.rm01222). (b) Takasu, K.; *Synlett* **2009**, 1905.
- (15) Origin of the diastereoselectivity was discussed in the Supporting Information.
- (16) (a) Ma, S.; Han, X.; Krishnan, S.; Virgil, S. C.; Stoltz, B. M. *Angew. Chem. Int. Ed.* **2009**, *48*, 8037. (b) Fang, C.-L.; Horne, S.; Taylor, N.; Rodrigo, R. *J. Am. Chem. Soc.* **1994**, *116*, 9480.
- (17) Yoshida, A.; Oda, K.; Kassai, T.; Koga, T.; Hasegawa, K. United States Patent, US005563169A, 1996.
- (18) Tellitu, I.; Serna, S.; Dominguez, E. *Arkivoc*, **2010**, *iii*, 7.