

This document is downloaded from DR-NTU, Nanyang Technological University Library, Singapore.

Title	Functional interplay among the flavivirus NS3 protease, helicase, and cofactors(Article)
Author(s)	Li, Kuohan; Phoo, Wint Wint; Luo, Dahai
Citation	Li, K., Phoo, W. W., & Luo, D. (2014). Functional interplay among the flavivirus NS3 protease, helicase, and cofactors. <i>Virologica Sinica</i> , 29(2), 74-85.
Date	2014
URL	http://hdl.handle.net/10220/19288
Rights	© 2014 WIV, CAS and Springer-Verlag Berlin Heidelberg. This is the author created version of a work that has been peer reviewed and accepted for publication by <i>Virologica Sinica</i> , WIV, CAS and Springer-Verlag Berlin Heidelberg. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [http://dx.doi.org/10.1007/s12250-014-3438-6].

DOI: 10.1007/s12250-014-3438-#

Received: 21 January 2014, Accepted: ## March 2014, Published online: ## March 2014

✉Correspondence.

Phone: 65-65869705, E-mail: luodahai@ntu.edu.sg

Running title: Flavivirus NS3 protein structure and function

Research Article

Functional interplay among the flavivirus NS3 protease, helicase, and cofactors

Kuohan Li, Wint Wint Phoo, Dahai Luo✉

Lee Kong Chian School of Medicine, Nanyang Technological University, 61 Biopolis Drive, Proteos Building, #07-03, Singapore 138673, Republic of Singapore

Abbreviations: DENV4, dengue virus serotype 4; NS3pro, nonstructural protein 3 protease; NS3hel, nonstructural protein 3 helicase protein; RdRP, RNA-dependent RNA polymerase

Flaviviruses are positive-sense RNA viruses, and many are important human pathogens. Nonstructural protein 2B and 3 of the flaviviruses (NS2BNS3) form an endoplasmic reticulum (ER) membrane-associated hetero-dimeric complex through the NS2B transmembrane region. The NS2BNS3 complex is multifunctional. The N-terminal region of NS3, and its cofactor NS2B fold into a protease that is responsible for viral polyprotein processing, and the C-terminal domain of NS3 possesses NTPase/RNA helicase activities and is involved in viral RNA replication and virus particle formation. In addition, NS2BNS3 complex has also been shown to modulate viral pathogenesis and the host immune response. Because of the essential functions that the NS2BNS3 complex plays in the flavivirus life cycle, it is an attractive target for antiviral development. This review focuses on the

recent biochemical and structural advances of NS2BNS3 and provides a brief update on the current status of drug development targeting of this viral protein complex.

KEYWORDS crystal structures; antiviral drug target; serine protease; RNA helicase

MANY FLAVIVIRUSES ARE IMPORTANT HUMAN PATHOGENS

Emerging and re-emerging viral infections threaten hundreds of thousands of human lives every year and cause serious global public health problems. Several well-known viral pathogens include Dengue virus (DENV), West Nile virus (WNV), Yellow fever virus (YFV), Japanese encephalitis virus (JEV) and Tick-borne encephalitis virus (TBE). They are classified into the flavivirus genus in the *flaviviridae* family (Lindenbach B D, et al., 2007). Some of these virus infections have become increasingly severe, frequently causing epidemics across the world. Taking dengue as an example, there are over 100 million cases of dengue fever occurring in the tropical and sub-tropical regions of the world annually. A number of the cases develop life-threatening conditions, such as dengue hemorrhagic fever (DHF) or dengue shock syndrome (DSS) (WHO, 2009). Despite global efforts to develop therapeutics, there are no effective antiviral drugs approved against any of these viral infections (Lim S P, et al., 2013). Vaccine development against DENV has also been slow (Sabchareon A, et al., 2012), although there are vaccines available against YFV (Verma R, et al., 2013), JEV (Yun S I, et al., 2013) and TBEV (Rendi-Wagner P, 2008). To facilitate the development of diagnostics, antiviral drugs, and effective treatments, basic research aims to better understand the mechanisms of viral replication, pathogenesis and virulence factors, transmission patterns, and the varying outcomes of host immune response.

Flavivirus are positive-sense, single-stranded RNA viruses (Lindenbach B D, et al., 2007). The virus particle has an icosahedral shape and is enveloped, with a diameter about 40-60 nm (Lindenbach B D, et al., 2007). The flavivirus genome is about 11 kilobases and has a 5' end cap similar to that of the cellular mRNA but no 3' end polyadenylation tail (Figure 1A) (Lindenbach B D, et al., 2007). There are also long untranslated regions at both ends, which play important roles in virus replication, viral protein translation, and virus particle assembly (Alvarez D E, et al., 2006; Filomatori C V, et al., 2006; Markoff L, 2003; Polacek C, et al., 2009; Villordo S M, et al., 2008; Yu L, et al., 2008). Viral protein translation results in a single polypeptide precursor spanning across the endoplasmic reticulum (ER)-derived membrane, which is processed into three structural proteins, namely the capsid protein (C), envelope protein (E), and

membrane protein (M) and seven non-structural proteins, NS1, NS2A, NS2B, NS3, NS4A NS4B, and NS5 (Figure 1A) (Lindenbach B D, et al., 2007; Mackenzie J, 2005). Both host ER-derived proteases and viral NS2BNS3 protease specifically cleave the precursor to release the individual viral proteins (Figure 1A) (Lindenbach B D, et al., 2007). The C-terminal region of NS3 is a NTPase/RNA helicase (Xu T, et al., 2005). NS5 possesses N-terminal RNA methyl-transferase (Egloff M P, et al., 2002; Ray D, et al., 2006) and C-terminal RNA-dependent RNA polymerase (RdRP) activities (Lescar J, et al., 2008; Lindenbach B D, et al., 2007). Together with the viral RNA, viral cofactors, and host cofactors, NS3 and NS5 form the virus replication complex on the intracellular membrane to amplify the viral genome (Lindenbach B D, et al., 2007; Murray C L, et al., 2008; Paul D, et al., 2013). Therefore, functional inhibition of the viral nonstructural proteins and disruption of the viral replication complex form the rationale behind anti-flavivirus drug development (Bollati M, et al., 2010; Lescar J, et al., 2008; Lim S P, et al., 2013; Malet H, et al., 2008; Sampath A, et al., 2009).

MULTIFUNCTIONAL ENZYME NS3

Flavivirus NS3 (69 kDa) is the second-largest viral protein (behind NS5) in the flavivirus genome and plays essential roles in the viral life cycle (Figure 1B). NS3 has two functional domains: the N-terminal protease cleaves the viral polyprotein precursor to release individual nonstructural proteins, and the C-terminal's nucleoside-triphosphatase (NTPase)-dependent RNA helicase activity is involved in genome replication and viral RNA synthesis (Lescar J, et al., 2008; Lindenbach B D, et al., 2007; Luo D, et al., 2012). In addition, a recent report described the ATP-independent RNA annealing activity of the DENV NS3 helicase (Gebhard L G, et al., 2012). The two domains are connected by a flexible linker that is about 10 amino acids (aa) in length (Figure 1C) (Luo D, et al., 2010). It is not known why a single gene encodes two distinct domains, but some studies have shown that these two domains are functionally coupled in DENV (Luo D, et al., 2012; Luo D, et al., 2010; Xu T, et al., 2006; Yon C, et al., 2005). Similarly, HCV NS3 is also a protease-ATPase/helicase fusion protein (Lindenbach B D, et al., 2007) and its activity has also been shown to be regulated by the protease domain (Frick D N, et al., 2004). Recently, several anti-HCV drugs targeting the NS3 protease received U.S. Food & Drug Administration (FDA) approval, and a few promising candidates are undergoing intensive clinical trials (Salam K A, et al., 2013). This will further stimulate the development of antiviral drugs that target the flavivirus NS3 protein. Therefore, we

present the recent structural and functional studies of this multifunctional viral enzyme from the flavivirus genus and compare it with the NS3 protein of hepatitis C virus belonging to hepacivirus genus.

NS2B-NS3 PROTEASE (NS2BNS3PRO)

The N-terminal domain of NS3 (aa 1–169) is a chymotrypsin-like serine protease and is able to perform both cis- and trans- viral polyprotein cleavage (Chambers T J, et al., 1990; Gorbalenya A E, et al., 1989; Li J, et al., 2005). To function as an active enzyme, NS3 protease requires the cofactor NS2B (Falgout B, et al., 1991; Jan L R, et al., 1995; Yusof R, et al., 2000; Zhang L, et al., 1992). The NS2B protein is a 14-kDa integral membrane protein that contains three domains: two putative hydrophobic transmembrane segments at both N and C termini and a central region (aa 49–96, 47 aa) that is the essential cofactor of the NS3 protease (Figure 1B) (Clum S, et al., 1997). It is known that the flavivirus NS3 protease domain is not very soluble or catalytically active *in vitro*. Full-length NS3 is not very soluble either, suggesting that the NS3 protease may not fold properly without the NS2B cofactor (Keller T H, et al., 2006; Xu T, et al., 2005). Indeed, Leung et al. engineered a fusion gene of the NS2B cofactor region that was connected to the NS3 protease domain via a flexible nine aa (Gly4-Ser-Gly4) linker; and this fusion protein became soluble when expressed in bacteria and was catalytically active *in vitro* (Leung D, et al., 2001). This idea of protein engineering greatly facilitated subsequent research on the NS2B-NS3 protease domain and the full-length proteins and promoted advances in NS3 structural biology (Table 1) (Arakaki T L, et al., 2002; Erbel P, et al., 2006; Lescar J, et al., 2008; Luo D, et al., 2008; Nall T A, et al., 2004; Robin G, et al., 2009). Furthermore, the central cofactor region of NS2B was revealed to have dual functions. The presence of the N-terminus (aa 49–67) from the NS2B cofactor region renders NS3 protease soluble but enzymatically inactive, which indicates that NS2B is responsible for the correct folding of the NS3 protease domain (Erbel P, et al., 2006; Luo D, et al., 2008). The C-terminal cofactor region of NS2B (aa 68–96) forms a conserved β -turn hairpin and binds to the hydrophobic S2 and S3 pockets in the substrate binding site of NS3. This peptide of NS2B directly interacts with the bound substrate or substrate-based-inhibitor, supporting the catalytic role of NS2B (Figure 2AB) (Erbel P, et al., 2006; Noble C G, et al., 2012; Robin G, et al., 2009). Thus, the NS2B cofactor is able to regulate NS3pro activity by stabilizing the correct folding of the core structure and by directly participating in the catalysis of substrate cleavage (Chappell K J, et al., 2008; Erbel P, et al., 2006; Luo D, et al., 2012; Robin G, et al., 2009). This is in contrast to HCV NS3pro, which only requires a short sequence from NS4A (~11 aa) to form the

active protease. Furthermore, the NS4A cofactor has been shown to play a structural role but does not directly participate in substrate recognition or catalysis directly (Kim J L, et al., 1996; Tomei L, et al., 1996; Urbani A, et al., 1998).

The protease catalytic triad (His51, Asp75, Ser135) of NS3pro is found in the central cleft (Figure 2AB) (Erbel P, et al., 2006; Noble C G, et al., 2012; Robin G, et al., 2009). In general, the protease recognizes positively charged Arg/Lys at the P1 and P2 positions followed by a small or polar amino acid at P1', although there are exceptions (e.g., Glu is found at the P2 position of the NS2B-NS3 junction) (Gouvea I E, et al., 2007; Li J, et al., 2005; Shiryaev S A, et al., 2007). Fluorogenic peptide substrates are usually used to study protease activities and for screening purposes (Li J, et al., 2005; Niyomrattanakit P, et al., 2006; Yusof R, et al., 2000). Structures of the NS2BNS3pro in complex with substrate-derived peptide inhibitors or aprotinin (bovine pancreatic trypsin inhibitor, BPTI) have revealed the substrate specificity and the catalysis mechanism (Erbel P, et al., 2006; Noble C G, et al., 2012; Robin G, et al., 2009).

NS3 protease inhibitors are currently designed by interfering either with the substrate binding cleft or with the interaction between the NS2B and NS3 protease domain (Lescar J, et al., 2008; Lim S P, et al., 2013). Despite structural and biochemical information available on the NS2BNS3pro substrate binding pocket, no compound has progressed to the preclinical stage to date. Researchers are still facing multiple challenges in identifying effective drugs that target NS2BNS3pro. The substrate binding pocket is shallow and solvent exposed, rendering interactions between mimics and enzyme labile. Moreover, dibasic residues at the P1 and P2 positions hinder peptidomimetic permeability and stability. More recently, researchers began to study the structure and function of the native-state-like NS2BNS3pro in the membrane context, which could lead to better screening platforms for protease inhibitors (Choksupmanee O, et al., 2012; Huang Q, et al., 2013).

NS3 NTPASE/RNA HELICASE (NS3HEL)

The C-terminal domain of the multifunctional NS3 protein (aa 180–618) was identified as a member of the superfamily 2 (SF2) helicase (Fairman-Williams M E, et al., 2010; Gorbalenya A E, et al., 1993). The overall structure comprises three subdomains (Figure 2C). Domain 1 and 2 adopt the RecA-like fold (Rao S T, et al., 1973; Story R M, et al., 1992) and contain eight conserved motifs that are essential for RNA binding and ATP hydrolysis activities and their coordination (Fairman-Williams M E, et al., 2010; Lescar

J, et al., 2008; Pyle A M, 2008). The third subdomain is unique to help form the single-stranded RNA binding groove. There is also evidence suggesting that domain 3 mediates the interaction with NS5 (Brooks A J, et al., 2002). In addition, NS3 also has RNA 5' triphosphatase activity (RTPase), which shares the same active site for ATP binding and hydrolysis (Figure 2C) (Balistreri G, et al., 2007). RNA 5' triphosphate hydrolysis is believed to be the first step of viral RNA capping (Decroly E, et al., 2012). Viruses carrying a defective or impaired NS3 helicase gene cannot replicate properly, indicating that NS3 is essential for virus replication (Matusan A E, et al., 2001). However, the exact function of the flavivirus NS3hel in the virus replication cycle is unknown. It is speculated that NS3hel could resolve secondary structures of the genomic RNA, displace transacting protein cofactors, and/or separate the dsRNA intermediate that is transiently formed during the polymerization reaction catalyzed by NS5 RdRP into single-strand form (Malet H, et al., 2007; Yu I M, et al., 2008; Yu L, et al., 2008).

Like most eukaryotic DExx helicase proteins, NS3 recognizes RNA largely in a sequence-independent manner (Fairman-Williams M E, et al., 2010; Luo D, et al., 2008; Pyle A M, 2008). During the duplex RNA unwinding process, one strand is inserted into the RNA binding groove of NS3hel, and the sugar-phosphate backbone makes direct or water-mediated contacts to the helicase residues, whereas the other strand is separated by a hydrophobic β -hairpin protruding from domain 2. This β -hairpin acts either actively or passively as a "helix opener" to disrupt base stacking and stabilize the unwound duplex, while the basic concave region between subdomain 2 and 3 works as "the translocator" by binding the 3' overhang (Buttner K, et al., 2007; Luo D, et al., 2008; Myong S, et al., 2007; Pyle A M, 2008). NS3hel binds to the 3' overhang sequence of the duplex RNA and unwind in the 3' to 5' direction (Figure 2C) (Benarroch D, et al., 2004; Li H, et al., 1999; Xu T, et al., 2005). NTP hydrolysis provides the energy to power the translocation and unwinding process, although the mechanism coupling these two enzymatic activities remains unclear. Single-molecule kinetic studies and structural biology work on HCV NS3hel collectively suggest that the basic step of translocation and unwinding is one base per ATP hydrolysis (Appleby T C, et al., 2010; Dumont S, et al., 2006; Gu M, et al., 2010; Myong S, et al., 2007). Comparative studies on flavivirus NS3 are not yet available and will be of great interest. To date, several flaviviral NS3hel domain structures have been reported (Table 1), providing the structural basis for its enzymatic activities, including basal and RNA-stimulated ATP hydrolysis activity, binding and translocation on the single-stranded RNA, and unwinding duplex RNA.

Helicase inhibitors could serve as common anti-viral therapy against flaviviruses (Lim S P, et al., 2013; Shadrick W R, et al., 2013). HCV NS3hel has been extensively studied, and compound inhibitors have been reported (Li K, et al., 2010; Maga G, et al., 2005; Salam K A, et al., 2013). However, lack of specificity at the RNA binding site and the NTP binding pocket might cause toxicity, as many cellular proteins carry similar helicase/NTPase activities (Linder P, et al., 2011; Shadrick W R, et al., 2013; Steimer L, et al., 2012). The intrinsic flexibility of motor proteins also makes it difficult to identify high-affinity and high potency compounds, although arguably allosteric inhibition remains an attractive concept for inhibitor design (Luo D, et al., 2008; Pyle A M, 2008; Saalau-Bethell S M, et al., 2012). Interesting and potent NS3hel inhibitors have been reported (Byrd C M, et al., 2013; Mastrangelo E, et al., 2012), but no helicase inhibitors have been approved for clinical trials or usage.

FULL-LENGTH NS3 IS MORE THAN NS2BNS3PRO PLUS NS3HEL

We do not fully understand why the two independent enzymatic activities hosted by two protein domains – protease and NTPase/helicase – are assembled into the single NS3 protein from *flaviviridae*. However, there are potential advantages for viruses to encode such a fusion gene. Polyprotein processing occurs co- and post-translationally in ER-derived intracellular membranes. NS3 may help bring NS4A-NS4B-NS5 in proximity to NS2BNS3pro for the efficient cleavage and release of individual viral proteins. In addition, NS2B anchors NS3 in the membrane, which is a prerequisite for viral replication complex maturation. It has been shown that flavivirus RNA replication is performed inside a vesicular-like compartment made of remodeled intracellular membranes (Miorin L, et al., 2013; Paul D, et al., 2013; Welsch S, et al., 2009). Therefore, NS2B together with other small membrane proteins might provide the membrane platform for the replicative enzymes (NS3 and NS5) and the viral RNA to assemble into the high-order structures of the flaviviral replication complex (Miorin L, et al., 2013; Paul D, et al., 2013; Welsch S, et al., 2009). Interestingly, YFV NS3 has been suggested to be involved in virus assembly independent of its enzymatic functions. A W349A mutation within NS3hel did not result in viral replication defects. However, no infectious viruses but only capsidless subviral particles could be detected from the mutant virus infected cells (Patkar C G, et al., 2008).

Molecular biologists studied the enzymatic activities of the two domains in the context of full-length NS3. Earlier work compared the enzymatic activities of full-length DENV2 NS3 (without NS2B cofactor) with NS3 helicase (aa171-618) and demonstrated that full-length NS3 has much higher unwinding activity

but a lower ATP hydrolysis rate (Xu T, et al., 2005; Yon C, et al., 2005). The affinity of dengue virus serotype 4 (DENV4) NS2B₁₈NS3 to ATP analogs is 10-fold higher than that of the truncated helicase (Luo D, et al., 2008). Intriguingly, no influence of the protease on helicase activity has been observed (or vice versa) in an MVEV NS2B₄₇NS3 construct (Assenberg R, et al., 2009). More recently, the Gamarnik group also reported comparable ATPase and helicase activities for his₆-tagged DENV2 NS2B₄₇NS3 and NS3hel, respectively (Gebhard L G, et al., 2012). This discrepancy may be due to the variations in protein constructs, enzyme preparation, and assay setup. Furthermore, the correct folded protease domain might assist NS3 to select the substrate specificity to RNA, as WNV NS3hel, but not NS2B₄₈NS3, unwinds both DNA and RNA (Chernov A V, et al., 2008). Studying the biochemistry of the native-state-like NS2BNS3 in the membrane context could clarify this issue and shed more light on the roles of NS3 in virus replication (Choksupmanee O, et al., 2012; Huang Q, et al., 2013).

Intriguingly, structural studies of full-length NS3 revealed three distinct configurations that differ from each other in the relative positioning of the NS3pro and NS3hel via a flexible inter-domain linker (Figure 3ABC) (Assenberg R, et al., 2009; Luo D, et al., 2012; Luo D, et al., 2008; Luo D, et al., 2010). We reported two different conformations of the DENV4 NS2B₁₈NS3 protein where the protease domain has rotated by approximately 161 degrees with respect to the helicase domain (Luo D, et al., 2012; Luo D, et al., 2008; Luo D, et al., 2010). The structure of the MVEV NS2B₄₇NS3 presented a third and a more radically different conformation of the NS3 protease-helicase (Assenberg R, et al., 2009). Interestingly, small-angle X-ray scattering (SAXS) experiments showed that both DENV4 NS2B₁₈NS3 protein and Kunjin NS3 adopted similar elongated shapes in solution (Luo D, et al., 2008; Mastrangelo E, et al., 2007). Given the close fit of the crystal structures and the hydrated envelope determined *ab initio* from SAXS data, it is evident that the flaviviral NS3 protease-helicase enzymes maintain an elongated conformation while the two domains are loosely connected through the linker (Figure 3). In contrast, the HCV NS3NS4A protease-helicase has been shown to adopt a globular conformation (Figure 3D) (Yao N, et al., 1999). In the first crystal structure of the similarly engineered NS4A₁₁NS3 fusion protein of HCV genotype 1b, the protease domain binds the C terminus of the NS3hel, mimicking the post-*cis*-cleavage state at the NS3-NS4A junction (Figure 3D) (Yao N, et al., 1999). NS4A₁₁NS3 fusion proteins from the same or different genotypes have also been crystallized in the similar globular conformation as *apo* enzyme (Appleby T C, et al., 2011), in complex with ssRNA and ATP analogs (Appleby T C, et al.,

2011), in complex with a macrocyclic protease inhibitor (Schiering N, et al., 2011), and in complex with allosteric inhibitors (Saalau-Bethell S M, et al., 2012). Sequence analysis revealed that the linker (aa 169–179) is less conserved compared to the two functional domains of the flavivirus NS3 (Figure 1C). The flavivirus NS3 linker is also rich in acidic residues (Luo D, et al., 2008). Conversely, the linker region of HCV NS3 (aa 181–194) is very conserved and rich in proline residues (Figure 1C). The sequence divergence between flavivirus and hepatitis virus NS3 seems to correlate with the distinct configurations of the protease-helicase: the flexible and less conserved linker is found in the extended conformations of flavivirus NS3 proteins, and the conserved and proline-rich linker is found in the globular conformations of HCV NS3 (Figure 1C and 3). In flaviviruses, the NS3 linker may have an optimum length and flexibility for polyprotein processing and RNA replication (Luo D, et al., 2010). It will be interesting to assess the impact of the HCV NS3 linker on virus replication. These studies on full-length NS3 facilitate a better understanding of the flavivirus life cycle and the exploration of new avenues for the development of antivirals and vaccines (Figure 4) (Luo D, et al., 2010; Saalau-Bethell S M, et al., 2012).

FUTURE PERSPECTIVE

Recent work utilizing yeast two-hybrid systems revealed the flavivirus NS proteins interactome (Khadka S, et al., 2011; Le Breton M, et al., 2011; Zou G, et al., 2011). More than 100 human proteins were identified to interact with NS3, NS5, or both, and many are involved in transcription regulation and host immune response (Krishnan M N, et al., 2014). Type I interferon (IFN) production in human monocyte-derived dendritic cells was inhibited as the cytoplasmic adaptor protein – stimulator of interferon genes (STING) (Ishikawa H, et al., 2008) or mediator of IRF3 activation (MITA) (Zhong B, et al., 2008) – was found cleaved and thus inactivated by DENV NS2BNS3pro (Aguirre S, et al., 2012). WNV NS2BNS3 has been shown to induce apoptosis through the activation of caspases 3 and 8 (Ramanathan M P, et al., 2006). Another study reported that JEV NS2BNS3pro inhibits the signaling pathway of activator protein 1 (AP-1), probably through proteolysis, and the authors suggested that NS2BNS3 protease may contribute to JEV-induced neurotropic pathogenesis (Lin C W, et al., 2006). Better and deeper understanding of how different enzymatic and non-enzymatic activities of NS3 are regulated is needed. The molecular and cellular biology of this protein and its structural and functional roles in viral replication complex formation definitely deserve more attention (Figure 4) as the dynamic nature of the host-pathogen interactions remains unclear (Suthar M S, et al., 2013). Advances in these

research areas will inevitably promote the rational design of novel inhibitors and vaccines against diseases caused by flavivirus infections.

ACKNOWLEDGMENTS

The author would like to thank Moon Tay and Julien Lescar for their critical reading of the manuscript.

COMPLIANCE WITH ETHICS GUIDELINES

All the authors declare that they have no competing interest. This article does not contain any studies with human or animal subjects performed by the any of the authors.

REFERENCES

- Aguirre S, Maestre A M, Pagni S, Patel J R, Savage T, Gutman D, Maringer K, Bernal-Rubio D, Shabman R S, Simon V, Rodriguez-Madoz J R, Mulder L C, Barber G N, and Fernandez-Sesma A. 2012. DENV inhibits type I IFN production in infected cells by cleaving human STING. *PLoS Pathog*, 8: e1002934.
- Aleshin A E, Shiryayev S A, Strongin A Y, and Liddington R C. 2007. Structural evidence for regulation and specificity of flaviviral proteases and evolution of the *Flaviviridae* fold. *Protein Sci*, 16: 795-806.
- Alvarez D E, Lodeiro M F, Filomatori C V, Fucito S, Mondotte J A, and Gamarnik A V. 2006. Structural and functional analysis of dengue virus RNA. *Novartis Found Symp*, 277: 120-132; discussion 132-125, 251-123.
- Appleby T C, Anderson R, Fedorova O, Pyle A M, Wang R, Liu X, Brendza K M, and Somoza J R. 2010. Visualizing ATP-dependent RNA translocation by the NS3 helicase from HCV. *J Mol Biol*, 405: 1139-1153.
- Arakaki T L, Fang N X, Fairlie D P, Young P R, and Martin J L. 2002. Catalytically active Dengue virus NS3 protease forms aggregates that are separable by size exclusion chromatography. *Protein Expr Purif*, 25: 241-247.
- Assenberg R, Mastrangelo E, Walter T S, Verma A, Milani M, Owens R J, Stuart D I, Grimes J M, and Mancini E J. 2009. Crystal structure of a novel conformational state of the flavivirus NS3 protein: implications for polyprotein processing and viral replication. *J Virol*, 83: 12895-12906.
- Balistreri G, Caldentey J, Kaariainen L, and Ahola T. 2007. Enzymatic defects of the nsP2 proteins of Semliki Forest virus temperature-sensitive mutants. *J Virol*, 81: 2849-2860.

- Benarroch D, Selisko B, Locatelli G A, Maga G, Romette J L, and Canard B. 2004. The RNA helicase, nucleotide 5'-triphosphatase, and RNA 5'-triphosphatase activities of Dengue virus protein NS3 are Mg²⁺-dependent and require a functional Walker B motif in the helicase catalytic core. *Virology*, 328: 208-218.
- Bollati M, Alvarez K, Assenberg R, Baronti C, Canard B, Cook S, Coutard B, Decroly E, de Lamballerie X, Gould E A, Grard G, Grimes J M, Hilgenfeld R, Jansson A M, Malet H, Mancini E J, Mastrangelo E, Mattevi A, Milani M, Moureau G, Neyts J, Owens R J, Ren J, Selisko B, Speroni S, Steuber H, Stuart D I, Unge T, and Bolognesi M. 2010. Structure and functionality in flavivirus NS-proteins: perspectives for drug design. *Antiviral Res*, 87: 125-148.
- Brooks A J, Johansson M, John A V, Xu Y, Jans D A, and Vasudevan S G. 2002. The interdomain region of dengue NS5 protein that binds to the viral helicase NS3 contains independently functional importin beta 1 and importin alpha/beta-recognized nuclear localization signals. *J Biol Chem*, 277: 36399-36407.
- Buttner K, Nehring S, and Hopfner K P. 2007. Structural basis for DNA duplex separation by a superfamily-2 helicase. *Nat Struct Mol Biol*, 14: 647-652.
- Byrd C M, Grosenbach D W, Berhanu A, Dai D, Jones K F, Cardwell K B, Schneider C, Yang G, Tyavanagimatt S, Harver C, Wineinger K A, Page J, Stavale E, Stone M A, Fuller K P, Lovejoy C, Leeds J M, Hruby D E, and Jordan R. 2013. Novel benzoxazole inhibitor of dengue virus replication that targets the NS3 helicase. *Antimicrob Agents Chemother*, 57: 1902-1912.
- Chambers T J, Hahn C S, Galler R, and Rice C M. 1990. Flavivirus genome organization, expression, and replication. *Annu Rev Microbiol*, 44: 649-688.
- Chandramouli S, Joseph J S, Daudenarde S, Gatchalian J, Cornillez-Ty C, and Kuhn P. 2010. Serotype-specific structural differences in the protease-cofactor complexes of the dengue virus family. *J Virol*, 84: 3059-3067.
- Chappell K J, Stoermer M J, Fairlie D P, and Young P R. 2008. Mutagenesis of the West Nile virus NS2B cofactor domain reveals two regions essential for protease activity. *J Gen Virol*, 89: 1010-1014.
- Chernov A V, Shiryaev S A, Aleshin A E, Ratnikov B I, Smith J W, Liddington R C, and Strongin A Y. 2008. The two-component NS2B-NS3 proteinase represses DNA unwinding activity of the West Nile virus NS3 helicase. *J Biol Chem*, 283: 17270-17278.

- Choksupmanee O, Hodge K, Katzenmeier G, and Chimnaronk S. 2012. Structural platform for the autolytic activity of an intact NS2B-NS3 protease complex from dengue virus. *Biochemistry*, 51: 2840-2851.
- Clum S, Ebner K E, and Padmanabhan R. 1997. Cotranslational membrane insertion of the serine proteinase precursor NS2B-NS3(Pro) of dengue virus type 2 is required for efficient in vitro processing and is mediated through the hydrophobic regions of NS2B. *J Biol Chem*, 272: 30715-30723.
- Decroly E, Ferron F, Lescar J, and Canard B. 2012. Conventional and unconventional mechanisms for capping viral mRNA. *Nat Rev Microbiol*, 10: 51-65.
- Dumont S, Cheng W, Serebrov V, Beran R K, Tinoco I, Jr., Pyle A M, and Bustamante C. 2006. RNA translocation and unwinding mechanism of HCV NS3 helicase and its coordination by ATP. *Nature*, 439: 105-108.
- Egloff M P, Benarroch D, Selisko B, Romette J L, and Canard B. 2002. An RNA cap (nucleoside-2'-O)-methyltransferase in the flavivirus RNA polymerase NS5: crystal structure and functional characterization. *EMBO J*, 21: 2757-2768.
- Erbel P, Schiering N, D'Arcy A, Renatus M, Kroemer M, Lim S P, Yin Z, Keller T H, Vasudevan S G, and Hommel U. 2006. Structural basis for the activation of flaviviral NS3 proteases from dengue and West Nile virus. *Nat Struct Mol Biol*, 13: 372-373.
- Fairman-Williams M E, Guenther U P, and Jankowsky E. 2010. SF1 and SF2 helicases: family matters. *Curr Opin Struct Biol*, 20: 313-324.
- Falgout B, Pethel M, Zhang Y M, and Lai C J. 1991. Both nonstructural proteins NS2B and NS3 are required for the proteolytic processing of dengue virus nonstructural proteins. *J Virol*, 65: 2467-2475.
- Filomatori C V, Lodeiro M F, Alvarez D E, Samsa M M, Pietrasanta L, and Gamarnik A V. 2006. A 5' RNA element promotes dengue virus RNA synthesis on a circular genome. *Genes Dev*, 20: 2238-2249.
- Frick D N, Rypma R S, Lam A M, and Gu B. 2004. The nonstructural protein 3 protease/helicase requires an intact protease domain to unwind duplex RNA efficiently. *J Biol Chem*, 279: 1269-1280.

- Gebhard L G, Kaufman S B, and Gamarnik A V. 2012. Novel ATP-independent RNA annealing activity of the dengue virus NS3 helicase. *PLoS One*, 7: e36244.
- Gorbalenya A E, and Koonin E V. 1993. Helicases: amino acid sequence comparisons and structure-function relationships. *Curr Opin Struc Biol*, 3: 419-429.
- Gorbalenya A E, Donchenko A P, Koonin E V, and Blinov V M. 1989. N-terminal domains of putative helicases of flavi- and pestiviruses may be serine proteases. *Nucleic Acids Res*, 17: 3889-3897.
- Gouvea I E, Izidoro M A, Judice W A, Cezari M H, Caliendo G, Santagada V, dos Santos C N, Queiroz M H, Juliano M A, Young P R, Fairlie D P, and Juliano L. 2007. Substrate specificity of recombinant dengue 2 virus NS2B-NS3 protease: influence of natural and unnatural basic amino acids on hydrolysis of synthetic fluorescent substrates. *Arch Biochem Biophys*, 457: 187-196.
- Gu M, and Rice C M. 2010. Three conformational snapshots of the hepatitis C virus NS3 helicase reveal a ratchet translocation mechanism. *Proc Natl Acad Sci USA*, 107: 521-528.
- Huang Q, Li Q, Joy J, Chen A S, Ruiz-Carrillo D, Hill J, Lescar J, and Kang C. 2013. Lyso-myristoyl phosphatidylcholine micelles sustain the activity of Dengue non-structural (NS) protein 3 protease domain fused with the full-length NS2B. *Protein Expr Purif*, 92: 156-162.
- Ishikawa H, and Barber G N. 2008. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature*, 455: 674-678.
- Jan L R, Yang C S, Trent D W, Falgout B, and Lai C J. 1995. Processing of Japanese encephalitis virus non-structural proteins: NS2B-NS3 complex and heterologous proteases. *J Gen Virol*, 76: 573-580.
- Keller T H, Chen Y L, Knox J E, Lim S P, Ma N L, Patel S J, Sampath A, Wang Q Y, Yin Z, and Vasudevan S G. 2006. Finding new medicines for flaviviral targets. *Novartis Found Symp*, 277: 102-114; discussion 114-109, 251-103.
- Khadka S, Vangeloff A D, Zhang C, Siddavatam P, Heaton N S, Wang L, Sengupta R, Sahasrabudhe S, Randall G, Gribskov M, Kuhn R J, Perera R, and LaCount D J. 2011. A physical interaction network of dengue virus and human proteins. *Mol Cell Proteomics*, 10: M111 012187.
- Kim J L, Morgenstern K A, Lin C, Fox T, Dwyer M D, Landro J A, Chambers S P, Markland W, Lepre C A, O'Malley E T, Harbeson S L, Rice C M, Mureko M A, Caron P R, and Thomson J A. 1996. Crystal structure of the hepatitis C virus NS3 protease domain complexed with a synthetic NS4A cofactor peptide. *Cell*, 87: 343-355.

- Krishnan M N, and Garcia-Blanco M A. 2014. Targeting host factors to treat west nile and dengue viral infections. *Viruses*, 6: 683-708.
- Le Breton M, Meyniel-Schicklin L, Deloire A, Coutard B, Canard B, de Lamballerie X, Andre P, Rabourdin-Combe C, Lotteau V, and Davoust N. 2011. Flavivirus NS3 and NS5 proteins interaction network: a high-throughput yeast two-hybrid screen. *BMC Microbiol*, 11: 234.
- Lescar J, Luo D, Xu T, Sampath A, Lim S P, Canard B, and Vasudevan S G. 2008. Towards the design of antiviral inhibitors against flaviviruses: the case for the multifunctional NS3 protein from Dengue virus as a target. *Antiviral Res*, 80: 94-101.
- Leung D, Schroder K, White H, Fang N X, Stoermer M J, Abbenante G, Martin J L, Young P R, and Fairlie D P. 2001. Activity of recombinant dengue 2 virus NS3 protease in the presence of a truncated NS2B co-factor, small peptide substrates, and inhibitors. *J Biol Chem*, 276: 45762-45771.
- Li H, Clum S, You S, Ebner K E, and Padmanabhan R. 1999. The serine protease and RNA-stimulated nucleoside triphosphatase and RNA helicase functional domains of dengue virus type 2 NS3 converge within a region of 20 amino acids. *J Virol*, 73: 3108-3116.
- Li J, Lim S P, Beer D, Patel V, Wen D, Tumanut C, Tully D C, Williams J A, Jiricek J, Priestle J P, Harris J L, and Vasudevan S G. 2005. Functional profiling of recombinant NS3 proteases from all four serotypes of dengue virus using tetrapeptide and octapeptide substrate libraries. *J Biol Chem*, 280: 28766-28774.
- Li K, Frankowski K J, Hanson A M, Ndjomou J, Shanahan M A, Mukherjee S, Kolli R, Shadrick W R, Sweeney N L, Belon C A, Neuenswander B, Ferguson J, Aube J, Schoenen F J, Blagg B S J, and Frick D N. 2010. Hepatitis C Virus NS3 Helicase Inhibitor Discovery, Probe Reports from the NIH Molecular Libraries Program, Bethesda (MD).
- Lim S P, Wang Q Y, Noble C G, Chen Y L, Dong H, Zou B, Yokokawa F, Nilar S, Smith P, Beer D, Lescar J, and Shi P Y. 2013. Ten years of dengue drug discovery: progress and prospects. *Antiviral Res*, 100: 500-519.
- Lin C W, Lin K H, Lyu P C, and Chen W J. 2006. Japanese encephalitis virus NS2B-NS3 protease binding to phage-displayed human brain proteins with the domain of trypsin inhibitor and basic region leucine zipper. *Virus Res*, 116: 106-113.

- Lindenbach B D, Thiel H J, and Rice C M. 2007. *Flaviviridae: the viruses and their replication*. Philadelphia: Lippincott-Raven Publishers.
- Linder P, and Jankowsky E. 2011. From unwinding to clamping - the DEAD box RNA helicase family. *Nat Rev Mol Cell Biol*, 12: 505-516.
- Luo D, Lim S P, and Lescar J. 2012. Chapter 5: The Flavivirus NS3 Protein: Structure and Functions *Molecular Virology and Control of Flaviviruses.*, ISBN: 978-1-904455-92-9. .
- Luo D, Xu T, Hunke C, Gruber G, Vasudevan S G, and Lescar J. 2008. Crystal structure of the NS3 protease-helicase from dengue virus. *J Virol*, 82: 173-183.
- Luo D, Wei N, Doan D N, Paradkar P N, Chong Y, Davidson A D, Kotaka M, Lescar J, and Vasudevan S G. 2010. Flexibility between the protease and helicase domains of the dengue virus NS3 protein conferred by the linker region and its functional implications. *J Biol Chem*, 285: 18817-18827.
- Luo D, Xu T, Watson R P, Scherer-Becker D, Sampath A, Jahnke W, Yeong S S, Wang C H, Lim S P, Strongin A, Vasudevan S G, and Lescar J. 2008. Insights into RNA unwinding and ATP hydrolysis by the flavivirus NS3 protein. *EMBO J*, 27: 3209-3219.
- Mackenzie J. 2005. Wrapping things up about virus RNA replication. *Traffic*, 6: 967-977.
- Maga G, Gemma S, Fattorusso C, Locatelli G A, Butini S, Persico M, Kukreja G, Romano M P, Chiasserini L, Savini L, Novellino E, Nacci V, Spadari S, and Campiani G. 2005. Specific targeting of hepatitis C virus NS3 RNA helicase. Discovery of the potent and selective competitive nucleotide-mimicking inhibitor QU663. *Biochemistry*, 44: 9637-9644.
- Malet H, Masse N, Selisko B, Romette J L, Alvarez K, Guillemot J C, Tolou H, Yap T L, Vasudevan S, Lescar J, and Canard B. 2008. The flavivirus polymerase as a target for drug discovery. *Antiviral Res*, 80: 23-35.
- Malet H, Egloff M P, Selisko B, Butcher R E, Wright P J, Roberts M, Gruez A, Sulzenbacher G, Vonnrhein C, Bricogne G, Mackenzie J M, Khromykh A A, Davidson A D, and Canard B. 2007. Crystal structure of the RNA polymerase domain of the West Nile virus non-structural protein 5. *J Biol Chem*, 282: 10678-10689.
- Mancini E J, Assenberg R, Verma A, Walter T S, Tuma R, Grimes J M, Owens R J, and Stuart D I. 2007. Structure of the Murray Valley encephalitis virus RNA helicase at 1.9 Angstrom resolution. *Protein Sci*, 16: 2294-2300.

- Markoff L. 2003. 5'- and 3'-noncoding regions in flavivirus RNA. *Adv Virus Res*, 59: 177-228.
- Mastrangelo E, Pezzullo M, De Burghgraeve T, Kaptein S, Pastorino B, Dallmeier K, de Lamballerie X, Neyts J, Hanson A M, Frick D N, Bolognesi M, and Milani M. 2012. Ivermectin is a potent inhibitor of flavivirus replication specifically targeting NS3 helicase activity: new prospects for an old drug. *J Antimicrob Chemother*, 67: 1884-1894.
- Mastrangelo E, Milani M, Bollati M, Selisko B, Peyrane F, Pandini V, Sorrentino G, Canard B, Konarev P V, Svergun D I, de Lamballerie X, Coutard B, Khromykh A A, and Bolognesi M. 2007. Crystal structure and activity of Kunjin virus NS3 helicase; protease and helicase domain assembly in the full length NS3 protein. *J Mol Biol*, 372: 444-455.
- Matusan A E, Pryor M J, Davidson A D, and Wright P J. 2001. Mutagenesis of the Dengue virus type 2 NS3 protein within and outside helicase motifs: effects on enzyme activity and virus replication. *J Virol*, 75: 9633-9643.
- Miorin L, Romero-Brey I, Maiuri P, Hoppe S, Krijnse-Locker J, Bartenschlager R, and Marcello A. 2013. Three-dimensional architecture of tick-borne encephalitis virus replication sites and trafficking of the replicated RNA. *J Virol*, 87: 6469-6481.
- Murray C L, Jones C T, and Rice C M. 2008. Architects of assembly: roles of Flaviviridae non-structural proteins in virion morphogenesis. *Nat Rev Microbiol*, 6: 699-708.
- Myong S, Bruno M M, Pyle A M, and Ha T. 2007. Spring-loaded mechanism of DNA unwinding by hepatitis C virus NS3 helicase. *Science*, 317: 513-516.
- Nall T A, Chappell K J, Stoermer M J, Fang N X, Tyndall J D, Young P R, and Fairlie D P. 2004. Enzymatic characterization and homology model of a catalytically active recombinant West Nile virus NS3 protease. *J Biol Chem*, 279: 48535-48542.
- Niyomrattanakit P, Yahorava S, Mutule I, Mutulis F, Petrovska R, Prusis P, Katzenmeier G, and Wikberg J E. 2006. Probing the substrate specificity of the dengue virus type 2 NS3 serine protease by using internally quenched fluorescent peptides. *Biochem J*, 397: 203-211.
- Noble C G, Seh C C, Chao A T, and Shi P Y. 2012. Ligand-bound structures of the dengue virus protease reveal the active conformation. *J Virol*, 86: 438-446.
- Patkar C G, and Kuhn R J. 2008. Yellow Fever virus NS3 plays an essential role in virus assembly independent of its known enzymatic functions. *J Virol*, 82: 3342-3352.

- Paul D, and Bartenschlager R. 2013. Architecture and biogenesis of plus-strand RNA virus replication factories. *World J Virol*, 2: 32-48.
- Polacek C, Foley J E, and Harris E. 2009. Conformational changes in the solution structure of the dengue virus 5' end in the presence and absence of the 3' untranslated region. *J Virol*, 83: 1161-1166.
- Pyle A M. 2008. Translocation and unwinding mechanisms of RNA and DNA helicases. *Annu Rev Biophys*, 37: 317-336.
- Ramanathan M P, Chambers J A, Pankhong P, Chattergoon M, Attatippaholkun W, Dang K, Shah N, and Weiner D B. 2006. Host cell killing by the West Nile Virus NS2B-NS3 proteolytic complex: NS3 alone is sufficient to recruit caspase-8-based apoptotic pathway. *Virology*, 345: 56-72.
- Rao S T, and Rossmann M G. 1973. Comparison of super-secondary structures in proteins. *J Mol Biol*, 76: 241-256.
- Ray D, Shah A, Tilgner M, Guo Y, Zhao Y, Dong H, Deas T S, Zhou Y, Li H, and Shi P Y. 2006. West Nile virus 5'-cap structure is formed by sequential guanine N-7 and ribose 2'-O methylations by nonstructural protein 5. *J Virol*, 80: 8362-8370.
- Rendi-Wagner P. 2008. Advances in vaccination against tick-borne encephalitis. *Expert Rev Vaccines*, 7: 589-596.
- Robin G, Chappell K, Stoermer M J, Hu S H, Young P R, Fairlie D P, and Martin J L. 2009. Structure of West Nile virus NS3 protease: ligand stabilization of the catalytic conformation. *J Mol Biol*, 385: 1568-1577.
- Saalau-Bethell S M, Woodhead A J, Chessari G, Carr M G, Coyle J, Graham B, Hiscock S D, Murray C W, Pathuri P, Rich S J, Richardson C J, Williams P A, and Jhoti H. 2012. Discovery of an allosteric mechanism for the regulation of HCV NS3 protein function. *Nat Chem Biol*, 8: 920-925.
- Sabchareon A, Wallace D, Sirivichayakul C, Limkittikul K, Chanthavanich P, Suvannadabba S, Jiwariyavej V, Dulyachai W, Pengsaa K, Wartel T A, Moureau A, Saville M, Bouckennooghe A, Viviani S, Tornieporth N G, and Lang J. 2012. Protective efficacy of the recombinant, live-attenuated, CYD tetravalent dengue vaccine in Thai schoolchildren: a randomised, controlled phase 2b trial. *Lancet*, 380: 1559-1567.
- Salam K A, and Akimitsu N. 2013. Hepatitis C virus NS3 inhibitors: current and future perspectives. *Biomed Res Int*, 2013: 467869.

- Sampath A, and Padmanabhan R. 2009. Molecular targets for flavivirus drug discovery. *Antiviral Res*, 81: 6-15.
- Schiering N, D'Arcy A, Villard F, Simic O, Kamke M, Monnet G, Hassiepen U, Svergun D I, Pulfer R, Eder J, Raman P, and Bodendorf U. 2011. A macrocyclic HCV NS3/4A protease inhibitor interacts with protease and helicase residues in the complex with its full-length target. *Proc Natl Acad Sci USA*, 108: 21052-21056.
- Shadrick W R, Ndjomou J, Kolli R, Mukherjee S, Hanson A M, and Frick D N. 2013. Discovering new medicines targeting helicases: challenges and recent progress. *J Biomol Screen*, 18: 761-781.
- Shiryaev S A, Kozlov I A, Ratnikov B I, Smith J W, Lebl M, and Strongin A Y. 2007. Cleavage preference distinguishes the two-component NS2B-NS3 serine proteinases of Dengue and West Nile viruses. *Biochem J*, 401: 743-752.
- Speroni S, De Colibus L, Mastrangelo E, Gould E, Coutard B, Forrester N L, Blanc S, Canard B, and Mattevi A. 2008. Structure and biochemical analysis of Kokobera virus helicase. *Proteins*, 70: 1120-1123.
- Steimer L, and Klostermeier D. 2012. RNA helicases in infection and disease. *RNA Biol*, 9.
- Story R M, and Steitz T A. 1992. Structure of the recA protein-ADP complex. *Nature*, 355: 374-376.
- Suthar M S, Aguirre S, and Fernandez-Sesma A. 2013. Innate immune sensing of flaviviruses. *PLoS Pathog*, 9: e1003541.
- Tomei L, Failla C, Vitale R L, Bianchi E, and De Francesco R. 1996. A central hydrophobic domain of the hepatitis C virus NS4A protein is necessary and sufficient for the activation of the NS3 protease. *J Gen Virol*, 77: 1065-1070.
- Urbani A, Bazzo R, Nardi M C, Cicero D O, De Francesco R, Steinkuhler C, and Barbato G. 1998. The metal binding site of the hepatitis C virus NS3 protease. A spectroscopic investigation. *J Biol Chem*, 273: 18760-18769.
- Verma R, Khanna P, and Chawla S. 2013. Yellow fever vaccine: An effective vaccine for travelers. *Hum Vaccin Immunother*, 10.
- Villordo S M, and Gamarnik A V. 2009. Genome cyclization as strategy for flavivirus RNA replication. *Virus Res*, 139: 230-239.

- Welsch S, Miller S, Romero-Brey I, Merz A, Bleck C K, Walther P, Fuller S D, Antony C, Krijnse-Locker J, and Bartenschlager R. 2009. Composition and three-dimensional architecture of the dengue virus replication and assembly sites. *Cell Host Microbe*, 5: 365-375.
- WHO. 2009. Fact sheet: Dengue and dengue haemorrhagic fever.
<http://www.who.int/csr/disease/dengue/en/>
- Wu J, Bera A K, Kuhn R J, and Smith J L. 2005. Structure of the Flavivirus helicase: implications for catalytic activity, protein interactions, and proteolytic processing. *J Virol*, 79: 10268-10277.
- Xu T, Sampath A, Chao A, Wen D, Nanao M, Chene P, Vasudevan S G, and Lescar J. 2005. Structure of the Dengue virus helicase/nucleoside triphosphatase catalytic domain at a resolution of 2.4 Å. *J Virol*, 79: 10278-10288.
- Xu T, Sampath A, Chao A, Wen D, Nanao M, Luo D, Chene P, Vasudevan S G, and Lescar J. 2006. Towards the design of flavivirus helicase/NTPase inhibitors: crystallographic and mutagenesis studies of the dengue virus NS3 helicase catalytic domain. *Novartis Found Symp*, 277: 87-97; discussion 97-101, 251-103.
- Yamashita T, Unno H, Mori Y, Tani H, Moriishi K, Takamizawa A, Agoh M, Tsukihara T, and Matsuura Y. 2008. Crystal structure of the catalytic domain of Japanese encephalitis virus NS3 helicase/nucleoside triphosphatase at a resolution of 1.8 Å. *Virology*, 373: 426-436.
- Yao N, Reichert P, Taremi S S, Prosser W W, and Weber P C. 1999. Molecular views of viral polyprotein processing revealed by the crystal structure of the hepatitis C virus bifunctional protease-helicase. *Structure*, 7: 1353-1363.
- Yon C, Teramoto T, Mueller N, Phelan J, Ganesh V K, Murthy K H, and Padmanabhan R. 2005. Modulation of the nucleoside triphosphatase/RNA helicase and 5'-RNA triphosphatase activities of Dengue virus type 2 nonstructural protein 3 (NS3) by interaction with NS5, the RNA-dependent RNA polymerase. *J Biol Chem*, 280: 27412-27419.
- Yu I M, Zhang W, Holdaway H A, Li L, Kostyuchenko V A, Chipman P R, Kuhn R J, Rossmann M G, and Chen J. 2008. Structure of the immature dengue virus at low pH primes proteolytic maturation. *Science*, 319: 1834-1837.
- Yu L, Nomaguchi M, Padmanabhan R, and Markoff L. 2008. Specific requirements for elements of the 5' and 3' terminal regions in flavivirus RNA synthesis and viral replication. *Virology*, 374: 170-185.

- Yun S I, and Lee Y M. 2013. Japanese Encephalitis: The virus and vaccines. *Hum Vaccin Immunother*, 10.
- Yusof R, Clum S, Wetzel M, Murthy H M, and Padmanabhan R. 2000. Purified NS2B/NS3 serine protease of dengue virus type 2 exhibits cofactor NS2B dependence for cleavage of substrates with dibasic amino acids in vitro. *J Biol Chem*, 275: 9963-9969.
- Zhang L, Mohan P M, and Padmanabhan R. 1992. Processing and localization of Dengue virus type 2 polyprotein precursor NS3-NS4A-NS4B-NS5. *J Virol*, 66: 7549-7554.
- Zhong B, Yang Y, Li S, Wang Y Y, Li Y, Diao F, Lei C, He X, Zhang L, Tien P, and Shu H B. 2008. The adaptor protein MITA links virus-sensing receptors to IRF3 transcription factor activation. *Immunity*, 29: 538-550.
- Zou G, Chen Y L, Dong H, Lim C C, Yap L J, Yau Y H, Shochat S G, Lescar J, and Shi P Y. 2011. Functional analysis of two cavities in flavivirus NS5 polymerase. *J Biol Chem*, 286: 14362-14372.

Titles and legends to figures




Figure 1. Schematic representation of the flaviviral genome, polyprotein, and functional domain partition along the NS2BNS3 gene. (A) Flavivirus genome and polyprotein. The viral genome contains a 5' cap and untranslated regions at both the 5' and 3' termini. The polyproteins are processed by both NS2BNS3 protease indicated as filled arrows and the host proteases by unfilled arrows. (B) The schematic representation of the NS2B and NS3 genes. The central cofactor region of NS2B is in red, and the putative

membrane associate regions of *NS2B* are displayed as blocks. (C) The sequence alignment of the linker regions between the protease and helicase domains of NS3.




Figure 2. Structures of NS2BNS3_{pro} and NS3_{hel}. (A) Overall fold of NS2B₄₇NS3_{pro} from WNV. NS3_{Pro} is shown in cyan, the NS2B₄₇ region in red, and the peptide inhibitor is in yellow. PDB code: 2FP7. (B) Surface view of the substrate binding site of WNV NS2BNS3_{pro}. (C) Ternary complex of DENV4 NS3_{hel}-ssRNA-AMPPNP. PDB code: 2JLV.




Figure 3. Structural views of the full-length NS3 protein. Structures of (A) and (B) NS2B₁₈NS3 from DENV4, (C) NS2B₄₇NS3 from MVEV, and (D) NS3NS4A₁₁ from HCV. NS2B (or NS4A) cofactor peptide is in red; NS3pro in green; linker in purple; and NS3hel is in blue, yellow, gray, and orange. In (A-D), the helicase domains are aligned, and the black dots represent the ssRNA binding tunnel. (E) When the protease domains are superimposed, the helicase domains display various orientations.




Figure 4. Putative model of the membrane-associated state of the NS2BNS3 complex. The full length of NS2B is modeled based on the predicted topology of NS2B (Luo D, et al., 2010). Numbering of NS2B residues: residues 49–66–96 are the central cofactor region of NS2B colored in red; residues 83–85 directly participate in substrate binding; membrane-associated regions include N-49 and 96-C. NS3pro is colored in green, NS3hel in cyan, and ssRNA in purple. The components of the viral replication complex, including NS5, NS1, NS4A, and NS4B, are drawn across the ER membrane. Lacking information about the precise mapping of the intermolecular interfaces, these individual components are only approximately modeled on the ER membrane.

Table 1. List of flavivirus NS2BNS3 structures

Domains	Virus	PDB ID	Ligands	Reference	
NS2B NS3pro	DENV1	3L6P	-	(Chandramouli S, et al., 2010)	
		3LKW	-		
	DENV2	2FOM	-	(Erbel P, et al., 2006)	
	DENV3	3U1I	Bz-Nle-Lys-Arg-Arg-H	(Noble C G, et al., 2012)	
		3U1J	Aprotinin / Bovine Pancreatic Trypsin Inhibitor (BPTI)		
	WNV	2YOL	3,4-dichlorophenylacetyl-Lys-Lys-GCMA	(Hammamy M Z, et al., 2013)	
			3E90	2-naphthoyl-Lys-Lys-Arg-H	(Robin G, et al., 2009)
		2IJO	Aprotinin	(Aleshin A E, et al., 2007)	
		2FP7	Bz-Nle-Lys-Arg-Arg-H	(Erbel P, et al., 2006)	
		2GGV	-	(Aleshin A E, et al., 2007)	
NS3hel	DENV2	2BHR	SO ₄	(Xu T, et al., 2005)	
		2BMF	-		
	DENV4	2JLQ	-	(Luo D, et al., 2008)	
		2JLR	AMPPNP_Mn ²⁺		
		2JLU	ssRNA ₁₂		
		2JLV	ssRNA ₁₂ , AMPPNP_Mn ²⁺		
		2JLW	ssRNA ₁₃		
		2JLX	ssRNA ₁₂ , ADP-Vandate_Mn ²⁺		
		2JLY	ssRNA ₁₂ , ADP-PO ₄ _Mn ²⁺		
		2JLZ	ssRNA ₁₂ , ADP_Mn ²⁺		
	Kokobera Virus	2V6I	-	(Speroni S, et al., 2008)	
		2V6J	-		
	JEV	2Z83	-	(Yamashita T, et al., 2008)	
	MVEV	2V8O	-	(Mancini E J, et al., 2007)	
	Kunjing Virus	2QEQ	-	(Mastrangelo E, et al., 2007)	
	YFV	1YKS	-	(Wu J, et al., 2005)	
		1YMF	ADP		
	NS2B ₁₈ NS3	DENV4	2WHX	ADP_Mn ²⁺	(Luo D, et al., 2010)
			2WZQ	-	
2VBC			-	(Luo D, et al., 2008)	
NS2B ₄₇ NS3	MVEV	2WV9	-	(Assenberg R, et al., 2009)	