

This document is downloaded from DR-NTU, Nanyang Technological University Library, Singapore.

Title	A comparison and optimization of methods and factors affecting the transformation of Escherichia coli
Author(s)	Chan, Weng-Tat; Verma, Chandra Shekhar; Lane, David P.; Gan, Samuel Ken-En
Citation	Chan, W., Verma, C., Lane, D., & Gan, S. (2013). A comparison and optimization of methods and factors affecting the transformation of Escherichia coli. Bioscience Reports, 33(6), 931-937.
Date	2013
URL	http://hdl.handle.net/10220/39215
Rights	© 2013 The Author(s) This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC-BY) (http://creativecommons.org/licenses/by/3.0/) which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.



OPEN ACCESS

A comparison and optimization of methods and factors affecting the transformation of *Escherichia coli*

Weng-Tat CHAN*, Chandra S. VERMA*†‡, David P. LANE‡§ and Samuel Ken-En GAN*¹

*Bioinformatics Institute, Agency for Science, Technology, and Research (A*STAR), Singapore 138671

†Department of Biological Sciences, National University of Singapore (NUS), Singapore 119077

‡School of Biological Sciences, Nanyang Technological University (NTU), Singapore 639798

§p53 Laboratory, Agency for Science, Technology, and Research (A*STAR), Singapore 138648

Synopsis

DNA manipulation routinely requires competent bacteria that can be made using one of numerous methods. To determine the best methods, we compared four commonly used chemical methods (DMSO, MgCl₂-CaCl₂, CaCl₂ and Hanahan's methods) on frequently used *Escherichia coli* (*E. coli*) strains: DH5 α , XL-1 Blue, SCS110, JM109, TOP10 and BL21-(DE3)-PLysS. Hanahan's method was found to be most effective for DH5 α , XL-1 Blue and JM109 strains ($P < 0.05$), whilst the CaCl₂ method was best for SCS110, TOP10 and BL21 strains ($P < 0.05$). The use of SOB (super optimal broth) over LB [Luria-Bertani (broth)] growth media was found to enhance the competency of XL-1 Blue ($P < 0.05$), dampened JM109's competency ($P < 0.05$), and had no effect on the other strains ($P > 0.05$). We found no significant differences between using 45 or 90 s heat shock across all the six strains ($P > 0.05$). Through further optimization by means of concentrating the aliquots, we were able to get further increases in transformation efficiencies. Based on the optimized parameters and methods, these common laboratory *E. coli* strains attained high levels of TrE (transformation efficiency), thus facilitating the production of highly efficient and cost-effective competent bacteria.

Key words: CaCl₂ method, competent bacteria, DMSO method, Hanahan's method, heat shock

Cite this article as: Chan, W.T., Verma, C.S., Lane, D.P and Gan, S.K. (2013) A comparison and optimization of methods and factors affecting the Transformation of *Escherichia coli*. Biosci. Rep. **33**(6), art:e00086.doi:10.1042/BSR20130098

INTRODUCTION

Bacterial transformation, the process whereby bacteria are able to take up foreign DNA was first demonstrated by Griffith with *Streptococcus pneumoniae* [1], and is now routinely used in laboratories. Although reported to occur naturally in bacteria such as *Bacillus subtilis* [2], such phenomenon is generally uncommon in *Escherichia coli*, which require induction by artificial methods, such as those first demonstrated by Mandel and Higa [3]. This chemical method involved treating the bacteria with bacteriophage λ DNA in the presence of Ca²⁺ ions, followed by a brief heat shock [4]. This was referred to as the 'calcium chloride' (CaCl₂) method [5] and was subsequently adopted to impart antibiotic resistance into the competent *E. coli*. However, it was discovered that only 1–2% of the transformants survived in this method [6], making it inefficient for routine DNA manipu-

lation. Later, modifications such as the use of multiple heat-shock cycles [7] and variations of chemicals were tested. These resulted in several optimized versions of the original CaCl₂ method [8,9] that typically yield 'TrE' (transformation efficiencies) of 5×10^6 – 2×10^7 colonies forming units per microgram (cfu/ μ g) of DNA.

In 1983, Douglas Hanahan proposed a new method that yielded competency levels of 1 – 5×10^8 cfu/ μ g across many *E. coli* strains [10,11]. However, his method was quite complex, combining several parameters such as the nature of chemicals, plasmids and labware. Subsequently, a simpler method based on the use of DMSO and PEG (polyethylene glycol) (replacing Ca²⁺ ions) and a short incubation of both bacteria and DNA on ice (replacing the heat-shock) was proposed. This method, commonly known as the DMSO method [12], produced up to 1×10^7 – 10^8 cfu/ μ g. However, to date there has not yet been a comprehensive study comparing the different methods across various strains of *E. coli*

Abbreviations used: CaCl₂, calcium chloride; CfU, colony-forming units; *E. coli*, *Escherichia coli*; FSB, frozen storage buffer; LB, Luria-Bertani (broth); PEG, polyethylene glycol; SOB, super optimal broth; SOC, super optimal broth with catabolite repression; TrE, transformation efficiency; TSB, transformation storage buffer.

¹ To whom any correspondence should be addressed (email samuelg@bii.a-star.edu.sg).

to determine the optimal conditions for maximizing the competency of the bacteria. Comparisons are complicated by issues such as the lack of consistency in the methods used to calculate TrE.

In order to address these issues, we examined the following four methods of producing competent bacteria: Hanahan's method [10], the DMSO method [12] and two variants of the CaCl₂ method [4,13]. Care was taken to ensure that conditions including aliquot volumes, TrE calculations, growth media, heat-shock duration, final resuspension volumes and plasmid were standardized for comparisons. These resulted in establishing the best method for each strain (we tested six different strains of *E. coli*), followed by optimization of other parameters such as growth media and heat-shock incubation times to produce bacteria with TrE levels that are statistically comparable with those reported commercially. To our knowledge, this is the first report of such a comprehensive study since that of Hanahan's (1983 and 1991).

EXPERIMENTAL

E. coli starting cultures:

Colonies of *E. coli* strains (DH5 α – Invitrogen, Cat no. 12297-016, XI-1 Blue – Stratagene, Cat no. 200247, SCS110 – Stratagene, Cat no. 200249, JM109 – Promega, Cat no. L2001, TOP10 – Invitrogen Cat no. C4040-10, BL21-(DE3)-PLyS – Invitrogen, Cat no. C6060-03) were isolated from commercial sources through plating on LB [Luria–Bertani (broth)] agar (Biopolis Shared Facilities, A*STAR). Single colonies were inoculated into 2 ml of relevant media overnight. Approximately, 100 μ l of the cultures were inoculated into 50 ml of pre-warmed respective media to allow growth to early log phase [OD₆₀₀ (optical density) reading of 0.3–0.5] [8,9] (Supplementary Table S1 at <http://www.biosciencerep.org/bsr/033/bsr033e086add.htm>), followed by immediate placement on ice for 15 min and centrifugation. The pellets were resuspended in method-specific buffers (see CaCl₂ [4,13], DMSO [12] and Hanahan's method [10] or supplementary data). All bacteria were grown at 37 °C with vigorous shaking at 200–220 rpm (Certomat BS-1, Satorius Stedim Biotech). All centrifugation steps were performed at 3200 g using Eppendorf, Model 5804R at 4 °C.

Comparison of the four methods on six *E. coli* strains

For comparisons, the following parameters were standardized: (1) Use of 100 μ l bacterial aliquots for transformation and TrE calculation; (2) Use of PUC18 plasmid only; (3) Use of LB media for growth and post heat-shock growth; (4) Use of 45 s heat-shock for transformation; and (5) Final resuspension volume of 4 ml from initial 50 ml of bacteria cultures.

MgCl₂–CaCl₂ method: adapted from the method of Sambrook and Russell [13]

Pelleted bacteria from 50 ml cultures were resuspended with gentle pipetting in 15 ml of 0.1 M MgCl₂ (formulated in de-

ionized water and autoclaved) (BDH, VWR International) and incubated on ice for 10 min. The bacteria were pelleted at 4k rpm at 4 °C for 10 min, resuspended in 15 ml of 0.1 M CaCl₂, and incubated on ice for 30 min. After spinning down, the supernatant were discarded, and the pellets gently resuspended in 4 ml (standardized volume) of 0.1 M CaCl₂ with 20% (v/v) glycerol solution and stored in 100 μ l aliquots at – 80 °C.

CaCl₂ method: adapted and modified from the method of Mandel and Higa [4]

Original protocols are in Supplementary Data; at <http://www.biosciencerep.org/bsr/033/bsr033e086add.htm>.

Pelleted bacterial strains from 50 ml cultures were resuspended with gentle pipetting in 25 ml (half the volume of the initial culture) of ice-cold 0.1 M CaCl₂ (BDH) (formulated in de-ionized water and autoclaved) and incubated on ice for 1 h. The bacteria suspensions were pelleted at 4k rpm at 4 °C for 10 min followed by gentle resuspension in 4 ml of 0.1 M CaCl₂ + 15% (v/v) glycerol and stored in 100 μ l aliquots at – 80 °C.

DMSO method: adapted and modified from the method of Chung and Miller [12]

Original protocols are in Supplementary Data; at <http://www.biosciencerep.org/bsr/033/bsr033e086add.htm>.

Pelleted bacteria strains from 50 ml cultures were resuspended in 4 ml of ice-cold TSB (transformation storage buffer: LB broth at pH 6.1, 10% (w/v) PEG4450, 5% (v/v) DMSO, 10 mM MgCl₂ and 10 mM MgSO₄, filter sterilized with 0.45 μ m filter) and incubated on ice for 30 min. The bacteria were then subsequently stored in 100 μ l aliquots at – 80 °C.

Hanahan's method: adapted and modified from Hanahan [10]

Detailed original protocols and FSB (frozen storage buffer) preparation are in Supplementary Data at <http://www.biosciencerep.org/bsr/033/bsr033e086add.htm>.

Pelleted bacterial strains from 50 ml culture were resuspended gently in 16.5 ml of FSB (10 mM CH₃CO₂K at pH 7.5, 45 mM MnCl₂, 10 mM CaCl₂, 0.1 M KCl, 3 mM [Co(NH₃)₆]Cl₃, 10% glycerol) and incubated on ice for 15 min. The bacteria were then pelleted at 4k rpm at 4 °C for 10 min, and resuspended in 4 ml of FSB. 140 μ l of DMSO was added twice in intervals of 5 min to the centre of the suspension with gentle swirling. The bacteria suspensions were stored in 200 μ l aliquots at – 80 °C.

Transformation protocol for the standardized methods

45 s heat-shock transformation – (modified from Stratagene's recommended protocol) – used for all methods except the DMSO method

100 μ l of competent bacteria were mixed with 1 μ l of control pUC18 plasmid [0.1 ng/ μ l in nuclease-free water (Agilent, 200231-42)] in cold 14 ml round bottomed tubes (BD, Product no. 352059) and incubated on ice for 30 min. A 42 °C heat-shock




Figure 1 Comparison of the four standardized chemical methods on six strains of *E. coli*

Bar chart showing the means and standard errors of the TrE obtained from the six strains of *E. coli* produced using the four different methods. Controls were performed by transforming competent bacteria with water, which did not yield any transformants.

of 45 s was performed, followed by immediate placement on ice for 2 min. 100 μ l of LB media were added to the bacterial suspensions before incubations at 37°C for 1 h. The entire aliquot was plated out on 1.5% (w/v) LB agar plates with 100 μ g/ml ampicillin (Goldbio, A-301-5) at 37°C overnight.

Transformation of competent bacterial (DMSO method):

100 μ l of thawed DMSO competent cells were transferred to ice-cold 14 ml round bottomed tubes (BD, Product no. 352059) and incubated with 0.1 ng/ μ l of control pUC18 plasmid (Agilent, 200231-42) on ice for 30 min. The cell suspensions were then allowed to grow in 0.9 ml of TSB with 20 mM of glucose at 37°C in vigorous shaking (speed 200–220) for 1 h. The cells were then plated on LB agar plates with 100 μ g/ml ampicillin and incubated overnight at 37°C.

Comparison of the culture media:

Optimization of culture media used for starter cultures

The various strains of *E. coli* were induced to be competent using the established best methods as described above, and varying the use of SOB (super optimal broth) [10] or LB [14] as the starting culture media. Heat shock was standardized to 45 s.

Comparison of the heat-shock incubation times for transformation

The incubation times of the heat shock for all the six strains were made using the exact transformation protocol as described

by Hanahan [10] (Supplementary data at <http://www.bioscirep.org/bsr/033/bsr033e086add.htm>), with the exception of varying the incubation to either 45 or 90 s in the 42°C heat bath.

Data collection and computation of TrE

All bacterial colonies on the plated agar were counted manually. The TrE were calculated according to the formula provided by Stratagene, and adjusted to aliquot volumes 100 μ l.

Fourfold concentration of optimally induced bacteria

The bacterial strains were concentrated prior to freezing and storage by virtue of a four-fold reduction in the volume of the final storage buffer used to resuspend the strains as per their optimized protocols.

Statistical computation and analysis

For comparison of the four methods on the six strains, both ANOVA and independent *t* tests were used. ANOVA test was performed to determine the reproducibility of the TrE within each method (Supplementary Table S2A at <http://www.bioscirep.org/bsr/033/bsr033e086add.htm>), as well as the differences between the four methods (Supplementary Table S2B at <http://www.bioscirep.org/bsr/033/bsr033e086add.htm>). Independent *t* test was used for the comparison of pair-wise method comparisons (Supplementary Table S2C at <http://www.bioscirep.org/bsr/>




Figure 2 Comparison of the use of LB or SOB as growth media

Bar chart showing the means, standard errors and *t* tests of TrE obtained from both CaCl_2 and Hanahan methods of chemical induction for the respective strains. (Top panel) Bar chart representing the three strains that responded best to Hanahan's method. (Bottom panel) Bar chart representing the three strains that responded best to CaCl_2 method. * denotes that $P < 0.05$; ** denotes that $P < 0.001$ for the *t* tests comparing the means. Refer to Supplementary Table S2 at <http://www.bioscirep.org/bsr/033/bsr033e086add.htm> for detailed statistical analysis. The *t* test showed that there was no significant difference between the use of SOB or LB media for DH5 α , SCS110, TOP10 and BL21 strains despite higher average TrE for SOB media.

033/bsr033e086add.htm), media (Figure 2), heat-shock incubation times (Figure 3), and four-fold concentration and neat (Figure 4). All statistical analysis was performed using SPSS ver. 17 (IBM) at a 95% confidence interval.

RESULTS AND DISCUSSION

Comparison of four different chemical methods of producing competent bacterial strains

TrE for the four different chemical methods mentioned above were tested using 100 μl aliquots, LB growth media, 4 ml final resuspension volumes and 45 s heat-shock duration (with the

exception of the DMSO method which uses 10–15 min ice incubation). All the methods yielded TrE that were significantly different from each other ($P < 0.05$) (see ANOVA results and independent *t* test results in Supplementary Tables S2B and S2C, respectively) with the exception of the DH5 α strain. To test for reproducibility within each method, ANOVA tests (see Supplementary Table S2A) were also performed. The results showed that all strains, again with the exception of DH5 α strain (produced using DMSO method; $F = 19.331$, $P = 0.004$, Supplementary Table S2A), were reproducible ($P > 0.05$).

We found that the standardized Hanahan's and CaCl_2 methods consistently produced higher TrE than the MgCl_2 – CaCl_2 or the DMSO methods across all the six strains of *E. coli* tested (Figure 1 and Supplementary Table S2). When transformed with pUC18 plasmid, Hanahan's method was most effective for DH5 α , XL-1




Figure 3 Comparison of the 45 and 90 s heat shock

Bar chart showing the means, standard error and t test results of 45 and 90 s incubation times performed across all strains of *E. coli* made using the optimized method with SOC. The absence of significance differences indicate that the 45 or 90 s heat-pulse yielded similar transformation efficiencies across all strains tested.




Figure 4 Comparison of four-fold concentration with the default final resuspension method in the optimized methods across the six *E. coli* strains

Bar chart showing the means, standard errors and t test results of the TrE obtained from optimally induced competent bacteria that were concentrated four-fold with that of default aliquots. * denotes that $P < 0.05$; ** denotes that $P < 0.001$ for the t tests comparing the means.

Blue and JM109 strains, while the CaCl_2 method was most effective for SCS110, TOP10 and BL21 strains (Figure 1). Surprisingly, repeated attempts (see Supplementary Table S1 for batches made) were still unsuccessful in reproducing the 1×10^7 – 10^8 cfu/ μg TrE reported by Chung and Miller [15] for the DMSO method. In fact, neither the SCS110 nor the BL21 strains yielded significant TrE levels for either the DMSO or the MgCl_2 – CaCl_2 methods (Figure 1). The DH5 α strain, produced with the DMSO method, failed to reach TrE levels (10^5 cfu/ μg) suitable for sub-cloning, but did produce better yields ($> 10^5$ cfu/ μg) with both the CaCl_2

methods, and these yields were statistically similar within the two variant methods ($P = 0.276$).

In agreement with reports suggesting genetic associations with sensitivity to the different methods of chemical induction [11,16], our analysis suggested that modifications of the *gal* operon were associated with the strains that responded better to the CaCl_2 method. In contrast, modifications of the *relA1* and *gyrA96* genes were associated with increased sensitivity to Hanahan's method (Supplementary Table S3 at <http://www.bioscierep.org/bsr/033/bsr033e086add.htm>). However, establishment of a

Table 1 Summary Table of the optimized methods and parameter for chemical induction of competency of the six *E. coli* strains

Table summarizing the various parameters for chemical induction of competency used.

<i>E. coli</i> strain	Best chemical competency induction method		P values			P values			Average TrE obtained after optimization	
	Growth media	t	df	(1-tailed)	Heat-shock duration	t	df	(2-tailed)		
DH5 α	Hanahan's method	No significant difference	0.038	10	0.485	No significant difference	1.258	6	0.255	9.31×10^7
XL-1 Blue	Hanahan's method	SOB > LB	4.280	16	<0.001	No significant difference	0.800	6	0.454	9.23×10^7
SCS110	CaCl ₂ method	No significant difference	1.330	6	0.116	No significant difference	0.243	6	0.816	3.03×10^5
JM109	Hanahan's method	LB > SOB	7.913	10	<0.001	No significant difference	1.622	6	0.156	2.26×10^7
TOP10	CaCl ₂ method	No significant difference	1.156	14	0.133	No significant difference	1.811	6	0.120	4.34×10^7
BL21	CaCl ₂ method	No significant difference	0.728	14	0.239	No significant difference	2.258	6	0.65	2.57×10^7

firm correlation between the genotype and sensitivity to the chemical induction methods would require more detailed studies.

Comparison of the use of LB and SOB media and their effects on TrE

We next examined the effect of LB [14] and SOB [10] as the starting growth media prior to competency induction. Both media are highly similar in composition, with the exception that SOB and its variant SOC (super optimal broth with catabolite repression) additionally contained MgCl₂, MgSO₄, KCl and glucose. Figure 2 shows that using SOB or LB for both growing and the final plated media yielded similar TrE for all the strains studied ($P > 0.05$, Figure 2), with the exception of JM109 and XL1-Blue. XL1-Blue was the only strain that responded significantly better to SOB ($P < 0.05$, Figure 2, top panel). Interestingly, JM109 achieved better TrE when grown in LB than in SOB [$t(12) = 3.130$, $P = 0.009$, Figure 2 top panel; the Hanahan method is normally associated with SOB]. This suggests that the additional chemicals in Hanahan's method affected the chemical induction methods for particular strains depending on their genotypes. Although Mg²⁺ ions have been demonstrated to be associated with improved competency [10], K⁺ ions present in SOB may also have a contributory effect.

Comparison of 45 and 90 s heat-shock protocols

The 45 s (Stratagene's recommended timing for their bacteria) was next compared with the 90 s heat-shock method [10]. We did not find significant differences between the two incubation timings for all the strains ($P > 0.05$, Figure 3), suggesting that 45 s incubations were sufficient for effective pUC18 transformation for all the six strains tested.

Boosting TrE levels through reduction of final resuspension buffer

Having established the optimum conditions (including growth media and heat-shock durations) for the six strains, we attempted

to increase TrE by increasing the density of bacteria in each aliquot. Our very first attempt, while following Hanahan's protocol exactly ([10], using 200 μ l aliquots for transformation), involved a four-fold reduction in volume of the final resuspension buffer (Figure 4). While not linear with concentration, we did obtain the minimum of a four-fold increase across all methods and strains tested (calculated in 100 μ l aliquot volumes for comparisons in Figure 4). To ensure comparability, all transformations were performed using pUC18 plasmid, and ANOVA tests were carried out to ensure reproducibility (Supplementary Table S4 at <http://www.bioscirep.org/bsr/033/bsr033e086add.htm>).

Issues with respect to the current formulae for TrE

One of the issues that make comparisons across methods difficult is the fact that there is no provision for inclusion of bacteria aliquot volumes used for transformations in the estimations of TrE (cfu/ μ g of DNA). This may affect the estimates since the actual bacterial numbers, as measured using OD_λ for log-phase determination, may vary. In addition, variations in bacterial numbers will also likely arise from the differences in final resuspension volumes. Thus, our study has demonstrated that increased bacterial density in the fixed volume aliquots can significantly boost the average TrE (Figure 4).

In summary, we propose that comparisons of TrE values across methods and strains need to ensure standardization of the following: (1) the aliquot volume used for transformation i.e. standardized to a fixed volume (e.g. 100 μ l); (2) the resuspension volumes of the final freezing buffer of the various protocols taking into account the initial culture volume; (3) the use of the same plasmid (e.g. pUC 18); and (4) the growth media used.

Conclusion

From our study comparing four commonly used chemical induction methods for producing competent bacteria, we have determined the optimal set of parameters for the most effective transformation strategy. We have obtained yields of competent

bacteria that are of high cloning TrE (see Table 1 for a summary of recommended parameters and methods for each of the strains). We have also demonstrated the need for the standardization of multiple factors when comparing methods including resuspension volume ratios, transformation aliquot volumes, plasmids and growth media.

AUTHOR CONTRIBUTION

Weng-Tat Chan performed the majority of the laboratory work, analysis and statistical tests and drafted the paper. Chandra Verma guided the intellectual content and critical revision for the paper. David Lane provided all of the facilities required for the study, the final revision of the paper, and advice on the intellectual content. Samuel Gan is the direct supervisor of Weng-Tat Chan, he conceived the study, the overall directions of the study, the formatting, and directed the data collection and writing of the paper.

ACKNOWLEDGEMENTS

We thank Elfin Boon-Swee Lim for his help in making some of the batches of CaCl₂ method.

FUNDING

This work was funded by the Joint Council Office, Agency of Science, Technology and Research, Singapore.

REFERENCES

- Griffith, F. (1928) The significance of pneumococcal types. *J. Hyg. (Lond.)* **27**, 113–159
- Michod, R. E., Wojciechowski, M. F. and Hoelzer, M. A. (1987) DNA repair and the evolution of transformation in the bacterium *Bacillus subtilis*. *Genetics* **118**, 31–39
- Bergmans, H. E., van Die, I. M. and Hoekstra, W. P. (1981) Transformation in *Escherichia coli*: stages in the process. *J. Bacteriol.* **146**, 564–570
- Mandel, M. and Higa, A. (1970) Calcium-dependent bacteriophage DNA infection. *J. Mol. Biol.* **53**, 159–162
- Cohen, S. N., Chang, A. C. and Hsu, L. (1972) Nonchromosomal antibiotic resistance in bacteria: genetic transformation of *Escherichia coli* by R-factor DNA. *Proc. Natl. Acad. Sci. U. S. A.* **69**, 2110–2114
- Weston, A., Humphreys, G. O., Brown, M. G. and Saunders, J. R. (1979) Simultaneous transformation of *Escherichia coli* by pairs of compatible and incompatible plasmid DNA molecules. *Mol. Gen. Genet.* **172**, 113–118
- Roychoudhury, A., Basu, S. and Sengupta, D. N. (2009) Analysis of comparative efficiencies of different transformation methods of *E. coli* using two common plasmid vectors. *Indian J. Biochem. Biophys.* **46**, 395–400
- Li, X., Sui, X., Zhang, Y., Sun, Y., Zhao, Y., Zhai, Y. and Wang, Q. (2010) An improvised calcium chloride method preparation and transformation of competent cells. *Afr. J. Biotechnol.* **9**, 8549–8554
- Tu, Z., He, G., Li, K. X., Chen, M. J., Chang, J., Chen, L., Yao, Q., Liu, D. P., Ye, H., Shi, J. and Wu, X. (2005) An improvised system for competent cell preparation and high efficiency plasmid transformation using different *Escherichia coli* strains. *Electron. J. Biotechnol.* **8**
- Hanahan, D. (1983) Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**, 557–580
- Hanahan, D., Jessee, J. and Bloom, F. R. (1991) Plasmid transformation of *Escherichia coli* and other bacteria. *Methods Enzymol.* **204**, 63–113
- Chung, C. T. and Miller, R. H. (1988) A rapid and convenient method for the preparation and storage of competent bacterial cells. *Nucleic Acids Res.* **16**, 3580
- Sambrook, J. and Russell, D. W. (2001) *Molecular Cloning: a Laboratory Manual*, Cold Spring Harbor Laboratory Press, Volume 1
- Bertani, G. (1951) Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J. Bacteriol.* **62**, 293–300
- Chung, C. T., Niemela, S. L. and Miller, R. H. (1989) One-step preparation of competent *Escherichia coli*: transformation and storage of bacterial cells in the same solution. *Proc. Natl. Acad. Sci. U. S. A.* **86**, 2172–2175
- Ryu, J. and Hartin, R. J. (1990) Quick transformation in *Salmonella typhimurium* LT2. *BioTechniques* **8**, 43–45

Received 27 September 2013/13 November 2013; accepted 14 November 2013

Published as Immediate Publication 14 November 2013, doi 10.1042/BSR20130098



OPEN ACCESS

SUPPLEMENTARY DATA

A comparison and optimization of methods and factors affecting the transformation of *Escherichia coli*

Weng-Tat CHAN*, Chandra S. VERMA*†‡, David P. LANE‡§ and Samuel Ken-En GAN*¹

*Bioinformatics Institute, Agency for Science, Technology, and Research (A*STAR), Singapore 138671, †Department of Biological Sciences, National University of Singapore (NUS), Singapore 119077, ‡School of Biological Sciences, Nanyang Technological University (NTU), Singapore 639798, and §p53 Laboratory, Agency for Science, Technology, and Research (A*STAR), Singapore 138648

EXPERIMENTAL

Detailed methods of chemical induction of competency

Original CaCl₂ method (from Mandel and Higa [1])

Pelleted bacteria were resuspended with gentle pipetting in 25 ml (half the volume of the initial culture) of 0.1 M CaCl₂ (formulated in de-ionized water and autoclaved) (BDH, VWR International) and incubated on ice for 1 h. The bacteria suspension were pelleted at 4k rpm at 4 °C for 8 min (Eppendorf, Model 5804R), and the supernatants were discarded. Following which, the pellets were resuspended gently in 4 ml (standardized volume) of 0.1 M CaCl₂ with 15% (v/v) glycerol solution and stored in 100 μl aliquots at – 80 °C.

Original MgCl₂–CaCl₂ method (from Sambrook and Russell [2])

Pelleted bacteria were resuspended with gentle pipetting in 15 ml of 0.1 M MgCl₂ (formulated in de-ionized water and autoclaved) (BDH, VWR International) and incubated on ice for 10 min. The bacteria were pelleted at 4k rpm at 4 °C for 8 min (Eppendorf, Model 5804R), and the supernatants were discarded. Following which, the pellets were resuspended in 15 ml of 0.1 M CaCl₂, and incubated on ice for 30 min. After spinning down, the supernatant were discarded, and the pellets resuspended in 4 ml (standardized volume) of 0.1 M CaCl₂ with 20% (v/v) glycerol solution, and stored in 100 μl aliquots at – 80 °C.

Original DMSO method (from Chung and Miller [3])

Pelleted bacteria were gently resuspended in 10% volume of the initial culture of ice-cold TSB [LB broth at pH 6.1, 10% (w/v) PEG3350, 5% (v/v) DMSO, 10 mM MgCl₂ and 10 mM MgSO₄, filter sterilized with 0.45 μm filter] and incubated on ice for 30 min. The bacteria were stored in 100 μl aliquots at – 80 °C.

Original Hanahan's method (from Hanahan [4])

Pelleted bacteria were resuspended gently in 1/3 volume of initial starting culture (50 ml) of FSB [10 mM CH₃CO₂K at pH 7.5, 45 mM MnCl₂, 10 mM CaCl₂, 0.1 M KCl, 3 mM [Co(NH₃)₆]Cl₃, 10% (v/v) glycerol] and incubated on ice for 15 min. The bacteria were then pelleted at 4k rpm at 4 °C for 8 min using Eppendorf, Model 5804R centrifuge and resuspended with 4 ml of FSB. 3.5% (v/v) of DMSO was added twice in intervals of 5 min [final concentration at 7% (v/v)] to the center of the suspension with gentle swirling. The bacteria suspensions were stored in 200 μl aliquots at – 80 °C.

Preparation of FSB

1 M CH₃CO₂K stock solution was prepared using Milli-Q grade water, and equilibrated to pH 7.5 using KOH and filtered using 0.22 μm pore-size filters prior to freezing for storage. A 10 mM CH₃CO₂K solution was prepared from the stock solution with 10% (v/v) of glycerol added. The rest of the chemicals were added into the 10 mM CH₃CO₂K solution, and the pH was adjusted to 6.4 using 0.1 N of HCL. Extreme care was taken to ensure that the pH values did not fall below 6.4 or a new buffer using new reagents was made. The pH of the buffer was allowed to drift from 6.4 for a ~1–2 days before finally settling at 6.1–6.2. The buffer was sterilized using a 0.22 μm pore-size filter and stored at 4 °C in the dark. Care was taken to ensure that the glassware used for competent bacteria production were autoclaved and clean, and that the chemicals used were uncontaminated and recently purchased.

Detailed transformation protocols

DMSO method

100 μl of thawed DMSO competent cells were transferred to cold 14 ml round bottomed tubes (BD, Product no. 352059) and incubated with pUC18 DNA (Agilent, 200231-42) on ice for 30 min. The cell suspensions were then allowed to grow in 0.9 ml of

¹ To whom any correspondence should be addressed (email samuelg@bii.a-star.edu.sg).

**Table S1 OD readings of the batches of competent cells produced for comparison of the four methods**

Table showing the various OD₆₀₀ readings obtained from the initial starting culture grown for each batch of bacteria produced using the respective methods. The number of different batches produced using the different methods are represented by *n*.

Method	Strain	<i>n</i>	Batch 1 OD ₆₀₀ readings	Batch 2 OD ₆₀₀ readings	Batch 3 OD ₆₀₀ readings	Batch 4 OD ₆₀₀ readings
MgCl₂-CaCl₂	DH5α	4	0.404	0.297	0.337	0.337
	XL-1 Blue	4	0.364	0.357	0.32	0.32
	SCS110	4	0.301	0.367	0.301	0.301
	JM109	4	0.444	0.444	0.481	0.481
	TOP10	4	0.387	0.387	0.3	0.3
	BL21	4	0.443	0.443	0.358	0.358
DMSO	DH5α	4	0.337	0.306	0.337	0.337
	XL-1 Blue	4	0.32	0.356	0.32	0.32
	SCS110	4	0.301	0.355	0.301	0.031
	JM109	4	0.444	0.444	0.481	0.481
	TOP10	4	0.387	0.387	0.3	0.3
	BL21	4	0.443	0.443	0.412	0.412
Hanahan's (standardized)	DH5α	4	0.327	0.31	0.396	0.337
	XL-1 Blue	4	0.318	0.31	0.324	0.32
	SCS110	4	0.301	0.355	0.301	0.301
	JM109	4	0.444	0.444	0.481	0.481
	TOP10	4	0.387	0.387	0.3	0.3
	BL21	4	0.443	0.443	0.412	0.412
CaCl₂	DH5α	4	0.363	0.363	0.363	0.363
	XL-1 Blue	4	0.367	0.367	0.367	0.367
	SCS110	4	0.322	0.322	0.322	0.322
	JM109	4	0.368	0.368	0.368	0.368
	TOP10	4	0.324	0.324	0.324	0.324
	BL21	4	0.357	0.357	0.357	0.357

TSB with 20 mM of glucose, at 37 °C in vigorous shaking (speed 200–220) for 1 h. The cells were then plated on LB agar plates with 100 μg/ml ampicillin (Goldbio, A-301-5) and incubated overnight at 37 °C.

Recommended 45 s heat-shock protocol by Stratagene

100 μl of competent bacteria were mixed with control pUC18 plasmid DNA (Agilent, 200231-42) in 14-ml BD Falcon Polypropylene round-bottom tubes (Falcon 2059) and incubated on ice for 30 min. A 42 °C heat shock of 45 s was applied to the tubes, followed by immediate placement on ice for 2 min. 900 μl of SOC (SOB + 20 mM glucose) media were added to the bacteria suspensions and subsequently incubated at 37 °C for 1 h. The entire suspension was plated out on LB agar plates with

100 μg/ml ampicillin (Goldbio, A-301-5) at 37 °C overnight. All transformations of commercially purchased competent bacteria utilized this protocol.

Hanahan method for transformation

200 μl of competent bacteria were mixed with control pUC18 plasmid DNA in 14-ml BD Falcon Polypropylene round-bottom tubes and incubated on ice for 30 min. A 42 °C heat shock of 90 s was applied to the bacteria, followed by immediate placement on ice for 2 min. 900 μl of SOC media were added to the bacteria suspensions and subsequently incubated at 37 °C for 1 h. The entire suspension was plated out on LB agar plates with 100 μg/ml ampicillin (Goldbio, A-301-5) using disposable spreaders at 37 °C overnight.

Table S2 Detailed statistical analysis of the reproducibility and differences within and between the four chemical methods of induction and six strains of *E. coli***(A) ANOVA test of the differences between the four chemical methods across the six strains of *E. coli* studied**

Strain	Sum of squares	df	Mean square	F	P value
DH5 α	1.514×10^{14}	3	5.046×10^{13}	6.947	0.001
XL-1 Blue	2.155×10^{14}	3	7.184×10^{13}	8.321	0.000
SCS110	8.440×10^{10}	3	2.813×10^{10}	32.303	0.000
JM109	3.296×10^{13}	3	1.099×10^{13}	78.713	0.000
TOP10	2.602×10^{14}	3	8.673×10^{13}	27.350	0.000
BL21	3.105×10^{11}	3	1.035×10^{11}	23.395	0.000

The tests were performed at a 95% confidence interval. $P < 0.05$ indicates that the methods were statistically different from one another within the particular strain.

(B) Independent t test of the four methods performed for each particular strain

<i>E. coli</i> strain	Comparison of methods	t	df	P (1-tailed)	Mean difference	95% confidence interval	
						Lower	Upper
DH5 α	Hanahan > MgCl ₂ -CaCl ₂	2.30	14	0.019	4.82×10^6	3.30×10^5	9.30×10^6
	Hanahan > DMSO	2.62	15	0.010	5.15×10^6	9.68×10^5	9.34×10^6
	Hanahan > CaCl ₂	2.92	19	0.004	4.73×10^6	1.34×10^6	8.12×10^6
	CaCl₂ > MgCl₂-CaCl₂	0.61	19	0.276	8.79×10^4	-2.15×10^5	3.91×10^5
	CaCl ₂ > DMSO	3.14	20	0.003	4.24×10^5	1.42×10^5	7.07×10^5
	MgCl ₂ -CaCl ₂ > DMSO	12.94	15	<0.001	3.37×10^5	2.81×10^5	3.92×10^5
XL1-Blue	Hanahan > MgCl ₂ -CaCl ₂	3.01	22	0.003	4.95×10^6	1.54×10^6	8.37×10^6
	Hanahan > DMSO	3.13	21	0.003	5.39×10^6	1.81×10^6	8.97×10^6
	Hanahan > CaCl ₂	2.45	22	0.011	4.11×10^6	6.30×10^5	7.58×10^6
	CaCl ₂ > MgCl ₂ -CaCl ₂	2.59	22	0.008	8.46×10^5	1.69×10^5	1.52×10^6
	CaCl ₂ > DMSO	3.83	21	<0.001	1.28×10^6	5.85×10^5	1.98×10^6
	MgCl ₂ -CaCl ₂ > DMSO	6.27	21	<0.001	4.35×10^5	2.91×10^5	5.79×10^5
SCS110	CaCl ₂ > MgCl ₂ -CaCl ₂	5.90	15	<0.001	1.17×10^5	7.47×10^4	1.59×10^5
	CaCl ₂ > DMSO	5.99	15	<0.001	1.19×10^5	7.64×10^4	1.61×10^5
	CaCl ₂ > Hanahan	5.14	15	<0.001	1.03×10^5	6.00×10^4	1.45×10^5
	Hanahan > MgCl ₂ -CaCl ₂	6.14	14	<0.001	1.44×10^4	9.35×10^3	1.94×10^4
	Hanahan > DMSO	6.98	14	<0.001	1.61×10^4	1.12×10^4	2.11×10^4
	MgCl ₂ -CaCl ₂ > DMSO	4.63	14	<0.001	1.75×10^3	9.39×10^2	2.56×10^3
JM109	Hanahan > MgCl ₂ -CaCl ₂	11.24	15	<0.001	2.37×10^6	1.92×10^6	2.82×10^6
	Hanahan > DMSO	12.47	16	<0.001	2.46×10^6	2.04×10^6	2.87×10^6
	Hanahan > CaCl ₂	6.23	15	<0.001	1.63×10^6	1.07×10^6	2.19×10^6
	CaCl ₂ > MgCl ₂ -CaCl ₂	4.83	16	<0.001	7.44×10^5	4.18×10^5	1.07×10^6
	CaCl ₂ > DMSO	5.81	17	<0.001	8.28×10^5	5.28×10^5	1.13×10^6
	MgCl ₂ -CaCl ₂ > DMSO	2.15	17	0.023	8.41×10^4	1.55×10^3	1.67×10^5
TOP 10	CaCl ₂ > MgCl ₂ -CaCl ₂	5.18	24	<0.001	5.14×10^6	3.09×10^6	7.18×10^6
	CaCl ₂ > DMSO	4.59	20	<0.001	5.40×10^6	2.94×10^6	7.85×10^6
	CaCl ₂ > Hanahan	5.85	26	<0.001	5.39×10^6	3.50×10^6	7.28×10^6
	MgCl ₂ -CaCl ₂ > DMSO	2.89	22	0.004	2.61×10^5	7.37×10^4	4.49×10^5
	MgCl ₂ -CaCl ₂ > Hanahan	3.56	28	0.001	2.53×10^5	1.07×10^5	3.98×10^5
	Hanahan > DMSO	2.00	24	0.029	8.75×10^3	-2.96×10^2	1.78×10^4
BL21	CaCl ₂ > MgCl ₂ -CaCl ₂	6.68	19	<0.001	2.11×10^5	1.45×10^5	2.77×10^5
	CaCl ₂ > DMSO	5.75	17	<0.001	2.11×10^5	1.33×10^5	2.88×10^5
	CaCl ₂ > Hanahan	2.20	23	0.038	7.24×10^4	4.41×10^3	1.40×10^5
	Hanahan > MgCl ₂ -CaCl ₂	5.20	18	<0.001	1.38×10^5	8.24×10^4	1.94×10^5
	Hanahan > DMSO	4.48	16	<0.001	1.38×10^5	7.28×10^4	2.04×10^5
	MgCl ₂ -CaCl ₂ > DMSO	NA	-	-	-	-	-

The tests were performed at 95% confidence. Absence of a P value indicates the absence of transformants. $P < 0.05$ showed that the strains responded differently to the various methods, with the exception of DH5 α , which responded similarly to both CaCl₂ and MgCl₂-CaCl₂ methods (bold).

**Table S2 Continue****(C) ANOVA test of reproducibility for the four methods for the six *E. coli* strains**

Methods	<i>E. coli</i> strains	Sum of squares	df	Mean square	F	P values
MgCl₂-CaCl₂ method	DH5 α	1.593 $\times 10^{10}$	3	5.312 $\times 10^9$	0.999	0.479
	XL1-Blue	1.371 $\times 10^{11}$	3	4.569 $\times 10^{10}$	0.832	0.513
	SCS110	4750000.000	3	1583333.333	1.949	0.264
	JM109	1.779 $\times 10^{10}$	3	5.931 $\times 10^9$	0.370	0.779
	TOP10	2.232 $\times 10^{11}$	3	7.439 $\times 10^{10}$	0.895	0.477
	BL21	0.000	3	0.000	.	.
DMSO method	DH5α	5.337 $\times 10^9$	3	1.779 $\times 10^9$	19.331	0.004
	XL1-Blue	5.879 $\times 10^8$	3	1.960 $\times 10^8$	1.075	0.419
	SCS110	0.000	3	0.000	.	.
	JM109	4.903 $\times 10^8$	3	1.634 $\times 10^8$	0.039	0.989
	TOP10	1.055 $\times 10^7$	3	3515625.000	0.783	0.546
	BL21	0.000	3	0.000	.	.
Hanahan's method	DH5 α	5.040 $\times 10^{11}$	3	1.680 $\times 10^{11}$	1.522	0.338
	XL1-Blue	8.225 $\times 10^{13}$	3	2.742 $\times 10^{13}$	0.798	0.529
	SCS110	1.174 $\times 10^8$	3	3.913 $\times 10^7$	0.862	0.530
	JM109	1.487 $\times 10^{12}$	3	4.956 $\times 10^{11}$	1.588	0.325
	TOP10	4.000 $\times 10^7$	3	1.333 $\times 10^7$	1.600	0.285
	BL21	1.577 $\times 10^{10}$	3	5.256 $\times 10^9$	0.926	0.471
CaCl₂ method	DH5 α	1.778 $\times 10^{11}$	3	5.926 $\times 10^{10}$	0.333	0.802
	XL1-Blue	3.090 $\times 10^{12}$	3	1.030 $\times 10^{12}$	0.792	0.532
	SCS110	1.452 $\times 10^{10}$	3	4.840 $\times 10^9$	2.321	0.192
	JM109	3.008 $\times 10^{11}$	3	1.003 $\times 10^{11}$	0.383	0.770
	TOP10	4.669 $\times 10^{13}$	3	1.556 $\times 10^{13}$	1.192	0.373
	BL21	2.581 $\times 10^{10}$	3	8.603 $\times 10^9$	1.876	0.204

The tests were performed at a 95% confidence level. $P > 0.05$ indicates that the various batches of bacteria were not different from one another (i.e. reproducible within the same method). Absence of P values indicates that there were no transformants observed. The bolded regions shows where $P < 0.05$ (i.e. the batches are not reproducible within the same method).

Table S3 Summary of the genotypes for the six *E. coli* strains

Genotypes were retrieved from: DH5 α – Invitrogen, Cat # 12297-016, XL-1 Blue – Stratagene, Cat # 200247, SCS110 – Stratagene, Cat # 200249, JM109 – Promega, Cat # L2001, TOP10 – Invitrogen, Cat # C4040-10, and BL21 – Invitrogen, Cat # C6060-03. Common modified genes were grouped accordingly to the methods that the strains responded best to.

<i>E. coli</i> strains	Hanahan's method			CaCl ₂ method		
	DH5 α	XL-1 Blue	JM109	TOP10	SCS110	BL21
Genotype modifications exclusive to <i>E. coli</i> strains sensitive to the method of production	<i>relA1</i>	<i>relA1</i>	<i>relA1</i>	<i>galU/galk</i>	<i>galk/galT</i>	<i>gal</i>
Genotype modification shared across the multiple strain of <i>E. coli</i> studied	<i>gyrA96</i> F-	<i>gyrA96</i>	<i>gyrA96</i> [F' <i>traD36, proAB, laqlqZ</i> Δ M15]	F-	[F' <i>traD36 proAB lac^qZ</i> Δ M15]	F-
	<i>recA1</i>	<i>recA1</i>	<i>recA1</i>	<i>recA1</i>		
	<i>endA1</i>	<i>endA1</i>	<i>endA1</i>	<i>endA1</i>	<i>endA1</i>	
	<i>hsdR17</i> (rk-, mk+)	<i>hsdR17</i>	<i>hsdR17</i> (rk-, mk+)	Δ (<i>mrr-hsdRMS-mcrBC</i>)		<i>hsdSB</i> (rB-, mB-)
	Δ (<i>lacZ</i>)M15	<i>lac</i>	Δ (<i>lac-proAB</i>)	Δ <i>lacX74</i>	<i>lacY</i>	
	<i>thi-1</i>	<i>thi-1</i>	<i>thi-1</i>		<i>thi-1</i>	
		<i>supE44</i>	<i>supE44</i>	(<i>StrR</i>)		(<i>StrR</i>)
					<i>supE44D</i>	
					<i>dcm</i>	<i>dcm</i>
	Φ 80		Φ 80 <i>lacZ</i> Δ M15			
Genotype modifications exclusive to individual strains	<i>fhuA2</i>			<i>mcrA</i>	<i>rpsL</i>	<i>ompT</i>
	Δ (<i>argF-lacZ</i>)			<i>araD139</i>	<i>thr</i>	(DE3)
	<i>U169</i>			Δ (<i>ara leu</i>)	<i>leu</i>	pLysS
	<i>phoA</i>			7697	<i>ara</i>	(CamR)
	<i>glnV44</i>			<i>rpsL</i>	<i>tonA</i>	
				<i>nupG</i>	<i>tsx</i>	
					<i>dam</i>	
					(<i>lac-proAB</i>)	

Table S4 ANOVA test for reproducibility of optimized competent bacteria

Table showing ANOVA of the different batches of bacteria produced using the optimized protocol with 4-fold concentration. Test was performed at 95 % confidence interval. $P > 0.05$ indicates that the batches were reproducible.

Optimized method for each strain	Sum of squares	df	Mean square	F	P values
DH5 α – Hanahan	2.632×10^{13}	1	2.632×10^{13}	0.160	0.728
XL-1 Blue – Hanahan	9.610×10^{12}	1	9.610×10^{12}	0.153	0.733
SCS110 – CaCl ₂	2.756×1^9	1	2.756×10^9	0.139	0.745
JM109 – Hanahan LB	1.232×10^{12}	1	1.232×10^{12}	0.006	0.945
TOP10 – CaCl ₂	5.595×10^{13}	1	5.595×10^{13}	0.168	0.722
BL21 – CaCl ₂	1.722×10^{13}	1	1.722×10^{13}	0.552	0.535



REFERENCES

- 1 Mandel, M. and Higa, A. (1970) Calcium-dependent bacteriophage DNA infection. *J. Mol. Biol.* **53**, 159–162
- 2 Sambrook, J. and Russell, D. W. (2001) *Molecular Cloning: a Laboratory Manual*, Cold Spring Harbor Laboratory Press, Volume 1
- 3 Chung, C. T. and Miller, R. H. (1988) A rapid and convenient method for the preparation and storage of competent bacterial cells. *Nucleic Acids Res.* **16**, 3580
- 4 Hanahan, D. (1983) Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**, 557–580

Received 27 September 2013/13 November 2013; accepted 14 November 2013

Published as Immediate Publication 14 November 2013, doi 10.1042/BSR20130098
