

This document is downloaded from DR-NTU, Nanyang Technological University Library, Singapore.

Title	Copper-catalyzed aerobic radical C-C bond cleavage of N-H ketimines
Author(s)	Tnay, Ya Lin; Ang, Gim Yean; Chiba, Shunsuke
Citation	Tnay, Y. L., Ang, G. Y., & Chiba, S. (2015). Copper-catalyzed aerobic radical C-C bond cleavage of N-H ketimines. <i>Beilstein Journal of Organic Chemistry</i> , 11, 1933-1943.
Date	2015
URL	<a href="http://hdl.handle.net/10220/40822">http://hdl.handle.net/10220/40822</a>
Rights	© 2015 The Author(s) (Published by Beilstein-Institut). This is an Open Access article under the terms of the Creative Commons Attribution License ( <a href="http://creativecommons.org/licenses/by/2.0">http://creativecommons.org/licenses/by/2.0</a> ), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Copper-catalyzed aerobic radical C–C bond cleavage of N–H ketimines

Ya Lin Tnay, Gim Yean Ang and Shunsuke Chiba\*§

## Full Research Paper

Open Access

### Address:

Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore

### Email:

Shunsuke Chiba\* - shunsuke@ntu.edu.sg

\* Corresponding author

§ Fax: +65-67911961; Tel: +65-65138013

### Keywords:

C–C bond cleavage; copper; ketimines; molecular oxygen; radical

*Beilstein J. Org. Chem.* **2015**, *11*, 1933–1943.

doi:10.3762/bjoc.11.209

Received: 23 July 2015

Accepted: 23 September 2015

Published: 19 October 2015

This article is part of the Thematic Series "Copper catalysis in organic synthesis".

Guest Editor: S. R. Chemler

© 2015 Tnay et al; licensee Beilstein-Institut.

License and terms: see end of document.

## Abstract


We report herein studies on copper-catalyzed aerobic radical C–C bond cleavage of N–H ketimines. Treatment of N–H ketimines having an  $\alpha$ -sp<sup>3</sup> hybridized carbon under Cu-catalyzed aerobic reaction conditions resulted in a radical fragmentation with C–C bond cleavage to give the corresponding carbonitrile and carbon radical intermediate. This radical process has been applied for the construction of oxaspirocyclohexadienones as well as in the electrophilic cyanation of Grignard reagents with pivalonitrile as a CN source.

## Introduction


Alkylideneaminy radicals (iminy radicals) have been utilized for the synthesis of azaheterocycles through an intramolecular N–C bond formation with the unsaturated systems [1–6]. As precursors of iminy radicals, readily available oximes and their derivatives have commonly been utilized. The generation of iminy radicals involves the homolysis of the N–O bond with radical initiators [7–15] (Scheme 1a) or using thermal [16–20] or photoreaction conditions [21–27] (Scheme 1b). An alternative route to iminy radicals is the single-electron reduction of oxime derivatives mediated by the appropriate lower valent transition

metals [28–32], electron-rich organic electron donors [33–38], or sensitized photolysis [39–42] (Scheme 1c). Although the oxidative generation of iminy radicals has also been reported, only iminoxyacetic acids have been used as the precursors when we started our studies on the oxidative reactions of N–H ketimines [43–49] (Scheme 1d).


The generation of iminy radicals by single-electron oxidation of N–H ketimines appears to be the most atom economical method, since only protons (H<sup>+</sup>) are produced along with the



iminyl radicals (Scheme 2). However, the instability of N–H ketimines [50] limits their use as starting materials.



Recently, we have studied the chemical reactivity of N–H ketimines towards copper-catalyzed aerobic reaction conditions [51–53]. In these studies, we employed the nucleophilic addition of Grignard reagents to carbonitriles followed by protonation as one of the methods for in situ generation of N–H ketimines, which were directly subjected to Cu-catalyzed aerobic reactions without further purification [54]. In this way, biaryl N–H ketimines generated from biaryl-2-carbonitriles were found to undergo copper-catalyzed aerobic aromatic C–H amination (Scheme 3a) [52] or 1,4-aminoxygenation (spirocyclization) (Scheme 3b) [51], affording phenanthridine deriva-



tives and azaspirocyclohexadienones, respectively, depending on the helical sense of the biaryl axis.

Herein we report applications of the copper-catalyzed aerobic C–C bond fission of iminyl radical species for the synthesis of oxaspirocyclohexadienones as well as the electrophilic cyana-tion of Grignard reagents using the readily available pivaloni-trile as a CN source.


## Results and Discussion

We further explored the reactivity of biaryl N–H ketimines under copper-catalyzed aerobic reaction conditions, aiming at the synthesis of 6-membered azaspirocycles such as **3a'** from carbonitrile **1a** having a quaternary sp<sup>3</sup>-hybridized carbon center at its  $\alpha$ -position. The reaction of *p*-tolylmagnesium bromide (**2a**) to carbonitrile **1a** proceeded smoothly in Et<sub>2</sub>O at 80 °C in a sealed tube, generating N–H ketimine **1aa** after protonation with MeOH. Subsequently, Cu(OAc)<sub>2</sub> (20 mol %), 1,10-phen (20 mol %) and DMF (to 0.1 M final concentration) were added and stirred at room temperature under an air atmosphere (Scheme 4). Interestingly, no formation of the desired 6-membered azaspirocycle **3a'** was observed, while oxaspirocyclohexadienone **3a**, biaryl alkene **4a**, and *p*-tolunitrile (**5a**) were isolated in 29%, 32%, and 86% yields, respectively (Scheme 4).

Oxaspirocyclohexadienone **3a** was formed through C–C bond cleavage from N–H ketimine intermediate **1aa** most likely via the corresponding iminyl radical, that undergoes radical frag-mentation to afford the corresponding C-radical and carboni-

trile **5a** [55]. Thus, it is deduced that oxaspirocyclohexadienone **3a** was formed through oxygenation of the putative C-radical. Moreover, in this transformation, the cyano group of carboni-trile **1a** is transferred to Grignard reagent **2a** to afford *p*-toluni-trile (**5a**). Considering the importance of carbonitriles in organic synthesis [56], we postulated that the cyano group transfer from simple carbonitriles onto Grignard reagents could be realized using this strategy.

Based on these preliminary results (Scheme 4), we next started to investigate the synthesis of oxaspirocyclohexadienones as the target product. Using carbonitrile **1a** and *p*-tolylmagnesium bromide (**2a**), an optimization of the reaction conditions was conducted (Table 1). Increasing the amount of the additive 1,10-phenanthroline to 40 mol % slightly improved the yield, giving **3a** in 40% yield (Table 1, entry 1). The use of 2,2'-bipyridine (bpy) provided a comparable result (Table 1, entry 2), while performing the reaction in the presence of 1,4-diazabicyclo[2.2.2]octane (DABCO) led to lower yields of spirodienone **3a** (Table 1, entry 3). We therefore decided to proceed with 1,10-phenanthroline as the optimal ligand and subsequently tested different Cu(II) and Cu(I) salts (Table 1, entries 4–7). The best results were obtained using 40 mol % CuI which provided **3a** in 46% isolated yield (Table 1, entry 7). A reduction of the catalyst loading to 20 mol % slightly lowered the yield of **3a** (Table 1, entry 9) and a stoichiometric amount of catalyst did not significantly improve the yield of **3a** (Table 1, entry 10). Performing the reaction under an O<sub>2</sub> atmosphere did also not increase the yield of **3a** (Table 1, entry 8).



**Table 1:** Optimization of reaction conditions: oxaspirocyclohexadienone synthesis.<sup>a</sup>




Entry	Cu salt [mol %]	Additive [mol %]	Atmosphere	Yield [%] <sup>b</sup>		
				3a	4a	5a
1	Cu(OAc) <sub>2</sub> [20]	1,10-phen [40]	air	35 (40)	32 (31)	85 (86)
2	Cu(OAc) <sub>2</sub> [20]	bpy [40]	air	32	36	80
3	Cu(OAc) <sub>2</sub> [20]	DABCO [40]	air	10	21	87
4	CuCl <sub>2</sub> [20]	1,10-phen [40]	air	27	22	78
5	CuCl [20]	1,10-phen [40]	air	37	28	96
6	[20]	1,10-phen [40]	air	40 (40) <sup>c</sup>	34 (27) <sup>c</sup>	95 (90) <sup>c</sup>
7	CuI [20]	1,10-phen [40]	air	42 (46) <sup>c</sup>	34 (28) <sup>c</sup>	92 (78) <sup>c</sup>
8	CuI [20]	1,10-phen [40]	O <sub>2</sub>	43	37	99
9	CuI [10]	1,10-phen [20]	air	37	34	95
10	CuI [100]	1,10-phen [100]	air	46	30	93

<sup>a</sup>All reactions were carried out using 0.5 mmol of biaryl carbonitrile **1a** with 1.3 equiv of Grignard reagent **2a** in Et<sub>2</sub>O (0.5 mL) at 80 °C (sealed tube) for 4 h followed by the addition of MeOH (60 μL, 3.0 equiv), DMF (5 mL), Cu catalyst and additive and subsequent stirring for 4 h at rt: 1,10-phen = 1,10-phenanthroline; bpy = 2,2'-bipyridine; DABCO = 1,4-diazabicyclo[2.2.2]octane. <sup>b</sup>Crude yields determined by <sup>1</sup>H NMR based on 1,1,2,2-tetrachloroethane as an internal standard. <sup>c</sup>Isolated yields are given in parentheses.

A proposed reaction mechanism for the formation of oxaspirocyclohexadienone **3a**, alkene **4a**, and *p*-tolunitrile (**5a**) are depicted in Scheme 5. Single-electron oxidation of N–H ketimine **1aa** with higher valent Cu(II) species generated under the aerobic reaction conditions forms iminyl radical species **A**, that undergoes β-carbon fragmentation to give *p*-tolunitrile (**5a**) and biaryl-2-isopropyl radical **B** (Scheme 5a). The aerobic oxygenation of C-radical **B** affords peroxy radical **C**, that is presumably reduced by Cu(I) species through the Fenton-type mechanism [57] to give alkoxy radical **D** [58]. Subsequent spirocyclization of the alkoxy radical **D** onto the benzene ring affords cyclohexadienyl radical **F**, oxygenation of which followed by C=O bond formation finally provides the oxaspirocyclohexadienone product **3a**. Whereas, the oxidation of the benzylic radical **B** by the existing Cu(II) species to carbocation **G** and subsequent E1-type elimination of a proton provides biaryl alkene **4a** (Scheme 5b). The presence of alkoxy radical **D** in the reaction process could be further supported by the reaction of biaryl hydroperoxide **6**, which could be converted into the alkoxy radical **D** under copper-catalyzed aerobic reaction conditions [58]. Indeed treatment of hydroperoxide **6** under the standard reaction conditions afforded **3a** in 24% yield along with biaryl alcohol **7** in 25% yield (Scheme 5c).

While formation of biaryl alkene **4a** could not be avoided at this moment, it is still appealing to probe the potential utility of the current transformation toward the synthesis of oxaspirocyclohexadienones **3**. Therefore, the reactions of carbonitriles **1b–d** were examined under the current best reaction conditions (Table 1, entry 8) using *p*-tolyl Grignard reagent **2a** (Table 2). Carbonitrile **1b**, having methyl groups in ortho-positions on both aryl rings of the biaryl moiety, underwent the C–C bond fission process smoothly to afford oxaspirocyclized product **3b** and biaryl alkene **4b** in 48% and 35% yields, respectively, along with *p*-tolunitrile (**5a**) in 90% yield (Table 2, entry 1). The reactions of N–H ketimines having cyclopentyl and tetrahydropyranyl rings derived from nitriles **1c** and **1d**, respectively, afforded tricyclic oxaspirocyclohexadienones **3c** and **3d** in moderate yields along with the corresponding alkenes **4c** and **4d** as well as *p*-tolunitrile (**5a**) (Table 2, entries 2 and 3).

On the other hand, the reaction starting from carbonitrile **1e** having a strained cyclobutane ring did not form the desired oxaspirocyclohexadienone (Scheme 6). Instead, γ-bromoketone **8e** was isolated in 44% yield along with nitrile **5a** in 80% yield. The formation of γ-bromoketone **8e** is most likely caused by radical ring opening from the transient cyclobutoxy radical **I**,

(a) Formation of oxospirocyclohexadienone **3a**(b) Formation of alkene **4a**(c) The reaction of hydroperoxide **6****Scheme 5:** Proposed reaction mechanisms for the formation of **3a**, **4a** and **5a**, and the reaction of hydroperoxide **6**.

which is driven by releasing ring strain of the cyclobutyl ring. The resulting  $\gamma$ -keto radical **J** subsequently undergoes radical bromination to form **8e** [59]. This result unambiguously supports the presence of the alkoxy radical intermediate during oxaspirocyclohexadienone formation in our mechanistic proposal.

Next, we turned our attention to apply the present copper-catalyzed aerobic C–C bond fission process in the electrophilic

cyanation of Grignard reagents. As carbonitriles are omnipresent components in various natural products, dyes and potent pharmaceutical drugs [60–62], new and versatile routes towards this substance class are always desirable. Conventional methods to install the cyano group on aryl rings such as the Rosenmund–von Braun reaction [63] or the Sandmeyer reaction [64] require the use of stoichiometric amounts of toxic metal cyanides (such as  $\text{CuCN}$ ) as the “CN” anion source [65]. Therefore, an employment of aliphatic carbonitriles which are

**Table 2:** Substrate scope: oxaspirocyclohexadienone synthesis.<sup>a</sup>

Entry	Substrate	Products <sup>b</sup>
1		
2		
3		




<sup>a</sup>All reactions were carried out using 0.5 mmol of biaryl carbonitrile **1** with 1.3 equiv of Grignard reagents **2a** in Et<sub>2</sub>O (0.5 mL) at 80 °C (sealed tube) for 4 h followed by the addition of MeOH (60 μL, 3.0 equiv), DMF (5 mL), CuI (20 mol %) and 1,10-phen (40 mol %) under ambient air at rt. 1,10-phen = 1,10-phenanthroline. <sup>b</sup>Isolated yields. <sup>c</sup><sup>1</sup>H NMR crude yields based on 1,1,2,2-tetrachloroethane as an internal standard.

less toxic and easier to handle (such as acetonitrile [66-70], benzyl cyanide [71-76], and malononitrile [77,78]) for the “CN” surrogates has been developed more recently [79]. We envisioned that the readily available pivalonitrile (**1f**) could be a potential CN source for the electrophilic cyanation [80-86] of Grignard reagents using the present protocol (Scheme 7).

Thus pivalonitrile (**1f**) was reacted with 2-naphthylmagnesium bromide (**2b**) for formation of the corresponding N–H ketimine **1fb**, which was subsequently treated with 10 mol % of Cu(OAc)<sub>2</sub> under an O<sub>2</sub> atmosphere (Scheme 8). As expected, formation of 2-naphthonitrile (**5b**) was observed in 79% yield. On the other hand, the reaction in the absence of O<sub>2</sub> (under an

Ar atmosphere) provided only 2-pivaloylnaphthalene (**9b**) in 83% yield formed through hydrolysis of unreacted N–H imine **1fb** during the aqueous work-up. Therefore molecular oxygen is indispensable to achieve the present cyanation through the C–C bond cleavage of the N–H ketimine. It was found that use of CuBr<sub>2</sub> as the catalyst resulted in formation of **5b** in higher yield (86%).

We next examined the substrate scope using different Grignard reagents (Table 3). The electrophilic cyanation proceeded smoothly even with sterically bulky Grignard reagents such as 1-naphthyl- and 2,4,6-trimethylphenyl Grignard reagents (for **5c** and **5d**) (Table 3, entries 1 and 2). Electron-rich aryl Grignard

Scheme 6: Formation of bromoketone **6e**.Scheme 7: Electrophilic cyanation of Grignard reagents with pivalonitrile (**1f**).

copper salts	atmosphere	time	<b>5b</b>	<b>9b</b>
$\text{Cu(OAc)}_2$	$\text{O}_2$	8 h	79%	0%
$\text{Cu(OAc)}_2$	Ar	24 h	0%	83%
$\text{CuBr}_2$	$\text{O}_2$	6 h	86%	0%

Scheme 8: Electrophilic cyanation with pivalonitrile (**1e**).



**Table 3:** Scope of the reaction using different Grignard reagents.<sup>a</sup>

Entry	R-MgBr (2)	Time x/y (h)	Product 5 <sup>b</sup>
1	 2c	12/18	 5c 76%
2 <sup>c</sup>	 2d	56/19	 5d 81%
3 <sup>c</sup>	 2e	50/18	 5e 67%
4	 2f	22/10	 5f 74%
5	 2g	48/24	 5g 81%
6	 2h	48/55	 5h 70%
7	 2i	48/24	 5i 63% <sup>d</sup>
8	 2j	48/48	 5j 47%

<sup>a</sup>Unless otherwise noted, the reactions were carried out using 1 mmol of pivalonitrile (**1f**) with 1.3 equiv of Grignard reagents **2** in Et<sub>2</sub>O (1 mL) at 60 °C (sealed tube) for the time x followed by the addition of MeOH (120 μL), DMF (10 mL), and CuBr<sub>2</sub> (10 mol %), and the mixture was stirred at 80 °C for time y under an O<sub>2</sub> atmosphere. <sup>b</sup>Isolated yields. <sup>c</sup>The reaction was conducted using Cu(OAc)<sub>2</sub> (10 mol %) as the catalyst. <sup>d</sup><sup>1</sup>H NMR yield.

reagents could also be used to give the corresponding benzonitriles **5e**, **5f** and **5g** in good yields (Table 3, entries 3–5). The reaction also proceeded with chlorinated substrates leaving the C–Cl bond intact (for **5h**) (Table 3, entry 6). Thiophen-2-

carbonitrile (**5i**) was also prepared in 63% yield. The present method could also be applied for the cyanation of the primary alkyl Grignard reagent, phenethylmagnesium bromide (for **5j**), albeit the product yield was moderate.

## Conclusion

In summary, we have demonstrated a copper-catalyzed aerobic generation of iminyl radicals from the corresponding N–H ketimines and their radical C–C bond fission. These processes could be applied for synthesis of oxaspirocyclohexadienones through spirocyclization of the transient alkoxy radicals generated by aerobic oxygenation of the resulting carbon radicals. With the present protocol, the electrophilic cyanation of Grignard reagents was also established using readily available pivalonitrile as a simple CN source.

## Supporting Information

### Supporting Information File 1

Full experimental details and analytical data.

[<http://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-11-209-S1.pdf>]

## Acknowledgements

This work was supported by funding from Nanyang Technological University and Singapore Ministry of Education (Academic Research Fund Tier 2: MOE2012-T2-1-014).

## References

- Kitamura, M.; Narasaka, K. *Bull. Chem. Soc. Jpn.* **2008**, *81*, 539–547. doi:10.1246/bcsj.81.539
- Narasaka, K.; Kitamura, M. *Eur. J. Org. Chem.* **2005**, 4505–4519. doi:10.1002/ejoc.200500389
- Stella, L. Nitrogen-Centered Radicals. In *Radicals in Organic Synthesis*; Renaud, P.; Sibi, M. P., Eds.; Wiley-VCH: Weinheim, Germany, 2001; Vol. 2, pp 407–426. doi:10.1002/9783527618293.ch45
- Mikami, T.; Narasaka, K. Generation of radical species by single electron transfer reactions and their application to the development of synthetic reactions. In *Advances in Free Radical Chemistry*; Zard, S. Z., Ed.; JAI: Stanford, 1999; Vol. 2, pp 45–88. doi:10.1016/S1874-5237(99)80004-X
- Fallis, A. G.; Brinza, I. M. *Tetrahedron* **1997**, *53*, 17543–17594. doi:10.1016/S0040-4020(97)10060-6
- Zard, S. Z. *Synlett* **1996**, *12*, 1148–1154. doi:10.1055/s-1996-5698
- El Kaim, L.; Meyer, C. *J. Org. Chem.* **1996**, *61*, 1556–1557. doi:10.1021/jo952173q
- Callier-Dublanquet, A.-C.; Quiclet-Sire, B.; Zard, S. Z. *Tetrahedron Lett.* **1995**, *36*, 8791–8794. doi:10.1016/0040-4039(95)01878-L
- Boivin, J.; Callier-Dublanquet, A.-C.; Quiclet-Sire, B.; Schiano, A.-M.; Zard, S. Z. *Tetrahedron* **1995**, *51*, 6517–6528. doi:10.1016/0040-4020(95)00319-4
- Callier, A.-C.; Quiclet-Sire, B.; Zard, S. Z. *Tetrahedron Lett.* **1994**, *35*, 6109–6112. doi:10.1016/0040-4039(94)88089-1
- Boivin, J.; Schiano, A.-M.; Zard, S. Z. *Tetrahedron Lett.* **1994**, *35*, 249–252. doi:10.1016/S0040-4039(00)76523-3
- Boivin, J.; Fouquet, E.; Zard, S. Z. *Tetrahedron* **1994**, *50*, 1757–1768. doi:10.1016/S0040-4020(01)80850-4
- Boivin, J.; Fouquet, E.; Zard, S. Z. *J. Am. Chem. Soc.* **1991**, *113*, 1055–1057. doi:10.1021/ja00003a057
- Boivin, J.; Fouquet, E.; Zard, S. Z. *Tetrahedron* **1994**, *50*, 1745–1756. doi:10.1016/S0040-4020(01)80849-8
- Boivin, J.; Fouquet, E.; Zard, S. Z. *Tetrahedron Lett.* **1990**, *31*, 3545–3548. doi:10.1016/S0040-4039(00)94438-1
- Cai, Y.; Jalan, A.; Kobosumi, A. R.; Castle, S. T. *Org. Lett.* **2015**, *17*, 488–491. doi:10.1021/ol5035047
- Portela-Cubillo, F.; Scott, J. S.; Walton, J. C. *Chem. Commun.* **2007**, 4041–4043. doi:10.1039/b712582h
- Portela-Cubillo, F.; Scott, J. S.; Walton, J. C. *J. Org. Chem.* **2008**, *73*, 5558–5565. doi:10.1021/jo800847h
- Blake, J. A.; Pratt, D. A.; Lin, S.; Walton, J. C.; Mulder, P.; Ingold, K. U. *J. Org. Chem.* **2004**, *69*, 3112–3120. doi:10.1021/jo049927y
- Lin, X.; Stien, D.; Weinreb, S. M. *Org. Lett.* **1999**, *1*, 637–640. doi:10.1021/ol990720e
- Faulkner, A.; Race, N. J.; Scott, J. S.; Bower, J. F. *Chem. Sci.* **2014**, *5*, 2416–2421. doi:10.1039/c4sc00652f
- Bingham, M.; Moutrille, C.; Zard, S. Z. *Heterocycles* **2014**, *88*, 953–960. doi:10.3987/COM-13-S(S)94
- Portela-Cubillo, F.; Lymer, J.; Scanlan, E. M.; Scott, J. S.; Walton, J. C. *Tetrahedron* **2008**, *64*, 11908–11916. doi:10.1016/j.tet.2008.08.112
- Alonso, R.; Campos, P. J.; Rodríguez, M. A.; Sampedro, D. *J. Org. Chem.* **2008**, *73*, 2234–2239. doi:10.1021/jo7025542
- Alonso, R.; Campos, P. J.; García, B.; Rodríguez, M. A. *Org. Lett.* **2006**, *8*, 3521–3523. doi:10.1021/ol061258i
- Gagosz, F.; Zard, S. Z. *Synlett* **1998**, 1978–1980.
- Boivin, J.; Fouquet, E.; Zard, S. Z. *Tetrahedron Lett.* **1991**, *32*, 4299–4302. doi:10.1016/S0040-4039(00)92153-1
- Zhao, M.-N.; Liang, H.; Ren, Z.-H.; Guan, Z.-H. *Synthesis* **2012**, *44*, 1501–1506. doi:10.1055/s-0031-1290779
- Tanaka, K.; Kitamura, M.; Narasaka, K. *Bull. Chem. Soc. Jpn.* **2005**, *78*, 1659–1664. doi:10.1246/bcsj.78.1659
- Koganemaru, Y.; Kitamura, M.; Narasaka, K. *Chem. Lett.* **2002**, *31*, 784–785. doi:10.1246/cl.2002.784
- Boivin, J.; Schiano, A.-M.; Zard, S. Z.; Zhang, H. *Tetrahedron Lett.* **1999**, *40*, 4531–4534. doi:10.1016/S0040-4039(99)00720-0
- Boivin, J.; Schiano, A.-M.; Zard, S. Z. *Tetrahedron Lett.* **1992**, *33*, 7849–7852. doi:10.1016/S0040-4039(00)74760-5
- Kitamura, M.; Yoshida, M.; Kikuchi, T.; Narasaka, K. *Synthesis* **2003**, *15*, 2415–2426. doi:10.1055/s-2003-42439
- Yoshida, M.; Kitamura, M.; Narasaka, K. *Bull. Chem. Soc. Jpn.* **2003**, *76*, 2003–2008. doi:10.1246/bcsj.76.2003
- Yoshida, M.; Kitamura, M.; Narasaka, K. *Chem. Lett.* **2002**, *31*, 144–145. doi:10.1246/cl.2002.144
- Uchiyama, K.; Hayashi, Y.; Narasaka, K. *Tetrahedron* **1999**, *55*, 8915–8930. doi:10.1016/S0040-4020(99)00453-6
- Uchiyama, K.; Ono, A.; Hayashi, Y.; Narasaka, K. *Bull. Chem. Soc. Jpn.* **1998**, *71*, 2945–2955. doi:10.1246/bcsj.71.2945
- Ono, A.; Uchiyama, K.; Hayashi, Y.; Narasaka, K. *Chem. Lett.* **1998**, *27*, 437–438. doi:10.1246/cl.1998.437
- Jiang, H.; An, X.; Tong, K.; Zheng, T.; Zhang, Y.; Yu, S. *Angew. Chem., Int. Ed.* **2015**, *54*, 4055–4059. doi:10.1002/anie.201411342
- McBurney, R. T.; Slawin, A. M. Z.; Smart, L. A.; Yu, Y.; Walton, J. C. *Chem. Commun.* **2011**, *47*, 7974–7976. doi:10.1039/c1cc12720a
- Kitamura, M.; Mori, Y.; Narasaka, K. *Tetrahedron Lett.* **2005**, *46*, 2373–2376. doi:10.1016/j.tetlet.2005.02.062
- Mikami, T.; Narasaka, K. *Chem. Lett.* **2000**, *29*, 338–339. doi:10.1246/cl.2000.338

43. Forrester, A. R.; Gill, M.; Meyer, C. J.; Sadd, J. S.; Thomson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1979**, 606–611. doi:10.1039/p19790000606
44. Forrester, A. R.; Gill, M.; Sadd, J. S.; Thomson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1979**, 612–615. doi:10.1039/p19790000612
45. Forrester, A. R.; Gill, M.; Thomson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1979**, 616–620. doi:10.1039/p19790000616
46. Forrester, A. R.; Gill, M.; Thomson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1979**, 621–631. doi:10.1039/p19790000621
47. Forrester, A. R.; Gill, M.; Napier, R. J.; Thomson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1979**, 632–636. doi:10.1039/p19790000632
48. Forrester, A. R.; Gill, M.; Meyer, C. J.; Thomson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1979**, 637–642. doi:10.1039/p19790000637
49. Forrester, A. R.; Napier, R. J.; Thomson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1981**, 984–987. doi:10.1039/p19810000984
50. Griller, D.; Mendenhall, G. D.; Van Hoof, W.; Ingold, K. U. *J. Am. Chem. Soc.* **1974**, *96*, 6068–6070. doi:10.1021/ja00826a018
51. Tnay, Y. L.; Chen, C.; Chua, Y. Y.; Zhang, L.; Chiba, S. *Org. Lett.* **2012**, *14*, 3550–3553. doi:10.1021/ol301583y
52. Zhang, L.; Ang, G. Y.; Chiba, S. *Org. Lett.* **2010**, *12*, 3682–3685. doi:10.1021/ol101490n
53. Chiba, S.; Zhang, L.; Lee, J.-Y. *J. Am. Chem. Soc.* **2010**, *132*, 7266–7267. doi:10.1021/ja1027327
54. Pickard, P. L.; Tolbert, T. L. *J. Org. Chem.* **1961**, *26*, 4886–4888. doi:10.1021/jo01070a025
55. Chiba, S.; Zhang, L.; Ang, G. Y.; Hui, B. W.-Q. *Org. Lett.* **2010**, *12*, 2052–2055. doi:10.1021/ol100522z
56. Paquette, L. A. *Comprehensive Organic Transformations*; VCH Publishers, Inc./VCH Verlagsgesellschaft: New York/Weinheim, 1989. doi:10.1002/ange.19901020837
57. Rachmilovich-Calis, S.; Masarwa, A.; Meyerstein, N.; Myerstein, D.; van Eldik, R. *Chem. – Eur. J.* **2009**, *15*, 8303–8309. doi:10.1002/chem.200802572
58. Too, P. C.; Tnay, Y. L.; Chiba, S. *Beilstein J. Org. Chem.* **2013**, *9*, 1217–1225. doi:10.3762/bjoc.9.138
59. Sanjaya, S.; Chiba, S. *Tetrahedron* **2011**, *67*, 590–596. doi:10.1016/j.tet.2010.11.060
60. Fatiadi, A. J. Preparation and Synthetic Applications of Cyano Compounds. In *Triple-Bonded Functional Groups*; Patai, S.; Rappaport, Z., Eds.; John Wiley & Sons, Ltd.: Chichester, UK, 1983; Vol. 2. doi:10.1002/9780470771709.ch9
61. Kleemann, A.; Engel, J.; Kutscher, B.; Reichert, D. *Pharmaceutical Substances: Synthesis, Patents, Applications*, 4th ed.; Thieme: Stuttgart, Germany, 2001.
62. Miller, J. S.; Manson, J. L. *Acc. Chem. Res.* **2001**, *34*, 563–570. doi:10.1021/ar0000354
63. Lindley, J. *Tetrahedron* **1984**, *40*, 1433–1456. doi:10.1016/S0040-4020(01)91791-0
64. Galli, C. *Chem. Rev.* **1988**, *88*, 765–792. doi:10.1021/cr00087a004
65. Wen, Q.; Jin, J.; Zhang, L.; Luo, Y.; Lu, P.; Wang, Y. *Tetrahedron Lett.* **2014**, *55*, 1271–1280. doi:10.1016/j.tetlet.2014.01.032
66. Zhu, Y.; Zhao, M.; Lu, W.; Li, L.; Shen, Z. *Org. Lett.* **2015**, *17*, 2602–2605. doi:10.1021/acs.orglett.5b00886
67. Pan, C.; Jin, H.; Xu, P.; Liu, X.; Cheng, Y.; Zhu, C. *J. Org. Chem.* **2013**, *78*, 9494–9498. doi:10.1021/jo4014904
68. Kou, X.; Zhao, M.; Qiao, X.; Zhu, Y.; Tong, X.; Shen, Z. *Chem. – Eur. J.* **2013**, *19*, 16880–16886. doi:10.1002/chem.201303637
69. Song, R.-J.; Wu, J.-C.; Liu, Y.; Deng, G.-B.; Wu, C.-Y.; Wei, W.-T.; Li, J.-H. *Synlett* **2012**, 2491–2496. doi:10.1055/s-0032-1317191
70. Luo, F.-H.; Chu, C.-I.; Cheng, C.-H. *Organometallics* **1998**, *17*, 1025–1030. doi:10.1021/om970842f
71. Wen, Q.; Jin, J.; Mei, Y.; Lu, P.; Wang, Y. *Eur. J. Org. Chem.* **2013**, *2013*, 4032–4036. doi:10.1002/ejoc.201300052
72. Zhang, L.; Wen, Q.; Jin, J.; Wang, C.; Lu, P.; Wang, Y. *Tetrahedron* **2013**, *69*, 4236–4240. doi:10.1016/j.tet.2013.03.089
73. Luo, Y.; Wen, Q.; Wu, Z.; Jin, J.; Lu, P.; Wang, Y. *Tetrahedron* **2013**, *69*, 8400–8404. doi:10.1016/j.tet.2013.07.063
74. Yuen, O. Y.; Choy, P. Y.; Chow, W. K.; Wong, W. T.; Kwong, F. Y. *J. Org. Chem.* **2013**, *78*, 3374–3378. doi:10.1021/jo3028278
75. Wen, Q.; Jin, J.; Hu, B.; Lu, P.; Wang, Y. *RSC Adv.* **2012**, *2*, 6167–6169. doi:10.1039/c2ra20770b
76. Jin, J.; Wen, Q.; Lu, P.; Wang, Y. *Chem. Commun.* **2012**, *48*, 9933–9935. doi:10.1039/c2cc35046g
77. Reeves, J. T.; Malapit, C. A.; Buono, F. G.; Sidhu, K. P.; Marsini, M. A.; Sader, C. A.; Fandrick, K. R.; Busacca, C. A.; Senanayake, C. H. *J. Am. Chem. Soc.* **2015**, *137*, 9481–9488. doi:10.1021/jacs.5b06136
78. Jiang, Z.; Huang, Q.; Chen, S.; Long, L.; Zhou, X. *Adv. Synth. Catal.* **2012**, *354*, 589–592. doi:10.1002/adsc.201100836
79. Wen, Q.; Lu, P.; Wang, Y. *RSC Adv.* **2014**, *4*, 47806–47826. doi:10.1039/C4RA08675A
80. Anbarasan, P.; Neumann, H.; Beller, M. *Chem. – Eur. J.* **2011**, *17*, 4217–4222. doi:10.1002/chem.201003388
81. Anbarasan, P.; Neumann, H.; Beller, M. *Chem. – Eur. J.* **2010**, *16*, 4725–4728. doi:10.1002/chem.201000086
82. Hughes, T. V.; Cava, M. P. *J. Org. Chem.* **1999**, *64*, 313–315. doi:10.1021/jo981924w
83. Koo, J. S.; Lee, J. I. *Synth. Commun.* **1996**, *26*, 3709–3713. doi:10.1080/00397919608003787
84. Klement, I.; Lennick, K.; Tucker, C. E.; Knochel, P. *Tetrahedron Lett.* **1993**, *34*, 4623–4626. doi:10.1016/S0040-4039(00)60640-8
85. van Leusen, A. M.; Jagt, J. C. *Tetrahedron Lett.* **1970**, *11*, 967–970. doi:10.1016/S0040-4039(01)97880-3
86. Foulger, N. J.; Wakefield, B. J. *Tetrahedron Lett.* **1972**, *13*, 4169–4170. doi:10.1016/S0040-4039(01)94266-2

## License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<http://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at:  
[doi:10.3762/bjoc.11.209](https://doi.org/10.3762/bjoc.11.209)