

This document is downloaded from DR-NTU, Nanyang Technological University Library, Singapore.

Title	Transient free-surface waves due to impulsive motion of a submerged source
Author(s)	Shu, Jian Jun
Citation	Shu, J. J. (2006). Transient free-surface waves due to impulsive motion of a submerged source. <i>Underwater Technology</i> , 26(4), 97-101.
Date	2006
URL	http://hdl.handle.net/10220/7134
Rights	© 2006 Society for Underwater Technology. This is the author created version of a work that has been peer reviewed and accepted for publication by Underwater Technology, Society for Underwater Technology. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at DOI: http://dx.doi.org/10.3723/175605406782725023 .

This document is downloaded from DR-NTU, Nanyang Technological University Library, Singapore.

Title	[SCG] Transient free-surface waves due to impulsive motion of a submerged source
Author(s)	Shu, Jian Jun
Citation	
Date	2006
URL	http://hdl.handle.net/null
Rights	

Transient free-surface waves due to impulsive motion of a submerged source

Dr JIAN-JUN SHU

School of Mechanical & Aerospace Engineering, Nanyang Technological University, Singapore

Abstract

The problem of a viscous flow past a submerged source starting from rest and moving with a constant velocity, below and parallel to a free surface, is examined and asymptotic expressions for long-time evolution of free-surface elevation are obtained. The results show explicitly the viscous effect on the free-surface elevation and no surface tension effect on the asymptotic wave.

Keywords: Transient free-surface waves, Kelvin wakes, wavefronts, surface tension

1. Introduction

The impulsive motion of a submerged source generates transient waves on the free surface of a viscous fluid. Lord Kelvin¹ ignored the free-surface shear stress and developed a theory to determine the kinematics and dynamics of the steady free-surface waves generated by a moving source with constant velocity in an inviscid fluid of infinite depth. It was found that the steady free-surface wave pattern consists of a series of so-called diverging and transverse waves. The diverging waves spread on each side of the moving source at an acute angle in relation to the source's moving direction, whereas the transverse waves move in the same direction as the moving source. The two wave systems intersect along the so-called 'cusp locus' on both sides of the moving source. The angle between this line and the source's moving direction can be calculated as $19^{\circ}28'$. The lines constitute the outer edge of the so-called Kelvin wake. It was also deduced that the diverging waves have a propagation direction of $35^{\circ}16'$ compared to the moving direction at a certain distance from the source's navigation route. Havelock² studied the long-time evolution of a free surface due to an impulsive motion of a cylinder with a constant velocity in an inviscid fluid. Wehausen and Laitone,³ Chan and Chwang⁴ and the author⁵ enriched the theory with the consideration of viscous effect.

In the present paper the analytical approach is extended to the transient interaction of impulsive motion of a submerged source with a free surface. It is interesting to study the viscous and surface tension

effects on a point force moving beneath a free surface. The point force solution can be used to construct solutions for a submerged source of any shape.

2. Mathematical formulation

It is considered that a point force is submerged in a viscous incompressible fluid that occupies initially the lower half space $z < 0$ in a fixed Cartesian coordinate system. The point force is located at a distance z_0 below the free surface of the fluid, being suddenly started from rest and made to move with uniform velocity $U^* \vec{e}_x$, where \vec{e}_x denotes the unit vector along the x direction. It is meaningful to nondimensionalise the time by

$\frac{U^*}{g}$, the distance by $\frac{U^{*2}}{g}$, the characteristic speed by

U^* and the pressure by ρU^{*2} , where g is the gravitational constant and ρ the density of the fluid. Because only the wave profile at large distances downstream for high Reynolds numbers is investigated, the nondimensional, linearised Navier-Stokes equations describing the fluid flow, induced by an external body force with dimensionless strength $\vec{F} H(t) \delta(\vec{x} - \vec{x}_0)$, are given by

$$\nabla \cdot \vec{u}^* = 0, \quad (1)$$

$$\frac{\partial \vec{u}^*}{\partial t} + \frac{\partial \vec{u}^*}{\partial x} = -\nabla p^* + \varepsilon \nabla^2 \vec{u}^* + \vec{F} H(t) \delta(\vec{x} - \vec{x}_0). \quad (2)$$

At negative time $t < 0$ everything is at rest,

$$\vec{u}^* = \vec{0}, \quad p^* = 0, \quad \eta = 0, \quad \text{for } t < 0. \quad (3)$$

On the free surface, the tangential stresses are assumed to vanish and the normal stress has a jump proportional to the surface tension and mean curvature. On $z < 0$ these free-surface conditions are linearised together with the kinematic boundary condition to yield

$$\frac{\partial u^*}{\partial z} + \frac{\partial w^*}{\partial x} = 0, \quad \frac{\partial v^*}{\partial z} + \frac{\partial w^*}{\partial y} = 0, \quad (4)$$

$$\eta - p^* + 2\varepsilon \frac{\partial w^*}{\partial z} = \sigma \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \eta, \quad (5)$$

$$\left(\frac{\partial}{\partial t} + \frac{\partial}{\partial x} \right) \eta = w^*. \quad (6)$$

The fluid velocity and pressure vanish at infinity,

$$\vec{u}^* \rightarrow \vec{0}, \quad p^* \rightarrow 0, \quad \text{as } z \rightarrow -\infty. \quad (7)$$

Here the variables $\vec{u}^* = (u^*, v^*, w^*)^T$ and p^* represent the non-dimensional disturbed velocity and disturbed pressure in the fluid, $\vec{x} = (x, y, z)^T$ and $\vec{x}_0 = (0, 0, -z_0)^T$ are the field point and source point, η is the free-surface elevation, and the dimensionless parameters $\varepsilon = \frac{1}{R_e}$ and $\sigma = \frac{1}{W_e}$ can be regarded as the reciprocal of the Reynolds number

$$R_e = \frac{\rho U^{*3}}{g \times (\text{dynamic viscosity})}$$

and the Weber number $W_e = \frac{\rho U^{*4}}{g \times (\text{surface tension})}$ respectively.

$H(\bullet)$ and $\delta(\bullet)$ are Heaviside's step function and Dirac delta function respectively. The solution to equations (1) and (2) for an unbounded fluid domain is given by the author⁶ as

$$\begin{aligned} \vec{u}_0 &= -\frac{H(t)}{4\pi} \vec{F} \bullet (\mathbf{I} \nabla^2 - \nabla \nabla) \int_0^t \frac{erf(r^*/2\sqrt{\varepsilon\tau})}{r^*} d\tau, \\ p_0 &= \frac{H(t) \vec{F} \bullet \vec{x}}{4\pi r^3}, \end{aligned} \quad (8)$$

where $\vec{x}^* = \vec{x} - \tau \vec{e}_z$, $r^* = \|\vec{x}^* - \vec{x}_0\|$, and $r = \|\vec{x} - \vec{x}_0\|$.

Now let the entire solution be written as

$$\vec{u}^* = \vec{u}_0 + \vec{u}, \quad p^* = p_0 + p. \quad (9)$$

To reduce the number of variables involved, the motion can be represented as a potential flow plus a rotational flow. Thus, two new functions ϕ and $\vec{\omega} = (\omega_x, 0, \omega_z)^T$ are defined⁸ by

$$\vec{u} = \nabla \phi + \nabla \times \vec{\omega} \quad (10)$$

such that

$$\nabla^2 \phi = 0, \quad (11)$$

$$\frac{\partial \vec{\omega}}{\partial t} + \frac{\partial \vec{\omega}}{\partial x} = \varepsilon \nabla^2 \vec{\omega}, \quad (12)$$

and

$$p = -\frac{\partial \phi}{\partial t} - \frac{\partial \phi}{\partial x} \quad (13)$$

subject to the condition $\eta = 0$, for $t = 0$. By means of the Laplace transform in t and the Fourier transforms in x and y ,

$$\begin{aligned} [\bar{\phi}, \bar{\vec{\omega}}](s, \alpha, \beta, z) &= \\ &\int_0^\infty \int_{-\infty}^\infty \int_{-\infty}^\infty [\phi, \vec{\omega}](t, x, y, z) \\ &\exp(-st - i\alpha x - i\beta y - [A, B](z + z_0)) dt dx dy, \end{aligned} \quad (14)$$

$$\begin{aligned} \bar{\eta}(s, \alpha, \beta) &= \int_0^\infty \int_{-\infty}^\infty \int_{-\infty}^\infty \eta(t, x, y) \\ &\exp(-st - i\alpha x - i\beta y) dt dx dy, \end{aligned} \quad (15)$$

it then follows from equations (11) to (13) that A and B must satisfy

$$A = \sqrt{\alpha^2 + \beta^2}, \quad B = \sqrt{\alpha^2 + \beta^2 + \frac{s + i\alpha}{\varepsilon}}. \quad (16)$$

The free-surface conditions (4) to (6) can be expressed in terms of $\bar{\phi}$, $\bar{\omega}_x$, $\bar{\omega}_z$ and $\bar{\eta}$ as

$$\mathbf{C} \vec{V} = \vec{C}^{\{A\}} \exp(-A(z + z_0)) + \vec{C}^{\{B\}} \exp(-B(z + z_0)) \quad (17)$$

where superscripts $\{A\}$ and $\{B\}$ denote contributions by the potential flow and the rotational flow respectively, $\vec{V} = (\bar{\phi}, \bar{\omega}_x, \bar{\omega}_z, \bar{\eta})^T$,

$$\mathbf{C} = \begin{bmatrix} 2i\alpha A(s+i\alpha)\varepsilon^{3/2} \\ 2i\beta A(s+i\alpha)\varepsilon^{3/2} \\ (s+i\alpha)^2 + A[1+A^2\sigma + 2A(s+i\alpha)\varepsilon] \\ A \\ \alpha\beta(s+i\alpha)\varepsilon^{3/2} & i\beta B(s+i\alpha)\varepsilon^{3/2} & 0 \\ (\beta^2 + B^2)(s+i\alpha)\varepsilon^{3/2} & -i\alpha B(s+i\alpha)\varepsilon^{3/2} & 0 \\ -i\beta[1+A^2\sigma + 2B(s+i\alpha)\varepsilon] & 0 & 0 \\ -i\beta & 0 & -(s+i\alpha) \end{bmatrix} \quad (18)$$

and

$$\begin{aligned} \vec{C}^{\{A\}} &= \frac{\alpha F_x + i A F_z}{2s(s+i\alpha)A} \\ &\begin{bmatrix} -2\alpha A(s+i\alpha)\varepsilon^{3/2} \\ -2\beta A(s+i\alpha)\varepsilon^{3/2} \\ i[A - (s+i\alpha)^2 + A^3\sigma - 2A^2(s+i\alpha)\varepsilon] \\ 0 \end{bmatrix}, \end{aligned} \quad (19)$$

$$\vec{C}^{\{B\}} = -\frac{i\alpha(1+A^2\sigma)F_x}{2s(s+i\alpha)} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} + O(\sqrt{\varepsilon}). \quad (20)$$

The solution may be expressed as

$$\bar{\eta} = \frac{i(\alpha F_x + i A F_z)[(s+i\alpha)^2 - A(1+A^2\sigma)] \exp(-A z_0)}{2sA[(s+i\alpha)^2 + A(1+A^2\sigma) + 4A^2(s+i\alpha)\varepsilon]} + O(\varepsilon^{3/2}). \quad (21)$$

Because of the exponential term $\exp(-B(z+z_0))$ involved in equation (17), the expansion of $\vec{C}^{\{B\}}$ in equation (20) will not affect the final result in equation (21).

⁸ As three new independent variables, ϕ , ω_x and ω_z replace u , v and w (three components of \vec{u}), the three independent replacements only allow having two independent components. Without loss of generality, say $\omega_y = 0$.

3. Results

The wave number vector $(\alpha, \beta)^T$ identifies a free-surface wave in the far wake of a point force so long as it points to a curve in the (α, β) -plane defined by setting one factor of the denominator in equation (21) to zero,

$$(s + i\alpha)^2 + A(1 + A^2\sigma) + 4A^2(s + i\alpha)\varepsilon = 0. \quad (22)$$

The curve consists of three or two branches depending on whether $\|s\| < 1/4$ or $\|s\| \geq 1/4$. In the steady limit ($s=0$), one closed branch shrinks to zero, while the other two open branches represent the Kelvin wave pattern¹, which consists of a series of so-called diverging and transverse waves.

The problem posed here is that of finding the far-field asymptotic behaviour of free-surface waves, induced by impulsive motion of a submerged point force $\vec{F} = (F_x, 0, F_z)$. F_x and F_z represent dimensionless drag and lift forces respectively. For the unsteady case, ie, $\|s\| \neq 0$, the cylindrical coordinates (R, θ) are introduced on the free surface through

$$x = R \cos\theta, \quad y = R \sin\theta. \quad (23)$$

To obtain the leading terms in the far-field asymptotic representation for small ε and s , Lighthill's two-stage scheme⁷ shall be employed, which in essence involves calculating the α -integration by residues⁸ and the R -integration by the method of steepest descent.⁹

For the first stage of Lighthill's scheme, the roots of the pole equation are considered

$$(s + i\alpha)^2 + A(1 + A^2\sigma) + 4A^2(s + i\alpha)\varepsilon = 0. \quad (24)$$

For small ε and s , the roots $(\alpha^j, j=1,2)$ take the form of

$$\alpha^{(j)} = (-1)^{j-1} A_2 - \frac{2iA_1^2(s + 2A_1^2\varepsilon)}{3A_2^2 - 2A_1(A_1 + 1)} + O(\varepsilon^{3/2} + \varepsilon s + s^2), \quad (25)$$

where $A_2 = \sqrt{A_1(1 + A_1^2\sigma)}$ and A_1 satisfies a cubic equation $\sigma A_1^3 - A_1^2 + A_1 + \beta^2 = 0$, that is,

$$A_1 = \begin{cases} \frac{1}{3\sigma} \left[1 + 2\sqrt{1-3\sigma} \cos\left(\frac{\pi+\psi}{3}\right) \right], \\ \text{no real positive root} \end{cases}$$

$$\cos\psi = \frac{27\beta^2\sigma^2 + 9\sigma - 2}{2(1-3\sigma)^{3/2}}, \quad 0 \leq \psi \leq \pi \quad \text{if } \sigma \leq \frac{1}{4}$$

$$\text{if } \sigma > \frac{1}{4} \quad (26)$$

Using the residue theorem of a meromorphic

function with respect to α , the leading terms contributing significantly to the asymptotic expressions about $\varepsilon=0$ and $s=0$ of the free-surface elevation can be written as

$$\int_0^\infty \eta \exp(-st) dt = -\frac{1}{2\pi s} \sum_{j=1}^2 \int_{-\infty}^\infty \frac{[A_2 F_x + (-1)^{j-1} i A_1 F_z] A_1 A_2}{3A_2^2 - 2A_1(A_1 + 1)} \exp(-A_1 z_0 + i R h_j) [1 + O(\varepsilon + s)] d\beta, \quad (27)$$

$$\text{where } h_j(\beta|\alpha^{(j)}) = \alpha^{(j)} \cos\theta + \beta \sin\theta.$$

For the second stage of Lighthill's scheme, the saddle points are considered to satisfy the derivative of the exponent of the Fourier kernel,

$$\frac{\partial}{\partial \beta} h_j(\beta|\alpha^{(j)}) = \frac{\partial \alpha^{(j)}}{\partial \beta} \cos\theta + \sin\theta = 0 \quad (28)$$

For small ε and s , the roots $\beta_\pm^{(j)}$ take the form of

$$\beta_\pm^{(j)} = (-1)^{j-1} \frac{3A_4^2 - 2(A_3 + 1)A_3}{3A_4^2 - 2A_3} \left\{ 1 + (-1)^{j-1} \frac{4iA_4^3[(3A_4^2 - 4A_3)s - 2(3A_4^2 - 4A_3^2)A_3^2 e]}{3A_4^6 - 6A_3^2 A_4^4 + 14A_3 A_4^4 - 28A_3^2 A_4^2 + 8A_3^4 + 8A_3^3} \right\} \times A_4 \tan\theta + O(\varepsilon^{3/2} + \varepsilon s + s^2), \quad (29)$$

where $A_4 = \sqrt{A_3(1 + A_3^2\sigma)}$ and A_3 satisfies a sextic equation

$$(1 + A_3^2\sigma)(3A_3^2\sigma - 2A_3 + 1)^2 \tan^2\theta + (A_3^2\sigma - A_3 + 1)(3A_3^2\sigma + 1)^2 = 0. \quad (30)$$

The sextic equation (30) cannot be solved by rational operations and root extraction on coefficients. For small σ , the root A_3 takes the form of

$$A_3 = M_\pm \left[1 + \frac{(5G_\pm - 6)M_\pm^2\sigma}{G_\pm - 2} + O(\sigma^2) \right] \quad (31)$$

with the G_\pm and M_\pm written as

$$M_\pm = \frac{G_\pm + 1}{2}, \quad G_\pm = \frac{1 \pm \sqrt{1 - 8 \tan^2\theta}}{4 \tan^2\theta}. \quad (32)$$

Hence

$$A_1 = M_\pm + (-1)^j \frac{4i(G_\pm - 1)M_\pm^{1/2}s}{G_\pm(G_\pm - 2)} + (-1)^{j-1} \frac{8i(G_\pm - 1)(2G_\pm - 1)M_\pm^{5/2}\varepsilon}{G_\pm(G_\pm - 2)} + \frac{(5G_\pm - 6)M_\pm^3\sigma}{G_\pm - 2} + O(\varepsilon^{3/2} + \varepsilon\sigma + \sigma^2 + \varepsilon s + \sigma s + s^2), \quad (33)$$

$$\begin{aligned}
 A_2 = & M_{\pm}^{1/2} + (-1)^j \frac{2i(G_{\pm}-1)s}{G_{\pm}(G_{\pm}-2)} \\
 & + (-1)^{j-1} \frac{4i(G_{\pm}-1)(2G_{\pm}-1)M_{\pm}^2 \varepsilon}{G_{\pm}(G_{\pm}-2)} \\
 & + \frac{(3G_{\pm}-4)M_{\pm}^{5/2} \sigma}{G_{\pm}-2} \\
 & + O(\varepsilon^{3/2} + \varepsilon \sigma + \sigma^2 + \varepsilon s + \sigma s + s^2), \quad (34)
 \end{aligned}$$

$$A_4 = M_{\pm}^{1/2} \left[1 + \frac{(3G_{\pm}-4)M_{\pm}^2 \sigma}{G_{\pm}-2} + O(\sigma^2) \right], \quad (35)$$

$$\begin{aligned}
 \alpha_{\pm}^{(j)} = & (-1)^{j-1} M_{\pm}^{1/2} \left\{ 1 + (-1)^{j-1} \frac{i(G_{\pm}-3)s}{(G_{\pm}-2)M_{\pm}^{1/2}} \right. \\
 & \left. + (-1)^{j-1} \frac{2i(5G_{\pm}-7)M_{\pm}^{3/2} \varepsilon}{G_{\pm}-2} + \frac{(3G_{\pm}-4)M_{\pm}^2 \sigma}{G_{\pm}-2} \right\} \\
 & + O(\varepsilon^{3/2} + \varepsilon \sigma + \sigma^2 + \varepsilon s + \sigma s + s^2), \quad (36)
 \end{aligned}$$

$$\begin{aligned}
 \beta_{\pm}^{(j)} = & (-1)^j G_{\pm} M_{\pm}^{1/2} \left\{ 1 + (-1)^j \frac{4is}{(G_{\pm}-2)M_{\pm}^{1/2}} \right. \\
 & \left. + (-1)^{j-1} \frac{8i(2G_{\pm}-1)M_{\pm}^{3/2} \varepsilon}{G_{\pm}-2} + \frac{(5G_{\pm}-2)M_{\pm}^2 \sigma}{G_{\pm}-2} \right\} \tan \theta \\
 & + O(\varepsilon^{3/2} + \varepsilon \sigma + \sigma^2 + \varepsilon s + \sigma s + s^2). \quad (37)
 \end{aligned}$$

Using the method of steepest descent, the leading term is obtained as

$$\begin{aligned}
 \int_0^{\infty} \eta \exp(-st) dt = & \frac{1}{s} \sqrt{\frac{1}{2\pi R \cos \theta}} (1-8 \tan^2 \theta)^{-1/4} \\
 & \sum_{j=1}^2 \sum_{\pm} M_{\pm}^{-1/4} \left[M_{\pm} F_x + (-1)^{j-1} i M_{\pm}^{3/2} F_z \right] \\
 & \times \left\{ \exp \left(-M_{\pm} z_0 + (-1)^{j-1} i R \frac{M_{\pm}^{3/2}}{G_{\pm}} \right. \right. \\
 & \left. \left. \left[1 + (-1)^{j-1} 4i M_{\pm}^{3/2} \varepsilon + M_{\pm}^2 \sigma \right] \cos \theta \pm (-1)^j i \frac{\pi}{4} \right) \right. \\
 & \left. + O \left(\varepsilon + \sigma + s + \frac{1}{R} \right) \right\} \times \exp \left(-\frac{2RM_{\pm} \cos \theta}{G_{\pm}} s \right). \quad (38)
 \end{aligned}$$

Upon substituting and other mathematical manipulations, the free-surface elevation can formally be expressed as

$$\begin{aligned}
 \int_0^{\infty} \eta \exp(-st) dt = & \frac{1}{s} \sqrt{\frac{2}{\pi R \cos \theta}} (1-8 \tan^2 \theta)^{-1/4} \\
 & \sum_{\pm} \left[M_{\pm}^{3/4} P_{\pm} (F_x \cos \gamma_{\pm} - M_{\pm}^{1/2} F_z \sin \gamma_{\pm}) \right. \\
 & \left. + O \left(\varepsilon + \sigma + s + \frac{1}{R} \right) \right] \times \exp \left(-\frac{2RM_{\pm} \cos \theta}{G_{\pm}} s \right) \quad (39)
 \end{aligned}$$

where

$$\begin{aligned}
 P_{\pm} = & \exp \left(-\frac{M_{\pm} (z_0 G_{\pm} + 4 R M_{\pm}^2 \varepsilon \cos \theta)}{G_{\pm}} \right), \quad (40) \\
 \gamma_{\pm} = & \frac{R M_{\pm}^{3/2} \cos \theta}{G_{\pm}} \mp \frac{\pi}{4}.
 \end{aligned}$$

Using the inverse Laplace transform with respect to s , the asymptotic expression of the free-surface elevation can be written as

$$\begin{aligned}
 \eta = & \sqrt{\frac{2}{\pi R \cos \theta}} (1-8 \tan^2 \theta)^{-1/4} \\
 & \sum_{\pm} M_{\pm}^{3/4} P_{\pm} (F_x \cos \gamma_{\pm} - M_{\pm}^{1/2} F_z \sin \gamma_{\pm}) \\
 & \times H \left(t - \frac{2 R M_{\pm} \cos \theta}{G_{\pm}} \right) + O \left(\varepsilon + \sigma + \frac{1}{R} \right). \quad (41)
 \end{aligned}$$

4. Conclusions

The Taylor expansion has been performed for the transient free-surface flow due to impulsive motion of a submerged point force. The new asymptotic expressions of free-surface elevations and wavefronts for large time are obtained including the effects of viscosity and surface tension. The presence of viscosity is found to reduce the free-surface wave amplitude, while the surface tension does not affect the asymptotic wave.

It is worth to mention to this end that the present new result, equation (41), is dealing with the Oseen problem⁶ of large Reynolds number

$$\left(\varepsilon = \frac{1}{R_e} \right)$$

and far field (asymptotic wave), in contrast to the classical result¹⁰ based on unsteady Stokes solution, which is only valid for small Reynolds numbers and near field around the submerged source.

References

1. Kelvin B. (1906). Deep sea ship-waves. *The London, Edinburgh and Dublin Philosophical Magazine and Journal of Science*, 11(6):1-25.
2. Havelock TH. (1949). The wave resistance of a cylinder started from rest. *The Quarterly Journal of Mechanics and Applied Mathematics*, 2(3): 325-334.
3. Wehausen JV and Laitone EV. (1960). Surface waves. In *Flügge S and Truesdell C (editors) Encyclopedia of Physics, Volume IX, Fluid Dynamics III*, 446-778. Springer-Verlag.
4. Chan AT and Chwang AT. (1996). Interaction of laminar far wake with a free surface. *The Physics of Fluids A*, 8(2): 421-429.
5. Shu J-J. (2004). Transient marangoni waves due to impulsive motion of a submerged body. *International*

- Applied Mechanics*, 40(6): 709-714.
6. Shu J-J and Chwang AT. (2001). Generalised fundamental solutions for unsteady viscous flows. *Physical Review E*, 63(5): 051201.
 7. Lighthill MJ. (1960). Studies on magneto-hydrodynamic waves and other anisotropic wave motions. *Philosophical Transactions of the Royal Society of London, Series A Mathematical and Physical Sciences*, 252: 397-430.
 8. Jeffrey A. (1992). *Complex Analysis and Applications*. CRC Press Inc.
 9. Murray JD. (1984). *Asymptotic Analysis*. Springer-Verlag, New York Inc.
 10. Pozrikidis C. (1997). *Introduction to Theoretical and Computational Fluid Dynamics*. Oxford University Press, New York.
-