

This document is downloaded from DR-NTU, Nanyang Technological University Library, Singapore.

Title	Involvement of PKC-alpha in PMA-induced facilitation of exocytosis and vesicle fusion in PC12 cells( Main article )
Author(s)	Xue, Renhao; Zhao, Yanying; Chen, Peng
Citation	Xue, R., Zhao, Y., & Chen, P. (2009). Involvement of PKC-alpha in PMA-induced facilitation of exocytosis and vesicle fusion in PC12 cells. <i>Biochemical and Biophysical Research Communications</i> , 380(2), 371-376.
Date	2009
URL	<a href="http://hdl.handle.net/10220/7524">http://hdl.handle.net/10220/7524</a>
Rights	© 2009 Elsevier. This is the author created version of a work that has been peer reviewed and accepted for publication by <i>Biochemical and Biophysical Research Communications</i> , Elsevier. It incorporates referee's comments but changes resulting from the publishing process, such as copyediting, structural formatting, may not be reflected in this document. The published version is available at: [DOI: <a href="http://dx.doi.org/10.1016/j.bbrc.2009.01.105">http://dx.doi.org/10.1016/j.bbrc.2009.01.105</a> ].

**Involvement of PKC $\alpha$  in PMA-induced facilitation of  
exocytosis and vesicle fusion in PC12 cells**

Renhao Xue, Yanying Zhao, Peng Chen\*

Division of Bioengineering, Nanyang Technological University, Singapore 637457

\* Corresponding author: Fax: +65-67911761; E-mail: [chenpeng@ntu.edu.sg](mailto:chenpeng@ntu.edu.sg)

## **Abstract**

Phorbol-12-myristate-13-acetate, a stable analogue of the important signaling membrane lipid diacylglycerol (DAG), is known to potentiate exocytosis and modulate vesicle fusion kinetics in neurons and endocrine cells. The exact mechanisms underlying the actions of PMA, however, is often not clear, largely because of the diversity of the DAG/PMA receptors involved in the exocytotic process, which include, most notably, various isoforms of protein kinase C (PKC). In the present study, the roles of PKC $\alpha$  in PMA-mediated regulation of exocytosis were investigated by over-expressing wild-type PKC $\alpha$  (wt-PKC $\alpha$ ) or dominant negative PKC $\alpha$  (dn-PKC $\alpha$ ). Amperometric measurements based on carbon fiber microelectrodes demonstrated that PKC $\alpha$  has a key role in the PMA-mediated facilitation of exocytosis and vesicle fusion in neuroendocrine PC12 cells.

**Keywords:** exocytosis; protein kinase C; amperometry; PMA; DAG

Regulated secretion, or exocytosis, of hormones and neurotransmitters is regulated by a plethora of proteins and lipids through their specific actions at distinct stages in the exocytotic cascade, involving vesicle trafficking, docking, priming, and final  $\text{Ca}^{2+}$ -dependent fusion [1]. A large body of experiments has demonstrated that exocytosis is regulated acutely by protein phosphorylation [2], most notably *via* protein kinase C (PKC) [3,4]. Several key secretory proteins, such as SNAP-25 [5,6], synaptotagmin [7], and Munc18 [8,9], are PKC substrates, suggesting the involvement of PKC in various aspects of exocytosis.

PKC functions are often examined using pharmacological tools, such as phorbol-12-myristate-13-acetate (PMA), a potent diacylglycerol (DAG) analogue that acts as a PKC activator, and bisindolylmaleimide (BIS), which blocks the ATP-binding sites in the catalytic domain of many PKCs and acts as a PKC inhibitor. PMA has been shown to potentiate exocytosis [10-13] and modulate vesicle fusion in various cell systems [14,15]. Many of these PMA effects are thought to be PKC-dependent, as they can be eliminated partially or completely by treatment with BIS. However, the exact mechanisms underlying the actions of PMA is often not clear, largely because of the diversity of PKC isoforms and the caveats of pharmacological assessment. The PKC family consists of conventional PKCs ( $\alpha$ ,  $\beta$ , and  $\gamma$ ), novel PKCs ( $\delta$ ,  $\epsilon$ ,  $\eta$ , and  $\theta$ ), and atypical PKCs ( $\zeta$ ,  $\tau$ ,  $\lambda$ ) [3]. To avoid these problems, over-expression of wild-type and dominant negative forms of individual PKC isoforms were used in this study to elucidate PKC-dependent PMA effects on exocytosis.

PKC $\alpha$  is expressed abundantly in neuron and neuroendocrine cells. In particular, it is the only conventional PKC expressed in bovine chromaffin cells [16], and it is the most

constitutively active PKC isoform in rat chromaffin cell-derived PC12 cells [17]. In this study, by over-expressing wild-type PKC $\alpha$  (wt-PKC $\alpha$ ) and dominant negative PKC $\alpha$  (dn-PKC $\alpha$ ), we demonstrated that PKC $\alpha$  is largely involved in the PMA-induced increase in the extent of exocytosis, the facilitation of fusion pore expansion, and the augment of quantal release in PC12 cells.

## **Materials and methods**

### *Cell culture and transfection*

PC12 cells (ATCC) were cultured in RPMI 1640 supplemented with 10% fetal bovine serum, 5% horse serum and 1% penicillin-streptomycin (Gibco) at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> / 95% air. PC12 cells were transfected with plasmid DNA using ESCORT III transfection reagent (Sigma), 1-2 days prior to the experiments. Cells were planted on poly-L-lysine pre-coated coverslips and treated with 5  $\mu$ M dexamethasone for 3-5 days as previously described [18,19] to enhance chromaffin cell phenotype.

### *Plasmids*

The cDNA plasmids coding for wt- and dn-PKC $\alpha$  were obtained from Addgene Plasmid Collection. The pIRES2-EGFP vector was purchased from Clontech. The wt- or dn-PKC $\alpha$ -IRES-EGFP was constructed by subcloning cDNA of PKC $\alpha$  or its mutation into pIRES2-EGFP vector at the sites of Nhe I and Sma I.

### *Amperometry measurement*

The bath solution contained (in mM): 150 NaCl, 5.5 KCl, 2 MgCl<sub>2</sub>, 2 CaCl<sub>2</sub>, 10 glucose, and 10 HEPES (titrated to pH 7.2 with NaOH); the high K<sup>+</sup> stimulation solution contained (in mM): 37 NaCl, 105 KCl, 6 CaCl<sub>2</sub>, 2 MgCl<sub>2</sub>, and 10 HEPES (titrated to pH 7.2). In some experiments, cells were treated with 0.1 μM PMA for 10 minutes or / and 2.5 μM BIS for 30 minutes.

Exocytosis was induced by perfusing the high K<sup>+</sup> solution through a glass micropipette (tip size of 2~3 μm) positioned 5~10 μm away from the cell. Amperometric spikes were recorded by a 5 μm-in-diameter carbon fiber electrode (ALA Scientific Instruments, USA) gently positioned onto the cell membrane. The microelectrode was biased at 700 mV and the amperometry signals were sampled at 4 kHz and filtered at 1 kHz using an EPC-10 double patch-clamp amplifier (HEKA Elektronik, Germany).

The parameters of amperometric spikes were obtained using an Igor program, Amperometric Spike Analysis [20]. Only spikes with amplitudes larger than 2 pA (5 times of the background noise) were considered as true amperometric events. Statistics of the spike parameters is represented by mean ± SEM. Statistical significance is evaluated with Student's *t*-test or Kolmogorov–Smirnov test.

## **Results and discussion**

In comparison to electrophysiological measurement of membrane capacitance [21,22], amperometry using carbon fiber microelectrodes is instrumental in the direct detection of exocytosis with single-vesicle sensitivity and revealing vesicle fusion kinetics with millisecond resolution. Upon Ca<sup>2+</sup> triggering, vesicle fusion starts with formation of a nanometric aqueous fusion pore through which secretory compounds escape from the

vesicular lumen into extracellular space, sometimes giving a small foot signal preceding the amperometric spike. The fusion pore then expands quickly to discharge the secretory compounds, causing a rapid increase in the amperometric signal. Finally, expansion of the fusion pore expansion may end in full collapse, or terminate prematurely in a so called kiss-and-run mode, which gives a more transient amperometric signal [23,24].

*PMA modulates both the extent and the fusion kinetics of vesicle exocytosis*

It has long been recognized that PMA enlarges the extent of  $\text{Ca}^{2+}$ -dependent exocytosis [12-14]. Fig. 1A presents typical amperometric recordings from PC12 cells in response to a membrane-depolarizing high- $\text{K}^+$  solution under different conditions. Each current spike corresponds to a single vesicle fusion event. PMA increased the number of exocytotic events per stimulus significantly ( $133.3 \pm 12.1\%$ , 11 cells, with respect to the average response from 22 control cells), and this enhancement was eliminated completely by BIS ( $90.5 \pm 10.2\%$ , 11 cells). Furthermore, BIS alone inhibited exocytosis severely compared to the control ( $38.1 \pm 5.5\%$ , 19 cells). Therefore, it is reasonable to conclude that the PMA enhancement is PKC-dependent and the basal activities of PKCs are important to support exocytosis.

The averaged amperometric spikes obtained from more than 400 fusion events in at least 11 cells are shown in Fig. 1B. In contrast to that from the control cells, the average amperometric signal from the PMA-treated cells has a steeper rising phase, indicating more rapid opening of the fusion pore. The rate of catecholamine release or the rate of pore expansion can be quantified by the slope of the rising amperometric signal between 35% and 90% of the peak amplitude. As shown in Fig. 1C, the average rise slope in the PMA-treated cells is significantly greater than that in the control cells ( $10.1 \pm 0.3$  pA/ms

vs.  $6.8 \pm 0.2$  pA/ms,  $P < 0.001$ ). The acceleration of pore expansion likely relies on PKC activation, because BIS eliminated this effect of PMA. Application of BIS alone, however, did not cause apparent reduction in the fusion rate. It seems that basal PKC activities are not required for opening the fusion pore. Similar PMA facilitation of fusion pore opening has been found in chromaffin cells [14,15], in which PMA also causes premature closure of the fusion pore, leading to kiss-and-run fusion with small quantal sizes [9,14]. By contrast, the quantal size of vesicle release in PC12 cells, or the total charge ( $Q$ ) from integration of the amperometric current signal, was enlarged in response to treatment with PMA. Since the distribution of  $Q$  is known to be severely skewed, we used the cube-root of the spike charge ( $Q^{1/3}$ ) for analysis. Student's  $t$ -test can be applied to determine statistically significant differences, since  $Q^{1/3}$  has a Gaussian distribution, and is proportional to the radius of the fusion vesicles [25]. As demonstrated in Fig. 1D, treatment with PMA increased  $Q^{1/3}$  appreciably. The ability of BIS to reverse this PMA-mediated enlargement again suggests the involvement of PKC. BIS application alone decreased the quantal size significantly (Fig. 1D), implying that basal PKC activities are important for maintaining the normal quantal size.

It is clear from the data presented here and as well as that reported by others that PMA plays multiple regulatory roles in the  $\text{Ca}^{2+}$ -dependent exocytotic process, usually through activation of PKCs. We further hypothesize that  $\text{PKC}\alpha$ , which is the abundant and most active conventional PKC isoform in PC12 cells activated by DAG and  $\text{Ca}^{2+}$ , has an important role in PMA-mediated regulation of  $\text{Ca}^{2+}$ -dependent exocytosis.

*PKC $\alpha$  contributes to PMA-induced enhancement of exocytosis*

The wt- or dn-PKC $\alpha$  created by a K368N mutation to permanently inactivate the ATP-binding site of PKC $\alpha$  catalytic domain was over-expressed in PC12 cells. The cDNA of PKC $\alpha$  is constructed together with enhanced green fluorescence protein (EGFP) in the same plasmid *via* an internal ribosome entry site (IRES) [26], which links two open reading frames. Thus, EGFP is co-expressed with PKC $\alpha$  as reporter.

Fig. 2 depicts the average time-courses and statistics of the amperometric responses to 60 s of high-K<sup>+</sup> stimulation of PC12 cells transfected with different plasmids, and with or without treatment with PMA. Only bright green cells, in which EGFP (hence PKC $\alpha$ ) was amply expressed, were chosen for recording. Over-expression of dn-PKC $\alpha$  largely reduced the extent of exocytosis by 34% ( $P < 0.01$ ) in comparison with over-expression of the empty vector (pIRES2-EGFP), indicating that competitive inhibition of endogenous PKC $\alpha$  activities impairs exocytosis. This is analogous to the effect of treatment with BIS. Compared to dn-PKC $\alpha$ , over-expression of dn-PKC $\epsilon$  was much less inhibitory (12.9% reduction,  $P < 0.05$ , 25 cells). PKC $\epsilon$  is a novel PKC isoform expressed in PC12 cells [17], and was found to be implicated in activity-dependent potentiation of exocytosis in chromaffin cells [27].

The over-expression of wt-PKC $\alpha$  gave only a marginal increase in exocytosis. Apparently, the abundance of PKC $\alpha$  is not sufficient to elevate exocytosis significantly. PMA together with PKC $\alpha$  over-expression, however, caused pronounced enhancement, greater than its potentiation in the vector cells (35% *vs.* 26%). On the other hand, PMA failed to rescue exocytosis in the dominant-negative cells. These observations provide strong support for the notion that PKC $\alpha$  plays a key role in PMA potentiation.

*PMA increases the rate of fusion pore expansion and enlarges the fusion quantal size in a PKC $\alpha$ -dependent manner*

Fig. 3 presents the average amperometric spikes and statistical analysis for different experimental conditions. As expected, in the empty vector-expressing cells, PMA increased the speed of pore expansion as evidenced by the steeper amperometric rising phase. In contrast, such PMA-induced acceleration was absent from the cells expressing dn-PKC $\alpha$ ; i.e., dn-PKC $\alpha$ , just like BIS, is able to abolish this PMA effect. This observation suggests the minor involvement of other PMA targets and the importance of PKC $\alpha$  activation in this phenomenon. Also similar to the BIS effect demonstrated in Fig. 1, over-expression of dn-PKC $\alpha$  without PMA did not affect the slope of the rising phase. Therefore, basal PKC $\alpha$  activities are probably not critically involved in normal fusion pore opening. The observations that over-expression of wt-PKC $\alpha$  gave a sharper rise in the amperometric signal, and that PMA could not further strengthen this effect in wt-PKC $\alpha$  over-expressing cells, suggest that the abundance of PKC $\alpha$ , but not that of DAG, is the limiting factor in activating the acceleration mechanism for fusion.

As anticipated from the earlier experiments (Fig. 1), PMA enlarged the quantal size in the vector cells. Again, analogous to BIS actions, dn-PKC $\alpha$  over-expression eliminated this PMA-induced augmentation and led to notably smaller quantal size in the absence of PMA. These observations clearly highlight the importance of PKC $\alpha$  activation in supporting normal vesicle fusion and PMA enhancement. Endogenous PKC $\alpha$  is adequate to ensure PMA-induced enhancement, as over-expression of wt-PKC $\alpha$  did not boost the effect of PMA further.

The results of this study are contrast to those of a recent study on rat chromaffin cells [28], in which the authors, mainly on the basis of pharmacological assessments, reported that it is novel PKC $\theta$  and not conventional PKC isoforms that is important in maintaining normal quantal size and the normal extent of exocytosis. The discrepancy between their findings and those presented here may be attributable to differences in the cell types used, and the stimulation protocols. Firstly, PC12 cells do not express PKC $\theta$  [29]. Secondly, we recorded the amperometric responses during 60 s of high-K<sup>+</sup> stimulation, while they recorded the responses after only a brief (3 s) high-K<sup>+</sup> stimulation. In addition, unlike chromaffin cells, which typically have hundreds of releasable pre-docked vesicles, the amperometric responses from PC12 cells are essentially resulted from the release of newly arrived vesicles [30].

We cannot completely rule out the possible involvement of other PKC isoforms in PMA regulations of vesicle fusion, due partly to cross-reactivity in the enzyme–substrate binding. To partially address this issue, we examined the effects of PKC $\epsilon$  on vesicle fusion and found that PKC $\epsilon$  was not able to modulate the rate of fusion pore expansion: wt ( $14.40 \pm 0.60$  pA/ms, 28 cells) *vs.* dn ( $12.63 \pm 0.75$  pA/ms, 25 cells),  $P > 0.05$ . Further, opposite to the effect of nt-PKC $\alpha$ , nt-PKC $\epsilon$  enlarged the quantal size ( $Q^{1/3}$ ): wt ( $3.55 \pm 0.02$  fC<sup>1/3</sup>) *vs.* dn ( $3.74 \pm 0.03$  fC<sup>1/3</sup>),  $P < 0.001$ .

PKC facilitation of vesicle fusion can be achieved by modulating the fusion protein machinery or by enzymatic alteration of the lipidic structures of the fusion pore. Phospholipase D (PLD), which can be activated by PKC [31], produces phosphatidic acids (PAs) at the inner leaflet of the cell membrane at the fusion site. Cone-shaped PAs energetically favor the formation of a negatively curved fusion pore and promote its

expansion [32]. We investigated the effects of 1-butanol, which blocks formation of PA by competitive binding to PLD [33], on the fusion kinetics. It was found that 1-butanol attenuated the spike slope from  $6.81 \pm 0.20$  pA/ms (22 untreated cells) to  $4.14 \pm 0.35$  pA/ms (11 treated cells)  $P < 0.001$ . But the application of PMA was able to rescue the amperometric rise slope of the cells treated with 1-butanol (6 cells) to a level even faster than that of the control ( $8.70 \pm 0.85$  pA/ms vs.  $6.81 \pm 0.20$  pA/ms;  $P < 0.001$ ). So, PLD is unlikely to be the target of PMA (thus PKC) in facilitating fusion pore expansion.

*Foot probability is lowered by PMA through PKC activities*

Fusion between vesicular and plasma membrane must overcome substantial energy barriers. When the driving force provided by the protein fusion machinery barely overwhelms these energy barriers, a small pre-spike foot in the amperometric signal occurs due to slow expansion of the fusion pore [34]. PMA accelerates fusion pore expansion, likely by lowering certain energy barriers of the fusion process. In line with this view, the probability of recording a foot signal was found to be reduced by treatment with PMA (Fig. 4). Foot probability is obtained by calculating the percentage of the events that have a preceding foot among all the recorded exocytotic events that have amplitude  $> 20$  pA (foot can be resolved definitely only for events with relatively large amplitude [35]). Not surprisingly, this PMA effect is also PKC-dependent, because BIS was able to inhibit it, while BIS application alone increased the foot probability. With reference to the vector-expressing cells, over-expression of dn-PKC $\alpha$  behaved just like BIS to increase the foot probability, whereas wt-PKC $\alpha$  decreased it. These experiments demonstrated that PKC activities, including, but maybe not exclusive to, the PKC $\alpha$

isoform, lowered the fusion energy barriers leading to a decrease in the number of footed events.

### **Acknowledgements**

This study was supported by an AcRF tier 2 (T206B3220) and tier 1 (RG41/05) grants.

## References


- [1] R.D. Burgoyne, A. Morgan, Secretory granule exocytosis, *Physiol. Rev.* 83 (2003) 581-632.
- [2] P.M. Jones, S.J. Persaud, Protein kinases, protein phosphorylation, and the regulation of insulin secretion from pancreatic beta-cells, *Endocr. Rev.* 19 (1998) 429-461.
- [3] A. Morgan, R.D. Burgoyne, J.W. Barclay, T.J. Craig, G.R. Prescott, L.F. Ciufo, G.J.O. Evans, M.E. Graham, Regulation of exocytosis by protein kinase C, *Biochem. Soc. Trans.* 33 (2005) 1341-1344.
- [4] Y. Yang, S. Udayasankar, J. Dunning, P. Chen, K.D. Gillis, A highly  $Ca^{2+}$ -sensitive pool of vesicles is regulated by protein kinase C in adrenal chromaffin cells, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 17060-17065.
- [5] Y. Shimazaki, T. Nishiki, A. Omori, M. Sekiguchi, Y. Kamata, S. Kozaki, M. Takahashi, Phosphorylation of 25-kDa synaptosome-associated protein - Possible involvement in protein kinase C-mediated regulation of neurotransmitter release, *J. Biol. Chem.* 271 (1996) 14548-14553.
- [6] G. Nagy, U. Matti, R.B. Nehring, T. Binz, J. Rettig, E. Neher, J.B. Sorensen, Protein kinase C-dependent phosphorylation of synaptosome-associated protein of 25 kDa at Ser(187) potentiates vesicle recruitment, *J. Neurosci.* 22 (2002) 9278-9286.
- [7] Y. Haberman, I. Ziv, Y. Gorzalczy, M. Fukuda, R. Sagl-Eisenberg, Classical protein kinase C(s) regulates targeting of synaptotagmin IX to the endocytic recycling compartment, *J. Cell Sci.* 118 (2005) 1641-1649.
- [8] Y. Fujita, T. Sasaki, K. Fukui, H. Kotani, T. Kimura, Y. Hata, T.C. Sudhof, R.H. Scheller, Y. Takai, Phosphorylation of Munc-18/n-Sec1/rbSec1 by protein kinase C -

- Its implication in regulating the interaction of Munc-18/n-Sec1/rbSec1 with syntaxin, *J. Biol. Chem.* 271 (1996) 7265-7268.
- [9] J.W. Barclay, T.J. Craig, R.J. Fisher, L.F. Ciufo, G.J.O. Evans, A. Morgan, R.D. Burgoyne, Phosphorylation of Munc18 by protein kinase C regulates the kinetics of exocytosis, *J. Biol. Chem.* 278 (2003) 10538-10545.
- [10] S.L. Pocotte, R.A. Frye, R.A. Senter, D.R. Terbush, S.A. Lee, R.W. Holz, Effects of phorbol ester on catecholamine secretion and protein-phosphorylation in adrenal-medullary cell-cultures, *Proc. Natl. Acad. Sci. U. S. A.* 82 (1985) 930-934.
- [11] M.L. Vitale, E.P. Seward, J.M. Trifaro, Chromaffin cell cortical actin network dynamics control the size of the release-ready vesicle pool and the initial rate of exocytosis, *Neuron* 14 (1995) 353-363.
- [12] K.D. Gillis, R. Mossner, E. Neher, Protein kinase C enhances exocytosis from chromaffin cells by increasing the size of the readily releasable pool of secretory granules, *Neuron* 16 (1996) 1209-1220.
- [13] Y. Shoji-Kasai, M. Itakura, M. Kataoka, S. Yamamori, M. Takahashi, Protein kinase C-mediated translocation of secretory vesicles to plasma membrane and enhancement of neurotransmitter release from PC12 cells, *Eur. J. Neurosci.* 15 (2002) 1390-1394.
- [14] M.E. Graham, R.J. Fisher, R.D. Burgoyne, Measurement of exocytosis by amperometry in adrenal chromaffin cells: Effects of clostridial neurotoxins and activation of protein kinase C on fusion pore kinetics, *Biochimie* 82 (2000) 469-479.
- [15] T. Fulop, C. Smith, Physiological stimulation regulates the exocytic mode through calcium activation of protein kinase C in mouse chromaffin cells, *Biochem. J.* 399 (2006) 111-119.

- [16] C.M. Sena, R.M. Santos, N.B. Standen, M.R. Boarder, L.M. Rosario, Isoform-specific inhibition of voltage-sensitive  $\text{Ca}^{2+}$  channels by protein kinase C in adrenal chromaffin cells, *FEBS Lett.* 492 (2001) 146-150.
- [17] M.W. Wooten, M.L. Seibenhener, Y. Soh, S.J. Ewald, K.R. White, E.D. Lloyd, A. Olivier, P.J. Parker, Characterization and differential expression of protein kinase C isoforms in PC12 cells - Differentiation parallels an Increase in PKC Beta(Ii), *FEBS Lett.* 298 (1992) 74-78.
- [18] R.H.S. Westerink, A.G. Ewing, The PC12 cell as model for neurosecretion, *Acta Physiol.* 192 (2008) 273-285.
- [19] A.S. Tischler, R.L. Perlman, G.M. Morse, B.E. Sheard, Glucocorticoids increase catecholamine synthesis and storage in PC12 pheochromocytoma cell cultures, *J. Neurochem.* 40 (1983) 364-370.
- [20] E.V. Mosharov, D. Sulzer, Analysis of exocytotic events recorded by amperometry, *Nature Methods* 2 (2005) 651-658.
- [21] P. Chen, K.D. Gillis, The noise of membrane capacitance measurements in the whole-cell recording configuration, *Biophys. J.* 79 (2000) 2162-2170.
- [22] P. Chen, T.C. Hwang, K.D. Gillis, The relationship between cAMP,  $\text{Ca}^{2+}$ , and transport of CFTR to the plasma membrane, *J. Gen. Physiol.* 118 (2001) 135-144.
- [23] R.D. Burgoyne, J.W. Barclay, Splitting the quantum: regulation of quantal release during vesicle fusion, *Trends Neurosci.* 25 (2002) 176-178.
- [24] M.B. Jackson, E.R. Chapman, Fusion pores and fusion machines in  $\text{Ca}^{2+}$ -triggered exocytosis, *Annu. Rev. Biophys. Biomol. Struct.* 35 (2006) 135-160.


- [25] R.A. Clark, A.G. Ewing, Quantitative measurements of released amines from individual exocytosis events, *Mol. Neurobiol.* 15 (1997) 1-16.
- [26] S.F. Bairstow, K. Ling, X.J. Su, A.J. Firestone, C. Carbonara, R.A. Anderson, Type I gamma 661 phosphatidylinositol phosphate kinase directly interacts with AP2 and regulates endocytosis, *J. Biol. Chem.* 281 (2006) 20632-20642.
- [27] Y.S. Park, E.M. Hur, B.H. Choi, E. Kwak, D.J. Jun, S.J. Park, K.T. Kim, Involvement of protein kinase C-epsilon in activity-dependent potentiation of large dense-core vesicle exocytosis in chromaffin cells, *J. Neurosci.* 26 (2006) 8999-9005.
- [28] R.G.W. Staal, A. Hananiya, D. Sulzer, PKC theta activity maintains normal quantal size in chromaffin cells, *J. Neurochem.* 105 (2008) 1635-1641.
- [29] P. Borgatti, M. Mazzoni, C. Carini, L.M. Neri, M. Marchisio, L. Bertolaso, M. Previati, G. Zauli, S. Capitani, Changes of nuclear protein kinase C activity and isotype composition in PC12 cell proliferation and differentiation, *Exp. Cell Res.* 224 (1996) 72-78.
- [30] E. Zhang, R. Xue, J. Soo, P. Chen, Effects of phorbol ester on vesicle dynamics as revealed by total internal reflection fluorescence microscopy, *Pflugers Arch.* 457 (2008) 211-222.
- [31] S.M. Hammond, J.M. Jenco, S. Nakashima, K. Cadwallader, Q.M. Gu, S. Cook, Y. Nozawa, G.D. Prestwich, M.A. Frohman, A.J. Morris, Characterization of two alternately spliced forms of phospholipase D1 - Activation of the purified enzymes by phosphatidylinositol 4,5-bisphosphate, ADP-ribosylation factor, and RHO family monomeric GTP-binding proteins and protein kinase C-alpha, *J. Biol. Chem.* 272 (1997) 3860-3868.

- [32] N. Vitale, A.S. Caumont, S. Chasserot-Golaz, G. Du, S. Wu, V.A. Sciorra, A.J. Morris, M.A. Frohman, M.F. Bader, Phospholipase D1: a key factor for the exocytotic machinery in neuroendocrine cells, *EMBO J.* 20 (2001) 2424-2434.
- [33] J. Klein, Functions and pathophysiological roles of phospholipase D in the brain, *J. Neurochem.* 94 (2005) 1473-1487.
- [34] R.H. Chow, L. Vonruden, E. Neher, Delay in vesicle fusion revealed by electrochemical monitoring of single secretory events in adrenal chromaffin cells, *Nature* 356 (1992) 60-63.
- [35] C.T. Wang, J.H. Bai, P.Y. Chang, E.R. Chapman, M.B. Jackson, Synaptotagmin- $\text{Ca}^{2+}$  triggers two sequential steps in regulated exocytosis in rat PC12 cells: fusion pore opening and fusion pore dilation, *J. Physiol.-London* 570 (2006) 295-307.




**Fig. 1.** Effects of PMA and BIS on exocytosis and vesicle fusion. **A.** Representative amperometric recordings from a control, PMA-treated, BIS+PMA-treated or BIS-treated cell, in response to 60 s of stimulation by a high- $K^+$  solution. **B.** Averaged amperometric spikes of all fusion events from 22 control cells, 11 PMA-treated cells, 11 BIS+PMA-treated cells or 19 BIS-treated cells. **C.** Statistics of the spike rise slope. The rise slope is


defined as the slope of the linear fit between 35% and 90% of the peak amplitude of the spike, as shown in the inset. Kolmogorov–Smirnov test: \*\*\*  $P < 0.001$ . **D**. Statistics of the cube-root of the spike charge,  $Q^{1/3}$ . The spike charge ( $Q$ ) was defined as the integration of the amperometric spike current as shown in the inset. Student's  $t$ -test: \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ . **C** and **D**, the data are shown as mean  $\pm$  SEM.



**Fig. 2.** Effects of PMA on exocytosis in PKC $\alpha$  over-expressing cells. Data were obtained by averaging the responses from 14 empty vector-expressing cells (Vector), 13 PMA-treated vector cells (Vector+PMA), 16 wt-PKC $\alpha$ -expressing cells (WT), 14 PMA-treated WT cells (WT+PMA), 21 dn-PKC $\alpha$ -expressing cells (DN) and 18 PMA-treated DN cells (DN+PMA), during 60 s high-K<sup>+</sup> stimulation. **A.** The cumulative spike number is plotted against time. **B.** The total numbers of the resulting spikes are shown as mean  $\pm$  SEM. Student's *t*-test: \*  $P < 0.05$ , \*\*  $P < 0.01$ .



**Fig. 3.** PMA modulates vesicle fusion *via* PKC $\alpha$ . The analysis is based on the experiments presented in Fig. 2. **A.** The average amperometric spikes of all events from differently treated cells. **B.** The rise slopes are shown as mean  $\pm$  SEM. Kolmogorov–Smirnov test: \*\*\*  $P < 0.001$ , not significant (N. S.) for  $P > 0.05$ . **C.** The cube-roots of spike charges ( $Q^{1/3}$ ) are shown as mean  $\pm$  SEM. Student’s  $t$ -test: \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .



**Fig. 4.** The effects of PMA, BIS and PKC $\alpha$  expression on foot probability.

Representative amperometric spikes with and without a preceding foot are illustrated in the inset. Foot probability is defined as the percentage of footed signals among all the events that were obtained from the same experiments presented in Figs. 1 and 2, and have amplitude > 20 pA (43 – 212 events).