

# Recent Advances in Alkenyl $sp^2$ C–H and C–F Bond Functionalizations: Scope, Mechanism and Applications

Ming-Zhu Lu,<sup>†,‡</sup> Manikantha Maraswami,<sup>‡</sup> Jeffrey Goh,<sup>‡</sup> Zhenhua Jia,<sup>†</sup> Jie-Sheng Tian,<sup>§</sup> Teck-Peng Loh<sup>\*,†,‡</sup>

<sup>†</sup>College of Advanced Interdisciplinary Science and Technology, Henan University of Technology, Zhengzhou, 450001, China

<sup>‡</sup>School of Chemistry, Chemical Engineering and Biotechnology, Nanyang Technological University, Singapore 637371

<sup>§</sup>School of Chemistry and Chemical Engineering, Northwestern Polytechnical University, Xi'an 710072, China

---

**ABSTRACT:** Alkenes and their derivatives are featured widely in a variety of natural products, pharmaceuticals, and advanced materials. Significant efforts have been made towards the development of new and practical methods to access this important class of compounds by selectively activating the alkenyl C( $sp^2$ )-H bonds in recent years. In this **comprehensive** review, we describe the state-of-the-art strategies for the direct functionalization of alkenyl  $sp^2$  C–H and C–F bonds **until June 2022**. Moreover, metal-free, photoredox, and electrochemical strategies are also covered. For clarity, this review is divided into two parts, the first part focuses on currently available alkenyl  $sp^2$  C–H functionalization methods using different alkene derivatives as the starting materials, and the second part describes the alkenyl  $sp^2$  C–F bond functionalization using easily accessible *gem*-difluoroalkenes as the starting material. This review includes the scope, limitations, mechanistic studies, stereoselective control (using directing groups as well as metal-migration strategies), and their applications to complex molecule synthesis where appropriate. Overall, this **comprehensive** review aims to comprehensively document the considerable advancements, current status, and emerging work by critically summarizing the contributions of researchers working in this fertile area, and is expected to stimulate novel, innovative, and broadly applicable strategies for alkenyl  $sp^2$  C–H and C–F bonds functionalization in the coming years.

---

## CONTENTS

1. Introduction
2. Alkenyl C–H Bond Functionalization of Simple Alkenes
  - 2.1. Direct Alkenyl C–H Bond Functionalization of Simple Alkenes
  - 2.2. Directed Alkenyl C–H Bond Functionalization of Aromatic Alkenes
  - 2.3. Migratory Alkenyl C–H Bond Functionalization of Aromatic Alkenes
  - 2.4. Annulation of Aromatic Alkenes via Alkenyl C–H Bond Functionalization
3. Alkenyl C–H Bond Functionalization of Aliphatic Alkenes Containing a Directing Group
4. Alkenyl C–H Bond Functionalization of Alkenes Containing an Electron-Withdrawing Directing Group
  - 4.1. Acrylic Acids
  - 4.2. Acrylic Esters
  - 4.3. Acrylic Amides
    - 4.3.1. Arylation
    - 4.3.2. Alkylation
    - 4.3.3. Alkenylation
    - 4.3.4. Allylation
    - 4.3.5. Alkynylation
    - 4.3.6. Fluoroalkylation
    - 4.3.7. Halogenation
    - 4.3.8. Annulation Reactions
    - 4.3.9. Other Useful Reactions
  - 4.4.  $\alpha,\beta$ -Unsaturated Ketones
  - 4.5.  $\alpha,\beta$ -Unsaturated Imines
  - 4.6.  $\alpha,\beta$ -Unsaturated Oximes and Derivatives.
  - 4.7. 2-Vinylpyridines
5. Alkenyl C–H Bond Functionalization of Alkenes Containing a Heteroatom
  - 5.1. Enamides
    - 5.1.1. Arylation
    - 5.1.2. Alkylation
    - 5.1.3. Fluoroalkylation
    - 5.1.4. Olefination
    - 5.1.5. Alkynylation
    - 5.1.6. Carbonylation
    - 5.1.7. Sulfonylation
    - 5.1.8. Phosphorylation
    - 5.1.9. Annulation Reaction
    - 5.1.10. Other Useful Reactions
  - 5.2. Enaminones

- 5.3. Enamines
- 5.4. Enolates
  - 5.4.1. Enol Acetates
  - 5.4.2. Enol Carbamates
  - 5.4.3. Enol Phosphates
  - 5.4.4. *N*-Enoxyphthalimides
- 5.5. Ketene Dithioacetals
- 6. Miscellaneous Alkenyl C–H Bond Functionalizations
- 7. Alkenyl C–F Bond Functionalizations of *gem*-Difluoroalkenes
  - 7.1. C–H Bond Formation
  - 7.2. C–C Bond Formation
    - 7.2.1. C–H/C–F Functionalization Reaction
    - 7.2.2. Alkylation
    - 7.2.3. Alkenylation and Allylation
    - 7.2.4. Arylation
    - 7.2.5. Alkynylation
    - 7.2.6. Carboxylation
    - 7.2.7. Other Useful Reactions
  - 7.3. C–Het Bond Formation
    - 7.3.1. Borylation and Silylation
    - 7.3.2. Miscellaneous
  - 7.4. Photoredox Catalysis
  - 7.5. Annulation Reactions
- 8. Conclusions and Outlook
- Author Information
  - Corresponding Authors
  - Authors
  - Notes
  - Biographies
- Acknowledgments
- References

## 1. INTRODUCTION


Alkenes and their derivatives belong to an important class of compounds in organic chemistry since they are key fragments featured widely in a myriad of pharmaceuticals, bioactive natural products, and advanced functional materials. Furthermore, they can also serve as extremely versatile building blocks in a diverse variety of synthetic transformations. Due to the importance of this class of compounds, numerous conceptually different methods have been developed for the efficient construction of alkenes and their derivatives including cyclic alkenes. Moreover, the stereoselective and intramolecular versions of these methods have also been achieved. Reported methods include the Nobel Prize winning reactions such as the Wittig reaction,<sup>1–3</sup> olefin metathesis,<sup>4–8</sup> and transition-metal catalyzed cross-coupling reactions.<sup>9</sup> While these reactions are extremely efficient and useful, there are still plagued with various problems. For example, the poor atom-economy of Wittig reaction, poor selectivity control for highly substituted alkenes as well as disposal problem associated with the phosphine oxide significantly limit their widespread applications for large production of many value-added alkenes. On the other hand, cross-coupling and olefin metathesis reactions often require the use of expensive transition-metal catalysts and environmentally unfriendly organohalides and/or

organometallics. This inevitably leads to the generation of halide wastes and the need to prepare the organometallics. Accordingly, there have been much effort directed towards the development of greener, atom-economical, and practical methods to obtain alkenes and their derivatives starting from simple alkenes or commercially available chemical feedstocks.

An alternative and sought-after strategy to construct alkene molecules is to apply an alkenyl  $sp^2$  C–H activation that can directly functionalize a simple alkene substrate without the pre-installation of a directing group.<sup>10,11</sup> It is often argued that a retrosynthetic scheme based on C–H functionalization is more straightforward and offers higher atom economy. Despite the high alkene  $sp^2$  C–H bond dissociation energy (110 kcal/mol, c.f. 101 kcal/mol of  $sp^3$  C–H bond dissociation energy),<sup>12</sup> we have witnessed a surge in research activities in the development of new, practical and broadly applicable methods for the alkenyl  $sp^2$  C–H bond functionalization to access structurally diverse alkenes and their derivatives that traditionally could only be obtained through tedious multistep synthesis.<sup>13–17</sup> Similar to alkenyl  $sp^2$  C–H bond, the alkenyl  $sp^2$  C–F bond functionalization of *gem*-difluoroalkenes have also been included in this review as it can be easily synthesized from carbonyl compounds and in many cases, the C–F bond functionalizations proceed through similar mechanisms with alkenyl C–H bond functionalizations. Over the past years, the alkenyl  $sp^2$  C–F bond functionalizations have attracted tremendous attention, especially in polymer synthesis. For example, teflon is readily synthesized via the homopolymerization of tetrafluoroethylene (TFE) which is an economical bulk organofluorine feedstock. Novel methods to selectively functionalize the alkenyl C–F bonds of multisubstituted fluorinated compounds such as tetrafluoroethylene (TFE) and its analogues will provide a wider access to many interesting fluorinated building blocks.<sup>18–26</sup> Arguably, the high bond dissociation energy poses challenges to selectively functionalize the alkenyl  $sp^2$  C–F bonds (130.6 kcal/mol for  $sp^2$  C–F bond dissociation energy of TFE),<sup>12</sup> especially in the presence of weaker C–H or C–X bonds, where X = Cl, Br, etc. However, many solutions have been successfully established such as the design of metal ligands with steric constraints. It should be mentioned that this review focuses on the alkenyl C–F bond functionalization of *gem*-difluoroalkenes, other C–X bond functionalization was not covered.<sup>27–30</sup>


Over the past several decades, transition metals such as Pd, Rh, Ir, Ru, Cu, Fe, Mn, and Co have been extensively employed to functionalize alkenyl  $sp^2$  C–H and C–F bonds. A large variety of different coupling partners such as  $\alpha,\beta$ -unsaturated systems, hypervalent iodine reagents, organometallics, etc., have been widely utilized in these reactions. In this review, we will focus on the latest advances in the development of such new and practical methods for the selective transition metal-catalyzed alkenyl  $sp^2$  C–H bond functionalization of alkene substrates and alkenyl  $sp^2$  C–F bond functionalization of *gem*-difluoroalkenes (Scheme 1).

### Scheme 1. Alkenyl $sp^2$ C–H and C–F Bond Functionalizations




Mechanism of alkenyl  $sp^2$  C–H activation by different transition metals has been extensively investigated by a myriad of research groups. In this section, we will highlight the general catalytic mechanism scheme for the reactions catalyzed by palladium and rhodium catalysts. For the Pd-catalyzed alkenyl C( $sp^2$ )–H activation, we can draw a generalized catalytic cycle as shown in Scheme 2. Initially, the palladium catalyst activates the electron-rich alkene substrate by the generation of vinyl palladium intermediate **A**. Then the incoming electron-deficient alkene coordinates to the intermediate **A**, which follows a migratory insertion into the palladium–C<sub>vinyl</sub> bond to produce the  $\sigma$ -Pd intermediate **B**. Finally,  $\beta$ -H elimination results in the formation of the expected coupling product **C**, while the oxidants employed in the reaction reactivates the catalyst for next catalytic cycle. It is important to note that the vinyl palladium species **A** can be either generated by a 1,2-addition process followed by deacetylation *or via* a direct C–H bond activation process. The difference in the reactivities of *Z*- and *E*-1,2-substituted  $sp^2$  C–H bonds of alkenes indicate that direct C–H activation is probably *involved* in this process.

### Scheme 2. General Mechanism of Palladium-Catalyzed Alkenyl $sp^2$ C–H Bond Functionalizations




The generalized mechanism of the directed alkenyl  $sp^2$  C–H activation of the reactions catalyzed by rhodium is depicted in Scheme 3. The rhodium catalyst activates the C–H bond by coordinating to the directing group to form the rhodium–substrate complex **A**, which in-turn inserts into the incoming functional groups to form the new C–FG bond **B**. The newly formed complex undergoes  $\beta$ -H elimination to furnish the coupling products **D**. Finally, the active rhodium catalyst is regenerated by the oxidants employed in the reaction.

### Scheme 3. General Mechanism of Rhodium-Catalyzed Directed Alkenyl $sp^2$ C–H Bond Functionalizations



### Scheme 4 The C–H bond functionalisation coupling processes involving photoredox catalysis or electrocatalysis promote radical reactions in the absence of radical initiators.

### Scheme 4. General Mechanism of Alkenyl $sp^2$ C–H Bond Functionalizations through Radical Processes



With respect to the mechanism of alkenyl  $sp^2$  C–F activation, activation of the  $sp^2$ -hybridized C–F bond mostly occurs via addition–elimination pathway since gem-difluoroalkenes in particular are highly reactive toward nucleophilic attack at the fluorinated  $sp^2$ -carbon.

Overall, this comprehensive review will be organised according to the types of olefins used in the reactions. In contrast to alkenyl  $sp^2$  C–H bond functionalization, we organize the alkenyl  $sp^2$  C–F bond functionalization section according to the cross-coupling partners, and methods published prior to the end of June 2022 are presented.


### 2. ALKENYL C–H BOND FUNCTIONALIZATION OF SIMPLE ALKENES

As mentioned above, alkenes and their derivatives are important compounds in organic synthesis. In this part of the review, we will focus on the development of the synthesis of complex alkenes and their derivatives starting from simple and easily accessible alkenes *via* alkenyl C(sp<sup>2</sup>)-H bond functionalization. It is organized based on the coupling reactions involving different types of simple alkenes. Strategies to control the stereochemistries (regio, *E/Z*, etc) as well as the intramolecular versions are also included.

### 2.1. Direct Alkenyl C-H Bond Functionalization of Simple Alkenes


In early 2000, Gusevskaya and co-workers reported a palladium-catalyzed dimerization of camphenes under aerobic oxidative conditions (Scheme 5).<sup>31,32</sup> In AcOH, employing Pd(OAc)<sub>2</sub>/Benzoquinone or LiNO<sub>3</sub>/O<sub>2</sub> (1.0 atm), the dimerized products were isolated.

#### Scheme 5. Palladium(II)-Catalyzed Dimerization of Camphenes




Although cross-coupling reaction between two different simple olefins is synthetically useful, it is significantly more difficult to realize. This is probably due to the difficulty to preferentially activate one alkenyl sp<sup>2</sup> C-H bond over the other. In search of new methods for the synthesis of conjugated dienes that are commonly encountered in numerous natural products and drug candidates with remarkable biological activities,<sup>33,34</sup> the first ever simple alkene-alkene cross-coupling reaction was disclosed by Loh's group in 2009. By exploiting the steric and electronic properties of different alkenes, they successfully established the first cross-coupling of this process by reacting  $\alpha$ -substituted styrenes with acrylates (Scheme 6a).<sup>35</sup> The direct cross-coupling reaction between simple alkenes and acrylates could be generally achieved in high efficiency with a catalytic amount of Pd catalyst under typically mild reaction conditions. In the mixture of DMSO/AcOH solvent, employing 20 mol% of Pd(OAc)<sub>2</sub> catalyst, 1 equivalent of Cu(OAc)<sub>2</sub> under O<sub>2</sub>, various conjugated 1,3-butadienes with different substitution patterns were smoothly synthesized in moderate to good yields (33–87%). Although an exceptional reaction was developed, the reported method has many limitations such as relatively high catalyst loading, low *E/Z* regioselectivities, and the need to use  $\alpha$ -substituted aryl alkenes, etc. In an attempt to render this strategy more general and convenient, Loh and colleagues in 2010 extended to establish a cross-coupling reaction between indenes and various electron-deficient alkenes using Pd(OAc)<sub>2</sub> (10 mol%) as the catalyst (Scheme 6b).<sup>36</sup>

### Scheme 6. Oxidative Cross-Coupling Reactions of 2-Substituted Alkenes with Acrylates



Exploring the generality of this strategy, Loh's group further disclosed a general protocol of the palladium-catalyzed cross-coupling reaction of readily available either TIPS-protected allylic or homoallylic alcohol substrates with a broad scope of acrylates, giving rise to a number of synthetically attractive conjugated dienyl alcohols in moderate to high yields with reasonable stereoselectivity. Gratifyingly, the presented strategy could be applied to the assembly of the key C<sub>13</sub>-C<sub>21</sub> fragment of palmerolide A (Scheme 7).<sup>37</sup>


#### Scheme 7. Palladium(II)-Catalyzed Cross-Coupling of Unactivated Alkenes with Acrylates



Encouraged by this success, Loh's group extended to describe a remarkable palladium-catalyzed direct cross-coupling reaction between simple diverse alkenes without

an  $\alpha$ -substitution and acrylates in the presence of a monoprotected amino acid (MPAA) as the ligand to afford the products with moderate to good *E/Z* selectivity (Scheme 8).<sup>38</sup> A broad substrate scope (33 examples) of this transformation was documented with yields of 35–86%. This atom- and step-economical strategy offers a direct method to construct new C–C bonds of synthetically useful dienes from readily available starting materials under benign conditions. It is interesting to note that Loh and co-workers also observed that the reactions with *Z*- and *E*-alkenes gave different results. In contrast to the *E*-alkenes, the *Z*-alkenes generates the products in good yields. On the basis of these results, they proposed that the alkenyl  $sp^2$  C–H activation mechanism is most probably operative in this type of reaction. An alternative mechanism involving 1,2-addition of alkene followed by facile elimination to generate the vinylic palladium was previously proposed as a possible catalytic mechanism. If this is the mechanism, the reaction should proceed similarly irrespective of the geometry of the alkenes.

### Scheme 8. Palladium(II)-Catalyzed Cross-Coupling Reactions of Simple Alkenes with Acrylates




Mechanistic studies provided further insights into the origins of this process. Later in 2016, Wu and co-workers combined DFT calculation and mass spectrometry to extensively elucidate on the possible mechanism of this Pd(II)/mono-*N*-protected amino acid (MPAA)-catalyzed alkene-alkene coupling reactions, and revealed that the strong binding of MPAA to Pd catalyst and basicity of the *N*-protecting group greatly facilitate the C–H activation step by stabilizing the active palladium catalyst, while the *E/Z* selectivity is mainly attributed to the interaction energy between the catalyst and alkene substrates.<sup>39</sup>

Due to the high potential for **polymerization**, unsaturated ketones such as methyl vinyl ketone (MVK) are rarely employed as the partners in cross-coupling reactions. In 2013, Loh's group introduced a general approach for the efficient synthesis of conjugated dienylic ketones enabled by palladium(II)-catalyzed cross-coupling reaction of simple alkenes with an array of vinyl ketones (Scheme 9).<sup>40</sup> Of note, the practicality of this strategy was illustrated by the straightforward synthesis of vitamin A1


and bornelone.

### Scheme 9. Palladium(II)-Catalyzed Oxidative Cross-Coupling of Alkenes and Vinyl Ketones



Since these initial reports, extensive studies by various research groups have been carried out to expand this alkene-alkene coupling reactions. For example, Liu and co-workers in 2012 developed a double vinylic C–H bond functionalization strategy to access diverse conjugated 1,3-dienes using Pd(OAc)<sub>2</sub> as the catalyst. The reaction occurred uneventfully with styrenes without an  $\alpha$ -substitution as well as a series of unactivated aliphatic alkenes, showcasing an elegant route for the synthesis of 1,3-butadienes (Scheme 10).<sup>41</sup> Nevertheless, this protocol has the disadvantages of having to use 15 mol% of Pd(OAc)<sub>2</sub> catalyst as well as 2.5 equivalent of AgOAc in order to obtain the coupling products in satisfactory yields.


### Scheme 10. Palladium(II)-Catalyzed C–H Olefination of Alkenes with Allylic Acetates and Acrylates



A more practical and efficient approach for the synthesis of conjugated dienes with low catalyst loading through biomimetic aerobic oxidative cross-coupling of two different C<sub>vinyl</sub>–H bonds has been developed by the Bäckvall group in 2013 (Scheme 11).<sup>42</sup> They employed only 5 mol% of Pd(OAc)<sub>2</sub> with catalytic amounts of electron-transfer mediators such as iron phthalocyanine and *p*-benzoquinone in presence of O<sub>2</sub> to achieve the synthesis of conjugated dienes with a broad set of alkene substrates.


### Scheme 11. Biomimetic Aerobic Oxidative Coupling of

## Simple Alkenes with Various Olefins



The palladium-catalyzed oxidative Heck reaction, a synthetically attractive transformation in which an olefinic C–H bond is generally converted into a C–C bond, has widespread applications in modern organic synthesis.<sup>43</sup> However, the overwhelming majority of these protocols require the use of electronically biased olefins such as acrylates or  $\alpha,\beta$ -unsaturated carbonyls in order to achieve high selectivity for the (*E*)-styrenyl products. In 2008, White et al. reported an efficient substrate chelate-controlled oxidative Heck reaction enabled by a versatile Pd(II)/bis-sulfoxide catalyst that proceeds smoothly with outstanding selectivities for non-resonance biased  $\alpha$ -olefins bearing proximal oxygen and nitrogen functionality, thus affording a wide range of coupling products in good yields (Scheme 12).<sup>44</sup>


## Scheme 12. Intermolecular Substrate Chelate-Controlled Oxidative Heck Reaction



Later in 2010, Sigman et al. developed a novel catalyst-controlled oxidative Heck reaction capable of generating a large variety of (*E*)-styrenyl products from electronically


nonbiased terminal alkenes with various arylboronic esters (Scheme 13).<sup>45</sup> The high selectivity observed under mild conditions is attributed to the Pd(*i*Pr)(OTs)<sub>2</sub> catalyst without the need to use base or high temperature. The authors performed preliminary mechanistic investigations, suggesting that the high selectivity may be mainly due to the catalyst's sensitivity to C–H bond strength in the key selectivity-determining  $\beta$ -H elimination step.

## Scheme 13. Intermolecular Catalyst-Controlled Oxidative Heck Reaction




As an extension of this approach, the same group further identified an efficient and simple reaction conditions for the (*E*)-styrenyl-selective Heck reactions of diverse electronically unbiased terminal olefins bearing a broad range of useful functionalities using Pd<sub>2</sub>dba<sub>3</sub> as the catalyst (Scheme 14).<sup>46</sup> Interestingly, the highly enantiomerically enriched olefin substrates that may be susceptible to racemization in many cases worked well under the mild conditions with no erosion of enantiomeric excess. Mechanistic investigations demonstrated that the  $\sigma$ -donating feature of DMA solvent is crucial for achieving high selectivity in this process. Meanwhile, Correia and co-workers also achieved a similar regio- and stereoselective substrate-directed arylations of allylamine derivatives with arenediazonium salts.<sup>47</sup> The application of this protocol is also illustrated by a short total synthesis of biologically active naftifine and abamine.

## Scheme 14. Highly (*E*)-Styrenyl-Selective Heck Reactions of Electronically Unbiased Olefins



By fine-tuning the steric and electronic factors of alkene substrates, Loh and co-workers were able to establish the divergent synthesis of indenes and  $\alpha$ -naphthols via an intramolecular rhodium(III)-catalyzed vinylic C–H bond functionalization (Scheme 15).<sup>48</sup> The intramolecular alkene-alkene coupling with acrylates as the internal coupling partners occurred smoothly to generate a series of highly decorated indenes with different substitution patterns in up to 93% yield, while the alkene-enolate coupling exclusively resulted in the formation of  $\alpha$ -naphthol derivatives. Detailed mechanistic investigations were also performed, and the results revealed that the reaction occurs via the vinylic  $sp^2$  C–H bond functionalization followed by the generation of a five-membered intermediate and possibly also a short-lived 3-membered intermediate which significantly depends on the coupling alkene functionalities.


**Scheme 15. Divergent Synthesis of Indenes and  $\alpha$ -Naphthols via  $Cp^*Rh(III)$ -Catalyzed Intramolecular Alkene-Alkene Coupling**



**2.2. Directed Alkenyl C–H Bond Functionalization of Aromatic Alkenes**


In addition to numerous examples of alkenyl C–H bond functionalizations of simple alkenes discussed above, remarkable efforts have been also devoted toward the chelation-assisted C–H bond functionalization of aromatic alkenes over the years. One of the major challenges in the  $sp^2$  C–H bond of aryl alkenes is to selectively control the  $sp^2$  C–H bond of the alkenes to yield either *E*- or *Z*-product. While it is generally easier to obtain the *E*-selective alkenyl metal complexes due to steric effect, the *Z*-selective product can be obtained via a chelating group (OH, NH, S, etc). Another appealing approach utilised the  $1,n$ -metal migration generated from an aryl halides or aryl metal reagents (Scheme 16).

**Scheme 16. General Scheme of Alkenyl C–H Bond Functionalizations of Aromatic Alkenes**




In 2015, Nachtsheim and colleagues reported the OH-assisted vinyl C–H alkynylation reaction of 2-vinylphenols with a modified hypervalent iodine(III) reagent (TIPS-EBX\*) under rhodium(III)-catalysis (Scheme 17).<sup>49</sup> The reactions proceeded smoothly under remarkably mild conditions with high functional group compatibility, excellent chemoselectivity, and exclusive (*Z*)-stereoselectivity, allowing the efficient synthesis of a diverse variety of highly substituted 1,3-enynes in 40–98% yields. The authors proposed that a base-assisted ligand exchange followed by the C–H activation step results in the formation of a rhodacycle intermediate. Subsequent regioselective insertion of the optimized hypervalent iodine (III) alkyne reagent (TIPS-EBX\*) produces the eight-membered rhodacycle species, which further undergoes facile  $\alpha$ -elimination and the extrusion of 2-iodo-6-methylbenzoate to give the key rhodium vinylidene complex. Finally, intramolecular concerted 1,2-migration accompanied by a ligand exchange releases the expected C–H alkynylation products, and regenerates the rhodium(III) catalyst.

**Scheme 17. Rhodium(III)-Catalyzed OH-Assisted C–H Alkynylation of 2-Vinylphenols with Ethynyl Benziodoxolones and Its Proposed Mechanism**




#### Proposed mechanism




Similarly, Zhang and co-workers carried out the vinylic C–H activation-desymmetrisation of 2-vinylphenols with diazabicycles (Scheme 18).<sup>50</sup> The products obtained were *tri*-substituted olefins with multiple functionalities. Mechanistically, the reaction was suggested to occur via alkenyl C–H activation and subsequent alkene insertion to produce the rhodacycle species. Then,  $\beta$ -N elimination occurs to deliver a key ring-opening *trans*-rhodium intermediate, which finally undergoes protonolysis to afford the alkenyl-substituted aminocyclopentenes.

#### Scheme 18. Rhodium-Catalyzed C–H Activation-Desymmetrisation of Diazabicycles with 2-Vinylphenols



Because of the distinct advantage of easy availability, and the fact that  $\text{NH}_2$  moieties indisputably serve as an ideal class of traceless directing groups for the synthesis of heterocyclic scaffolds, free amino functionality ( $\text{NH}_2$ ) has always been identified as one of the most appealing group in the area of alkenyl  $\text{C}(\text{sp}^2)\text{--H}$  activation reactions. Indeed, remarkable endeavors have been devoted towards the development of  $\text{NH}_2$ -assisted C–H bond functionalizations in recent years. Specifically, You and co-workers in 2009 reported their investigations on the amine-assisted vinylic C–H allylation of 2-aminostyrene derivatives with diverse allylic carbonates under  $[\text{Ir}(\text{COD})\text{Cl}]_2$ /Feringa's ligand catalysis, delivering a variety of (*Z/E*)-1,4-dienes with the formation of a *cis* double bond, which is complementary to Heck cross-coupling reaction (Scheme 19).<sup>51</sup> This protocol exhibited excellent functional group compatibility to both sides with the yields ranging from 51% to 99%. The authors also carried out mechanistic studies, and the results suggested that the leaving group of the allyl precursors is crucial for directing the reaction pattern in this case.


#### Scheme 19. Free Amine-Assisted Vinylic C–H Bond Allylation of Styrenes




Later in 2017, Nachtsheim et al. described an effective Ir(III)-catalyzed, free  $\text{NH}_2$ -directed C–H alkynylations of 2-vinylanilines by using alkynylbenziodoxolones as the efficient electrophilic alkyne transfer reagents, leading to the expedient synthesis of highly desirable 1,3-enynes in excellent yields of up to 97% with exclusive *Z*-stereoselectivity (Scheme 20a).<sup>52</sup> In the same year, they further extended to report the Ir(III)-catalysed, free  $\text{NH}_2$ -directed C–H alkenylation of 2-vinylanilines using alkenyl- $\lambda^3$ -iodanes as the electrophilic alkene-transfer reagents, enabling the synthesis of highly desirable 1,3-dienes in excellent yields (76–98%) with high to perfect (*Z,E*) stereoselectivity (Scheme 20b).<sup>53</sup>

#### Scheme 20. $\text{NH}_2$ -Directed C–H Bond Alkynylation and Alkenylation of 2-Vinylanilines

a) **Nachtsheim et al., 2017**





b) **Nachtsheim et al., 2017**



Despite considerable progress in alkenyl C(sp<sup>2</sup>)-H functionalizations of styrenes, the direct construction of atropisomeric axially chiral styrenes bearing a conjugated 1,3-diene moiety via asymmetric alkenyl C-H olefination strategy remains a great long-standing challenge to overcome over the past decades. Very Recently, Shi et al. pioneered to report a highly efficient and atom-economical synthesis of atropisomeric styrenes via an enantioselective palladium(II)-catalyzed thioether-directed alkenyl C-H olefination (Scheme 21).<sup>54</sup> The reaction occurred through the formation of an *endo*-palladacycle intermediate using chiral spiro phosphoric acid as the chiral ligand under mild conditions. Notably, this stereospecific strategy features high functional group tolerance, broad substrate scope (69 examples), excellent enantioselectivities (up to 99% ee), and complete *Z*-selectivity control. More importantly, diverse chiral sulfoxide derivatives could be easily obtained by the high diastereoselective oxidation of the resulting axially chiral styrenes, thus providing an alternative handle for the enantioselective synthesis of axial chiral sulfur-olefin ligands, which have been well recognized as efficient and versatile chiral ligands in many catalytic asymmetric synthesis.<sup>55-58</sup> In a very recent report, Xu and co-workers achieved an analogous tosylamine-group-directed transformation in the presence of a chiral tridentate Pybox ligand.<sup>59</sup>


### Scheme 21. Palladium-Catalyzed Thioether-Directed Atroposelective Alkenyl C-H Olefination




Despite of the indisputable advantages of chelating group strategies, the laborious pre-installation and subsequent removal of the directing group sometimes significantly compromise the overall efficiency of the transformations. To address this issue, a large number of transient directing group (TDG) strategies have been devised over the past years.<sup>60-65</sup> Quite recently, by taking advantage of the chiral transient directing group (CTDG) strategy,<sup>66,67</sup> Zhang and co-workers achieved the asymmetric vinylic C-H olefination of 2-vinyl benzaldehydes through a challenging seven-membered *endo*-cyclometallation process. Successfully, an amino acid-derived transient chiral auxiliary was identified in this case (Scheme 22a).<sup>68</sup> A broad range of 2-vinyl benzaldehydes reacted smoothly with various acrylates and styrenes to afford the corresponding coupling products in high yields (up to 93%) and excellent enantioselectivities (up to > 99% ee). However, electronically activated alkenes such as vinyl ketones, vinyl sulfones, and acrylamides were proven to be incompatible with the conditions. Unlike this asymmetric coupling reactions where *endo* metallacycles were generated to regioselectively functionalize the vinylic β-C(sp<sup>2</sup>)-H bonds of styrene derivatives, the same group expanded to establish the asymmetric α-C(sp<sup>2</sup>)-H alkenylations of styrenes using *L*-*t*-leucine as the transient chiral auxiliary through a six-membered *exo*-cyclopalladation process (Scheme 22b).<sup>69</sup>

### Scheme 22. Synthesis of Axially Chiral Aryl 1,3-Dienes by a Chiral Transient Directing Group

a) Zhang et al., 2022




b) Zhang et al., 2022



Almost simultaneously, Engle and co-workers also reported their exploration of a TDG approach to realize the feasibility of palladium(II)-catalyzed  $\alpha$ -C–H olefination of styrenes by the generation of an *exo* alkenyl palladacycle, providing access to diverse 2-aryl-substituted 1,3-dienes with high regio- and *E/Z*-selectivity (Scheme 23).<sup>70</sup> In this report, commercially available *tert*-leucine was identified as the optimal transient directing group, and alkenyl aldehydes with different substitution patterns could be readily olefinated to the expected coupling products. Notably, unbiased aliphatic alkenes were found to be viable substrates for this transformation, albeit with diminished efficiency. Mechanistic studies including synthesis of two catalytically relevant alkenyl palladacycle complexes and DFT calculations were also carried out to elucidate the plausible mechanism.


### Scheme 23. Pd(II)-Catalyzed Alkenyl C–H Olefination of Aldehydes by a Transient Directing Group



Around the same time, Zhang et al. investigated the *N,N*-bidentate-chelation-assisted  $\alpha$ -selective vinylic C–H bond functionalization of styrene derivatives (Scheme 24).<sup>71</sup> The 8-aminoquinoline (AQ) was readily employed as the bidentate directing group of best choice. By the combination of  $\text{Pd}(\text{OAc})_2$ /benzoquinone (BQ),  $\text{MnO}_2$ ,

oxidant, and PivOH additive in EtOH at 60 °C, a wide substrate scope (25 examples) of vinylic  $\alpha$ -C(sp<sup>2</sup>)–H olefination product was documented in satisfactory yields of up to 92%. Interestingly, using 2-pyrazinamide (PC) as the efficacious chelating group, the authors could establish an aerobic approach for  $\alpha$ - and  $\beta$ -bis-alkenylations of styrenes under identical conditions. As expected, a series of  $\beta$ -alkenylation products were formed with  $\alpha$ -substituted styrenes as substrates.



### Scheme 24. Highly Regio- and Stereoselective Vinylic C–H Functionalizations of Styrenes



### 2.3. Migratory Alkenyl C–H Bond Functionalization of Aromatic Alkenes

In addition to the aforementioned chelation-assisted strategy for the activation and further functionalization of styrene olefinic C–H bonds, strategies based on 1,*n*-metal shift allow the direct functionalization of specific distal C–H bonds which may be otherwise difficult to achieve in many cases. Inspired by the seminal work of Larock illustrating the possible vinyl to aryl palladium 1,4-migration as a reversible process in the migration as a plausible mechanism,<sup>72</sup> the group of Lin and Feng made a significant contribution to the field of migratory alkenyl C–H bond functionalization of aromatic alkenes. In 2016, they elegantly reported the first example of aryl to vinyl 1,4-palladium migration process (Scheme 25).<sup>73</sup> According to the proposed mechanism, the oxidative addition of aryl bromide to palladium(0) generates the alkenyl palladium(II) intermediate which further reacted with diboron reagents under Miyaura borylation conditions, affording a novel method to the rapid and diverse synthesis of remarkably useful  $\beta,\beta$ -disubstituted vinylboronates with broad scope and excellent regioselectivity.


### Scheme 25. Alkenyl C–H Bond Borylation via Aryl to Vinyl Palladium 1,4-Migration




In continuation, Lin, Feng, and co-workers expanded to apply their reliable migration strategy to the highly stereoselective synthesis of conjugated 1,3-butadienes via aryl to vinyl palladium 1,4-migration/Heck sequence of 2-bromostyrene substrates with diverse electron-deficient olefins (Scheme 26a),<sup>74</sup> and later styrenes (Scheme 26b).<sup>75</sup> Interestingly, this new methodology could also be applied to efficiently synthesize a series of trisubstituted conjugated 1,3,5-hexatriene derivatives with modest to decent yields (Scheme 26c),<sup>76</sup> which are ubiquitous structures encountered in many natural products and organic functional materials.<sup>77-79</sup>

### Scheme 26. Synthesis of Di- and Trienes via Aryl to Vinyl 1,4-Palladium Migration/Heck Sequences


a) Feng, Lin et al., 2018



b) Feng, Lin et al., 2021





c) Feng, Lin et al., 2020



In 2020, an efficient aryl to vinyl palladium 1,4-migration/Suzuki-Miyaura sequence of 2-bromostyrene substrates with organoboron reagents was also achieved by Lin, Feng, and co-workers (Scheme 27).<sup>80</sup> In this protocol,



various phenylboron reagents were tested, and five-membered ethylene glycol-derived boronate esters was found to be the best coupling partner to afford the olefinic C(sp<sup>2</sup>)-H arylation products. The coupling reaction with various commercially available arylenyl boronates enables the stereospecific synthesis of multisubstituted conjugated 1,3-butadienes. Of note, the reaction features easy scalability, extraordinarily broad substrate scope (>90 examples), excellent functional group tolerance, and versatile utility of the corresponding products.

### Scheme 27. Suzuki-Miyaura Coupling Enabled by Aryl to Vinyl 1,4-Palladium Migration





As an extension of this approach, Lin, Feng, and co-workers recently continued this 1,4-palladium migration strategy to accomplish a highly regioselective palladium-catalyzed three-component tandem C-H alkylation/cross-coupling reaction of 2-iodostyrenes with alkyl halides as the electrophile and various easily accessible nucleophiles as the coupling reagents (Scheme 28).<sup>81</sup> The reaction occurs through a cyclopalladation process enabled by the judicious choice of electron-rich 2-pyridone ligand with regiodetermining C-H alkylation of the key aryl-alkenyl-palladacycle species which was *in situ* captured and well-identified by NMR and X-Ray spectra.

### Scheme 28. Regioselective Three-Component Tandem C-H Alkylation/Cross-Coupling Reaction of *ortho*-Iodophenylethenes




Subsequently, Yu and co-workers illustrated an efficient palladium-catalyzed olefinic C–H bond alkylation of 2-bromostyrenes with cyclobutanols as the coupling partners enabled by a controllable aryl to vinyl 1,4-palladium migration/ring-opening C–C cleavage cascade strategy (Scheme 29).<sup>82</sup> This protocol features excellent regio- and stereoselectivity, wide substrate scope, and high functional group compatibility, providing access to  $\gamma$ -ketone olefins in decent yields (43–95%). Combined mechanistic investigations and DFT calculations demonstrated that the reaction occurs via oxidative addition, aryl to vinyl palladium 1,4-migration, ring-opening C–C bond cleavage, and reductive elimination sequences. It should be mentioned that the high efficiency of this protocol is mainly attributed to the thermodynamically favored aryl to vinyl 1,4-palladium migration effectively assisted by the 2-fluorophenol additive.

### Scheme 29. Alkenyl C–H Bond Alkylation through an Aryl to Vinyl 1,4-Palladium Migration/C–C Cleavage Cascade



The Huang group established a stereospecific olefinic C–H cyanation by using *o*-(alkenyl) substituted aryl boronic acids as the substrates combined with easily accessible *N*-cyano-*N*-phenyl-*p*-methylbenzenesulfonamide (NCTS) as the cyanating reagent. The transformation occurred through an aryl to vinyl 1,4-rhodium migration process, giving rise to a diverse variety of  $\beta,\beta$ -disubstituted acrylonitriles in up to 99% yield (Scheme 30).<sup>83</sup> Of note, the reaction features a broad substrate scope and excellent functional group tolerance. The proposed mechanism involves the facile formation of an alkenylrhodium intermediate that was readily generated in situ via 1,4-rhodium migration from aryl to vinyl, accompanied by an electrophilic cyanation step.

### Scheme 30. Stereospecific Olefinic C–H Cyanation Enabled by 1,4-Rhodium Migration





### 2.4. Annulation of Aromatic Alkenes via Alkenyl C–H Bond Functionalization

The annulation processes of styrenes via vinylic C–H bond functionalization are especially useful methods since they provide an easy and straightforward access to various

polycyclic compounds. Different types of annulation reaction to form pyrans, latones, lactams, as well as diverse spiro rings are covered in this section.



In 2017, Sheykhani, Abbasnia, and their co-workers reported an unprecedented annulation of simple styrenes with readily available electron-deficient maleimides and maleic anhydrides as coupling partners via a tandem activation of both aromatic and olefinic C–H bonds in a pseudo-Diels–Alder mode, allowing an economical and convenient synthesis of Benzo[*e*]isoindole-1,3-diones in 63–84% yield (Scheme 31).<sup>84</sup>

**Scheme 31. Synthesis of Benzo[*e*]isoindole-1,3-diones**





In an effort to expand the scope of 1,4-palladium migration strategy, the group of Lin and Feng further established a sequential cross-coupling/annulation of 2-bromostyrene derivatives with internal alkynes, allowing a convenient approach to the synthesis of multisubstituted naphthalenes via a controllable aryl to vinyl 1,4-palladium migration process (Scheme 32).<sup>85</sup> Later, they continued to apply this sequential cross-coupling/annulation strategy of 2-bromostyrenes with aromatic bromides to identify a direct approach for the facile synthesis of polycyclic aromatic compounds in yields ranging from 32% to 86% with excellent chemo- and regioselectivity (Scheme 33).<sup>86</sup> On the basis of SAESI-MS analysis, a vinyl-coordinated palladacycle species was tentatively proposed as the key intermediate for this 1,4-palladium migration and further annulation processes. The authors also performed the kinetic isotope effect (KIE) experiments, and an intermolecular KIE value of 1.68 unambiguously suggests that the alkenyl C–H bond cleavage was probably involved in the rate-determining step in this case.

**Scheme 32. Synthesis of Naphthalene via Benzannulation of (2-Aryl)vinyl Metal Species and Internal Alkynes**





**Scheme 33. Sequential Cross-Coupling/Annulation of 2-Bromostyrenes with Aromatic Bromides**




In 2019, a practical and step-economical approach to construct a diverse range of highly valuable arylidene  $\gamma$ -lactams and indanone derivatives through a palladium(o)-catalyzed intramolecular cascade reaction involving 1,4-palladium migration and subsequent C(sp<sup>3</sup>)–H activation has been reported by Baudoin's group (Scheme 34).<sup>87</sup> As a particular highlight, the applicability of this protocol was demonstrated as a key step with an appreciable 50% yield in the formal synthesis of a pyrrolizidine alkaloid (–)pyrrolam A.

**Scheme 34. Synthesis of  $\alpha$ -Arylidene  $\gamma$ -Lactams via 1,4-Palladium Shift/C(sp<sup>3</sup>)–H Activation Strategy**





#### Formal Synthesis of (-)-Pyrrolam A



Reaction conditions: (a) (COCl)<sub>2</sub> (1.5 equiv), Et<sub>3</sub>N (2.0 equiv), DCM, 20 °C. (b) Pd(PCy<sub>3</sub>)<sub>2</sub> (10 mol%), PivOH (30 mol%), Rb<sub>2</sub>CO<sub>3</sub> (1.5 equiv), mesitylene, 160 °C, 18 h.



Phenols are a highly conserved functional group in natural and synthetic chemistry.<sup>88</sup> Strategies to construct heterocycles have also been reported. For examples, methodologies that selectively functionalize a specific C–H bond of phenols can dramatically alter molecular diversity and its function.<sup>89</sup> Particularly, the direct OH-assisted olefinic C–H cyclocarbonylation of 2-vinylphenols could provide an efficient approach for the synthesis of coumarins which are a ubiquitous class of natural products with a broad variety of biological activities.<sup>90</sup> In this regard, Alper and colleagues in 2012 first illustrated an attractive Pd(II)-catalyzed direct oxidative cyclocarbonylation of 2-vinylphenols with remarkably low pressure of CO ( $\leq 100$  psi), enabling the preparation of coumarins bearing a diverse array of functional groups in up to 85% yield by using air or 1,4-benzoquinone (BQ) as the terminal oxidant (Scheme 35).<sup>91</sup>

#### Scheme 35. Palladium(II)-Catalyzed Oxidative Cyclocarbonylation of 2-Vinylphenols





In 2014, Gulías, Mascareñas, and co-workers achieved the rhodium(III)-catalyzed C–H activation/annulation of readily available 2-hydroxystyrenes with CO as the one-carbon coupling partner for the construction of coumarin scaffolds (Scheme 36).<sup>92</sup> Subsequently, the same group further expanded this annulation strategy to large-scale synthesis of coumarins. The process was straightforward and economical for the assembly of coumarins with different substitution patterns in 53–89% yield. Of note, the utility of this protocol was also demonstrated by synthesizing two biologically important coumarin derivatives (Scheme 36).<sup>93</sup>

#### Scheme 36. Synthesis of Coumarins via Rhodium(III)-Catalyzed C–H Activation/Annulation





Shortly after, the Wang research group reported the cyclocarbonylation of 2-alkenylphenols with CO under cost-effective Cp\*Co(III) catalysis (Scheme 37).<sup>94</sup> The reaction proceeded under typically mild conditions. Using only balloon pressure CO could achieve high reactivity in this case. The utility of this approach is highlighted by a practical gram-scale preparation and the total synthesis of three natural products herniarin, xanthyletin, and seselin.

**Scheme 37.** Cp\*Co(III)-Catalyzed Annulations of 2-Alkenylphenols with CO




The direct C–H bond carboxylation reaction using easily accessible, inexpensive, naturally abundant atmospheric CO<sub>2</sub> represents a straightforward method for the atom- and step-economical synthesis of diverse value-added carboxylic acids or their derivatives.<sup>95–100</sup> In 2013, Iwasawa and co-workers described the first example of alkenyl C–H carboxylation of 2-hydroxystyrenes in the presence of 5 mol% of Pd(OAc)<sub>2</sub> in diglyme at 100 °C under an atmospheric pressure of CO<sub>2</sub> with Cs<sub>2</sub>CO<sub>3</sub> as a base, producing the expected coumarins in high yields (Scheme 38).<sup>101</sup> In this case, the hydroxy group served as an efficacious directing group for the C–H activation step. The authors proposed a plausible reaction mechanism involving the following pathways: the six-membered alkenyl palladium(II) species is readily produced by the chelation-assisted alkenyl C–H bond cleavage of 2-hydroxystyrene with a Pd(OAc)<sub>2</sub> catalyst, along with the coordination of another molecule of 2-hydroxystyrene as its caesium salt. The alkenyl palladium(II) species then undergoes a reversible nucleophilic carboxylation process to generate the key palladium carboxylate intermediate, which finally reacts with another molecule of 2-hydroxystyrene as well as the base to deliver the coumarin derivatives with the regeneration of the cyclometalated intermediate.

**Scheme 38.** Pd(II)-Catalyzed Carboxylation of Alkenyl C–H Bonds with CO<sub>2</sub> and Its Proposed Mechanism




**Proposed mechanism**




In 2017, Zhi, Yu, and co-workers realized a transition-metal-free alkenyl C–H bond lactonization of 2-vinylphenols with CO<sub>2</sub> as the ideal one-carbon (C<sub>1</sub>) source to synthesize highly valuable coumarin derivatives under redox-neutral conditions (Scheme 39a).<sup>102</sup> Later in 2021, the Chen group also reported an analogous cascade radical addition/cyclization reaction of diverse 2-alkenylphenols with readily available CBr<sub>4</sub> as the one-carbon source in the presence of water under photoredox catalysis (Scheme 39b),<sup>103</sup> providing an efficient access to 4-arylcoumarins in a one-pot fashion with a wide substrate scope. Mechanistic investigations confirmed that the reaction should proceed through an oxidizing quenching radical-involving pathway.

**Scheme 39. Direct Alkenyl C–H Carboxylation of 2-Alkenylphenols**

a) Zhi, Yu et al. 2017





b) Chen et al. 2021




The formal [5 + 2] cycloaddition of 2-vinylphenols with diverse alkynes to readily assemble highly appealing benzoxepines in an atom-economical manner was first reported by the Gulías and Mascareñas group in 2014 by using a catalytic amount of [RhCp\*Cl<sub>2</sub>]<sub>2</sub> (2.5 mol%) as a catalyst in conjunction with 0.5 equiv of Cu(OAc)<sub>2</sub>•H<sub>2</sub>O as an oxidant. Notably, the reaction accommodated a variety of substituents in the aryl group of vinylphenol substrates (Scheme 40).<sup>92</sup>

**Scheme 40. Synthesis of Benzoxepines via Cp\*Rh(III)-Catalyzed Formal [5 + 2] Cycloadditions**





The same year, Gulías and Lam independently reported the dearomatizing oxidative annulation reaction of various 2-alkenylphenols with internal alkynes under Rh(III) catalysis, generating the highly functionalized spirocyclic enones with good yields and excellent regioselectivities (Scheme 41a and b).<sup>104,105</sup> Interestingly, conjugated 1,3-enynes were also proved to be viable substrates for the dearomatizing oxidative annulation in Lam's work. As expected, a wide range of spirocyclic products were obtained in high yields (72–95%), albeit the need to use a superstoichiometric amount of Cu(OAc)<sub>2</sub> (2.1 equiv) as the oxidant. The reaction was proposed to proceed via the cleavage of alkenyl C–H bond and the dearomatization of the phenol aromatic ring. Recently, Lu and co-workers performed density functional theory calculations to investigate the competing formal [5 + 2] annulation and dearomatizing [3 + 2] annulation pathways, and the results clearly showed the direct reductive elimination of eight-membered rhodacycles is disfavored, and instead the dissociation of the Rh–O bond followed by antarafacial nucleophilic attack is the most favorable pathway for the generation of seven-membered azacyclic and spirocyclic products.<sup>106</sup> In a related report, Gogoi et al. also realized a similar transformation under ruthenium(II) catalysis.<sup>107</sup>

**Scheme 41. Synthesis of Spirocyclic Enones by Rhodium(III)-Catalyzed Dearomatizing Oxidative Annulation**




More recently, the Ackermann group achieved the elegant synthesis of highly appealing seven-membered benzoxepine derivatives through electro-oxidative rhodium-catalyzed [5 + 2] annulation reactions (Scheme 42).<sup>108</sup> The reaction proceeds via dual activation of C–H and O–H bond with molecular hydrogen as the sole byproduct. The expected benzoxepine products were synthesized with a diverse range of functional group compatibility. Detailed mechanistic studies were also performed, and monitoring the reaction by NMR spectroscopy clearly revealed that a low-valent rhodium intermediate was likely the catalyst resting state. A key benzoxepine-coordinated rhodium(I) sandwich complex prepared and isolated by the reaction of Cp\*Rh(OAc)<sub>2</sub> with the substrates unambiguously provided a solid support for a fast C–H rhodation and a rhodium(III/I) regime in this annulative transformations.

**Scheme 42. Electrooxidative Rhodium-Catalyzed [5 + 2] Annulations of 2-Vinylphenol with Alkynes**





Apart from above-mentioned alkynes and enynes, ynamides as synthetically valuable subgroup of alkynes,<sup>109–111</sup> can also couple with 2-alkenylphenols in a highly regioselective manner. In 2018, Li and colleagues first reported the oxidative [5 + 2] annulation reaction of 2-vinylphenols with ynamides to fabricate valuable 2-aminobenzoxepines under earth-abundant cobalt catalysis via C–H/O–H bond functionalizations (Scheme 43).<sup>112</sup> Shortly after, Wang, Li, and their co-workers extended this method to a formal dearomative [3 + 2] spiroannulation reaction to generate highly decorated spiro[4,5]decane derivatives bearing a quaternary stereogenic centre entirely made of carbons. The reaction worked under especially mild conditions with a broad variety of functional group compatibility (Scheme 43).<sup>113</sup> The coordinating characteristic of the sulfonyl group in the ynamides may determine the regioselectivity of the reaction.

**Scheme 43. Cobalt-Catalyzed Oxidative [5 + 2] Annulation and Dearomative [3 + 2] Spiroannulation of 2-Alkenylphenols with Ynamides**





In 2015, the group of Gulías and Mascareñas reported a simple and atom-economical Rh(III)-catalyzed oxidative [5 + 1] annulation of 2-vinylphenols with allenes as one-carbon coupling partners to afford a series of biologically active, six-membered 2,2-disubstituted 2*H*-chromenes, formally involving the cleavage of the C–H and O–H bonds of 2-alkenylphenol substrates (Scheme 44a).<sup>14</sup> Similar to this transformation, Cheng and co-workers also realized the present phenolic OH-assisted C–H oxidative annulation reactions by using non-noble [CoCp\*(CO)<sub>2</sub>] as the catalyst under mild conditions (Scheme 44b).<sup>15</sup> This annulative reaction is proposed to proceed via the initial alkenyl C–H activation, allene insertion, and an unusual intramolecular regioselective phenoxide addition. The authors argued that the Co–O bond in the  $\pi$ -allylic moiety could have been selectively attacked during the oxidative [5 + 1] annulations to afford the phenoxide intermediate.

**Scheme 44. Synthesis of 2,2-Disubstituted 2*H*-Chromenes via Formal [5 + 1] Annulations Between 2-Alkenylphenols and Allenes**





Quite recently, Prabhu and co-workers studied the [5 + 1] annulation reactions of 2-alkenylphenols with various maleimides under Rh(III) catalysis in the presence of Cu(OAc)<sub>2</sub> (2.0 equiv) as the terminal oxidant, exclusively giving rise to highly functionalized spirocyclic scaffolds bearing an oxygen-containing spiro carbon (Scheme 45).<sup>16</sup>

**Scheme 45. Rhodium(III)-Catalyzed [5 + 1] Annulation of 2-Alkenylphenols with Maleimides**



In contrast to the above results obtained with Rh and Co catalysts which afforded chromene-like coupling products, Gulías and Mascareñas et al. demonstrated that 2-alkenylphenols could couple with allenes via formal [5 + 2] cycloadditions in the presence of Pd(OAc)<sub>2</sub> catalyst to produce the seven-membered benzoxepines in decent yields (60–97%) with good regio- and diastereoselectivities (Scheme 46).<sup>17</sup> Detailed computational studies demonstrated that the different outcome of the reaction by Pd(II) or Cp\*Rh(III) catalysts could be ascribed to the geometric requirements associated with their respective square planar or pseudo-octahedral geometries. The square planar geometry of palladium catalyst required in this protocol is generally favored to undergo a reductive elimination process to generate benzoxepine products.


**Scheme 46. Palladium(II)-Catalyzed Formal [5 + 2] Annulations of *ortho*-Alkenylphenols with Allenes**



Moreover, propargyl carbonates can also engage in the rhodium(III)-catalyzed oxidative [5 + 2] annulation with various 2-alkenylphenols, exclusively giving the 3-alkenylated benzoxepine derivatives with excellent regioselectivity and high functional group compatibility


(Scheme 47).<sup>118</sup> Combined DFT calculations and experimental mechanistic studies clearly revealed the essential role of the traceless assisting OBoc group in controlling the regio- and chemoselectivities of this transformation in terms of its steric hindrance and good leaving ability.

### Scheme 47. Rhodium(III)-Catalyzed Oxidative [5 + 2] Annulations of 2-Alkenylphenols with Propargyl Carbonates




Unlike the case with propargyl carbonates, Reddy and colleague demonstrated that unprotected propargyl alcohols could couple with a series of 2-alkenyl phenols to assemble diverse naphtho furan derivatives through a migratory three-point double annulation process (Scheme 48).<sup>119</sup> Under the optimal conditions, a broad range of fused naphtho furans were documented in reasonable yields up to 86%. Of note, this protocol was also applicable in a more complex setting. It should be mentioned that unsubstituted or disubstituted 2-alkenyl phenols failed to deliver the expected coupling products.

### Scheme 48. Rh(III)-Catalyzed Double Annulation of 2-Alkenylphenols with Unprotected Tertiary Propargyl Alcohols




An efficient approach for the concise construction of 2,3-diarylbenzofurans by coupling 2-hydroxystyrenes with iodobenzenes through a C–H activation/oxidation cascade reaction has been reported by Jia and co-workers (Scheme 49).<sup>120</sup> By using this strategy, the overall efficiency of the direct synthesis of natural products decursivine, serotobenine and their analogs was greatly improved without the need of protecting group.

### Scheme 49. Synthesis of 2,3-Diarylbenzofurans via Alkenyl C–H Activation/Oxidation Tandem Reaction of 2-Hydroxystyrenes with Iodobenzenes




Along with their novel strategy to synthesize coumarins via the direct oxidative cyclocarbonylation with carbon monoxide,<sup>91</sup> Alper and co-workers extended to disclose an oxidative C–H cyclocarbonylation of various *N*-monosubstituted-2-vinylanilines with CO, providing a general and practical approach for the synthesis of a variety of biologically active 2(1*H*)-quinolinone derivatives in up to 97% yield (Scheme 50).<sup>121</sup>

### Scheme 50. Palladium(II)-Catalyzed Direct Oxidative Cyclocarbonylation of 2-Vinylanilines with CO



More recently, the impressive example of transition metal-free carbonylative C–H [5 + 1] annulation of 2-alkenylanilines with environmental and user-friendly dioxazolones as the carbonylating reagents was realized by Nan and co-workers (Scheme 51).<sup>122</sup> This protocol allows the rapid synthesis of highly privileged quinolinone derivatives through an unprotected NH<sub>2</sub>-assisted olefinic C–H bond functionalizations. A diverse series of value-added products were obtained in appreciable to high yields (35–87%) with excellent functionality tolerance.





### Scheme 51. Metal-Free C–H [5 + 1] Carbonylation of 2-Alkenylanilines with Dioxazolones



In 2017, Yu and co-workers reported the first example of transition metal-free lactamization reaction of olefinic C–H bonds of unprotected 2-alkenylanilines with nontoxic and readily available carbon dioxide as the ideal carbonyl source in the presence of alkali metal *tert*-butoxides to synthesize electronically and sterically diverse 2-quinolinones in modest to excellent yields (Scheme 52).<sup>123</sup> This efficient and eco-friendly process exhibited a broad substrate scope, high functional group compatibility, and facile scalability, which represents an appealing method for the pharmaceutical industry to the rapid synthesis of drugs and bioactive molecules.



### Scheme 52. Transition-Metal-Free Lactamization of 2-

### Alkenylanilines with CO<sub>2</sub>





Later in 2019, the group of Ma and Nan disclosed an unprecedented Ru(II)-catalyzed [5 + 1] annulation of 2-alkenylanilines with readily available sulfoxonium ylides as one-carbon coupling partners through the NH<sub>2</sub>-assisted olefinic C–H functionalization strategy (Scheme 53).<sup>124</sup> This new ruthenium-catalyzed annulation reaction employs synthetically ideal free amino functionality as a traceless directing group to assist the vinylic C–H bond functionalizations under aerobic conditions, providing a straightforward construction of 2-acylquinolines which are widespread in biologically active pharmaceuticals. Quite recently, Yu, Wang, and their co-workers also achieved the same transformation by an operationally simple, photo-thermo-mechanochemical approach under iron(II) phthalocyanine catalysis that remarkably obviates the use of any solvent or harsh reaction conditions.<sup>125</sup>

### Scheme 53. Ru-Catalyzed Formal [5 + 1] Annulation of Alkenylanilines with Sulfoxonium Ylides





A general palladium-catalyzed oxidative coupling of 2-vinylanilines with isocyanides was achieved by Zeng and co-workers, providing an exceptionally efficient and straightforward approach for the preparation of a variety of highly decorated 2-aminoquinolines (Scheme 54a),<sup>126</sup> which represent an important class of heterocyclic motifs exhibiting a broad range of biological and medicinal activities.<sup>127</sup> Moreover, this isocyanide insertion reaction of 2-vinylanilines was also realized with comparable efficiency by the Ji and Wang group in the presence of cost-effective  $\text{Co}(\text{acac})_2$  as the catalyst and molecular oxygen as the oxidant (Scheme 54b).<sup>128</sup>

#### Scheme 54. Rapid Synthesis of 2-Aminoquinolines by Oxidative Isocyanide Insertion with 2-Vinylanilines




Recently, Ding's group reported a  $\text{Cu}(\text{II})$ -catalyzed [5 + 1] oxidative annulation of free 2-vinylanilines with glyoxylic acid for the synthesis of 4-arylated quinolines (Scheme 55).<sup>129</sup> Here, a diverse range of 4-arylated quinolines were synthesized in excellent yields. Of note, the reaction was typically finished within 1.5 h at 80 °C in DMSO. The authors proposed that the  $\text{Cu}(\text{II})$  catalyst mediated the nucleophilic addition of aniline to the glyoxylic acid to produce an intermediate, which was subsequently dehydrated to generate an imine intermediate. Then, a  $6\pi$ -electrocyclization of the imine afforded the corresponding 2,3-dihydroquinolines. Further oxidative aromatization and the expulsion of  $\text{CO}_2$  resulted in the formation of the desired quinoline products.

#### Scheme 55. Copper-Mediated Formal [5 + 1] Annulation of 2-Vinylanilines and Glyoxylic Acid




In 2017, Mascareñas' group developed an elegant  $\text{Pd}(\text{II})$ -catalyzed [5 + 2] annulation of *ortho*-alkenylanilides and allenes for the direct synthesis of 2,3-dihydro-1*H*-benzo[*b*]azepines (Scheme 56).<sup>130</sup> The reaction was highly regio- and diastereoselective, affording the expected products in decent yields. As to the reaction mechanism, preliminary mechanistic studies showed that the C–H activation of the anilides occurred via a traditional metalation-deprotonation pathway. Specifically, an initial ligand exchange with the  $\text{Pd}(\text{II})$  catalyst and the anilide kickstart the reaction to generate the  $\text{Pd}(\text{II})$  complex intermediate followed by a C–H activation to afford a six-membered palladacycle. Subsequent coordination and regioselective migratory insertion of the allene moiety produced a  $\pi$ -allylic palladacycle. Finally, reductive elimination afforded the corresponding annulation products. Almost at the same time, Zeng and co-workers also achieved this oxidative annulation reactions with excellent yields.<sup>131</sup>

#### Scheme 56. Pd-Catalyzed Formal [5 + 2] Annulation of *ortho*-Alkenylanilides with Allenes and Its Proposed Mechanism




#### Proposed mechanism



Quite recently, Volla et al. established a non-oxidative protocol for the [5 + 1] annulation reaction of 2-alkenylanilides (Scheme 57).<sup>132</sup> The reaction occurred smoothly at room temperature with allenyl acetates under Rh(III) catalysis, thereafter affording a number of highly functionalized 1,2-dihydroquinolines in excellent yields (76–91%). On the basis of mechanistic studies, the authors tentatively proposed a catalytic cycle involving highly regioselective 2,3-migratory insertion, facile  $\beta$ -oxygen elimination followed by intramolecular nucleophilic cyclization to elucidate the plausible pathway. However, it is worth noting that both 1,3-disubstituted allenes and aliphatic allenes were found to be incompatible with this protocol. Moreover, by slightly modifying the conditions, the authors successfully expanded this [5 + 1] annulation strategy to 2-alkenylphenol substrates. A series of allenyl acetates with different substitution patterns were coupled to deliver the appealing chromene derivatives.


#### Scheme 57. [5 + 1] Annulation of 2-Alkenylanilides and 2-Alkenylphenols with Allenyl Acetates



The groups of Gulías and Lam achieved the vinylic C–H activation on 2-alkenylphenols to give oxacyclic or


spirocyclic molecules.<sup>92,104</sup> To build on that discovery, they expanded to carry out C–H bond functionalization of styrene-type alkenes to produce 2-substituted indolines. By means of a modified Rh(III) catalyst featuring an electron-deficient  $\eta^5$ -cyclopentadienyl ligand ( $\text{Cp}^E$ ), the coupling reaction of alkynes with 2-alkenyl anilides bearing an Ns group proceeded smoothly in toluene at 80 °C (Scheme 58).<sup>133</sup> By separating out the intermediates and performing deuterium-labelling experiments, it can be concluded that the possible mechanism involves alkenylation of C–H bond to form an intermediate, followed by a 1,5-H shift simultaneously with migration of Rh to afford an intermediate which finally undergoes reductive elimination.

#### Scheme 58. Synthesis of 2-Substituted Indolines by Rhodium(III)-Catalyzed Annulation of 2-Alkenyl Anilides with Alkynes





In 2019, the same group also discovered another kind of annulation reaction between alkynes and 2-alkenyl anilides bearing a Tf group (Scheme 59).<sup>134</sup> Apart from obtaining the typical naphthylamide product, an unexpected isomer was identified. Mechanistic studies were performed to justify the production of the isomer when the naphthylamide of the alkenyl underwent migration. The results drawn were used in combination with DFT calculations to elucidate the likely mechanism. The C–H bond is activated, then alkyne coordination and insertion occur to produce a seven-membered rhodacycle. The rhodacycle then undergoes reductive elimination to give a Rh-spirocycle that can form two different kinds of cyclopropyl tricyclic intermediates, as supported by computational results. These two different intermediates would then proceed to form their respective products.

#### Scheme 59. Formation of Unexpected Naphthalene Adducts by Annulation of *ortho*-Alkenyl Anilides with Alkynes





However, the oxidative annulation between alkynes and 2-alkenyl anilides by a Pd(OAc)<sub>2</sub> catalyst exceptionally produced a rare class of highly appealing nitrogen-containing cyclopentaquinoline derivatives (Scheme 60).<sup>135</sup> Various 2-alkenylanilines with different substitution patterns have proven to be compatible. Mechanistically, this novel transformation undergoes sequential vinylic C–H alkenylation/amination/pyrindination processes in one pot under the same conditions.

### Scheme 60. Synthesis of Cyclopentaquinolines by 2-Vinylanilines with Alkynes




The ability to assemble functionalized benzo[*b*]azapines from a chemo- and regioselective [5 + 2] cloaddition of 2-alkenylanilines with propargylic esters is a highly sought-after goal. To this end, Zeng and co-workers established a palladium(II)/Lewis acid co-catalyzed annulation protocol to synthesize diverse synthetically valuable benzo[*b*]azapine derivatives by using commercially available 2-alkenylanilines with various propargylic esters (Scheme 61).<sup>136</sup> A broad scope of 35 examples of this transformation was presented with yields of 30–75%. Of note, this [5 + 2] annulation protocol was also applicable to the reaction between alkynes and 2-alkenylanilines.

### Scheme 61. Oxidative Annulation of 2-Alkenylanilines and Propargylic Esters





The readily accessible *gem*-dihaloalkenes are versatile synthons which could be used as alkyne precursors for cross-coupling reactions.<sup>137</sup> Quite recently, Tian and co-workers employed *gem*-dichloroalkenes for the Ni(o)-catalyzed reaction of 2-vinylanilines to construct polysubstituted quinolines (Scheme 62).<sup>138</sup> By the combination of Ni(cod)<sub>2</sub> catalyst, *N*-heterocyclic carbene ligand (IPr·HCl), and NaO<sup>t</sup>Bu base in mesitylene at 80 °C, a broad scope of 30 examples was documented in yield up to 80%. Notably, *gem*-dichloroalkenes bearing both aryl and alkyl substituents were well tolerated in this case. This reaction can be carried out on a preparative scale, and was applicable to late-stage derivatization of natural products.

### Scheme 62. Nickel-Catalyzed Cascade Reaction of 2-Vinylanilines with *gem*-Dichloroalkenes






As detailed in Scheme 63, a highly regioselective palladium(II)-catalyzed sequential vinylic C–H bond arylation/intramolecular amination of Ts-protected 2-vinylanilines with aryl boronic acids under oxidative conditions was reported by Zeng's group in 2018.<sup>139</sup> The scope of this protocol was found to be remarkably broad, delivering an efficient and straightforward platform to access synthetically valuable multifunctionalized indoles.

### Scheme 63. Sequential C–H Arylation/Amination of 2-Vinylanilines with Arylboronic Acids



In 2020, Yi, Zhou, and colleagues reported a unique intramolecular dehydrogenative cross-coupling reaction by taking advantage of a modified  $\text{Cp}^*\text{Rh(III)}$  catalyst, resulting in the direct synthesis of highly valuable eight-membered sultam/lactam derivatives with a wide substrate scope and excellent functional group compatibility (Scheme 64).<sup>140</sup> The presented reaction undergoes a redox-neutral pathway to furnish the intramolecular olefinic C–H arylation products with  $\text{H}_2$  as the major byproduct.

#### Scheme 64. Intramolecular Dehydrogenative Cross-Couplings for Building Eight-Membered Sultam/Lactam Scaffolds




Besides, the intramolecular vinylic C–H amination of 2-vinyl anilines provide a straightforward and atom-economical approach to construct the privileged indole skeletons. In this regard, a series of conceptually different strategies have been successively established over the past decades. Specifically, Stille and co-workers in 1988 reported the Pd(II)-catalyzed intramolecular cyclization


reaction of tosyl-protected 2-alkenylanilines by using *p*-benzoquinone (BQ) as a reoxidant (Scheme 65a).<sup>141</sup> In this case, 10 excess equivalent of lithium chloride was employed to obtain high efficiency. Recently, the Li group also illustrated an analogous transformation with comparable efficiency with unprotected 2-alkenylanilines as the suitable substrates enabled by a  $\text{FeF}_2/\text{K}_2\text{S}_2\text{O}_8$  catalytic system (Scheme 65b).<sup>142</sup>

#### Scheme 65. Synthesis of Indoles via Intramolecular Vinylic C–H Amination of 2-Alkenylanilines

a) Stille et al., 1988





b) Li et al., 2019



In 2017, Cheng et al. accomplished a copper-mediated intramolecular cyclization of various 2-alkenylaniline substrates (Scheme 66).<sup>143</sup> Under the optimal conditions, diverse 2-alkenylanilines bearing a broad range of functionalities could be smoothly converted into the expected products with yields of 33–96%. Remarkably, 2,3-disubstituted indole can be obtained uneventfully, albeit with diminished efficiency. The authors demonstrated the practicality of this protocol by a preparative scale synthesis.

#### Scheme 66. Synthesis of Indoles by Cu-Mediated Intramolecular Cyclization of 2-Alkenylanilines




Although these transition metal-based strategies have shown notable features, innovative strategies for the sustainable indole synthesis from 2-alkenylanilines are highly desirable. To this end, a myriad of efficient metal-

free methods, including dioxygen-promoted,<sup>144</sup> DDQ-mediated,<sup>145</sup> selenium-catalyzed,<sup>146,147</sup> iodine(III) reagent-mediated,<sup>148–150</sup> as well as NIS-mediated intramolecular aminations<sup>151</sup> have been successively established over the past years. These strategies greatly expanded the toolbox for the expedient synthesis of structurally diverse indole derivatives.

With the ever-increasing interests in photoredox catalysis, Zheng et al. in 2012 first explored the visible-light-mediated photocatalytic process for the synthesis of *N*-arylindoles by using [Ru(bpz)<sub>3</sub>](PF<sub>6</sub>)<sub>2</sub> as a competent photocatalyst (Scheme 67).<sup>152</sup> In this case, a variety of substrates with different substitution patterns are compatible with the open-air conditions. Nevertheless, a *p*-alkoxyphenyl group on the nitrogen atom is indispensable for the reaction, and no reaction was observed under the same conditions when it was replaced by a phenyl group. It is noteworthy that a broad array of *gem*-disubstituted styryl anilines could undergo a 1,2-carbon shift process to furnish 2,3-disubstituted *N*-arylindoles. Both aryl and alkyl groups can participate in this migratory transformation, and the former one is more preferentially migrated.

### Scheme 67. Visible-Light-Mediated Synthesis of *N*-Arylindoles




Another alternative sustainable strategy to synthesize indoles from 2-alkenylanilines is to employ the electrochemically driven dehydrogenative cyclization process. In this regard, the group of Zha and Wang recently established an iodine-mediated electrochemical protocol of vinylic C–H amination through an ionic process (Scheme 68a).<sup>153</sup> Using Pt plates as both the cathode and the anode in an easy-to-use undivided cell, diverse Ts-protected 2-alkenylanilines bearing different substituents reacted smoothly to afford the expected indoles in yield up to 92%. However, remarkable electronic and steric effects were observed in this case as 7-brominated substrate was proven to be incompatible with the electrochemical conditions. Later in 2021, Xu et al. disclosed an analogous process carried out in a Schlenk tube equipped with a reticulated vitreous carbon anode (RVC) and a Pt plate cathode in the presence of an organic phenothiazine-based redox catalyst. By virtue of anodic oxidation under metal-free conditions, this protocol produced a variety of 3-substituted and 2,3-disubstituted indoles with H<sub>2</sub> evolution (Scheme 68b).<sup>154</sup> In a related report, Chang and co-workers realized an operationally convenient, Ir(III)-catalyzed dehydrogenative cyclization of unprotected 2-alkenyl anilines under undivided electrolytic conditions. Combined computational studies and electrochemical


investigations revealed that the reaction probably proceeds through an electro-oxidation induced reductive elimination pathway (Scheme 68c).<sup>155</sup>

### Scheme 68. Synthesis of Indoles through an Electrocatalytic Dehydrogenative Cyclization of 2-Vinylanilides


a) Zha, Wang et al., 2020



b) Xu et al., 2021



c) Chang et al., 2021




## 3. ALKENYL C–H BOND FUNCTIONALIZATION OF ALIPHATIC ALKENES CONTAINING A DIRECTING GROUP

After discussing on alkenyl C–H bond functionalization of aryl styrenes, we next turned to focus on the reactions using simple aliphatic terminal alkenes. One of the major challenges in the sp<sup>2</sup> alkenyl C–H bond functionalization of alkenes is the control of the stereochemistries (regio, *E/Z*, etc). Similar to the reactions using aryl alkenes, the *E*-selective vinylic C–H bond can be controlled by steric effect. Similarly, most of the reported *Z*-selective C–H bond functionalizations made use of a directing group. Many different chelating groups forming 5–6 membered metal chelates have been successfully used to give *Z*-selective products. In this part of the review, we will organise the advances according to the types of chelating groups that form 5–6 membered ring chelated intermediate.

Conjugated dienes are found extensively in a variety of pharmaceutically relevant molecules and bioactive natural products, and are also important starting materials in organic synthesis. Alkene C–H alkenylation reactions


catalysed by different transition metal catalyst are an atom-economic way to achieve these conjugated dienes. In 2017, Loh's group reported a stereospecific olefination between two alkenes by using a native hydroxyl directing group (Scheme 69).<sup>156</sup> Good *E/Z* selectivity was obtained in the products. The hydroxyl group was found to serve as a chelating group not only to guide the activation of alkenyl C–H bonds, but to improve on the *Z*-stereoselectivity in this process. By tuning the protecting group in homoallylic alcohols, this protocol can be extended to the stereochemical synthesis of denoates.

### Scheme 69. Oxidative Cross-Coupling Reactions between Homoallylic Alcohols and Alkenes




As an extension of this strategy, Zhong and co-workers later in 2019 also discovered the functionalization of proximal alkenyl C–H bonds of simple aliphatic alkenes (Scheme 70).<sup>157</sup> Monodentate chelation was used for the C–H bond alkenylation, and the reaction was successful with broad substrates containing hydroxyl, carbamate, and amide functional groups. In this work, the proximal C–H bond was activated *via* an *o*-monodentate chelation, which occurs through four- to eight-membered *exo*-palladacycles, which is in contrast to the *N,N*-bidentate chelation that occurs through a six-membered *endo*-palladacycle as generally proposed in Engle's work. Notably, steroids and ricinoleates are also competent substrates to undergo late-stage functionalizations, which remarkably showcased the capability of this approach.

### Scheme 70. Geminal Group-Directed Alkenyl C–H Olefination



Almost simultaneously, Dong and co-workers reported a novel approach for the distal-selective alkenyl C–H arylation enabled by palladium/norbornene (NBE) cooperative catalysis (Scheme 71).<sup>158,159</sup> This reaction occurs by activating a proximal olefinic C–H bond, and can be conducted in air with a large variety of *cis*-alkenes, aryl iodides, MeI, and methyl bromoacetate. This reaction is found to be more probable to undergo the functionalization of alkenyl  $sp^2$  C–H bond rather than the classical Heck cross-coupling pathway as supported by the detailed mechanistic studies. Mechanistically, the proximal C–H bond of the olefin substrate is activated to form an intermediate, and this process is directed by the easily removable oxime ether directing group. Then, the NBE derivative is inserted into the intermediate, and the oxime ether de-coordinates to afford another intermediate which undergoes C–H activation again at the distal position. This results in the formation of a palladacycle, which reacts with ArI to obtain the final product.


### Scheme 71. Distal Olefinic C–H Functionalization via Palladium/Norbornene Cooperative Catalysis



The Dong group extended their distal C–H alkenylation through the palladium/norbornene cooperative catalysis strategy to arylate and alkylate the *cis*-olefins by introducing a selective donating group (Scheme 72).<sup>160</sup> The judicious choice of structurally modified norbornene mediator is critical to realise this transformation. The


authors successfully demonstrated the arylation of allylamines, primary and secondary derived alcohols.

### Scheme 72. Palladium/NBE-Catalyzed Distal Alkenyl C–H Arylation and Alkylation of *cis*-Olefins



In 2020, Miura and co-workers reported a regioselective palladium(II)-catalyzed phenanthroline-directed vinylic C–H alkenylation of allylic alcohols with electron-deficient alkenes (Scheme 73).<sup>161</sup> In this case, the proximal alkenyl C–H bond was activated over allylic C–O bond with the assistance of a bidentate directing group. Under the same Pd(II)/phenanthroline conditions, the authors utilized alkynyl bromides to alkynylate the allylic alcohols in a regioselective manner.


### Scheme 73. Phenanthroline-Directed, Palladium(II)-Catalyzed Regioselective C–H Alkenylation and Alkynylation of Allylic Alcohols



Lu and Zhao et al. reported an oxyacetamide-directed *Z*-type alkenyl C–H functionalization enabled by a Cp\**Rh*(III) catalyst through a rare *exo*-rhodacycle process (Scheme 74).<sup>162</sup> Multisubstituted alkenes and allenes were synthesized in modest to excellent yields. With the judicious choice of solvents and base, both alkenes and propargylic carbonates coupling partners reacted well to furnish the corresponding products with high regio- and


stereoselectivity. Moreover, unnatural  $\beta$ -amino acid can be synthesized using this strategy.

### Scheme 74. Oxyacetamide-Directed *Rh*(III)-Catalyzed Alkenyl C–H Functionalization of Allylic Alcohols



In the same year, Dethe and colleagues extended Loh's strategy,<sup>156</sup> and developed a direct Ru-catalyzed method involving an oxidative coupling of allyl alcohols with activated olefins by C(allyl)–H activation of allyl alcohols, providing an efficient and direct access to synthetically useful class of  $\alpha,\beta$ -unsaturated enones (Scheme 75).<sup>163</sup> The authors demonstrated the utility of this method by applying to the synthesis of bioactive natural products such as Hydroxy- $\beta$ -sanshool, ZP-amide I, Chondrillin, Plakorin, and (+)-*cis*-Solamin A.



### Scheme 75. Ru(II)-Catalyzed Dehydrogenative Cross-Coupling of Allyl Alcohols with Acrylates




Allylamine derivatives are highly valuable synthetic building blocks for the synthesis of various nitrogen-containing molecules. The direct activation of alkenyl C–H bonds of allylamines also attract tremendous interests in recent years. For example, Li's group in 2013 described a rhodium(III)-catalyzed oxidative vinylic C–H olefination of *N*-sulfonyl allylamines with acrylates and acrylonitriles (Scheme 76).<sup>164</sup> More interestingly, the reaction with electron-deficient *N,N*-dimethylacrylamide as the coupling partner exclusively resulted in the formation of 2,3-dihydropyrrole derivatives in modest yields (51–58%). Later on, Babu and co-workers disclosed the bidentate picolinamide-assisted  $\gamma$ -selective C–H arylation of

allylamines in the presence of the Pd(OAc)<sub>2</sub> catalyst and AgOAc additive to afford a series of *Z*-cinnamylamines with appreciable yields and up to 2:98 *E/Z* ratio (Scheme 77).<sup>165</sup>

### Scheme 76. Rhodium(III)-Catalyzed Oxidative C–H Olefination of *N*-Allyl Sulfonamides





### Scheme 77. Pd(II)-Catalyzed Picolinamide-Assisted Construction of *Z*-Cinnamylamines





Additionally, Loh's group also disclosed a general approach for the alkenyl C–H alkylation of allylamines by means of a bidentate isoquinoline-1-carboxamide (*iQA*)-assisted strategy (Scheme 78),<sup>166</sup> providing a general method to selectively produce (*Z*)-tri- and tetra-substituted olefins. Notably, both primary and secondary alkyl iodides reacted well under these conditions.

### Scheme 78. Stereospecific Vinylic C–H Alkylation of Allylamines with Alkyl Iodides



Undoubtedly, *C*-vinyl glycosides are a prominent class of carbohydrates widely encountered in many natural products and bioactive molecules.<sup>167–169</sup> Their efficient synthesis is of great importance, and a tremendous synthetic challenge as well.<sup>170–173</sup> More recently, the straightforward and stereoselective synthesis of *C*-vinyl glycosides was disclosed by the He and Chen group through a palladium(II)-catalyzed chelation-assisted C–H glycosylation of unbiased aliphatic alkenes bearing an easily removable isoquinoline-1-carboxamide (*iQA*) auxiliary<sup>166</sup> with glycosyl chlorides (Scheme 79).<sup>174</sup> This strategy can glycosylate both  $\gamma$  and  $\delta$  C–H bonds of various readily available alkene substrates in moderate to high yields with excellent regio- and stereoselectivity.


### Scheme 79. Stereoselective Synthesis of *C*-Vinyl Glycosides via C–H Glycosylation of Aliphatic Alkenes



Moreover, Young's group demonstrated an efficient, simple, and highly selective  $\gamma$ -arylation of unprotected allylamines through a stereo- and regiospecific olefin insertion mechanism (Scheme 80).<sup>175</sup> The authors employed both primary and secondary amines in this


transformation. Different functional groups such as ketones, esters and heterocyclic motifs are compatible with this method, and an extraordinarily broad range of 3,3'-diarylated allylamines were obtained in decent yields.

### Scheme 80. Regioselective $\gamma$ -Arylation of Unprotected Allylamines



Following this, the same group extended to report the efficient synthesis of *Z*-selective alkenyl amines by the  $\gamma,\gamma'$ -diarylation reaction of unprotected  $\beta$ -alkyl allylamines through an interrupted chain walking process (Scheme 81).<sup>176</sup> In this case, they demonstrated the choice of silver salt plays a significant role in arylating the less bulkier secondary and tertiary amines. Even the unprotected allylamines could react efficiently in this protocol.


### Scheme 81. Pd(II)-Catalyzed $\gamma,\gamma'$ -Diarylation of Free Alkenyl Amines



Moreover, Gulias and colleagues disclosed a palladium-catalyzed enantioselective annulation of allyltriflamides with allenes which relies on an asymmetric alkenyl C–H functionalization step. This protocol allowed an easy and straightforward access to enantioenriched allylamines and


piperidines with decent stereocontrol in a step- and atom-economical manner (Scheme 82).<sup>177</sup>

### Scheme 82. Kinetic Resolution of Allyltriflamides through a Palladium(II)-Catalyzed Vinylic C–H Bond Functionalization with Allenes



In addition to the example of  $\gamma$ -carbonylation<sup>178</sup> and silylation<sup>179</sup> of aliphatic amines (Scheme 83a), Zhao and co-workers also disclosed an example of olefinic  $\delta$ -C(sp<sup>2</sup>)-H alkylation via six-membered palladacycles using a readily available oxalyl amide directing group (Scheme 83b).<sup>180</sup> Another notable example of alkenyl  $\delta$ -C–H bond alkylation of unbiased alkenes with bromoalkyne was reported by Sarpong and co-workers using a  $\beta$ -carboline amide as inherent directing group (Scheme 83c).<sup>181</sup>


### Scheme 83. Various Olefinic C–H Functionalization Reactions of Unbiased Alkenes



Meanwhile, He and colleagues illustrated an efficient site-selective palladium(II)-catalyzed olefinic  $\delta$ -C(sp<sup>2</sup>)-H acetoxylation,<sup>182</sup> and later alkenylation<sup>183</sup> of unactivated


cyclic alkenes by using picolinamide as a remote bidentate directing group via a six-membered palladacycle intermediate (Scheme 84).

#### Scheme 84. Palladium-Catalyzed Olefinic $\delta$ -C(sp<sup>2</sup>)-H Acetoxylation and Alkenylation of Unactivated Cycloalkenes



Apart from the above intermolecular oxidative coupling reactions, Loh's research group in 2017 established an intramolecular approach of vinylic C–H amination for the synthesis of multisubstituted dihydropyrroles (Scheme 85).<sup>184</sup> In this report, the judicious use of chloranil instead of benzoquinone (BQ) greatly facilitated this intramolecular 5-*endo* cyclization to result in the formation of dihydropyrroles. Overall, a broad range of  $\gamma,\delta$ -unsaturated tosyl-protected amines delivered the expected products in excellent yields. However, diminished efficiency was observed for electron-rich substituents on the alkene moiety, and long-chain alkyl group completely inhibited this transformation.



#### Scheme 85. Synthesis of Dihydropyrroles via Pd(II)-Catalyzed Intramolecular Alkenyl C–H Amination



With the assistance of a removable bidentate 8-aminoquinoline auxiliary, Pan and co-workers were able to



regioselectively activate the  $\beta$ -C–H bonds of diverse  $\beta,\gamma$ -unactivated amides, enabling an expeditious route to a variety of conjugated 1,3-enynes (Scheme 86).<sup>185</sup> Notably, a wide scope with respect to both terminal and internal unactivated aliphatic alkenes were smoothly coupled with bromoalkynes to generate the alkylation products with overall high yields (58–79%). Bromoalkynes bearing TIPS, TBS and TES groups were appropriate coupling partners, while other substituents such as phenylacetylene and methyl propionate were proven to be incompatible with the conditions. On the basis of mechanistic investigations, a plausible catalytic cycle involving Pd(II)/Pd(IV) redox manifold was proposed to elucidate the mechanism.

#### Scheme 86. Synthesis of 1,3-Enynes via Palladium-Catalyzed Vinylic C–H Alkylation of Alkenes




It is commonly reported that neighbouring directing groups of alkenes always resulted in the activation of distal C–H bonds, and this is notably distinct from C–H activation in aryls.<sup>186</sup> However, the activation of proximal C–H bonds in olefin substrates has been not as well explored. In 2018, Engle and co-workers elegantly disclosed a landmark work on obtaining various highly substituted 1,3-dienes by using Pd(OAc)<sub>2</sub> to catalyse the alkenylation of proximal olefinic C–H bonds with a stoichiometric amount of MnO<sub>2</sub> or O<sub>2</sub> and catalytic Co(OAc)<sub>2</sub> (Scheme 87).<sup>187</sup> Internal non-conjugated alkenes reacted well under the typical reaction conditions. Combined experimental and computational investigations confirmed that proximal C–H alkenyl bond activation occurred in this protocol. The structurally well-defined alkenylpalladium(II) dimer was readily isolated and well-characterized, which can be also used as the catalyst.

#### Scheme 87. Synthesis of 1,3-Diene via Six-Membered Palladacycles



Subsequently, Zhong and colleagues disclosed an efficient approach to the synthesis of branched 1,4-dienes by the distal alkenyl C-H allylation of non-conjugated alkenyl amides with allyl carbonates using 8-aminoquinoline (AQ) as the bidentate chelating group (Scheme 88).<sup>188</sup> This reaction builds on their previous work in proximal alkenyl C-H functionalization reactions through an *exo*-palladacycle intermediate.<sup>157</sup> Their new protocol employs unactivated Z-olefins to couple with allyl carbonates under typically mild and oxidant-free conditions.


### Scheme 88. Synthesis of Branched 1,4-Dienes via Bidentate Auxiliary-Directed Alkenyl C-H Allylation



Alkenyl iodides represent a versatile class of building blocks frequently used in organic synthesis,<sup>189–192</sup> which significantly highlights the increasing need for efficient synthesis methods. A straightforward approach to prepare alkenyl iodides would be the direct olefinic C-H iodination of their alkene precursors. In 2019, Carreira and co-workers disclosed an elegant protocol of the regio- and stereoselective Pd(II)-catalyzed C-H iodination of electronically unbiased alkenes bearing the bidentate


picolinamide directing group, furnishing the iodination products in yields ranging from 41% to 85% with excellent functional group compatibility (Scheme 89).<sup>193</sup>

### Scheme 89. Synthesis of Alkenyl Iodides through the Regioselective Vinylic C-H Iodination of Unactivated Alkenes



As an extension of this approach, the same group later expanded to disclose a palladium-catalyzed remote C-H alkynylation of unactivated alkenes by using bromoalkynes as the alkynylating source (Scheme 90).<sup>194</sup> In this case, readily available picolinamide auxiliary enables the formation of putative 5- and 6-*exo*-metallacycles as key intermediates to produce a series of multisubstituted 1,3-enynes in up to 91% yield with excellent regio- and diastereoselectivity. As a particular highlight, the utility of this strategy has been showcased by late-stage modification of biologically important molecules.



### Scheme 90. Palladium-Catalyzed C-H Alkynylation of Electronically Unbiased Alkenes



Successfully, Fu, Bi, and co-workers in 2018 achieved a distinctive Cu-catalyzed Heck-type C-H alkylation of unbiased aliphatic alkenes with various alkyl halides by the assistance of a bidentate 8-aminoquinoline (AQ) auxiliary strategy (Scheme 91).<sup>195</sup> Alkene substrates bearing other common mono- or bidentate auxiliaries thoroughly failed



to furnish the desired products in this transformation. Notably, both 1°, 2°, 3° alkyl bromides and sterically diverse nonactivated alkenes were tolerated under the conditions to afford the expected products with excellent regio- and stereoselectivity. A wide scope of 41 examples of this transformation was presented with yields of 46–92%. The authors conducted preliminary mechanistic investigations and DFT calculations that clearly indicated a radical pathway involving a concerted H–Br elimination step of a putative conformationally strained Cu(III) intermediate, which is greatly assisted by dimethyl sulfoxide.

### Scheme 91. Intermolecular Heck-Type C–H Alkylation of Unactivated Olefins



Macrocyclic peptides are prevalent in nature and increasingly emerged as fascinating molecular scaffolds to explore the chemical space between small organic molecules and large biologicals for the development of therapeutics and chemical probes to interrogate many complicated biological systems.<sup>196</sup> In 2020, Chen and co-workers disclosed an intramolecular palladium-catalyzed picolinamide-directed alkenyl C–H arylation at the  $\gamma$  and  $\delta$  positions of *N*-terminal amino acids to the construction of a handful of aryl-alkene-braced macrocyclic peptides from readily accessible precursors, providing the arylated peptide macrocycles in decent yields (Scheme 92).<sup>197</sup>


### Scheme 92. Synthesis of Peptide Macrocycles by Intramolecular Alkenyl C–H Arylation



## 4. ALKENYL C–H BOND FUNCTIONALIZATION OF ALKENES CONTAINING AN ELECTRON-WITHDRAWING DIRECTING GROUP

In this section, we will focus on another class of alkene derivatives where the alkenes are attached to an electron-withdrawing groups such as acrylic acids, acrylic esters, acrylic amides,  $\alpha,\beta$ -unsaturated ketones,  $\alpha,\beta$ -unsaturated imines,  $\alpha,\beta$ -unsaturated oximes and their derivatives, as well as 2-vinylpyridines (Scheme 93). These electron-withdrawing groups also serve as directing group to control the stereochemistry of the coupling reactions. In addition, the coupling products contain these versatile functional groups also make them more useful building blocks in organic synthesis.

### Scheme 93. General Scheme of Alkenyl C–H Bond Functionalization of Alkenes Containing an Electron-Withdrawing Directing Group




#### 4.1. Acrylic Acids

Acrylic acids and their derivatives are ubiquitous building blocks in chemical synthesis. Transition metal-catalyzed alkenyl C–H activation of acrylic acids and their derivatives, which allows to selectively functionalize the molecule via employing the directing and coordinating groups, has drawn more attention in the last few decades. In C–H activation of acrylic derivatives, a plethora of reports using rhodium and palladium catalysts have dominated. Nonetheless, this section aims at discussing the alkenyl C(sp<sup>2</sup>)–H activation of acrylic derivatives using different transition-metal catalysts.


In 2008, Yu's group reported a direct palladium(II)-catalyzed C–H carboxylation by using C–H activation/CO insertion sequences to prepare dicarboxylic acids in modest to excellent yields. Gratifyingly, the alkenyl β-C–H bond of α,β-unsaturated 2-phenylacrylic acid could be carboxylated selectively to afford a *cis*-1,2-dicarboxylic acid in 68% yield (Scheme 94).<sup>198</sup>

#### Scheme 94. Pd(II)-Catalyzed Vinyl C–H Carboxylation of 2-Phenylacrylic Acid




Pioneering work of Miura and co-workers elegantly elaborated the oxidative vinylic C–H annulation reaction of acrylic acids with diverse alkynes and alkenes by using Cp<sup>E</sup>Rh(III) catalysts to synthesize various α-pyrone and butenolides (Scheme 95).<sup>199</sup>

#### Scheme 95. Oxidative Coupling of Acrylic Acids with Alkynes and Alkenes



Later in 2013, the same group expanded to report the efficient rhodium(III)-catalyzed decarboxylative and dehydrogenative coupling reaction of commercially available maleic acids with both alkynes and 1,3-dienes to afford a series of multisubstituted α-pyrone derivatives in decent yields (Scheme 96).<sup>200</sup> Moreover, the rhodium catalyst system is also applicable to the reaction with alkenes to exclusively produce dienoic acid derivatives.


#### Scheme 96. Decarboxylative and Dehydrogenative Coupling of Maleic Acids with Alkynes and Alkenes




Jiang and colleagues reported a practical palladium(II)-catalyzed oxidative [4 + 2] annulation of unsubstituted acrylic acids with diverse internal alkynes by using of O<sub>2</sub> (1 atm) as the oxidant (Scheme 97a).<sup>201</sup> Later in 2016, Tanaka's group also disclosed an analogous oxidative annulation process of acrylic acids with alkynes enabled by an electron-deficient Cp<sup>E</sup> rhodium(III) catalyst under mild conditions (Scheme 97b).<sup>202</sup> Following this, Zhao and co-workers also reported the synthesis of α-pyrone and co-workers also reported the synthesis of α-pyrone via oxidative annulation reaction catalyzed by Rh(III) and silver co-catalyst between various substituted acrylic acids and alkynes (Scheme 97c).<sup>203</sup> The protocol accommodated a diverse range of acrylic acids and alkynes, delivering the corresponding α-pyrone in good to excellent yields. Impressively, the sorbic acid was also competent substrate for this transformation.

#### Scheme 97. Oxidative [4 + 2] Annulation Reaction of Acrylic Acids with Alkynes


a) Jiang et al., 2014



b) Tanaka et al., 2016




c) Zhao et al., 2018



In 2015, Gogoi and co-workers established a general method for the preparation of diversely decorated α-pyrone derivatives through a Ru(II)-catalyzed C–H activation/annulation of cinnamic acids with various



disubstituted alkynes under cost-effective ruthenium catalysis (Scheme 98).<sup>204</sup> The oxidative annulation reaction features a wide substrate scope and high regioselectivity, affording the expected products in up to 92% yield with low catalyst loading.

### Scheme 98. Synthesis of $\alpha$ -Pyrone via Ru(II)-Catalyzed Alkene C–H Bond Functionalization




Mei and co-workers combined electrochemistry and Ir catalysis to perform alkenyl C–H activation/annulation under oxidative conditions (Scheme 99).<sup>205</sup> This reaction is tolerant towards a broad variety of substrates including complex molecules, giving moderate to excellent yields. The reaction mechanism was investigated in detail. By using stoichiometric reactions, an Ir(I) diene complex was isolated which was hypothesized to be one of the intermediates in the possible catalytic reaction. Further mechanistic experiments were performed to prove that common oxidants such as Ag(I) and Cu(II) were not as important as anodic oxidation.

### Scheme 99. Cp\*Ir(III)-Catalyzed Electrochemical C–H Annulation of Acrylic Acids with Alkynes



Su and co-workers achieved a novel Rh-catalyzed C–H alkylation of alkenyl carboxylic acids with diverse ketones (Scheme 100).<sup>206</sup> This protocol proceeded via the merging of Cu-catalyzed ketone dehydrogenative desaturation process and Rh-catalyzed carboxyl-directed vinyl *ortho*-C–H alkylation. In this case, TEMPO was proved to be essential for the dehydrogenation of ketones as well as the generation of the catalytically active Rh(III)(TEMPO)<sub>2</sub>L<sub>n</sub> catalyst for C–H activation. Besides alkenyl carboxylic acids, the aryl counterparts were also found to be suitable C–H sources for this transformation.


### Scheme 100. Carboxyl-Directed C–H Alkylation of Alkenyl Carboxylic Acids




Alkenyl C–H activation and intramolecular cyclization was successfully conducted to generate *N*-arylmaleimide using Rh catalyst by the Li group in 2016 (Scheme 101a),<sup>207</sup> with the intermediate being an amide. In the following year, Zhong and co-workers conducted the annulation reaction of substituted acrylic acids with electron-deficient alkenes to produce a broad range of synthetically valuable  $\gamma$ -alkylidenebutenolides (Scheme 101b).<sup>208</sup> Successfully, the reaction could be scaled up to gram-scale with only 1 mol % catalyst loading.

### Scheme 101. Rhodium(III)-Catalyzed C–H Activation of Acrylic Acids with Isocyanates and Alkenes

a) Li et al., 2016




b) Zhong et al., 2017




In 2017, Zhu's group reported the synthesis of furan-2(5*H*)-ones through the alkenyl C–H activation of cyclo-alkenecarboxylic acids with acrylates in the presence of a Cp<sup>\*</sup>Rh(III) catalyst (Scheme 102).<sup>209</sup> The mechanism involved the formation of 7-membered rhodacycle, followed by hydride elimination, and intramolecular cyclization through Michael addition to furnish the desired furan derivatives.

### Scheme 102. Synthesis of Furan-2(5*H*)-ones from cyclo-Alkenecarboxylic Acids and Acrylates





Ackermann's group demonstrated two examples of Rh(III)-catalyzed cross-dehydrogenative alkenylation of acrylic acids bearing various substitution patterns with electron-deficient alkenes via electrooxidation (Scheme 103).<sup>210</sup> The expected coupling products were synthesized with chemo- and stereoselectivity via environmentally benign oxidative two-fold C–H activation with H<sub>2</sub> as the sole byproduct.

### Scheme 103. Rhodium-Catalyzed Electrochemical C–H Alkenylation of Acrylic Acids





The formal [4 + 2] cycloaddition protocol of acrylic acids with various 1,3-dienes was achieved by Zhang and co-workers in 2018 (Scheme 104).<sup>211</sup> The reaction worked uneventfully with 10 mol% PdSO<sub>4</sub> as the catalyst in conjunction with stoichiometric Cu(OAc)<sub>2</sub> (2.0 equiv) as the oxidant. As a particular highlight, this approach was applicable to the short synthesis of natural product Clausamine B.

### Scheme 104. Palladium-Catalyzed Formal [4 + 2] Cycloaddition of Acrylic Acids with 1,3-Dienes




More recently, Jeganmohan and co-workers utilized alkylidenecyclopropanes in the vinylic C–H annulation reactions of acrylic acids under Cp<sup>\*</sup>Rh(III) catalysis (Scheme 105).<sup>212</sup> In this case, acrylic acids with α-substituted reacted uneventfully to produce α-pyrone in moderate yields (31–54%), while β-substituted acrylic acids such as cinnamic acid and crotonic acid were found to be incompatible with this protocol.

### Scheme 105. Rhodium(III)-Catalyzed Annulation of Acrylic Acids with Alkylidenecyclopropanes




Despite the remarkable advances in alkenyl C–H olefinations over the past decades, these strategies have been greatly restricted to electronically unbiased alkenes such as acrylates and styrenes. Establishing a general and practical procedure for the vinylic C–H olefination with unbiased alkenes is a long-standing challenge. This can be attributed to the intrinsic poor reactivity as well as the unbiased nature of aliphatic alkenes.<sup>213</sup> In this regard, Jeganmohan et al. in 2020 illustrated the Rh<sup>III</sup>-catalyzed COOH-assisted C–H olefination of diverse α-substituted acrylic acids with unbiased olefins at room temperature (Scheme 106),<sup>214</sup> generating *ortho*-alkenylated vinylic acids in modest yields (22–69%). Unfortunately, the reaction conditions were not compatible with simple acrylic acid and β-substituted acrylic acids.

### Scheme 106. Rhodium(III)-Catalyzed C–H Olefination of Vinyl Acids with Unactivated Olefins



The development of efficient chalcogenylation reactions for the assembly of unsymmetrical diaryl sulfides or diaryl selenides has attracted remarkable attention because of their prevalence in a diverse variety of potent drug candidates, biologically active natural products and advanced functional materials.<sup>215</sup> In this context, the group of Wang and Ji reported the first example of vinyl C–H thiolation of acrylic acids by using readily available inorganic rhodium salt  $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$  as a catalyst in 2019 (Scheme 107).<sup>216</sup> The efficiency of  $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$  as catalyst was better than that of  $[\text{RhCp}^*\text{Cl}_2]_2$  in this strategy, which produced a series of (*Z*)-alkenyl sulfides exclusively. Of note, the thiolation products could be easily converted into biologically and pharmacologically useful thioflavones.


### Scheme 107. Rhodium(III)-Catalyzed C–H Thiolation of Acrylic Acids



In 2018, Chou et al. established a straightforward method for the rapid synthesis of various *ortho*-alkylated vinylarenes from readily available benzoic acids employing Birch reductive alkylation followed by a tandem decarboxylative C–H olefination/rearomatization sequence (Scheme 108a).<sup>217</sup> Subsequently, the same group further disclosed that proaromatic 1,3-dienes can be synthesized through C–H activation of 1,4-cyclohexadiene by using free carboxylic acid as the directing group (Scheme 108b).<sup>218</sup> Based on the judicious choice of silver salt, direct and sequential bisolefinations of proaromatic 1,3-dienes were also achieved. The authors carried out


kinetic resolution studies to understand the mechanism of this process.

### Scheme 108. Palladium-Catalyzed Alkenyl C–H Bond Olefination of Proaromatic Acids




Quite recently, Gooßen et al. disclosed the highly *Z*-selective coupling reaction of 2-arylacrylic acids with aryl bromides by the combination of  $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$  catalyst,  $\text{PEt}_3 \cdot \text{HBF}_4$  ligand and  $\text{Li}_2\text{CO}_3$  base in NMP at 100 °C. A broad scope of 28 examples was documented in reasonable yields of up to 94% and (*Z/E*)-ratios of up to 99:1. Mechanistic investigations indicated that this carboxylate-directed vinylic C–H arylations probably proceeds via a base-assisted cyclometalation process rather than via a Heck-type mechanism (Scheme 109).<sup>219</sup>

### Scheme 109. Ru(II)-Catalyzed Vinylic C–H Arylation of Acrylic Acids with Aryl Bromides




Gao, Hu, and co-workers established a protocol for the amination/annulation of readily available acrylic acids with anthranils in the presence of a  $\text{Cp}^*\text{Rh(III)}$  catalyst (Scheme 110).<sup>220</sup> The carboxylic acid functional group serves as a traceless directing group in this protocol, producing diverse polysubstituted quinolones in modest to excellent yields with  $\text{H}_2\text{O}$  and  $\text{CO}_2$  as byproducts.

### Scheme 110. Rh-Catalyzed C–H Amination/Annulation of Acrylic Acids and Anthranils




The synthesis of conjugated dienes is mostly achieved by the oxidative cross-coupling reaction between two different olefins. In early 2019, Liu's group disclosed a novel redox-neutral method for the synthesis of dienes from acrylic acids and yndienones in the presence of a Rh(III) catalyst (Scheme 111).<sup>221</sup> This reaction is diastereospecific, and happens through activation of olefinic C–H bond, alkyne insertion, and Michael addition sequence.

### Scheme 111. Synthesis of *cis*-Hydrobenzofuranone via Cp\*Rh(III)-Catalyzed Coupling of Acrylic Acids and Ynenones




Moreover, Zhang and colleagues elegantly developed an efficient protocol to synthesize cyclopentenones by coupling acrylic acids with commercially available malonates and formaldehyde (Scheme 112).<sup>222</sup> The authors showcased the application of this method by the synthesis of various 5-alkylated cyclopentenones. Using excess malonate and formaldehyde resulted in the formation of multisubstituted cyclopentenones. In continuation of their work on vinylic C–H activation of acrylic acids with formaldehyde, the same group later extended to disclose a Co(III)-catalyzed carboxyl-assisted C–H functionalization of acrylic acids with formaldehyde to produce butenolides in up to 95% yield (Scheme 113).<sup>223</sup> Interestingly, by addition of Na<sub>2</sub>CO<sub>3</sub> to the catalytic system, a series of  $\gamma$ -hydroxymethylated butenolides could be obtained in up to 72% yield.

### Scheme 112. Synthesis of Cyclopentenones through Rhodium-Catalyzed C–H Annulation of Acrylic Acids with Formaldehyde and Malonates







### Scheme 113. Synthesis of Butenolides via Cobalt-Catalyzed Vinyl C–H Addition to Formaldehyde





Quite recently, Zhang and co-workers further employed  $\alpha$ -diazocarbonyls to couple with acrylic acids under Cp\*Rh(III) catalysis to synthesize  $\alpha$ -pyrones (Scheme 114).<sup>224</sup> Detailed optimization studies revealed that the addition of Zn(OAc)<sub>2</sub> was essential for this transformation, which remarkably accelerates this carboxyl-directed vinylic C–H annulation process. Electronically diverse acrylic acids reacted smoothly with a variety of  $\alpha$ -diazocarbonyl compounds. By slightly elevating the reaction temperature to 90 °C, this protocol also accommodates a series of benzoic acids to afford isocoumarins in modest yields (31–71%). Subsequently, the same group made further efforts to fabricate pyranone derivatives through a Ru(II)-catalyzed three-component cascade reaction of acrylic acids, ethyl glyoxylate, and *p*-toluenesulfonamide (Scheme 115).<sup>225</sup> Notably, an array of butenolides could be readily prepared in high yields under similar catalytic system.

### Scheme 114. Synthesis of $\alpha$ -Pyrones by Rh-Catalyzed Coupling of Acrylic Acids with $\alpha$ -Diazocarbonyls



### Scheme 115. Ru(II)-Catalyzed Cascade Reaction via Vinylic C–H Addition to Glyoxylate





A recent report by Li and co-workers documented a straightforward assembly of various cyclic skeletons enabled by rhodium(III) catalysis under redox-neutral conditions (Scheme 116).<sup>226</sup> In this work, readily available hypervalent iodonium ylides were identified as efficient carbene precursors. It should be emphasized that the catalyst loading could be sharply decreased to 0.5 mol% for the preparative-scale synthesis.

### Scheme 116. Iodonium Ylides as Carbene Precursors in Rhodium(III)-Catalyzed Vinylic C–H Activation



Beyond the intermolecular cross-coupling reaction, the Gulías group in 2019 disclosed a general approach to synthesise attractive bicyclic or tricyclic macrocycles containing pyran-2-one or isocoumarin from simple precursors via an intramolecular oxidative [4 + 2] annulation of acrylic acids with alkynes in the presence of a  $\text{Cp}^*\text{Rh}(\text{III})$  catalyst (Scheme 117).<sup>227</sup>

### Scheme 117. Rhodium(III)-Catalyzed Intramolecular Annulations of Acrylic Acids to Alkynes




The  $\text{Cp}^*\text{Rh}(\text{III})$ -catalyzed formal oxidative [4 + 1] annulation of vinylic carboxylic acids with allenes was reported by Cheng and co-workers in 2015 (Scheme 118).<sup>228</sup> Both symmetrical and unsymmetrical internal allenes were compatible with the conditions, leading to the synthesis of a variety of bioactive 5-vinyl-substituted 2-furanones in 47–93% yields.

### Scheme 118. Rh(III)-Catalyzed [4 + 1] Annulation of Vinylic Carboxylic Acids and Allenes



An enantioselective version of this annulation process was later established by Cramer and co-workers in 2020. They were able to enantioselectively activate the alkenyl C–H bonds in acrylic acids with allenes by using a modified Rh(III) catalyst bearing a novel rigid axial chiral cyclopentadienyl (Cp<sup>x</sup>) ligand featuring a semi-saturated H8-binaphthyl backbone.<sup>229</sup> This enantioselective [4 + 1] annulation process allows the construction of synthetically appealing enantioenriched  $\alpha,\beta$ -unsaturated- $\gamma$ -lactones in decent yields with enantioselectivity of up to 99% enantiomeric ratio (Scheme 119).<sup>230</sup>



#### Scheme 119. Synthesis of Chiral $\gamma$ -Lactones via Cp<sup>x</sup>Rh(III)-Catalyzed C–H Activation of Acrylic Acids with Allenes



#### 4.2. Acrylic Esters


Early in 1995, Trost's group first disclosed a Ru(o)-catalyzed C–H alkylation of diverse  $\alpha,\beta$ -unsaturated esters with vinylsilanes (Scheme 120).<sup>231</sup> As expected, a wide range of functional groups such as epoxide, alkyl bromide, ketals and thioketals were tolerated to deliver the alkylation products in decent yields. Ketone and amide moieties are suitable directing groups in this strategy. Notably, this protocol can be applied to functionalize a sugar derivative.

#### Scheme 120. Ruthenium-Catalyzed Vinylic C–H Alkylation of $\alpha,\beta$ -Unsaturated Esters





Later, Glorius and colleagues revealed a method of vinylic C–H bond cross-coupling with various directing or activating groups under Cp<sup>x</sup>Rh(III) catalysis. A variety of 1,1-, 1,2- and 1,1,2-substituted alkenes reacted uneventfully to give linear 1,3-butadiene products. However, modest yields and poor stereoselectivity were observed with esters as the chelating group (Scheme 121).<sup>232</sup>

#### Scheme 121. Cp<sup>x</sup>Rh(III)-Catalyzed Oxidative Vinylic C–H Olefination of Acrylates





Subsequently, Ishiyama and Ito reported an efficient vinylic C–H borylation of 1-cycloalkenecarboxylates with bis(pinacolato)diboron catalyzed by an in situ-generated Ir complex consisting of [Ir(OMe)(cod)]<sub>2</sub> and AsPh<sub>3</sub>, affording a range of synthetically useful alkenylboronates. A scope of 15 examples of this transformation was documented with yields of 20–96% (Scheme 122).<sup>233</sup>

#### Scheme 122. Ir(I)-catalyzed Vinylic C–H Borylation of 1-Cycloalkenecarboxylates





There have been some notable contributions on alkene C–H alkenylation reactions to obtain various dienes in the presence of a Ru(II) catalyst. Loh and co-workers were able to stereo- and chemo-selectively carry out cross-coupling between two electron-deficient acrylates (Scheme 123).<sup>234</sup> This procedure allows a direct and atom-economical method to obtain functionalized (*Z,E*)-muconates, and is tolerant towards a large variety of substrates. A diverse array of aliphatic and aryl substituted acrylates can be reacted smoothly to provide moderate to good yields with excellent stereoselectivity. The subtle difference in steric and electronic effect enables the preferential C–H functionalization of  $\alpha$ -alkylated acrylates over simple acrylates, and also allows the cross-coupling to proceed.

### Scheme 123. Ru-Catalyzed Cross-Coupling between Two Electron-Deficient Acrylates





Similarly, Zhang's group also reported an analogous regioselective cross-coupling reaction of acrylates under Cp<sup>\*</sup>Rh(III) catalysis, producing substituted 1,3-butadienes in satisfactory yields with high stereoselectivity (Scheme 124).<sup>235</sup> Notably, a diverse array of styrenes were also found to be competent substrates for this transformation.

### Scheme 124. Rhodium(III)-Catalyzed Ester-Directed Olefination of Acrylates



In 2018, Echavarren et al. described the vinylic C(sp<sup>2</sup>)–H alkenylation of  $\alpha,\beta$ -unsaturated esters with bromoalkynes (inverse-Sonogashira reaction) under rhodium(III) catalysis (Scheme 125).<sup>236</sup> The reaction proceeded smoothly using synthetically useful ether as the weakly directing group. Of note, the aryl counterparts were also found to be suitable C–H sources for this alkenylation process.



### Scheme 125. Rhodium(III)-Catalyzed Ester-Directed Alkenyl C–H Alkynylation



## 4.3. Acrylic Amides




**4.3.1. Arylation.** With the success of the alkenyl sp<sup>2</sup> C–H bond functionalization of acrylates, researcher started to turn their attention to acrylic amides.<sup>237</sup> In 2012, Glorius and co-workers reported a novel Cp<sup>\*</sup>Rh(III)-catalyzed dehydrogenative cross-coupling reaction of *N,N*-diisopropylmethacrylamides with bromoarenes with a stoichiometric amount of pivalic acid and a catalytic amount of cesium pivalate as additives, leading to the formation of diverse tri- and tetrasubstituted alkenes in moderate yields (27–65%). The reaction proceeded without any chelate assisting-directing group in arene substrates, thus providing a novel protocol for the highly *Z*-selective synthesis of arene-substituted olefins (Scheme 126).<sup>238</sup>

### Scheme 126. Cp<sup>\*</sup>Rh(III)-Catalyzed Dehydrogenative Cross-Coupling of Vinylic Amides with Bromoarenes





The Babu group in 2015 described the synthesis of  $\beta$ -arylated acrylamides and *Z*-cinnamamides by using 8-aminoquinoline as the bidentate directing group in presence of Pd(OAc)<sub>2</sub> catalyst (Scheme 127a).<sup>239</sup> The reaction was greatly promoted by AgOAc, through the directed *Z*-selective C–H activation followed by the  $\beta$ -arylation of the vinylic C(sp<sup>2</sup>)–H bond of *N*-(quinolin-8-yl)acrylamide systems. A broad array of aryl- and heteroaryl iodides bearing different substitution patterns were used as arylating source. The *Z*-selective  $\beta$ -arylation of *N*-(quinolin-8-yl)acrylamide systems was elucidated by proposing a plausible mechanism involving the bidentate ligand-assisted, chelation-based C–H functionalization. Subsequently, Jiang et al. also achieved a similar 8-aminoquinoline-directed vinylic C–H arylation of acrylamides with 0.5 equivalent of (BnO)<sub>2</sub>PO<sub>2</sub>H as the additive in conjunction with 1.0 equivalent of oxone as the oxidant (Scheme 127b).<sup>240</sup>

### Scheme 127. Palladium-Catalyzed 8-Aminoquinoline-Assisted Vinylic C–H Arylation of Acrylamides





Successfully, Ferry and co-workers demonstrated that the installation of a bidentate 8-aminoquinoline auxiliary at the C2 position of glycols is effective in a Pd(II)-catalyzed alkenyl C(sp<sup>2</sup>)–H arylation of the anomeric position to access a broad range of unsaturated C-aryl glucosides (Scheme 128).<sup>241</sup> Of note, the application of this novel methodology was showcased by the efficient synthesis of a dapagliflozin analogue.

### Scheme 128. Palladium(II)-Catalyzed Vinylic C–H Arylation of the Anomeric Position in Glycosides



In 2016, a novel cobalt-promoted regioselective arylation of alkenyl C(sp<sup>2</sup>)–H bond of acrylamides with various arylboronic acids has been achieved by using bidentate 8-aminoquinoline as the auxiliary by Zhu and co-workers (Scheme 129),<sup>242</sup> leading to the preparation of arylation products in decent yields ranging from 61 to 81%.


### Scheme 129. Cobalt-Promoted C(sp<sup>2</sup>)–H Arylation of Acrylamides with Arylboronic Acids



Ackermann and co-workers developed a novel protocol for the sp<sup>2</sup> C–H functionalization by using 1,2,3-triazole as the powerful auxiliary. The authors described the iron-catalyzed C–H arylations of acrylamides in chemo-, site-, and diastereoselective fashions with a broad substrate scope (Scheme 130).<sup>243</sup> This triazole-assisted C–H activation strategy occurred under remarkably mild reaction conditions, and the triazole auxiliary could be


easily removed in a traceless fashion. Notably, this versatile iron-catalyzed system also proved to be applicable for the challenging C(sp<sup>3</sup>)-H functionalizations, and proceeds by an organometallic mode of action.

### Scheme 130. Iron-Catalyzed Alkenyl C-H Arylation by Triazole Assistance





More Recently, Lu and co-workers accomplished the cross-coupling of acrylamides with organoboron reactants by employing [Cp\*Rh(MeCN)<sub>3</sub>](SbF<sub>6</sub>)<sub>2</sub> as the catalyst (Scheme 131).<sup>244</sup> The authors used monoprotected amino acids *N*-Ac-Leucine as the ligand for this transformation in MeCN at 60 °C. Silver pivalate and potassium carbonate were used as the oxidant and base, respectively. This reaction proceeds under moderate reaction conditions to afford a broad range of vinyl, aryl and heterocyclic substituted acrylamides in up to 93% yield.

### Scheme 131. Cp\*Rh(III)-Catalyzed C-H (Het)arylation of *N*-2,6-Difluoroaryl Acrylamides




**4.3.2. Alkylation.** In 2013, the Chatani group reported a bidentate chelation-assisted, nickel-catalyzed C(sp<sup>2</sup>)-H alkylation of acrylamides and benzamides with unactivated alkyl halides (Scheme 132).<sup>245</sup> They employed 10 mol% of Ni(OTf)<sub>2</sub> as the catalyst and 20 mol% of PPh<sub>3</sub> as the ligand in toluene solvent at 140 °C. The desired alkylated products were synthesized in high yields by using sodium carbonate as a base.

### Scheme 132. Nickel-Catalyzed Vinyl C-H Alkylation of Acrylamides with *n*-Butyl Bromide





In 2014, Nakamura et al. extended their inexpensive and benign iron-catalysis strategy to selectively alkylate C-H bonds of various arene, heteroarene and alkene molecules (Scheme 133).<sup>246</sup> By taking advantage of the efficient Fe(acac)<sub>3</sub>/diphosphine catalytic system, a diverse range of primary and secondary alkyl halides, mesylate, and tosylates alkylated the C-H bonds. ArZnBr was used as a base in this case. The reaction proceeds stereospecifically for alkene substrate. Around the same time, Cook and co-workers also realized an analogous iron-catalyzed C(sp<sup>2</sup>)-H alkylation of acrylamides with *n*-butyl bromide in 2-methyltetrahydrofuran (2-MeTHF).<sup>247</sup> The reaction does not require a co-oxidant, and proceeds in less than 10 min.

### Scheme 133. Iron-Catalyzed Directed C-H Alkylation of Olefinic Carboxamides







Later, Nakamura and co-workers further reported a stereospecific iron/diphosphine catalyzed C-H alkylation which utilized a bidentate 8-aminoquinoline auxiliary (Scheme 134).<sup>248</sup> Either primary or secondary alkylzinc halide was used as the alkylation source while a dichloroalkane (DCIB) was screened as an oxidant in this protocol. The authors proposed that the reaction may involve an organoiron(III) species, which selectively activates the alkenyl C-H bond of monosubstituted and unsubstituted acrylamides, to alkylate in a stereoselective fashion. It should be mentioned that homocoupling of the organometallic reagent and β-hydrogen elimination were not observed in this iron-catalyzed protocol.

### Scheme 134. Iron-Catalyzed Directed C-H Alkylation of Alkenes with Alkylzinc Halides




Loh and co-workers came up with a substrate-controlled strategy for alkenyl C(sp<sup>2</sup>)-H activation reactions. In 2017, they presented an efficient route to synthesize diverse  $\beta$ -alkylated acrylamides and dihydropyrro-2-ones by the activation of alkenyl C-H bonds of *N*-tosyl acrylamides with acrylosilanes as the viable coupling partners under Cp<sup>\*</sup>Rh(III) catalysis (Scheme 135),<sup>249</sup> By slightly modifying the conditions, either C-H alkylation or alkenylation products could be obtained in satisfactory yields. The distinct reactivity of acrylosilanes is probably attributed to their inherent electronic properties which are distinctive from other carbonyl compounds. On the basis of their mechanistic studies, the authors proposed that the catalytic process may proceed via *N*-tosylamide-assisted alkenyl C-H bond cleavage followed by C=C double bond insertion to produce a rhodacycle species. Finally, protodemetalation occurs under acidic conditions to generate the alkylated products. Alternatively, the annulated products were formed through facile  $\beta$ -H elimination followed by an intramolecular *aza*-Michael reaction.

**Scheme 135. Rhodium(III)-Catalyzed Switchable C-H Bond Functionalizations of *N*-Tosyl Acrylamides with Acryloylsilanes**




**Proposed mechanism**




The rhodium(III)-catalyzed vinylic C-H alkylation and further migration reaction of electron-deficient acrylamides with various maleimides was achieved by Kim's group in 2016 under redox-neutral conditions (Scheme 136a),<sup>250</sup> affording the biologically important exo-cyclised succinimide-containing amides with high regio- and stereoselectivity. Inspired by this precedential work, Jeganmohan et al. also established an analogous C-H alkylation of acrylamides with maleimides in the presence of pivalic acid additive to afford olefin-migrated alkylated products under cost-effective Cp<sup>\*</sup>Co(III) catalysis (Scheme 136b).<sup>251</sup> The reaction is compatible with a broad range of substituted acrylamides and maleimides. Under the same conditions, the linear-selective alkylated products were formed with acrylates as the coupling partners.

**Scheme 136. Redox-Neutral Coupling of Acrylamides with Activated Alkenes**

a) Kim et al., 2016




b) Jeganmohan et al., 2020





Besides the above-mentioned activated alkene coupling partners, readily available nitroalkenes could also participate in the alkenyl C–H alkylation of acrylamides. In 2017, Ellman's group disclosed the efficient  $Cp^*Rh(III)$ -catalyzed alkenyl C–H bond addition to electron-deficient nitroalkenes under redox-neutral conditions (Scheme 137).<sup>262</sup> furnishing the nitro-containing C–H alkylation products in 58–77% yield.

### Scheme 137. $Cp^*Rh(III)$ -Catalyzed Alkenyl C–H Bond Addition to Nitroalkenes





More recently, the highly enantioselective vinylic C–H alkylation of  $\alpha,\beta$ -unsaturated amides with readily available crotonates was elegantly reported by Shibata and colleagues (Scheme 138).<sup>253</sup> The coupling adducts were generally obtained with high yield and ee (up to 99% yield and up to 95% ee) by using a chiral Ir catalyst.

### Scheme 138. Enantioselective Cross-Coupling of $\alpha,\beta$ -Unsaturated Amides with Crotonate Derivatives




In 2017, López, Mascareñas and Gulías illustrated an intramolecular Ir(I)-catalyzed functionalization reactions of alkenyl C–H bonds followed by an hydrocarbonation process to construct a diverse number of versatile cyclic systems bearing quaternary stereocenters (Scheme 139).<sup>254</sup> The aryl and heteroaryl counterparts were also found to be suitable C–H sources for this strategy. More interestingly, the cyclic products can be easily transformed to unusual cyclic ketones. Mechanistically, the authors proposed that the Ir(I) catalyst first undergoes a C–H activation step with the olefin to form an Ir–H intermediate, followed by a selective *exo*-migratory insertion to generate a cyclic Ir–complex. A subsequent C–H reductive elimination generated the expected product.

### Scheme 139. Enantioselective Iridium(I)-Catalyzed Intramolecular Hydrocarbonation of Alkenes




**4.3.3. Alkenylation.** In 2012, Loh and co-workers proposed a general method for oxidative cross-coupling acrylamides with diverse electron-deficient olefins (Scheme 140).<sup>255</sup> It is notable that both Ru and Rh were competent catalysts for this transformation, generating the corresponding (*Z,E*)-dienamides products in high yields with excellent stereoselectivities.

### Scheme 140. Ruthenium- and Rhodium-Catalyzed Cross-Coupling of Acrylamides with Alkenes




Successfully, Nakamura and co-workers reported an iron-catalyzed alkenylation of olefinic, aromatic and heteroaromatic substrates under mild oxidative conditions with aryl and alkenyl boron compounds (Scheme 141).<sup>256</sup> The authors argued that the use of zinc(II) salt is essential for this transformation, as it significantly facilitates the transfer of organic group from boron to iron catalyst. The process of alkenyl C–H activation involved the formation of a reactive organoiron(III) intermediate in this case.

### Scheme 141. Iron-Catalyzed C(sp<sup>2</sup>)-H Olefination of Alkene Carboxamides with Alkenyl Boronates




Zhong's group employed Weinreb amide as the directing group for the C–H olefinations of alkenes under Cp\*Rh(III) catalysis (Scheme 142).<sup>257</sup> This reaction provided an easy route to obtain highly functionalized (*Z,E*)-butadienes. Both terminal and internal alkenes were tolerated. It is important to note that the Weinreb amide is easy to synthesise and could be readily converted into other useful functional groups.

### Scheme 142. Rhodium-Catalyzed Weinreb Amide Directed Cross-Coupling between Electron-Deficient Alkenes




As an extension of this strategy, the same group further reported a direct Ru(II)-catalyzed alkenylation of electron-deficient olefins with allyl acetate via alkenyl C–H activation (Scheme 143).<sup>258</sup> By employing *N,N*-disubstituted aminocarbonyl as the chelating group, this oxidant-free strategy features high efficiency and decent stereo- and regioselectivities, which unlocks a novel synthetic tool for the synthesis of (*2Z,4E*)-butadiene skeletons.

### Scheme 143. Ruthenium-Catalyzed C–H Olefination of Electron-Deficient Alkenes with Allyl Acetates




An efficient approach for the oxidant-free cross-coupling reaction of activated alkenes has been established under Ru(II) catalysis by the group of Zhong and Zhang (Scheme 144).<sup>259</sup> With the assistance of the oxidizing directing group CONH(OMe), this strategy presents a benign, direct and efficient approach for the construction of synthetically useful 1,3-butadiene skeletons with excellent *Z,E* selectivities.

### Scheme 144. Cross-Coupling of Electron-Deficient Alkenes Using an Oxidizing Directing Group




To exploit the potential of cost-effective cobalt catalysis, the same group further achieved the  $\text{Cp}^*\text{Co}(\text{III})$ -catalyzed cross-coupling reactions between electron-deficient alkenes by using readily synthesizable **Co(III)** complexes  $[\text{Cp}^*\text{Co}(\text{CO})\text{I}_2]$ , giving rise to regioselective and stereoselective transformation of a wide variety of *Z,E*-dienamides, along with  $\gamma$ -alkenyl ketones by prudent selection of coupling partners (Scheme 145).<sup>260</sup> The carbonyl group character and reaction conditions are responsible for determining the selectivity of the process, which results in the formation of the two kinds of products. Acrylates afforded conjugated dienes via  $\beta$ -H elimination, while  $\alpha,\beta$ -unsaturated ketones produced the C–H alkylated products by proto-demetalation.

### Scheme 145. Cobalt-Catalyzed Olefinic C–H Bond Alkenylation and Alkylation




Furthermore, the same group also demonstrated the  $\text{Cp}^*\text{Ir}(\text{III})$ -catalyzed cross-coupling of electron-deficient olefins by integration of chelation-assisted vinylic C–H alkenylation and transfer hydrogenation, which remarkably obviates the use of a metal oxidant, and instead employs a hydrogen acceptor such as chloranil, leading to a highly site- and stereoselective synthesis of (*Z,E*)-configured dienamides in 47–97% yields (Scheme 146).<sup>261</sup>

### Scheme 146. Iridium(III)-Catalyzed Cross-Coupling Reactions of Alkenes by Hydrogen Transfer




Tanaka and Shibata demonstrated that a highly electron-deficient  $\text{Cp}^*\text{Rh}(\text{III})$  complex, bearing two ester moieties on the Cp ring,  $[\text{Cp}^*\text{Rh}^{\text{III}}]$ , could catalyze the aerobic oxidative cross-coupling of substituted acrylamides with both activated and unactivated alkenes (Scheme 147),<sup>262</sup> leading to (*zZ*, *4E*)-dienamides at relatively low temperature (80 °C). The authors employed tertiary, secondary, and primary amide directing groups in this chemistry to obtain the expected products. Through mechanistic studies, the authors revealed that the electron-deficient nature of  $[\text{Cp}^*\text{Rh}^{\text{III}}]$  complex greatly facilitates the turnover-limiting vinylic C–H bond cleavage of the acrylamides.

### Scheme 147. Electron-Deficient $\text{Rh}^{\text{III}}$ -Catalyzed Cross-Coupling of Acrylamides with Alkenes





Despite of this indisputable advances, chelation-assisted alkene-alkene cross-coupling reactions are predominantly restricted to form (*Z,E*)- and (*E,E*)-butadienes. Sparse examples were reported on the alkene-alkyne coupling, giving rise to highly valuable (*Z,Z*)-butadienes. In 2009, Tanaka and co-workers achieved a regioselective amide-directed alkenylations of olefinic  $\text{sp}^2$  C–H bond by using a cationic rhodium(I)/BIPHEP complex (Scheme 148).<sup>263</sup> In this case, the sagacious choice of a 1-pyrrolidinecarbonyl group as the chelating group dramatically accelerates the reaction under especially mild conditions.

### Scheme 148. Cationic $\text{Rh}(\text{I})$ /BIPHEP-Catalyzed Amide-Directed C–H Alkenylation





Later in 2017, a highly regio- and stereoselective ruthenium(II)-catalyzed cross-coupling reaction between  $\alpha$ -substituted acrylamides and internal alkynes via directed vinylic C–H bond activation has been realized by the group of Zhang and Zhong (Scheme 149).<sup>264</sup> This atom-economic *syn*-hydrovinylation of alkynes resulted in good efficiency and excellent stereoselectivities under oxidant-free conditions, providing an attractive approach for the synthesis of diverse highly functionalized (*zZ*,*4Z*)-butadienes.  $\beta$ -Substituted acrylamides, however, were proven to be incompatible with this protocol.

### Scheme 149. Synthesis of (*Z,Z*)-Dienamides via Ruthenium-Catalyzed Cross-Coupling between Alkenes and Alkynes





Subsequently, Zhong's group continued to achieve an iridium-catalyzed alkene–alkyne cross-coupling under ligand- and additive-free conditions (Scheme 150).<sup>265</sup> A broad variety of acrylamides could couple with internal alkynes, generating a diverse array of branched (*Z,Z*)-butadienes in up to 98% yield with excellent site- and stereoselectivities. As a particular highlight, the synthetic practicality and versatility was demonstrated by late-stage C–H functionalization of biologically active perillid and artemisic acid derived amides.

### Scheme 150. Iridium-Catalyzed Cross-Coupling between Alkenes and Alkynes




In 2018, Mascareñas, López, and their co-workers reported an intramolecular Ir(I)-catalyzed carboxamide-assisted C–H olefination via the addition of vinylic C(sp<sup>2</sup>)–H bonds to the unsaturated moiety of alkynes (Scheme 151),<sup>266</sup> providing a diverse variety of cyclic scaffolds bearing an *exo*-dienyl moiety with well-defined stereochemistry in modest to excellent yields.

### Scheme 151. Intramolecular Hydroalkenylation of Alkynes





The group of Wang and Yu demonstrated the examples of C–H alkenylation through an efficient cost-effective copper-catalyzed reaction of acrylamides with both acyclic and cyclic vinyl boronates in DMSO at 45 °C (Scheme 152).<sup>267</sup> Of note, the substrate scope and functional group tolerance of this method are superior to Pd-catalyzed cross-coupling with vinyl borons.

### Scheme 152. Cu-Catalyzed C–H Alkenylation of Acrylic Acids with Vinyl Boronates




By means of vinylic C(sp<sup>2</sup>)-H activation strategy, the Jeganmohan group developed an aerobic oxidative cross-coupling of acrylamides with unactivated aliphatic alkenes catalyzed by a Cp\*Rh(III) catalyst (Scheme 153).<sup>268</sup> The authors carried out the reaction in 1,4-dioxane with 5 mol% Rh(III) catalyst, 20 mol% AgSbF<sub>6</sub> and Cu(OAc)<sub>2</sub>·H<sub>2</sub>O at 100 °C to provide diverse synthetically valuable amide-functionalized 1,3-butadienes. The authors tentatively proposed a plausible reaction mechanism involving the chelation-assisted vinylic C-H activation via a carboxylate-assisted deprotonation pathway.

### Scheme 153. Rhodium(III)-Catalyzed C-H Olefination of Acrylamides with Unbiased Alkenes





Following this, the same group extended their alkenyl C-H bond functionalization work by disclosing the chemodivergent chelation-directed C-H vinylation and alkenylation of structurally diverse acrylamides with easily accessible vinyl acetates as the coupling partners under Co(III) and Ru(II) catalysis, respectively (Scheme 154).<sup>269</sup>

### Scheme 154. Chemodivergent Cross-Coupling of Acrylamides with Vinyl Acetate




In 2013, Glorius and colleagues illustrated a novel and straightforward construction of highly unsaturated [3]dendralenes by a Cp\*Rh(III)-catalyzed alkenyl C-H activation and cross-coupling with allenyl carbinol carbonates (Scheme 155).<sup>270</sup> An assortment of dendralenes with a variety of substitution patterns can be accessed with decent efficiency. The method is highly stereoselective and suited with various directing groups as well as a number of functional groups.

### Scheme 155. Rh(III)-Catalyzed Alkenyl C-H Activation of Acrylamides with Allenyl Carbinol Carbonates




An efficient and stereoselective strategy to produce trifluoromethylated 1,3-butadienes through the vinylic C-H activation of acrylamides with inexpensive 2-bromo-3,3,3-trifluoropropene (BTP) has been developed by Bouillon and Poisson (Scheme 156).<sup>271</sup> Intriguingly, pentafluoroethylated dienes could be also synthesized by the reaction of 2-bromo-3,3,4,4,4-pentafluorobutene with  $\alpha$ -aryl-acrylamides under similar conditions.

### Scheme 156. Rapid Synthesis of 3-Trifluoromethyl-Substituted 1,3-Butadienes via Palladium-Catalyzed C-H Bond Functionalization




Macrocycles are essential building blocks in numerous bioactive natural products and pharmaceutically relevant molecules which are especially difficult to synthesize in some cases. Accordingly, it is crucial to develop general and practical strategies for macrocycle synthesis.<sup>272</sup> In 2017, Loh and co-workers elegantly reported the first example of intramolecular oxidative annulation reaction between two activated olefins in the presence of Cp<sup>\*</sup>Rh(III) catalyst. The intramolecular alkene-alkene coupling reaction produced macrolactams containing a 1,3-conjugated diene moiety with excellent chemo- and stereoselectivity (Scheme 157).<sup>273</sup>

#### Scheme 157. Macrolide Synthesis via Intramolecular Oxidative Cross-Coupling of Alkenes





Later in 2020, Loh's group extended their intramolecular oxidative annulation strategy to  $\alpha,\beta$ -unsaturated ketone fragments, and successfully reported an expeditious method for macrolactams synthesis through a Cp<sup>\*</sup>Rh(III)-catalyzed alkene-alkene cross-coupling (Scheme 158).<sup>274</sup> The ring-closing reaction proceeded via a Rh(III)-catalyzed vinylic C–H activation process, which opens a new pathway to access diverse macrocyclic molecules of different ring sizes in satisfactory yields (33–72%). Moreover, macrolactams featuring a conjugated 1,3-diene moiety could be easily prepared with high chemoselectivities and *Z,E*-stereoselectivities. The versatile nature of the 1,3-diene moiety was further converted into many useful functional groups, thus allowing the synthesis of many different types of macrocyclics.

#### Scheme 158. Macrolactams Synthesis via Ring-Closing Alkene-Alkene Cross-Coupling





**4.3.4. Allylation.** The allylation reaction is one of the most synthetically useful transformations in organic synthesis because the allyl moieties could be readily converted into many versatile functional groups.<sup>275</sup> In particular, the transition-metal-catalyzed vinylic C–H allylation of readily available starting materials with diverse allylic electrophiles provides a reliable route to incorporate allyl units into organic molecules for the synthesis of skipped 1,4-diene skeletons. For example, Loh's group in 2015 established an elegant synthetic procedure to easily produce 1,4-diene skeletons via Cp<sup>\*</sup>Rh(III)-catalyzed C–H allylation of electron-deficient acrylamides with allyl acetates (Scheme 159).<sup>276</sup> As a consequence of using a weakly coordinating group, this reaction provided high yields, and a diverse variety of functional groups were tolerated while maintaining exceptional  $\gamma$ -selectivity.

#### Scheme 159. Cp<sup>\*</sup>Rh(III)-Catalyzed C–H Allylation of Electron-Deficient Alkenes with Allyl Acetates



The same year, Glorius et al. disclosed an amide-directed C–H allylation of aromatics, heteroaromatics, and acrylamides with allyl carbonates as reaction partners under cost-effective Cp<sup>\*</sup>Co(III) catalysis (Scheme 160).<sup>277</sup> A number of allyl carbonates can be readily employed to incorporate this synthetically useful functional group, allowing for the construction of skipped 1,4-dienes. Moreover, Kim and co-workers also achieved an analogous transformation under ruthenium catalysis to generate the allylation products in modest yields.<sup>278</sup>



#### Scheme 160. Cobalt(III)-Catalyzed C–H Allylation of Acrylamides with Allyl Carbonates





The highly chemo- and stereoselective Cp<sup>\*</sup>Rh(III)-catalyzed C–H allylation of acrylic acid derivatives with 4-vinyl-1,3-dioxolan-2-ones as coupling partners has been reported by Wang and co-workers (Scheme 161a),<sup>279</sup> allowing an efficient synthesis of highly functionalized allylic alcohols, which are the most synthetically useful building blocks in organic synthesis.<sup>280</sup> The reaction occurred uneventfully to produce the skipped dienes with remarkable efficiency and excellent *E/Z* ratios under ambient conditions. Meanwhile, the same group also illustrated the synthesis of 1,4-dienes through a Cp<sup>\*</sup>Rh(III)-catalyzed sequential vinylic C–H activation and C–C activation reaction with vinylcyclopropanes as a versatile coupling partner under identical conditions (Scheme 161b).<sup>281</sup>

### Scheme 161. Cp<sup>\*</sup>Rh(III)-Catalyzed Vinylic C–H Allylation of Acrylamides

a) Wang et al., 2015





b) Wang et al., 2015



The Ji group was able to efficiently synthesise a broad range of skipped 1,4-diene skeletons by a Ru(II)-catalyzed C–H allylation of electronically activated olefins with the assistance of a *N*-methoxycarbonyl directing group (Scheme 162).<sup>282</sup> This procedure utilizes easily available



allyl alcohols to couple with various electron-deficient acrylamides in the presence of inexpensive [RuCl<sub>2</sub>(*p*-cymene)<sub>2</sub>]<sub>2</sub> catalyst in aqueous solution. A variety of functional groups were tolerated to generate the expected products with excellent regio- and stereoselectivity.

### Scheme 162. Ruthenium-Catalyzed C–H Allylation of Acrylamides with Allyl Alcohols



Ackermann's group expanded their sustainable C–H activation strategy to disclose a highly regioselective iron-catalyzed C–H allylation of (hetero)arenes and alkenes with allyl chloride by the assistance of a removable bidentate triazole auxiliary (Scheme 163).<sup>283</sup> This triazole-assisted C–H allylation tolerates with both primary and secondary halides. Notably, the triazole chelating group could be easily removed in a traceless fashion. The authors carried out preliminary mechanistic investigations suggesting that a single-electron-transfer (SET) pathway is probably involved in this process.


### Scheme 163. Iron-Catalyzed Alkenyl C–H Allylation of Acrylamides with Allyl Chloride



In 2019, Zhang and Zhong elaborated an iridium-catalyzed olefinic C–H allylation of acrylamides with diverse conjugated 1,3-dienes, exclusively affording a variety of branched 1,4-diene skeletons (Scheme 164).<sup>284</sup> This atom-economic C–H allylation took place with the assistance of a NH-Ts amide group under additive-and


ligand-free conditions. Successfully, they further expanded to carry out this olefinic C–H allylation of NH-Ts acrylamides in water, efficiently affording the corresponding skipped 1,4-dienes in excellent yields (52–99%).<sup>285</sup>

#### Scheme 164. Iridium-Catalyzed Olefinic C–H Allylation of Acrylamides with Conjugated 1,3-Dienes



Moreover, Krische's group in 2009 established a chelation-directed C–H allylation of  $\alpha,\beta$ -unsaturated carboxamides via the addition of vinylic  $\text{C}(\text{sp}^2)$ –H bonds to 1,1-dimethylallenes catalyzed by a cationic iridium complex assembled from  $[\text{Ir}(\text{cod})_2]\text{BAR}^{\text{F}}_4$  and *rac*-BINAP (Scheme 165),<sup>286</sup> resulting in the adduct products of C–H prenylation in good yields as single isomers.

#### Scheme 165. Iridium-Catalyzed Alkenyl C–H Oxidative Addition–Allene Insertion of $\alpha,\beta$ -Unsaturated Carboxamides




**4.3.5. Alkynylation.** Alkyne functional groups are among the most commonly used motifs in a diverse variety of bioactive organic molecules. They can also serve as extremely versatile synthetic building blocks, and could be easily converted into other synthetically useful functional groups which dramatically leads to an increase in molecular complexity. It is well-known that the

Sonogashira cross-coupling is one of the most robust and powerful strategies for the straightforward synthesis of various functionalized alkynes. Considering the remarkable atom- and step-economy, transition-metal-catalyzed directed C–H bond alkynylation has been significantly advanced in recent years with the efficacious assistance of a suitable directing group, which greatly enables the rapid and straightforward assembly of diverse alkynes.<sup>287,288</sup>



Over the past decades, a diverse variety of olefinic C–H alkynylations of acrylamides have been realized by different transition metal catalysts, providing concise and efficient methods to highly functionalized conjugated 1,3-enynes. Specifically, the Chatani group described a palladium(II)-catalyzed chelation-assisted  $\text{C}(\text{sp}^2)$ –H alkynylation of aromatic amides by using bidentate 8-aminoquinoline as a removable auxiliary with 1-bromoalkyne (Scheme 166a).<sup>289</sup> Notably, This directed C–H alkynylation is applicable to vinylic C–H bond of  $\alpha,\beta$ -unsaturated amide, allowing the straightforward construction of a conjugated (*Z*)-enyne in 66% yield. Following this bidentate chelation strategy, Li and co-workers subsequently achieved the same vinylic C–H alkynylation by a nonprecious Ni/BDMAE catalyst system (Scheme 166b).<sup>290</sup> In this case, the identified flexible bis(2-dimethylaminoethyl) ether (BDMAE) ligand was crucial to achieve remarkable catalytic efficiency. Furthermore, Balaraman's group later demonstrated that the use of benzoic acid (10 mol%) as a ligand could also achieve comparable reactivity in the  $\text{Ni}(\text{OTf})_2$ -catalyzed olefinic C–H alkynylation reactions (Scheme 166c).<sup>291</sup>

#### Scheme 166. Palladium- and Nickel-Catalyzed Vinylic C–H Alkynylation of $\alpha,\beta$ -Unsaturated Amides


a) Chatani et al., 2012



b) Li et al., 2015





c) Balaraman et al., 2016



The straightforward protocol for the efficient assembly of *gem*-diethynylethenes would be through a sequential double vinylic C–H activation process. In this regard, Gao's laboratory recently published the first example of nickel-catalyzed 3,3-dialkynylation of 2-aryl acrylamides with 1-bromotriisopropylsilylacetylene (Scheme 167).<sup>292</sup> Of note, the sagacious choice of bidentate 8-aminoquinoline as the auxiliary was indispensable for this transformation. However, alkynylating reagents including phenylacetylene, 3-methylbut-1-yne, and *tert*-butyl(ethynyl)dimethylsilane were all found to be incompatible with the conditions.



### Scheme 167. Nickel-Catalyzed 3,3-Dialkynylation of 2-Aryl Acrylamides



Afterward, Ackermann and co-workers published an earth-abundant, environmentally benign iron-catalyzed C–H alkylation of various arenes and heteroarenes enabled by the assistance of a triazole auxiliary (Scheme 168).<sup>293</sup> Of note, the reaction was typically finished within 1



h at 65 °C in THF. This sustainable methodology was applicable to  $\alpha,\beta$ -unsaturated amide substrates, furnishing the corresponding 1,3-enynes in moderate yields (54–63%).

### Scheme 168. Iron-Catalyzed Alkenyl C–H Alkynylation of Alkenes through Triazole Assistance




In 2020, the Li group successfully realized a highly regioselective palladium/monoprotected amino acid (MPAA)-catalyzed C–H alkylation of weak coordination nitrogen functionality including diverse benzylamines, arylethyl amines, and benzedrines. In this report, readily available  $\alpha,\beta$ -unsaturated cinnamamides were also competent substrates to afford the conjugated 1,3-enynes in decent yields (Scheme 169).<sup>294</sup>

### Scheme 169. Palladium-Catalyzed Vinylic C–H Alkynylation of Cinnamamides with 1-Bromoalkyne




Loh and co-workers continued their research on alkylation reaction, and established an efficient approach for olefinic C–H alkylation of electron-deficient alkenes with ethynylbenziodoxole (TIPS-EBS) as the alkynylating reagent in presence of a Cp\*Rh(III) catalyst (Scheme 170).<sup>295</sup> Tosyl-imide group was employed as the directing group for this transformation. The weak coordinating ability of this directing group was responsible for the highly efficient and stereospecific C–H alkylation of olefinic C–H bonds. Operational simplicity, excellent functional group compatibility as well as especially mild reaction conditions were the key features of this strategy. Hence, this method represented an efficient route for the synthesis of synthetically valuable 1,3-enyne moieties. To showcase the potential of this protocol, the authors derivatized the obtained products into a series of pyridinone and triazole moieties.

### Scheme 170. Rhodium(III)-Catalyzed Olefinic C–H Alkynylation of Acrylamides



Almost simultaneously, Glorius and co-workers also established a direct olefinic C–H bond alkynylation of  $\alpha,\beta$ -unsaturated amides with hypervalent iodonium reagents (TIPS-EBS) under  $\text{Cp}^*\text{Rh(III)}$  catalysis (Scheme 171).<sup>296</sup> This experimentally simple C–H alkynylation reaction occurred smoothly under additive-free conditions, eventually providing an electronically inverted retrosynthetic disconnection of conjugated 1,3-enynes when compared to the classical Sonogashira cross-coupling. Unfortunately,  $\alpha,\beta$ -disubstituted acrylamides precluded alkynylation product formation in this process. Moreover, this protocol could be applied to late-stage C–H alkynylation of natural product piperine, albeit with 30% yield.


### Scheme 171. Synthesis of Conjugated 1,3-Enynes by $\text{Cp}^*\text{Rh(III)}$ -Catalyzed Alkynylation of Alkenes



In the same year, Li and co-workers illustrated an efficient chelation-assisted C–H alkynylation of a broad range of (hetero)arenes by means of  $\text{Cp}^*\text{Rh(III)}$  or  $\text{Cp}^*\text{Ir(III)}$  catalysis with hypervalent iodine-alkyne reagents under typically mild conditions (Scheme 172).<sup>297</sup> Olefin substrates


were also proved to be compatible in the presence of  $[\text{IrCp}^*\text{Cl}_2]_2$  (4 mol%) as a catalyst, enabling the formation of 1,3-enynes in 48–75% yields.

### Scheme 172. $\text{Cp}^*\text{Ir(III)}$ -Catalyzed C–H Alkynylation of $\alpha,\beta$ -Unsaturated Amides



Besides the preactivated alkynylating reagents such as ethynylbenziodoxolones (EBX) and haloalkynes, the use of commercially available terminal alkynes as the alkynylating reagents represents an atom-economical strategy for the C–H alkynylation reactions. In this regard, You and co-workers successfully reported a general and inexpensive  $\text{Co}(\text{acac})_3$ -catalyzed oxidative cross-coupling between a broad range of acrylamides and triisopropylsilylacetylene with the assistance of the bidentate 8-aminoquinoline auxiliary (Scheme 173).<sup>298</sup> Notably, a broad scope of 36 examples of this transformation was documented with yields of 33–82%. Moreover, this protocol was applicable to late-stage alkynylation of a derivative of steroid drug Epristeride.


### Scheme 173. $\text{Co(III)}$ -Catalyzed C–H Alkynylation of Alkenes with Triisopropylsilylacetylene



Meanwhile, Li's group elaborated an efficient example of C–H alkynylation of acrylamides by taking advantage of


terminal alkynes as the readily available alkynylating reagents under Cp\*Ir(III) catalysis (Scheme 174),<sup>299</sup> thus affording a variety of **multisubstituted** 1,3-enynes with excellent stereoselectivity.

#### Scheme 174. Cp\*Ir(III)-Catalyzed Stereoselective C–H Bond Alkynylation of $\alpha,\beta$ -Unsaturated Amides



**4.3.6. Fluoroalkylation.** The incorporation of a fluoroalkyl group into an organic molecule sometimes dramatically enhances its chemical stability, bioavailability, lipophilicity and so on. Accordingly, remarkable efforts have been devoted towards the development of efficient and straightforward approaches for the synthesis of fluoroalkyl-substituted molecules in past decades. In this context, the direct vinylic C–H trifluoromethylation has drawn significant attentions from synthetic organic chemists.<sup>300</sup> For example, Loh and co-workers in 2013 disclosed an efficient copper-catalyzed olefinic C–H trifluoromethylation of acrylamides with Togni's reagent (Scheme 175).<sup>301</sup> The Ts-protected directing group plays an important role in achieving highly *cis*-selective  $\beta$ -CF<sub>3</sub>-functionalized acrylamides in this strategy.


#### Scheme 175. Chelation-Assisted Copper-Catalyzed Olefinic C–H Trifluoromethylation of Acrylamides



Almost at the same time, Besset's group disclosed a regioselective copper-mediated C–H trifluoromethylation of acrylamides with commercially available Umemoto's reagent (Scheme 176).<sup>302</sup> The reaction occurred in the presence of CuI (1.1 equiv) under acidic conditions, thus


resulting in the formation of *Z*-trifluoromethylated  $\alpha,\beta$ -unsaturated amides in 11–69% yield.

#### Scheme 176. Copper-Mediated Vinylic C–H Bond Trifluoromethylation of $\alpha,\beta$ -Unsaturated Amides




Successfully, Tan and Liu reported an elegant transition-metal-free *E*-selective C–H  $\beta$ -trifluoromethylation of  $\alpha,\beta$ -unsaturated carbonyls with Togni's reagent as CF<sub>3</sub> radical precursor (Scheme 177).<sup>303</sup> In this protocol, the authors employed 1.5 equiv of tetrabutylammonium iodide as an initiator activate Togni's reagent to in situ generate the highly electrophilic iodine(III) species.<sup>304–306</sup> Interestingly, the judicious choice of sterically bulky  $\alpha$ -alkyl acrylamides could dramatically alter the reaction pattern of this strategy, and exclusively afforded the allylic C–H trifluoromethylated products with high efficiency under identical conditions.

#### Scheme 177. Transition-Metal-Free Vinylic C–H Trifluoromethylation of $\alpha,\beta$ -Unsaturated Amides




Besides the  $\beta$ -C-H trifluoromethylation of  $\alpha,\beta$ -unsaturated carbonyl compounds, Bi's group in 2014 successfully achieved a highly regioselective copper-catalyzed vinylic C-H  $\alpha$ -trifluoromethylation with Togni's reagent (Scheme 178).<sup>307</sup> Likewise, a broad range of alkene substrates including  $\alpha,\beta$ -unsaturated amides, esters, thioesters as well as enones could engage this transformation to stereospecifically furnish the (*E*)- $\alpha$ -trifluoromethylated products in modest to good yields (30–92%). The authors highlighted the value of this protocol by late-stage vinylic C-H trifluoromethylation of biologically active molecules.

### Scheme 178. Copper-Catalyzed Vinylic C-H $\alpha$ -Trifluoromethylation of $\alpha,\beta$ -Unsaturated Carbonyls




Later, Dai's group also described a regioselective C-H  $\alpha$ -trifluoromethylation of  $\alpha,\beta$ -unsaturated acrylamides with readily available, inexpensive Ruppert's reagent (TMSCF<sub>3</sub>) (Scheme 179).<sup>308</sup> The reaction was typically finished within 0.5 h at 120 °C in DMSO. A large variety of  $\beta$ -substituted acrylamide derivatives are well compatible with the conditions, thereafter providing the corresponding (*E*)-trifluoromethylated products in satisfactory yields.

### Scheme 179. Copper-Catalyzed $\alpha$ -Selective C-H Trifluoromethylation of Acrylamides with TMSCF<sub>3</sub>




Moreover, Xu and his co-workers expanded their sustainable electrochemical strategy,<sup>309</sup> and elaborated the efficient 1,2-hydroxydifluoromethylation and vinylic C-H bond difluoromethylation of various acrylamides by using CF<sub>2</sub>HOSO<sub>2</sub>NHNHBoc as the CF<sub>2</sub>H source under electrochemical catalysis. These electricity-powered oxidative olefin functionalizations do not require any metal catalyst or oxidant. The reaction outcome, 1,2-difunctionalization or C-H functionalization reaction, is greatly dictated by the substituents on the amide nitrogen atom of the acrylamides as compared to the reaction conditions. (Scheme 180).<sup>310</sup>

### Scheme 180. Electrochemical Difluoromethylation of Electron-Deficient Alkenes



Beset and co-workers reported a straightforward *Z*-selective 2,2,2-trifluoroethylation of acrylamides by Pd-catalyzed vinyl C-H activation with a fluorinated hypervalent iodine reagent as the CH<sub>2</sub>CF<sub>3</sub> source (Scheme 181).<sup>311</sup> Indeed, the reaction occurred smoothly at room temperature under additive-free conditions, enabling the direct synthesis of various *Z*-2,2,2-trifluoroethylated acrylamides in a stereoselective manner.

### Scheme 181. Pd-Catalyzed 2,2,2-Trifluoroethylation of Acrylamides at Room Temperature




**4.3.7. Halogenation.** The carbon-halogen bond is one of the central functional groups in organic chemistry. With the ever-increasing efforts devoted towards transition-metal-catalyzed C-H functionalizations, the direct C-H halogenation has also gained considerable momentum in recent years.<sup>312–315</sup> For example, Glorius and co-workers in 2013 demonstrated a remarkable Cp\*Rh(III)-catalyzed alkenyl C(sp<sup>2</sup>)-H bond halogenation of various  $\alpha,\beta$ -unsaturated amides with readily available *N*-Bromosuccinimide and *N*-Iodosuccinimide to obtain



diverse *Z*-haloacrylamide derivatives in DCE at 60 °C (Scheme 182a).<sup>316</sup> Indeed, a large variety of synthetically useful and versatile functional groups are tolerated in this case. Subsequently, they further extended to report the earth-abundant, inexpensive Cp\*Co(III)-catalyzed alkenyl C–H iodination of acrylamides to selectively afford the mono-iodinated products in modest yields (Scheme 182b).<sup>317</sup>

### Scheme 182. Synthesis of Haloacrylamides via C–H Halogenation of $\alpha,\beta$ -Unsaturated Acrylamides

a) Glorius et al., 2013





b) Glorius et al., 2014





In 2020, Besset's group disclosed a palladium-catalyzed *Z*-selective chlorination of  $\alpha,\beta$ -unsaturated acrylamides at room temperature with the inexpensive and commercially available *N*-chlorosuccinimide as the chlorinating agent (Scheme 183).<sup>318</sup> 8-Aminoquinoline directing group was readily employed to access a diverse variety of value-added chlorinated olefins as a single *Z* stereoisomer.

### Scheme 183. Palladium(II)-Catalyzed Vinylic C–H Chlorination of Acrylamides



More recently, Morrill et al. elaborated a sustainable electrochemical strategy for the highly *Z*-selective C–H chlorination of various acrylamides by taking advantage of MgCl<sub>2</sub> as both the chloride source and the electrolyte (Scheme 184).<sup>319</sup> Of note, this protocol obviates the use of transition-metal catalyst and organic oxidant which tolerates a variety of substituted acrylamides, providing an efficient access to an array of synthetically useful *Z*- $\beta$ -chloroacrylamides in an environmental-friendly manner.

### Scheme 184. Electrochemical Oxidative *Z*-Selective Vinylic C–H Chlorination of Acrylamides




**4.3.8. Annulation Reactions.** The efficient synthesis of substituted 2-pyridone scaffold is a contemporary field of great interest because a number of bioactive molecules, pharmaceuticals, and advanced functional materials contain this privileged moiety. Accordingly, remarkable efforts have been devoted towards the construction of 2-pyridone derivatives through the vinylic C–H functionalization over the past decade.<sup>320</sup> Specifically, Li and colleagues in 2010 described a Cp\*Rh(III)-catalyzed oxidative cross-coupling of acrylamides with alkynes through C–H/N–H bond functionalizations (Scheme 185a),<sup>321</sup> affording a variety of synthetically valuable 2-pyridones in appreciable to high yields (48–94%). The reaction occurred smoothly by using a low 0.5 mol% catalyst loading of [RhCp\*Cl<sub>2</sub>]<sub>2</sub> in conjunction with stoichiometric amounts of Cu(OAc)<sub>2</sub> (2.2 equiv) as the terminal oxidant. Unfortunately, the use of acrylamides bearing electron-deficient *N*-substituents led to unsatisfactory selectivities, and the  $\beta$ -substituents of the acrylamide substrates showed pronounced effects, furnishing a substituted indole as the major product in this protocol. Additionally, alkyl- and aryl-substituted unsymmetrical internal alkynes gave mixed products of regioisomers. Understandably, ligand is a key element to increase the reactivity and regioselectivity in C–H functionalization reactions. Later in 2011, the Rovis group successfully developed a new ligand, 1,3-di-*tert*-butylcyclopentadienyl (termed Cp<sup>t</sup>),<sup>322</sup> that could greatly


result in an improved regioselectivity in the alkyne insertion step (Scheme 185b).<sup>323</sup> Internal alkynes with unsymmetrical alkyl and aryl substitution engaged this reaction with good yields and satisfactory regioselectivity.

### Scheme 185. Oxidative [4 + 2] Annulation Reactions of Acrylamides with Alkynes

a) Li et al., 2010





b) Rovis et al., 2011



The same year, Ackermann's group disclosed an inexpensive ruthenium-catalyzed oxidative [4 + 2] annulation reactions of alkynes with broad substrate scopes (Scheme 186).<sup>324</sup> Notably, both electron-rich and electron-deficient acrylamides coupled well with diverse unsymmetrically substituted alkynes in this protocol to produce a series of 2-pyridones with remarkably high regioselectivity.



### Scheme 186. Oxidative [4 + 2] Annulation Reactions of Acrylamides with Alkynes



In 2016, Nakamura and Ilies achieved an efficient assembly of 2-pyridones from acrylamides and alkynes enabled by a sustainable iron catalyst under typically mild conditions (Scheme 187).<sup>325</sup> In this report, both acyclic and cyclic alkenes bearing 8-aminoquinoline directing group tolerated smoothly with diverse internal alkynes. A diverse

variety of synthetically useful functional groups including alkenyl, allyl, alkynyl, silyl and thienyl were readily converted, thus showing the robustness of this strategy. More importantly, unsymmetrical internal alkyne produces the 2-pyridones with excellent regioselectivity.


### Scheme 187. Iron-Catalyzed Cross-Coupling of Amides with Alkynes




Gratifyingly, transition-metal-catalyzed direct oxidative cyclization reactions of acrylamides with alkynes are not restricted to monoannulation. In 2017, the Sahoo group demonstrated an elegant example of Ru-catalyzed traceless sulfoximine DG-assisted oxidative double annulation of acrylamides with internal alkynes (Scheme 188a).<sup>326</sup> This one-pot protocol was able to assemble four bonds in a naked acrylamide substrate through a single operation, thereafter generating a variety of  $\pi$ -extended pyrido-fused-isoquinolinones which are widely encounter in numerous pharmaceutically relevant molecules and functional materials. Subsequently, the same group expanded this cascade annulation strategy to the annulation of acrylamides with alkynes and naphthoquinone, giving rise to an array of unusual spiro-fused-isoquinolones (Scheme 188b).<sup>327</sup>

### Scheme 188. Ru-Catalyzed Double Annulation of MPS-Enabled Acrylamides with Alkynes

a) Sahoo et al., 2017




b) Sahoo et al., 2018




Almost simultaneously, Fan and Zhang disclosed a  $\text{Cp}^*\text{Rh}$ -catalyzed cascade  $[4 + 2]$  annulation/lactonization reaction between acryloyl hydroxamates and ynoates bearing a tertiary propargyl alcohol, which opened a rapid route to assemble  $\gamma$ -lactone ring-fused pyridones (Scheme 189).<sup>328</sup> This redox-neutral protocol employed readily available starting materials, and occurred with high regioselectivity which was greatly facilitated by the chelation assistance of  $\text{Cp}^*\text{Rh(III)}$  catalyst with the hydroxyl group in the ynoates.

**Scheme 189.  $\text{Cp}^*\text{Rh(III)}$ -Catalyzed Cascade Reactions of Vinylic *N*-Alkoxyamides with 4-Hydroxy-2-alkynoates**



A similar regioselective oxidative annulation of acrylamides with 2-alkynoates through the Ru-catalyzed vinylic C–H activation was established by Mhaske and co-workers in 2019 (Scheme 190).<sup>329</sup> Contrary to the well-known Ru(II)- or Rh(III)-catalyzed annulation protocols, this strategy provided a complete reverse regioselectivity mainly due to the electron-withdrawing nature of the ester group in the 2-alkynoate substrates.


**Scheme 190. Ru-Catalyzed Regioselective Annulation of Acrylamides with 2-Alkynoates**



In 2014, Daugulis and colleagues illustrated the cobalt-catalyzed oxidative  $[4 + 2]$  annulation of  $\text{C}(\text{sp}^2)\text{-H}$  bond by using either picolinamide or aminoquinoline directing groups (Scheme 191a).<sup>330</sup> They used readily available cobalt(II) acetate tetrahydrate catalyst,  $\text{Mn}(\text{OAc})_2$  co-catalyst in conjunction with atmospheric oxygen as the terminal oxidant. Both internal and external alkynes were


tolerated in this protocol. Moreover, the Daugulis group also reported a cobalt-catalyzed carbonylation of aminoquinoline benzamides with carbon monoxide by using atmospheric oxygen as an oxidant in trifluoroethanol solvent at room temperature (Scheme 191b).<sup>331</sup>  $\text{Mn}(\text{OAc})_2$  was used as an efficient cocatalyst in this process. The corresponding imides were synthesized in good yields from acrylic and benzylic acid derivatives. Shortly after, they also extended their 8-aminoquinoline-directed, cobalt-catalyzed  $\text{sp}^2$  C–H bond of cinnamic and methacrylic amides with alkenes to produce nitrogen-containing heterocycles (Scheme 191c).<sup>332</sup>

**Scheme 191. Directed Cobalt-Catalyzed Oxidative Annulation Reaction of Vinylic C–H Bonds with Carbon Monoxide, Alkenes, and Alkynes**




Incorporating fluorine atom into organic molecules is highly valuable because fluorine can provide special characteristics, including increasing the metabolic stability of the molecule and improving the lipophilicity. In 2020, Cramer and co-workers successfully managed to produce 4-fluoro-2-pyridones that are particularly difficult to construct with excellent regioselectivity and efficiency (Scheme 192).<sup>333</sup> This reaction took advantage of alkenic C–H activation, alkynyl triazene insertion, then Lossen rearrangement to afford a key intermediate. Subsequent addition occurs to generate a triazenyl molecule that can be further modified using Wallach reaction to generate the desired product. Additionally, the triazenyl moiety can be modified into fluorinated alkoxy and trifluoromethyl groups.

**Scheme 192. Regioselective Access to 4-Fluoro-2-pyridones by a  $\text{Cp}^*\text{Rh(III)}$ -Catalyzed C–H Activation–Lossen Rearrangement–Wallach Reaction**




Mei and co-workers continued on their electrochemical transition metal catalysis strategy,<sup>334</sup> and elaborated the synthesis of  $\alpha$ -pyridones and  $\alpha$ -pyrones via electrochemical C–H activation (Scheme 193).<sup>335</sup> In this report, the authors employed a series of acrylamides and alkynes. Depending on the substituents on nitrogen atom and reaction temperature, a variety of structurally diverse  $\alpha$ -pyridones and  $\alpha$ -pyrones were produced accordingly in excellent yields. It should be mentioned that the undivided electrolytic conditions employed in this protocol ultimately avoided the use of any external oxidant.

### Scheme 193. Divergent Rh-Catalyzed Electrochemical Vinylic C–H Annulation of Acrylamides with Alkynes




Moreover, a notable cascade oxidative vinylic C–H/N–H annulation/alkenylation sequence of fumaramides with various internal alkynes was achieved by means of ruthenium(II) catalysis (Scheme 194).<sup>336</sup> Alkynes bearing a wide range of electron-donating functionalities were well compatible, while electron-withdrawing alkynes furnished the respective product with diminished efficiency. Asymmetric terminal alkynes, however, were found to be incompatible in this protocol. Gratifyingly, in the presence of  $K_2CO_3$  base, the reaction of fumaramides with alkynes proceeded efficiently under identical conditions, giving rise to 2,6-naphthyridine-1,5-diones as the major products.

### Scheme 194. Divergent Synthesis of Pyridones and Naphthyridinediones via Ru(II)-Catalyzed Double C(sp<sup>2</sup>)–H of Fumaramides with Alkynes



Apart from the intermolecular oxidative annulation reactions, Gulías, Mascareñas, and their co-workers in 2013 achieved a rhodium(III)-catalyzed the formal [4 + 2] annulation of alkyne-tethered acrylamides to readily generate tricyclic indolizinones derivatives with 62–73% yield in an intramolecular fashion (Scheme 195).<sup>337</sup> The authors performed DFT studies which suggested that the favoured migratory insertion of the alkyne moiety into the resulting rhodacycle generated from the initial C–H activation takes place into the Rh–N instead of the Rh–C bond in this process.


### Scheme 195. Intramolecular Cycloaddition Reaction of Acrylamides and Alkynes




However, this intramolecular protocol requires a superstoichiometric amounts of  $Cu(OAc)_2$  as the external oxidant, leading to the generation of undesired waste. In 2012, the Park group successfully reported an atom-economical oxidizing-directing-group strategy that make use of alkynetethered hydroxamic esters as the substrates, affording diverse hydroxyalkyl-substituted isoquinolones with excellent regioselectivity and broad substrate scope (Scheme 196a).<sup>338</sup> Notably, the utility of this redox-neutral methodology is elegantly demonstrated by the facile total synthesis of phenanthroindolizidine alkaloids using this  $Cp^*Rh(III)$ -catalyzed C–H activation as the key step. Inspired by this work, Meyer, Cossy, and co-workers later in 2019 further expanded this strategy toward the vinylic C–H functionalization of cyclobutenyl hydroxamates, enabling the rapid synthesis of cyclobuta[c]pyridines in decent yields (Scheme 196b).<sup>339</sup>

## Scheme 196. Rhodium(III)-Catalyzed Intramolecular Annulation of Alkynyl-ethered Hydroxamic Esters

a) Park et al., 2012




b) Meyer, Cossy et al., 2019





Another elegant example of intramolecular annulative reaction through the  $Cp^*Rh(III)$ -catalyzed vinylic C–H activation strategy was disclosed by Meyer and Cossy. They employed a series of  $\omega$ -alkynyl  $\alpha$ -substituted acrylic hydroxamates to undergo macrocyclization to construct a large variety of structurally diverse 12- to 21-membered macrocyclic pyridones. The judicious choice of a *O*-pivaloyl hydroxamate as the effective oxidizing-chelating group was crucial for this strategy to achieve high efficiency (Scheme 197).<sup>340</sup>

## Scheme 197. Synthesis of Macrocyclic Pyridones via Rh(III)-Catalyzed C–H Activation/Heterocyclization





Due to the absence of suitable bias, the vinylic C–H olefination with aliphatic unbiased olefins often suffers from low reactivity and poor regioselectivity.<sup>213</sup> In 2019, Rovis and co-workers examined the formal oxidative [4 + 2] annulation of acrylamides with diverse both terminal and internal unbiased aliphatic alkenes (Scheme 198).<sup>341</sup> The regioselectivity of this reaction was greatly improved by the use of their modified bulky  $Cp^t$  ligand ( $Cp^t = 1,3$ -di-*tert*-butylcyclopentadienyl) on the  $Rh^{III}$  catalyst. A large variety of unprotected  $\alpha,\beta$ -unsaturated- $\delta$ -lactams bearing a diverse array of functional groups were produced with modest to good yields and synthetically satisfactory regio/diastereoselectivity.

## Scheme 198. Rh(III)-Catalyzed Oxidative [4 + 2] Annulation of Acrylamides with Unbiased Alkenes




More recently, Wang, Sun, and colleagues elaborated a highly regioselective C–H functionalization/annulation reaction of *N*-sulfonyl acrylamides with unactivated allylbenzenes (Scheme 199).<sup>342</sup> The reaction occurred in the presence of 10 mol%  $Pd(TFA)_2$  and 20 mol%  $Cu(OAc)_2$  in conjunction with air as the terminal oxidant, which undergoes a  $C(sp^2)$ –H allylation/aminopalladation/ $\beta$ -H elimination/isomerization sequence, leading to an array of highly functionalized pyridinone derivatives in appreciable yields (43–53%).

## Scheme 199. Palladium-Catalyzed Regioselective C–H Functionalization/Annulation of Acrylamides with Allylbenzenes for the Synthesis of Pyridinones




By taking advantage of the picolinamide (PA) directing group, Liu and co-workers were able to construct quinolinone scaffolds through an intramolecular oxidative *endo*-cyclization reaction of vinyl  $C(sp^2)$ –H and aryl  $C(sp^2)$ –H of *N*-aryl acrylamides (Scheme 200).<sup>343</sup> It is worth noting that the cyclic hypervalent iodine  $Oa(III)$  is proven to be crucial for the reaction, delivering the expected cyclization product in satisfactory yields. Interestingly, a spiro product was formed for the substrate with incorporation of OMe in this case.

### Scheme 200. Pd(II)-Catalyzed Intramolecular C(sp<sup>2</sup>)-H/C(sp<sup>2</sup>)-H Coupling of *N*-Aryl Acrylamides




Synthesising *tetra*-substituted carbons are typically difficult in many cases, especially via C–H activation. However, Lu and co-workers were able to use  $\alpha$ -allenols as the coupling partners to couple with a  $\gamma$ -lactam containing a tetrasubstituted carbon in the presence of Cp<sup>\*</sup>Rh(III) catalyst (Scheme 201).<sup>344</sup> In ensuring chemo- and regioselectivity, the coordination of hydroxyl group in  $\alpha$ -allenols was found to be crucial as revealed from the detailed mechanistic studies. This regiocontrolled annulative [4 + 1] protocol provided an efficient route to produce a broad range of 1,5-dihydro-pyrrol-2-ones in synthetically useful yields.

### Scheme 201. Regiocontrolled Cross-Coupling of Vinylic Amides with $\alpha$ -Allenols




The slight variation of electronic or steric properties of allenols has the potential to alter their reactivity patterns of C–H activation reactions in many cases. For example, Maiti and Volla in 2016 reported a cobalt-catalyzed heterocyclization of acrylamides with allenols at room temperature (Scheme 202).<sup>345</sup> Here, the reaction of acrylamides with phenylallene and sterical 1,1-dimethylallene by readily available Co(acac)<sub>2</sub> smoothly afforded the corresponding dihydropyridone products, while the reaction with cyclohexylallene, methoxyallene or electron-deficient allenols such as allenyl phosphonate exclusively resulted in the formation of pyridones. The authors argued that the different regioselectivity of the products may be mainly due to the back donation of lone pair electrons from the oxygen atom of allene substrates.

### Scheme 202. Synthesis of Dihydropyridones and Pyridones via Cobalt-Catalyzed Alkenyl C–H Activation of Acrylamides with Allenes



Besides the oxidative annulation with alkynes, alkenes and allenols, Ma and Xu in 2015 successfully achieved an example of Pd(II)-catalyzed oxidative [4 + 2] annulation between acrylamides and arynes (Scheme 203).<sup>346</sup> In this protocol, the authors made use of 2-(trimethylsilyl)phenyl trifluoromethanesulfonate as the benzyne precursors. Various aryne precursors bearing both electron-rich and electron-deficient groups could smoothly participate in this annulation reaction, producing the corresponding quinolinones in synthetically useful yields.



### Scheme 203. Synthesis of Quinolinones via Palladium-Catalyzed Oxidative Annulation Reaction between Acrylamides and Arynes



The Zhu group was able to construct the same scaffolds through an efficient Cp<sup>\*</sup>Rh(III)-catalyzed alkenyl C–H

functionalization of *N*-methoxycycloalkene carboxamides with aryl boronic acid pinacol esters (Scheme 204).<sup>347</sup> A variety of *N*-methoxycycloalkene carboxamide derivatives reacted smoothly with diverse aryl boronates in the presence of 5 mol% [RhCp\*Cl<sub>2</sub>]<sub>2</sub> catalyst and 2 equivalents of Ag<sub>2</sub>O oxidant to furnish the corresponding cycloalkaquinolinones in up to 92% yield. On the basis of detailed mechanistic investigations, the authors revealed that this annulative transformation may occur via a dual C–H activation pathway.


#### Scheme 204. Cp\*Rh(III)-Catalyzed Cross-Coupling of *N*-Methoxycycloalkene-1-carboxamides with Aryl Boronates



Even though asymmetric C–H activation is extremely useful for obtaining chiral compounds, controlling their stereochemistry is an extremely difficult task in some cases.<sup>348–352</sup> Enantioselective C–H functionalization often demands harsh reaction conditions and high temperature that are incompatible with many asymmetric reactions. While the asymmetric C–H activation in the presence of Pd catalyst has gained much traction, catalytic asymmetric C–H activation in aryls have seen advancements due to the development of chiral cyclopentadienyl ligands and chiral acid adducts in past decades. Unfortunately, not much discovery has been done on C–H activation in alkenes.


Lately, an innovative synthetic metalloenzymes (ArMs) system has been established by the Rovis and McNaughton group. This is achieved by inserting a Cp\*Rh unit in the mSav active site to asymmetrically produce  $\alpha,\beta$ -unsaturated- $\delta$ -lactams, which is done by alkenic C–H activation (Scheme 205).<sup>353</sup> This [4 + 2] annulative protocol made use of monomeric streptavidin (mSav)-Rh(III) to catalyze the reaction, instead of previous mechanisms that use biotin-tetrameric streptavidin (biotin-tSav) that was established by Whitesides,<sup>354</sup> and Ward.<sup>355</sup> This new concept is efficient and provides up to 99% yield and enantiomeric excess of 97% under mild and aqueous conditions. Notably, the products can also subsequently be easily converted into enantioenriched piperidines.

#### Scheme 205. Asymmetric $\delta$ -Lactam Synthesis with a Monomeric Streptavidin Artificial Metalloenzyme




Cramer and co-workers in 2019 revealed another strategy to enantioselectively activate the alkenyl C–H bonds to obtain enantioenriched  $\alpha,\beta$ -unsaturated- $\gamma$ -lactams bearing a quaternary stereocentre by means of a chiral Rh(III) catalyst (Scheme 206).<sup>356</sup> This procedure can be used on a wide variety of acrylamides and allenes to provide an excellent enantioselectivity of up to 97:3 er.

#### Scheme 206. Synthesis of $\alpha\beta$ -Pyrrol-2-ones through [4 + 1] Annulation of Acryl Amides and Allenes





In 2015, Wang et al. illustrated the general protocol for the Cp\*Rh(III)-catalyzed cyclization reaction of *N*-methoxymethacrylamides and diazo compounds to construct pyridone derivatives.<sup>357</sup> In the following year, Ma and co-workers presented the Cp\*Rh(III)-catalyzed cyclization of *N*-tosylacrylamides with diazoacetoacetates to readily construct a series of multisubstituted  $\alpha$ -pyrones under redox-neutral conditions (Scheme 207).<sup>358</sup> More interestingly, the reaction between *N*-tosylacrylamides and diazoacetoacetates could also selectively undergo a formal oxidative [2 + 3] cyclization in the presence AgOAc as the oxidant via sequential C–H and C–C cleavage to assemble highly functionalized furans. Of note, the authors employed readily available acylsulfonamide as an oxidizing or a traceless directing group in this controllable protocol.

#### Scheme 207. Rh(III)-Catalyzed Oxidative Cyclization of Acrylamides with Diazo Compounds



Later in 2019, Dai and co-workers reported a domino annulation strategy to construct spirooxindole pyrrolones using C–H activation reaction under Cp\*Rh(III) catalysis together with Lossen rearrangement (Scheme 208).<sup>359</sup> As a particular highlight, the annulation of *N*-pivaloyloxy benzamides with diazo compounds is also demonstrated by late-stage diversification of pentoxifylline, endofolliculina and pregnenolone. In this report, the vinylic C–H bond is activated to afford an intermediate followed by carbene migratory insertion to generate alkyl rhodium species. Subsequent Lossen rearrangement occurs on this intermediate to generate another intermediate which then goes through nucleophilic addition and protonation to unambiguously furnish the spirooxindole pyrrolones.


### Scheme 208. Synthesis of Spirooxindole Pyrrolones by Cp\*Rh(III)-Catalyzed Vinylic C–H Activation/Carbene Insertion/Lossen Rearrangement Sequence



Despite of the remarkable advances on the above-mentioned  $\beta$ -olefinic C–H activation reactions which involves five-membered cyclometalation intermediates, the  $\alpha$ -selective alkenyl C(sp<sup>2</sup>)–H functionalization of  $\alpha,\beta$ -disubstituted acrylamides has been seldom reported. In 2016, an elegant study by Yu and co-workers elaborated the palladium-catalyzed  $\alpha$ -olefinic C–H functionalization of

diverse  $\alpha,\beta$ -unsaturated olefins with *t*-BuNC (Scheme 209).<sup>360</sup> By activating of  $\alpha$ -olefinic C–H bonds, the synthesis of highly *cis*-stereoselective 4-imino- $\beta$ -lactam was readily achieved. This protocol tolerated a broad range of heterocycles at the  $\beta$ -position using air as the sole oxidant. Upon the treatment of BF<sub>3</sub>·OEt<sub>2</sub>, the obtained products can be easily converted into 2-aminoquinoline derivatives, which are privileged scaffolds widespread in many natural products and pharmaceutically relevant molecules.


### Scheme 209. Synthesis of 4-Imino- $\beta$ -Lactams via Pd-Catalyzed $\alpha$ -Selective C–H Functionalization of Olefins





Moreover, the nickel-catalyzed oxidative [4 + 1] annulation of  $\alpha,\beta$ -unsaturated amides bearing the 8-aminoquinoline auxiliary with elemental selenium has been reported by Nishihara and co-workers under aerobic conditions (Scheme 210a),<sup>361</sup> which provides an alternative route to isoselenazolones in synthetically acceptable yields (50–64%). Shortly afterwards, Besset's group realized the synthesis of functionalized isothiazolone derivatives from  $\alpha,\beta$ -unsaturated acrylamides and an electrophilic SCN reagent through an aerobic palladium-catalyzed vinylic C–H Bond activation (Scheme 210b).<sup>362</sup>

### Scheme 210. Synthesis of Isoselenazolones and Isothiazolones through Vinylic C–H Activation

a) Nishihara et al., 2017




b) Besset et al., 2019




Nitriles are the key structural unit widely found in numerous organic molecules, and the direct redox-neutral nucleophilic addition to nitriles undoubtedly exhibits a better atom- and step-economy. More recently, Chen, and co-workers investigated the inert C–H addition to nitriles, and accomplished the Cp\*Rh(III)-catalyzed vinylic C(sp<sup>2</sup>)-H addition to phenoxyacetone nitriles followed by an annulation sequence, giving rise to a broad set of 1,5-dihydro-2H-pyrrol-2-ones in a one-pot manner with decent efficiency (40–81%). Besides acrylamides, the aryl counterparts were also proven to be viable C–H sources in this strategy (Scheme 211).<sup>363</sup>

**Scheme 211. Synthesis of 1,5-Dihydro-2H-pyrrol-2-ones by Cp\*Rh(III)-Catalyzed Vinylic C(sp<sup>2</sup>)-H Addition to Aliphatic Nitriles**




**4.3.9. Other Useful Reactions.** In 2015, Zhang and co-workers studied the nickel-catalyzed direct vinylic C–H thiolation of alkenes and arenes with diverse diaryl disulfides (Scheme 212).<sup>364</sup> Here, the authors employed 10 mol% NiCl<sub>2</sub>, 4 equivalents of TBAI in conjunction with 2 equivalents of sodium carbonate in DMSO at 100 °C to produce alkenyl sulfides and diaryl sulfides in good yields with a broad range of functional group tolerance.

**Scheme 212. Nickel-Catalyzed Vinylic C–H Thiolation of Acrylamides**




Later in 2018, Wang and Ji disclosed an efficient approach for the vinylic C–H thiolation through the assistance of a weakly coordination *N*-tosylamide group in presence of 2 mol% [RhCp\*Cl<sub>2</sub>]<sub>2</sub> catalyst, 50 mol% of silver carbonate and copper acetate in DMF at 80 °C (Scheme 213).<sup>365</sup> A diverse range of (*Z*)-alkenyl sulfides were synthesized in excellent yields. Notably, this protocol tolerated a large variety of functional groups, and up to 7100 of turnover numbers were obtained for this transformation using as low as 0.01 mol% Cp\*Rh(III) catalyst. The authors showcased the applicability of this protocol for the synthesis of (*Z*)-β-alkenyl selenides under identical conditions. Quite recently, Yu and co-workers also achieved an analogous olefinic C–H thiolation of *N*-2,6-difluoroaryl acrylamides with comparable efficiency.<sup>366</sup>

**Scheme 213. Cp\*Rh(III)-Catalyzed C–H Thiolation of *N*-Tosyl Acrylamides**




In 2017, the Jana group investigated a dual copper/manganese-catalyzed selenation of both acrylic and benzoic acids using 8-aminoquinoline auxiliary as the directing group (Scheme 214).<sup>367</sup> In presence of 40 mol% Cu(OAc)<sub>2</sub>·H<sub>2</sub>O and 20 mol% Mn(OAc)<sub>2</sub> in DMSO at 90 °C, a variety of (*Z*)-vinyl selenides were synthesized in good yields. Lithium *tert*-butoxide was used as a base for this transformation while atmospheric oxygen is used as a terminal oxidant.

**Scheme 214. Copper/Manganese Cocatalyzed Vinylic C–H Selenation of Alkenes**




A highly regioselective Pd-catalyzed chelation-assisted vinylic C–H bond trifluoromethylthiolation of acrylamides bearing the 8-aminoquinoline auxiliary was achieved by Bouillon and Besset in 2017 (Scheme 215).<sup>368</sup> The reaction tolerates a diverse array of  $\alpha,\beta$ -unsaturated acrylamides bearing different substitution patterns, and affords the corresponding SCF<sub>3</sub>-containing alkenes with complete Z-selectivity. The authors tentatively proposed a plausible Pd(II)/Pd(IV) catalytic mechanism for this protocol.

### Scheme 215. Pd(II)-Catalyzed Directed Vinylic C–H Trifluoromethylthiolation of Acrylamides



In 2018, Zhang et al. reported a general oxidative C–H/O–H cross-coupling reaction enabled by a Co(OAc)<sub>2</sub> catalyst for the selective vinylic C–H acyloxylation (Scheme 216).<sup>369</sup> 8-Aminoquinoline directing group was installed and used as the directing group for the acyloxylation of alkenes and arenes. A wide range of esters were readily synthesized from corresponding acids in presence of 20 mol% Co(OAc)<sub>2</sub>, silver sulphate and sodium carbonate in DCE under N<sub>2</sub> atmosphere at 70 °C. The oxidant silver sulphate, although used in excess, could be recovered back. The recyclable directing group makes this protocol as a robust strategy for the acyloxylation for the alkenyl carboxamides.


### Scheme 216. Cobalt(II)-Catalyzed C–H Acyloxylation of Acrylamides



Sulfones are an important class of organic molecules which can be served as versatile building blocks in a number of synthetic transformations, including classical Julia olefination and Ramberg-Backlund reaction. The direct functionalization of inert C–H bonds represents an


atom- and step-economical strategy to construct structurally diverse sulfones.<sup>370</sup> Recently, a highly stereoselective and straightforward synthesis of (Z)- $\beta$ -alkenyl sulfones through the copper-catalyzed reaction of N-tosyl acrylamides, 1,4-diazabicyclo[2.2.2]octane-sulfur dioxide surrogate in conjunction with aryldiazonium tetrafluoroborates has been accomplished by Wu and co-workers (Scheme 217).<sup>371</sup> In the presence of copper trifluoroacetate, the vinylic C(sp<sup>2</sup>)-H arylsulfonylation of acrylamides with sulfur dioxide (SO<sub>2</sub>) as the sulfonyl source proceeds smoothly, leading to diverse (Z)- $\beta$ -alkenyl sulfones with excellent regio- and stereoselectivities.

### Scheme 217. Copper-Catalyzed C–H Arylsulfonylation of Acrylamides with Sulfur Dioxide



Meanwhile, the Wu group further extended to report the direct C(sp<sup>2</sup>)-H alkylsulfonylation of N-tosyl acrylamides (Scheme 218).<sup>372</sup> The reaction was conducted in DMF at 50 °C using Cu(TFA)<sub>2</sub> as the catalyst. Phenyliodine(III) dicarboxylates were employed as the alkyl radical precursors with N-tosyl amidyl group as a weakly coordinating directing group. This copper-catalyzed three-component reaction provided diverse (Z)- $\beta$ -alkenyl alkylsulfones in satisfactory yields.


### Scheme 218. Copper-Catalyzed Three-Component Decarboxylative Alkylsulfonylation



In 2015, Fu's group reported a practical and highly efficient Cp\*Rh(III)-catalyzed vinylic C–H cyanation


reaction of various acrylamides and ketoximes with readily prepared, environmental friendly *N*-Cyano-*N*-phenyl-*p*-methylbenzenesulfonamide (NCTS) as the cyanation reagent (Scheme 219).<sup>373</sup> The protocol tolerated a variety of synthetically useful functional groups, allowing for the synthesis of versatile alkenyl nitriles in up to 90% yield.

### Scheme 219. Rhodium(III)-Catalyzed Vinylic C–H Cyanation of Acrylamides



The direct synthesis of vinylsilanes attracts special attention from synthetic community as they are an important class of valuable building blocks widely used in synthetic chemistry (Hiyama–Denmark coupling, Tamao–Fleming oxidation, etc.), polymers, and medicines.<sup>374–375</sup> In 2017, Zhang and co-workers elaborated a stereoselective palladium(II)-catalyzed vinylic  $C(sp^2)$ –H silylation of acrylamides with disilanes as **the silicon source** (Scheme 230).<sup>376</sup> The bidentate 8-aminoquinoline amide directing group was readily employed to exclusively produce *Z*-vinylsilanes in synthetically satisfactory yields. Unboubtedly, the use of environmentally benign 1,4-benzoquinone (BQ) as a nonmetal oxidant significantly enables this strategy synthetically appealing for the synthesis of *Z*-stereoselective vinylsilanes from readily available starting materials.


### Scheme 230. Pd(II)-Catalyzed Vinylic C–H Silylation of Acrylamides with Disilanes




The formation of C–O bonds is one of the fundamental reactions in synthetic chemistry. In 2015, Niu and Song successfully realized an inexpensive, earth-abundant cobalt-catalyzed vinylic  $C(sp^2)$ –H alkoxylation of both aromatic and olefinic carboxamides using their 2-aminopyridine-1-oxide as a removable *N,O*-bidentate auxiliary (Scheme 231a).<sup>377</sup> Indeed, the reaction worked smoothly under typically mild conditions and tolerated a broad range of both alcohols and amide substrates. Following this, Ackermann's group in 2017 devised an electrochemical cobalt-catalyzed alkenyl C–H alkoxylation of acrylamides which substantially avoided the use of stoichiometric silver(I) oxidants, providing a sustainable strategy to the synthesis of enoether derivatives (Scheme 231b).<sup>378</sup>

### Scheme 231. Direct Alkoxylation of Acrylamides through Vinylic C–H Activation

a) Niu, Song et al., 2015




b) Ackermann et al., 2017



In 2013, Chang et al. exploited the  $Cp^*Ir(III)$ -catalyzed olefinic C–H amidation of  $\alpha,\beta$ -unsaturated acrylamides by means of acyl azides as the nitrogen source (Scheme 232).<sup>379</sup> A broad range of acyl azides bearing various functional groups could readily be employed under mild conditions. This protocol obviates the use of external oxidants and releases  $N_2$  gas as the only byproduct, affording an environmentally benign access to a variety of *Z*-enamides with excellent regio- and stereoselectivity. Subsequently, the same group expanded to report a vinylic


C–H amidation with azidoformates as an easily deprotectable amino source, which eventually gives rise to *N*-protected enamines (Scheme 232).<sup>38o</sup>

### Scheme 232. Ir(III)-Catalyzed Vinylic C–H Amidation of Acrylamides with Acyl Azides and Azidoformates



Li's group was able to assemble synthetically useful enamides via a cost-effective  $\text{Cp}^*\text{Co(III)}$ -catalyzed alkenyl C–H amidation of *N*-methoxy acrylamides with dioxazolones as an amidating reagent (Scheme 233).<sup>38i</sup> Interestingly, elevating the reaction temperature to 110 °C, the resulting enamides could further undergo a dehydrative cyclization to afford pyrimidones, which can act as a ready directing group for subsequent second aryl C–H amidation to generate diamidated products in high yields.

### Scheme 233. $\text{Cp}^*\text{Co(III)}$ -Catalyzed Amidation of Olefinic C–H Bonds




#### 4.4. $\alpha,\beta$ -Unsaturated Ketones

In 1995, Trost and colleagues first elaborated the  $\text{Ru(0)}$ -catalyzed alkenyl C–H alkylation of 1-acetylcyclohexene with vinylsilanes, giving rise to the linear products in 94% and 76% yield, respectively (Scheme 234a).<sup>23i</sup> Around the same time, Murai's research group also achieved this transformation with a broader olefin scope (Scheme 234b).<sup>38j</sup> In this report, both five- and six-membered  $\alpha,\beta$ -enones worked well with diverse alkenes such as vinylsilanes, styrenes, and vinylcyclohexane to furnish the vinylic C–H alkylation products in 20% to quantitative yields. Later in 1998, Murai and co-workers further expanded to establish a stereodivergent C–H alkylation by


using acyclic  $\alpha,\beta$ -enones as substrates (Scheme 234c).<sup>38j</sup> Specifically, the reaction between phenylsubstituted *trans*-enone and styrene exclusively delivered the linear alkylated product with retention of the olefin geometry, while the methyl-substituted *trans*-enone generated a branched product with inversion of the olefin geometry. The authors tentatively outlined a C–H activation and a hydrometallation pathway to elucidate this discovery.

### Scheme 234. $\text{Ru(0)}$ -Catalyzed Vinylic C–H Alkylation of $\alpha,\beta$ -Unsaturated Ketones


a) Trost et al., 1995



b) Murai et al., 1995





c) Murai et al., 1998





Vinyl ketones were seldom utilized as coupling partners in cross-coupling reactions due to their potential to undergo homodimerization or polymerization. Quite recently, Dethe et al. investigated the viability of direct coupling reactions between two different vinyl ketones. They exploited the cost-effective ruthenium(II) catalysis for the stereoselective oxidative coupling of vinyl ketones (Scheme 235).<sup>38k</sup> Markedly, a broad range of vinyl ketones were identified as viable substrates in this protocol, giving rise to highly functionalized (*E,E*)-1,6-dioxo-2,4-dienes in appreciable to good yields. Nevertheless, appreciable amounts of homo-coupling product were observed in most cases. Moreover, this strategy was applicable to the direct synthesis of some bioactive natural products.

### Scheme 235. Ruthenium(II)-Catalyzed Stereoselective Oxidative Cross-Coupling of Vinyl Ketones



Alkene annulations have been getting more recognition in recent years. Cheng and co-worker disclosed a multi-component reaction to produce highly substituted pyridinium salts in 2015 (Scheme 236).<sup>385</sup> The authors proposed that the imine was produced *in situ*, and underwent oxidative coupling with alkynes to obtain highly-substituted pyridinium salts.



### Scheme 236. Synthesis of Pyridinium Salts via Rhodium(III)-Catalyzed Alkenyl C–H Activation



Your's group elaborated a general method for producing ploysubstituted furans by ketone-directed vinylic C–H annulation of  $\alpha,\beta$ -unsaturated ketones with alkynes under synergistic Rh/Cu catalysis (Scheme 237).<sup>386</sup> A series of tri- and tetra-substituted furans can be synthesized through the vinylic C–H activation followed by [4 + 2] annulation and ring contraction processes. This reaction mechanism is supported by <sup>18</sup>O-labelling experiments and intermediate verification reactions. The series of reactions begin when the alkene C–H bond is activated, allowing the alkyne to be inserted and hence forming a seven-membered rhodacycle. This process is directed by the ketone group. The intermediate pyrylium salt is then



produced via reductive elimination to give the [4 + 2] O-annulation intermediate. The Rh(I) salt is also produced. The superoxide radical then undergoes nucleophilic attack to give a hydroperoxide intermediate, and then heterolysis of the O–O bond occurs with intramolecular cyclisation to give O-acylated furanium. The final desired product is then produced via hydrolysis.

### Scheme 237. Synthesis of Ploysubstituted Furans via Cp\*Rh(III)-Catalyzed Alkenyl C–H Activation $\alpha$ -Aryl Enones with Alkynes





Following this, the same group further published an efficient synthesis of 1*H*-benzo[*f*]chromene derivatives by the reaction between exocyclic  $\alpha,\beta$ -enones and alkynes under Cp\*Rh(III) catalysis (Scheme 238).<sup>387</sup> This strategy underwent the formal [4 + 2] vinylic C–H O-annulation to afford the active pyrylium salts as the intermediates, which subsequently participated in a base-promoted 1,5-H shift process. By using this protocol, a scope of 25 examples was documented in decent yields.

### Scheme 238. Cp\*Rh(III)-Catalyzed [4 + 2] Vinylic C–H O-Annulation of Exocyclic Enones with Alkynes and 1,5-H Shift





In addition to these annulation reactions, significant advances on the vinylic  $\alpha$ -C-H functionalization of  $\alpha,\beta$ -enones has been made over the decades. Specifically, when exposure of cyclic enones or enals to  $\text{BiAr}_3\text{Cl}_2$  and Hunig's base (DIPEA) in the presence of a catalytic amount of tributylphosphine at room temperature, an array of  $\alpha$ -arylated enones and enals were smoothly produced in decent yields (44–93%) (Scheme 239).<sup>388</sup>

### Scheme 239. Phosphine-Catalyzed $\alpha$ -Arylation of Enones and Enals Using Hypervalent Bismuth Reagents





Recently, Wengryniuk and co-workers established an alternative approach for the synthesis of  $\alpha$ -arylated enones through a metal-free alkenyl C-H arylation mediated by hypervalent iodine(III) reagents (Scheme 240).<sup>389</sup> The reaction occurred via a novel reductive iodonium Claisen rearrangement of the in situ-generated  $\beta$ -pyridinium silyl enol ethers. A broad scope of 27 examples was documented with yield up to 90%. Interestingly, the incorporated arenes maintained the synthetically valuable iodine functional handle, enabling further elaboration by traditional cross-coupling reactions.

### Scheme 240. Metal-Free Vinylic C-H Arylation of Enones Mediated by Hypervalent Iodine Reagents



Huang and collaborators in 2015 developed an efficient method to assemble functionalized  $\alpha$ -benzylated enones (Scheme 241).<sup>390</sup> The authors discovered that a reaction between enones and simple toluenes in the presence of  $\text{Cu}(\text{tfacac})_2$  ( $\text{tfacac} = \text{CF}_3\text{COCHCOCH}_3$ ) catalyst generated a series of  $\alpha$ -substituted enones containing different groups. In this report, a broad scope of 27 examples was presented with yield up to 98%. This oxidative coupling protocol made use of di-*tert*-butyl peroxide (TBP) as the oxidant to trigger the generation of benzyl carbon radical. Notably, methyl cinnamate could be also used as the substrate in this case, albeit with a lower yield (28%).


### Scheme 241. Copper-Catalyzed Regioselective Alkenyl C-H Benzoylation of Enones with Toluenes



Cho's group elaborated a practical route for the easy preparation of  $\alpha$ -trifluoromethylthio- $\alpha,\beta$ -unsaturated carbonyls in decent yields via a highly efficient DABCO-mediated alkenyl C-H trifluoromethylthiolation of diverse  $\alpha,\beta$ -unsaturated carbonyls by using easily available *N*-trifluoromethylthio-dibenzenesulfonimide as the  $\text{SCF}_3$  source (Scheme 242).<sup>391</sup> Impressively, the reaction occurred smoothly at room temperature, and was well compatible with a myriad of  $\alpha,\beta$ -unsaturated esters to afford the corresponding products in satisfactory yields.


### Scheme 242. DABCO-Mediated Electrophilic Alkenyl C-H Trifluoromethylthiolation of $\alpha,\beta$ -Unsaturated

## Carbonyl Compounds



An impressive work by Hwang and Ryu uncovered the catalytic carbon insertion into the  $\beta$ -vinyl C–H bond of cyclic enones with diazoacetates as the coupling partner under boron Lewis acid catalysis (Scheme 243).<sup>392</sup> The authors utilized readily available  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  or a novel oxazaborolidinium ion as the efficacious catalyst. Meanwhile, the asymmetric version of this process was also achieved by using a chiral oxazaborolidinium ion catalyst (Scheme 243).<sup>393</sup> A variety of enantioenriched  $\beta$ -substituted cyclic enones were obtained in up to 99% yield along with excellent enantiomeric excess (up to 99% ee). As a particular highlight, the potential of this protocol was demonstrated by the formal synthesis of (+)-epijuabione.

### Scheme 243. Catalytic Carbon Insertion into Vinyl $\beta$ -C–H Bond of Cyclic Enones and Its Enantioselective Version




Moreover, Wang, Pan, and their colleagues disclosed a regioselective vinylic C–H amination for the robust synthesis of  $\beta$ -aryl enaminones from chalcones and benzyl azides (Scheme 244a).<sup>394</sup> By means of a catalytic amount of


$\text{Ce}(\text{OTf})_3$  (5 mol%), the reaction of chalcone with benzyl azides furnished a diverse array of multisubstituted  $\beta$ -aryl enaminones in appreciable to excellent yields with complete *Z*-selectivity. The authors rationally speculated that the initial regioselective 1,3-dipolar cycloaddition of chalcone with azides produced the unstable triazoline intermediate, which underwent decomposition to afford a zwitterionic species. Subsequent 1,2-H migration resulted in the formation of the imine intermediate, followed by 1,3-hydrogen shift to afford the  $\beta$ -enaminones. A couple of years later, Phan's group also elaborated a similar synthesis of  $\beta$ -enaminones through trisulfur-radical-anion ( $\text{S}_3^-$ )-triggered alkenyl C–H amination of  $\alpha,\beta$ -unsaturated carbonyls with simple amines (Scheme 244b).<sup>395</sup>

### Scheme 244. Alkenyl $\beta$ -C–H Amination of Enaminones

#### a) Wang, Pan et al., 2014





#### b) Wang, Phan et al., 2020



## 4.5. $\alpha,\beta$ -Unsaturated Imines



Over the past years, transition-metal-catalyzed alkenyl C–H bond functionalizations of  $\alpha,\beta$ -unsaturated imines has also emerged rapidly as a versatile tool to the synthesis of various complex nitrogen-containing molecules. Early in 2002, pioneering work of Miura and co-workers elegantly elaborated the Ru(0)-catalyzed carbonylation of olefinic C–H bonds of  $\alpha,\beta$ -unsaturated imines with alkenes and carbon monoxide (CO) to synthesize  $\beta,\gamma$ -unsaturated  $\gamma$ -butyrolactams (Scheme 245).<sup>396</sup> Various aryl and alkyl groups at  $\beta$ -position of the imine substrates reacted uneventfully under the conditions to produce the corresponding lactams in appreciable to good yields. The authors tentatively proposed that the reaction occurred via the initial formation of ketone derivatives generated by the direct carbonylation at the vinylic  $\beta$ -C–H bond of  $\alpha,\beta$ -unsaturated imines. An intramolecular attack by the imino nitrogen on the carbonyl group followed by a 1,2-shift of the ethyl group then afforded the expected product.

### Scheme 245. $\text{Ru}_3(\text{CO})_{12}$ -Catalyzed Reaction of $\alpha,\beta$ -Unsaturated Imines with CO and Alkenes





Later, the Ellman group uncovered the rhodium-catalyzed C–C bond formation protocol through a stereoselective alkylation of diverse  $\alpha,\beta$ -unsaturated imines via alkenyl C–H activation (Scheme 246).<sup>397</sup> They achieved the highly stereoselective synthesis of the tri- and tetrasubstituted  $\alpha,\beta$ -unsaturated imines by means of the electron-donating (dicyclohexylphosphinyl)ferrocene ligand. The *Z*-selective imines could be further hydrolyzed to obtain the  $\beta$ -alkylated  $\alpha,\beta$ -unsaturated aldehydes.

#### Scheme 246. Rh-Catalyzed C–H Alkylation of $\alpha,\beta$ -Unsaturated Imines




In continuation with their C–H bond functionalization investigations, Ellman's group further elaborated the synthesis of highly substituted pyridines from  $\alpha,\beta$ -unsaturated imines and diverse alkynes via alkenyl C–H alkylation/electrocyclization/aromatization sequence (Scheme 247).<sup>398</sup> In this case, the authors developed new class of ligands, and extensively investigated the possible catalytic mechanism of transformation through kinetic simulation studies.

#### Scheme 247. One-Pot Synthesis of Pyridines from $\alpha,\beta$ -Unsaturated Imines and Alkynes




Following this, the same group expanded their rhodium-catalyzed olefinic C(sp<sup>2</sup>)–H activation strategy for the asymmetric synthesis of (–)-Incarvilleine via an intramolecular alkenylation of  $\alpha,\beta$ -unsaturated imines (Scheme 248).<sup>399</sup> By loading 2.5 mol% of the Rh catalysts in conjunction with employing the mist selective ligand (DMAPh)PEt<sub>2</sub>, they synthesized the key intermediate with moderate diastereoselectivity.

#### Scheme 248. Asymmetric Synthesis of Fragment of (–)-Incarvilleine




Subsequently, Kuninobu, Takai, and their colleagues elegantly demonstrated the synthesis of cyclopentadienyl-rhenium complexes through olefinic C–H activation (Scheme 249).<sup>400</sup> The ketimines reacted with  $\alpha,\beta$ -unsaturated carbonyl compounds in the presence of a rhenium(o) complex, Re<sub>2</sub>(CO)<sub>10</sub>, affording a series of cyclopentadienyl–rhenium complex in 47–94% yield.

#### Scheme 249. Synthesis of Cyclopentadienyl-Rhenium Complexes from Ketimines, $\alpha,\beta$ -Unsaturated Carbonyls and Rhenium Complex





#### Proposed mechanism




Tetrahydropyridines are present in numerous bioactive natural products. Moreover, they are also extremely versatile intermediates for alkene addition reactions to produce privileged piperidine frameworks, which are the most prevalent nitrogen-containing heterocycles widely found in a number of drug molecules.<sup>401,402</sup> As an extension of their continued interest in olefinic C–H activation of  $\alpha,\beta$ -unsaturated imines, the Ellman group in 2012 presented the Rh(I)-catalyzed highly diastereoselective synthesis of tetrahydropyridines (Scheme 250).<sup>403</sup> The established method tolerated a diverse set of internal alkynes bearing alkyl and aryl substituents on the imine-Nitrogen to yield multisubstituted piperidine derivatives in 52–95% yield with high diastereoselectivity. The authors rationalized that the observed stereochemical outcome may be due to kinetically controlled protonation followed by face-selective borohydride reduction.

#### Scheme 250. Rh(I)-Catalyzed Diastereoselective Synthesis of Tetrahydropyridines



Followed this, the Ellman group further elaborated an extraordinary work to synthesize highly diastereoselective piperidine derivatives (Scheme 251).<sup>404</sup> By initial rhodium-catalyzed olefinic C–H activation, they synthesized multisubstituted piperidine derivatives from alkyne and  $\alpha,\beta$ -unsaturated imines, which then subjected to distereo- and regioselective protonation under either thermodynamic or kinetic control to provide the distinct iminium ion intermediate. Final nucleophilic additions occurred in an highly diastereoselective manner.


#### Scheme 251. Synthesis of Substituted Piperidine, Tropane and 2-Azabicyclo[3.1.0] Systems




In 2013, Yoshikai and co-workers illustrated an efficient synthesis of polysubstituted dihydropyridines through an annulative reaction between internal alkynes and  $\alpha,\beta$ -unsaturated imines enabled by a cobalt-triarylphosphine catalyst (Scheme 252).<sup>405</sup> The authors concluded that the

reaction involves of the cobalt-mediated olefinic C–H bond alkenylation followed by a  $6\pi$  electrocyclization process of the resulting azatriene intermediate. By means of this strategy, a variety of polysubstituted piperidines were synthesized in good yields.

### Scheme 252. Cobalt-Catalyzed Annulation of $\alpha,\beta$ -Unsaturated Imines with Alkynes




#### Proposed mechanism




Subsequently, The Ellman group expanded their Rh-catalyzed olefinic C–H bond functionalization strategy in combination with electrocyclization to synthesize densely substituted 2-silyl 1,2-dihydropyridines from  $\alpha,\beta$ -unsaturated imines and TMS acetylenes (Scheme 253).<sup>406</sup> The authors treated the piperidine intermediates with acids to produce unstabilized azomethane ylides which were used to synthesize various nitrogen containing heterocycles. In this case, polysubstituted piperidines, azabicyclo[3.1.0] systems and tropanes were synthesized from a common intermediate in good to excellent yields with high diastereoselectivity.

### Scheme 253. Synthesis of Substituted Piperidine, Tropane and 2-Azabicyclo[3.1.0] Systems





Successfully, the same group prepared a robust bench stable Rh catalyst from commercially available  $[\text{Rh}(\text{cod})\text{Cl}]_2$ . With the new catalyst, the reaction could be performed outside the glovebox under air atmosphere. The authors synthesized polysubstituted tetrahydropyridines via Rh(I)-catalyzed olefinic C–H activation/alkenylation/electrocyclization cascade followed by reduction (Scheme 254).<sup>407</sup> The practicality of this method was demonstrated through a >100 mol scale cascade reaction.

### Scheme 254. Synthesis of Multisubstituted Tetrahydropyridines via Rh(I)-Catalyzed C–H Bond Functionalization Sequence



A diverse array of highly functionalized pyridines were prepared from a reaction of  $\text{Cp}^*\text{Rh}(\text{III})$ -catalyzed alkenyl C–H activation of *N*-sulfonyl ketimines with diverse internal alkynes by Dong's group in 2014 (Scheme 255).<sup>408</sup> The N–S bond of *N*-sulfonyl ketimine substrates served as an internal oxidant for this transformation which involved new C–C/C–N bond formation, and S–C/S–N bond cleavage followed by desulfonylation under mild conditions.



### Scheme 255. Synthesis of Functionalized Pyridines from *N*-Sulfonyl Ketimines and Alkynes



#### 4.6. $\alpha,\beta$ -Unsaturated Oximes and Derivatives

$\alpha,\beta$ -Unsaturated oximes are an easily prepared and transformed synthetic scaffolds, which are commonly employed in synthetic organic chemistry.<sup>409,410</sup> Over the decades, remarkable advances have been achieved in the alkenyl C–H functionalizations of  $\alpha,\beta$ -unsaturated oximes and their derivatives. Specifically, Cheng's group in 2008 elegantly illustrated a ketoximes-assisted C–H activation and further intermolecular cyclization of  $\alpha,\beta$ -unsaturated oximes with internal alkynes under  $\text{RhCl}(\text{PPh}_3)_3$  catalysis, enabling an efficient route to assemble multisubstituted pyridine scaffolds (Scheme 256).<sup>411</sup> By using this strategy, the same group subsequently expanded to report a gram-scale synthesis of pentasubstituted pyridines (C2–C6) in a one-pot fashion. However, unsymmetrical alkynes afforded the annulation products with low regioselectivity while requiring high reaction temperature.<sup>412</sup>

#### Scheme 256. Rh(I)-Catalyzed C–H Annulation of $\alpha,\beta$ -Unsaturated Ketoximes with Alkynes





Following this, Rovis and co-workers elaborated a similar synthesis of pyridines via the Rh(III)-catalyzed reaction of  $\alpha,\beta$ -unsaturated oximes with alkynes under typically benign conditions (Scheme 257a).<sup>413</sup> Notably, it was established that sterically different ligands led to



different product selectivities in this protocol. Almost at the same time, Li and Chiba described an analogous transformation for the construction of highly substituted pyridine scaffolds with comparable yields (40–97%) under redox-neutral conditions (Scheme 257b).<sup>414</sup>

#### Scheme 257. Rh(III)-Catalyzed Synthesis of Pyridines from $\alpha,\beta$ -Unsaturated Oximes and Alkynes

a) Rovis et al., 2011





b) Li, Chiba et al., 2011




Apart from internal alkynes, the group of Bergman and Ellman also employed terminal alkynes as the coupling partner to engage in a Rh(I)-catalyzed one-pot C–H functionalization/electrocyclization/dehydration for the construction of multisubstituted pyridines with moderate to excellent regioselectivities (Scheme 258).<sup>415</sup> In this protocol, an electron-poor phosphite ligand was readily used to prevent the dimerization of the alkynes.

#### Scheme 258. Synthesis of Substituted Pyridines From $\alpha,\beta$ -Unsaturated Ketoximes and Terminal Alkynes



Subsequently, Ellman and co-workers extended to demonstrate a novel  $Cp^*Rh(III)$ -catalyzed C–H activation of  $\alpha$ -fluoro- $\alpha,\beta$ -unsaturated oximes with both internal and terminal alkynes for the synthesis of multisubstituted 3-fluoropyridines (Scheme 259).<sup>416</sup> It was established that structurally diverse oximes bearing aryl, heteroaryl, and alkyl  $\beta$ -substituents were efficacious substrates for this reaction. Notably, terminal alkynes participated in this annulative process exclusively affording 3-fluoropyridines as a single regioisomer.


### Scheme 259. $Cp^*Rh(III)$ -Catalyzed Synthesis of Fluorinated Pyridines



Moreover, Glorius and co-workers described an impressive  $Cp^*Rh(III)$ -catalyzed cyclization reaction between  $\alpha,\beta$ -unsaturated oximes and diazos through the dual activation of both aryl and vinyl C–H bonds, allowing the construction of a series of multisubstituted isoquinolines and pyridine  $N$ -oxides in high yields (Scheme 260).<sup>417</sup> This annulative strategy incorporated a tandem C–H activation, ring formation, and condensation


steps. The reaction also avoided the need for any external oxidant, while releasing  $N_2$  and  $H_2O$  as the byproducts.

### Scheme 260. Direct Synthesis of Multisubstituted Isoquinoline and Pyridine $N$ -Oxides from $\alpha,\beta$ -Unsaturated Oximes and Diazo Compounds



Later in 2020, Samanta et al. investigated the annulation reaction of unsaturated oxime ethers with fluorinated diazomalonates to construct 2-pyridone derivatives (Scheme 261).<sup>418</sup> Various unsaturated oxime ethers bearing with different substitution patterns were compatible, and a broad scope of 30 examples was documented in yield up to 92%. Notably, keto-oximes bearing five-, six-, seven and eight member rings proceeded uneventfully in this case.


### Scheme 261. Synthesis of 2-Pyridones by Rhodium(III)-Catalyzed Annulation of Unsaturated Oxime Ethers with Fluorinated Diazomalonates



In addition to these annulation examples, Ellman's group also elaborated a novel annulation strategy for the robust assembly of substituted furans and pyrroles. Specifically, the reaction between  $O$ -methyl  $\alpha,\beta$ -unsaturated oximes with both aldehydes and  $N$ -tosyl imines yields secondary alcohols and amine intermediates, which could readily undergo subsequent cyclization and aromatization to furnish biologically important


heterocyclic compounds in modest to excellent yields (Scheme 262).<sup>419</sup>

### Scheme 262. Rh(III)-Catalyzed Annulation Reaction for the Synthesis of Substituted Furans and Pyrroles



Encouraged by this work, the same group further identified a novel air-stable cationic Co(III)-catalyst, and expanded to evaluate the alkenyl C–H functionalization of  $\alpha,\beta$ -unsaturated oxime ethers with aldehydes, followed by a subsequent cyclization and aromatization, providing a cost-effective synthesis of multisubstituted furans in a single step (Scheme 263).<sup>420</sup> Interestingly, only a catalytic amount of AcOH additive was required for achieving high efficiency. This protocol was the first illustration of a Co(III)-catalyzed additions to aldehydes for the synthesis of furans, and was compatible with a diverse range of both (hetero)aromatic and aliphatic compounds. Moreover, a diverse array of indazoles can be obtained by this strategy with azobenzenes as the substrates.


### Scheme 263. Co(III)-Catalyzed Synthesis of Furans from $\alpha,\beta$ -Unsaturated Oxime Ethers



The group of Zhou and Li reported a  $\text{Cp}^*\text{Rh}(\text{III})$ -catalyzed activation of the alkenyl C–H bond to isocyanates, followed by a subsequent annulation to afford biologically important 5-ylidene pyrrol-2(5H)-ones (Scheme 264).<sup>421</sup> The reaction was atom-economic, and operated under benign and redox-neutral conditions without the need for any additive, while no


environmentally hazardous waste was generated.

### Scheme 264. Synthesis of 5-Ylidenepyrrol-2(5H)-ones from $\alpha,\beta$ -Unsaturated Oxime Ethers and Isocyanates



More recently, Ravikumar and co-workers also achieved a redox-neutral  $\text{Cp}^*\text{Co}(\text{III})$ -catalyzed intermolecular cyclization of  $\alpha,\beta$ -unsaturated oxime ethers with internal alkynes (Scheme 265).<sup>422</sup> A variety of multisubstituted pyridines were obtained in decent yields. Remarkably, the strategy exhibited good functional group compatibility, and was applicable for late-stage diversification of a bioactive molecule-dehydroepigenolone.

### Scheme 265. Redox-Neutral Co(III)-Catalyzed Alkenyl C–H Activation of $\alpha,\beta$ -Unsaturated Oxime Ethers with Internal Alkynes




In order to improve the regioselectivity and substrate scope, the Rovis group in 2013 developed a  $\text{Cp}^*\text{Rh}(\text{III})$ -catalyzed complementary approach for pyridine synthesis from  $\alpha,\beta$ -unsaturated oxime esters and simple olefins


(Scheme 266a).<sup>423</sup> The reaction was found to be highly regioselective, and affording a series of multisubstituted pyridines in decent yields when an activated alkene was employed. Mechanistic investigations revealed that the reaction proceeded through a reversible C–H activation, followed by the insertion of the olefin, a subsequent C–N bond forming/N–O bond breaking process afforded the corresponding pyridines. Recently, Tamura's group realized an analogous transformation enabled by Pd(II)-catalyzed vinylic C–H alkenylation of  $\alpha,\beta$ -unsaturated oxime ethers followed by aza- $6\pi$ -electrocyclization. A sterically hindered pyridine-based ligand was readily identified to facilitate this process (Scheme 266b).<sup>424</sup> Substrates with different degrees of substitution at the alkene moiety furnished the expected products in satisfactory yields (33–95%).

### Scheme 266. Regioselective Pyridine Synthesis from $\alpha,\beta$ -Unsaturated Oxime Esters and Alkenes

a) Rovis et al., 2013





b) Tamura et al., 2021





In the following year, the same group further illustrated an efficient decarboxylative cross-coupling of  $\alpha,\beta$ -unsaturated *O*-pivaloyl oximes with unsaturated carboxylic acids to assemble pyridine scaffolds (Scheme 267).<sup>425</sup> The carboxylic acid functioned as a traceless activating group, affording the expected pyridines in good yields and excellent regioselectivity. Further investigations into the mechanism dismissed the picolinic acid intermediate, while a rhodium complex intermediate was isolated providing clues into the mechanism of the process.

### Scheme 267. Decarboxylative Cross-Coupling of $\alpha,\beta$ -Unsaturated *O*-Pivaloyl Oximes with Unsaturated Carboxylic Acids




Almost at the same time, Glorius and colleagues elaborated a highly regioselective Cp<sup>\*</sup>Rh(III)-catalyzed C–H activation/cyclization/isomerization approach for the synthesis of multisubstituted pyridines via the reaction of  $\alpha,\beta$ -unsaturated oxime esters with 1,3-dienes (Scheme 268).<sup>426</sup> As a particular highlight, the protocol does not require an external oxidant, operated under redox-neutral conditions with a diverse substrate scope.

### Scheme 268. Cp<sup>\*</sup>Rh(III)-Catalyzed Pyridine Synthesis from $\alpha,\beta$ -Unsaturated Oxime Esters with 1,3-Dienes




In 2015, Rovis and co-workers investigated the Rh(III)-catalyzed C–H functionalization of  $\alpha,\beta$ -unsaturated oxime pivalates with 1,1-disubstituted alkenes in an effort to produce 2,3-dihydropyridines. In this report, the oxime pivalates go through a reversible alkenyl C–H insertion with Rh(III) complexes to afford the 5-membered rhodacycle. Subsequently, this intermediate readily undergoes irreversible migratory insertion with 1,1-disubstituted alkenes followed by a reductive elimination, affording the desired 2,3-dihydropyridines in 67–92% yield with excellent regioselectivities (Scheme 269).<sup>427</sup> Upon hydrogenation, the corresponding piperidines could be obtained, which are crucial core structures of numerous pharmaceutically relevant molecules.

### Scheme 269. Direct Synthesis of 2,3-Dihydropyridines through Rh(III)-Catalyzed C–H Functionalization of Unsaturated Oximes



Apart from the **annulation** examples discussed above, in 2014, Xu and co-workers reported a straightforward fluorination of vinylic C–H bonds by taking advantage of NFSI as the fluorinating agents under benign reaction conditions (ambient conditions in most examples). In the presence of a catalytic amount of inexpensive AgNO<sub>3</sub>, the alkenyl C(sp<sup>2</sup>)–H bonds bearing a diversity of functional groups were selectively fluorinated (Scheme 270).<sup>428</sup> On the basis of their mechanistic investigations, the authors tentatively proposed a Pd(II)/Pd(IV) catalytic cycle kickstarted by the formation of the cationic [Pd(NO<sub>3</sub>)]<sup>+</sup> species. Of note, the aryl counterparts were also found to be suitable C–H sources for this transformation.


#### Scheme 270. Nitrate-Mediated Alkenyl C–H Bond Fluorination Under Benign Conditions



Recently, Tamura and co-workers identified an alkyl aryl thioether ligand to enable the palladium(II)-catalyzed electrophilic C–H arylation of  $\alpha,\beta$ -unsaturated *O*-SEM oximes with arylboronic acids (Scheme 271).<sup>429</sup> Varying the structure of the ligand could greatly improve the efficiency of this protocol. The reaction was typically finished within 1 h at 90 °C in 1,4-dioxane, and a wide scope of 23 examples


of this transformation was documented with satisfactory yields of 22–96%.

#### Scheme 271. Pd(II)-Catalyzed C–H Arylation of $\alpha,\beta$ -Unsaturated *O*-SEM Oximes



Moreover, Cramer and co-workers expanded their chiral Cp\*Rh(III) catalysis strategy, and illustrated the first enantioselective alkenyl C–H activation/ring-opening between various  $\alpha,\beta$ -unsaturated oxime ethers and 2,3-diazabicyclo[2.2.1]hept-5-enes, enabling an efficient access to chiral cyclopentenylamines in decent yields and high enantioselectivities (Scheme 272).<sup>430</sup>

#### Scheme 272. Cp\*Rh(III)-Catalyzed Enantioselective C–H Activation/Ring-Opening to Synthesize Chiral Cyclopentenylamines




#### 4.7. 2-Vinylpyridines

In the early days of alkenyl C–H bond activation of 2-vinylpyridines, Murai and co-workers in 1998 first investigated the Ru(0)-catalyzed C–H carbonylation of alkenes. Reacting 2-vinylpyridines with carbon monoxide and ethene in a catalytic amount of Ru<sub>3</sub>(CO)<sub>12</sub> catalyst afforded propionylation at the alkenyl C–H bond in these

2-pyridylalkenes (Scheme 273).<sup>431</sup> The carbonylation took place regioselectively at the  $\gamma$ -position with respect to the nitrogen on the pyridine ring. Remarkably, this method can also be applied to *N*-(2-pyridyl)enamines, whereby the alkenyl moiety is segregated from the pyridine by an  $sp^3$ -nitrogen, generating the corresponding ethyl ketones as the product. Moreover,  $Rh_4(CO)_{12}$  also exhibited excellent catalytic activity for the reactions of *N*-(2-pyridyl)enamines with a diverse array of alkenes.


### Scheme 273. $Ru_3(CO)_{12}$ -Catalyzed Carbonylation of $C(sp^2)$ -H Bond of 2-Vinylpyridines



In 2005, the group of Oi and Inoue disclosed the regio- and stereoselective alkenyl C–H arylation reaction of 2-vinylpyridines with aryl bromides catalyzed by specific Ru(II)-phosphine complexes, delivering a series of  $\beta$ -arylated (*Z*)-2-alkenylpyridines with the aryl moiety *cis* to the pyridyl group, which is in sharp contrast to the  $Pd(OAc)_2$ -catalyzed Mizoroki-Heck coupling reaction (Scheme 274a).<sup>432</sup> Later in 2007, Ackermann's group developed an efficacious diastereoselective Ru(IV) carbene-catalyzed arylation of the vinylic  $C(sp^2)$ -H bonds of 2-vinylpyridines with aryl chlorides (Scheme 274b).<sup>433</sup> Of note, the direct arylation-hydrosilylation sequence was also presented in this report.


A couple of years later, the same group further extended to elaborate a chemo-, site-, and diastereoselective C–H functionalization of alkenes enabled by the well-defined air- and moisture-stable ruthenium(II) complexes bearing a trivalent phosphinous acid (PA) ligand (Scheme 274c).<sup>434</sup> Detailed mechanistic investigation revealed that a C–X bond breaking occurred via a single-electron transfer (SET) process, and proof of a phosphinic acid-assisted C–H ruthenation step was also reported. Following this, Gramage-Doria's group in 2018 also illustrated an analogous Ru(II)-catalyzed olefinic  $C(sp^2)$ -H bond (hetero)arylation by making use of diazines as the directing groups (Scheme 274d).<sup>435</sup>

### Scheme 274. Ru-Catalyzed Vinylic $C(sp^2)$ -H Bond Arylation of 2-Vinylpyridines




Afterward, Wang and co-workers achieved an efficient  $[Rh(cod)Cl]_2$ -catalyzed 2-pyridyl group-assisted alkenyl  $C(sp^2)$ -H functionalization reaction with aroyl- or acrylamides, giving rise to a series of arylated or alkenylated olefins in satisfactory yields with good functional group compatibility (Scheme 275).<sup>436</sup>

### Scheme 275. Rh-Catalyzed Vinylic $C(sp^2)$ -H Bond Functionalization with Aroyl- and Acrylamides




Kuninobu, Takai, and their co-workers reported on a Re-catalyzed insertion of  $\alpha,\beta$ -unsaturated carbonyls, alkynes, and aldehydes into alkenyl C(sp<sup>2</sup>)-H bonds to produce  $\gamma,\delta$ -unsaturated carbonyl compounds, dienes, and allyl silyl ethers, respectively (Scheme 276).<sup>437</sup> The reaction kickstarted with a vinylic C-H activation, further insertion of the unsaturated compounds to form the new Re-C bond, and a final reductive elimination (or transmetalation for aldehydes) to yield the expected products.

### Scheme 276. Re-Catalyzed Insertion of Unsaturated Molecules into Olefinic C-H Bonds



In a related study, Wang and co-workers elaborated a cost-effective Mn(I)-catalyzed nucleophilic addition of chemically inactive alkenyl C(sp<sup>2</sup>)-H bonds to aldehydes (Scheme 277).<sup>438</sup> The strategy exhibited a broad scope of substrates with excellent regio- and stereoselectivity. Detailed mechanistic investigation revealed a plausible mechanistic pathway of the reaction.


### Scheme 277. Mn(I)-Catalyzed Nucleophilic Addition of Alkenyl C(sp<sup>2</sup>)-H Bonds to Aldehydes



Nakamura and co-workers achieved an impressive iron-catalyzed stereospecific alkenyl C(sp<sup>2</sup>)-H bond functionalization with Grignard reagents for the synthesis of substituted alkenes (Scheme 278).<sup>439</sup> The reaction between aryl magnesium compounds and 2-vinylpyridines resulted in a stereospecific substitution at the C-H bond *syn* to directing group. The transformation operated with


ease and without isomerization of the alkene product. Mechanistic studies suggested that the strategy was consistent with a Fe-catalyzed C-H bond activation, instead of a Mizoroki-Heck reaction.

### Scheme 278. Fe(III)-Catalyzed Stereospecific Alkenyl C(sp<sup>2</sup>)-H Functionalization with Grignard Reagents



One year later, Shi's group uncovered a convenient approach of Cp\*Rh(III)-catalyzed alkenyl C(sp<sup>2</sup>)-H addition to both *N*-sulfonylaldimines and aryl aldehydes, eventually allowing an efficacious and atom-economical strategy to access a diverse array of allyl amines and allyl alcohols in decent yields (Scheme 279).<sup>440</sup>


### Scheme 279. Cp\*Rh(III)-Catalyzed Alkenyl C(sp<sup>2</sup>)-H Addition to *N*-Sulfonylaldimines and Aldehydes



Expanding their alkenyl C-H activation strategy, Loh's group in 2012 established a general approach for the alkenyl C-H sulfonylation of  $\alpha$ -methyl vinyl pyridines with sulfonyl chlorides (Scheme 280).<sup>441</sup> Various aryl sulfonyl chlorides were moderately efficient, giving rise to the expected products in reasonable yield (54–71%). Notably, aliphatic sulfonyl chloride was also a competent substrate for this transformation, albeit with diminished efficiency (25%). The authors tentatively proposed a plausible Pd(II)/Pd(IV) catalytic cycle to elucidate the possible


mechanism of this reaction.

### Scheme 280. Palladium(II)-Catalyzed Alkenyl C–H Bond Sulfonylation





Moreover, Yu and colleagues in 2016 presented a directed Cp\*Rh(III)-catalyzed C(sp<sup>2</sup>)-H Michael addition to CF<sub>3</sub>-substituted α,β-unsaturated ketones (Scheme 281).<sup>442</sup> Both aromatic and alkene substrates bearing a chelating group reacted smoothly with a wide range of β-trifluoromethyl-α,β-unsaturated ketones to deliver the respective products in an atom-economical manner with decent yields.

### Scheme 281. Cp\*Rh(III)-Catalyzed C(sp<sup>2</sup>)-H Michael Addition to β-Trifluoromethyl-α,β-Unsaturated Ketones




Furthermore, Wang's group elaborated the synthesis of β-alkenyl carbonyls through a Cp\*Rh(III)-catalyzed oxidative coupling of 2-vinylpyridines with readily available allylic alcohols (Scheme 282).<sup>443</sup> A series of allylic alcohols coupled effectively to afford the corresponding products in modest to good yields. Markedly, allylic alcohols reacted specifically at the γ-position while alkenylpyridines reacted at the cis-β-position in this case.

### Scheme 282. Rhodium(III)-Catalyzed C–H Oxidative Coupling of 2-Vinylpyridines with Allylic Alcohols




α,β-Unsaturated nitriles are one of the most common scaffolds embedded in pharmaceutically relevant molecules and bioactive natural products. Anbarasan's group in 2015 established an efficacious and regioselective Cp\*Rh(III)-catalyzed alkenyl C–H cyanation by exploiting environmentally friendly N-cyano-N-phenyl-p-methylbenzenesulfonamide (NCTS) as the cyanating agent (Scheme 283).<sup>444</sup> This strategy exhibited high functional group compatibility and enabled the construction of a diverse variety of substituted acrylonitriles in decent yields (41–88%). As a particular highlight, the authors showcased the practicality of this strategy by the synthesis of a chlorpheniramine-based antagonist.

### Scheme 283. Rhodium(III)-Catalyzed Vinylic C(sp<sup>2</sup>)-H Cyanation of Alkenes



Quite recently, Wang's group initiated their investigations to exploit the ruthenium(II)-catalyzed oxidative coupling reaction of 2-vinylpyridines with triisopropylsilylacetylene. In this report, silver oxide (Ag<sub>2</sub>O) was proven to be indispensable for achieving high efficiency. Gratifyingly, both 2-pyridyl and 2-pyrimidyl were competent directing groups for this strategy, enabling the synthesis of highly functionalized 3-enyne products in good yields with excellent Z-selectivity (Scheme 284).<sup>445</sup>


### Scheme 284. Ru-Catalyzed Cross-Coupling Reaction of Alkenes with Triisopropylsilylacetylene



## 5. ALKENYL C–H BOND FUNCTIONALIZATION OF ALKENES CONTAINING A HETEROATOM

In this section of the review, we will summarise the work related to alkenyl  $sp^2$  C–H bond of diverse 2-heteroatom substituted alkenes (Scheme 285). In most of these cases, there can serve as enolate equivalents or latent protecting groups for the corresponding atom. The use of photoredox and electrochemical strategies to carry out C–H bond functionalization of alkenes bearing a 2-heteroatom substituted auxiliary have recently gained ever-increasing popularity. Metal-free systems have also emerged in recent years which may provide greener approaches for C–H functionalization of alkenes and their derivatives. Moreover, the use temporary chiral amines to in situ generate enamides, enamines, and their equivalents provides novel opportunities for the asymmetric C–H bond functionalizations with coupling partners containing pro-chiral centers.

### Scheme 285. General Scheme of Alkenyl C–H Bond Functionalizations of Alkenes Containing a Heteroatom Directing Group




## 5.1 Enamides

Enamides are extremely important synthetic building blocks since they are stable enolate equivalents and can be used to couple with a wide variety of electrophiles.<sup>446</sup> Furthermore, they can be easily converted into amines including chiral amines via asymmetric hydrogenation. As a consequence, there have been many methods developed for the preparation of this important class of compounds. Among these established methods, the selective alkenyl C–H bond functionalization of simple and easily accessible enamides provides one of the most direct and practical methods to access highly functionalized enamides.<sup>447,448</sup> In this part of the review, we summarized the comprehensive advances on the alkenyl C–H bond functionalizations of enamides. Metal-catalysis, photoredox as well as electrochemical strategies will be included.


**5.1.1. Arylation.** The first  $\beta$ -C( $sp^2$ )-H arylation of enamides was investigated by Loh and co-workers in 2009. They reported an unprecedented arylation of cyclic enamides with various arylboronic acids through a C–H activation reaction enabled by  $Pd(OAc)_2$  catalyst (Scheme 286).<sup>449</sup> A diverse number of arylboronic acids bearing both electron-donating and electron-withdrawing groups undergo a smooth cross-coupling to yield the arylated products in modest to good yields, while electron-withdrawing groups, such as  $CO_2R$ , COR, CN,  $NO_2$ , etc. on the cyclic enamides were not scrutinized. Due to excessive oxidation, *N*-(2-arylnaphthalen-1-yl)acetamides were also observed as byproducts when X was methylene.

### Scheme 286. Pd(II)-Catalyzed Vinylic C–H Arylation of Cyclic Enamides




Subsequently, the same group extended to establish a palladium(II)-catalyzed C–H arylation of cyclic enamides by using environmentally friendly arylsilanes as the coupling reagents, producing the corresponding 2-arylated enamides in 19–95% yield (Scheme 287a).<sup>450</sup> In this silicon-based strategy, AgF was employed as both an oxidant and a de-silyl reagent. However, this strategy was found to be incompatible with acyclic enamides. To address this issue, Luo and co-workers very recently successfully developed an efficient Cp\*Rh(III) catalysis system to realize the highly stereoselective direct  $\beta$ -C(sp<sup>2</sup>)-H bond arylation of acyclic enamides with electronically and sterically (hetero)arylsilanes by the judicious utilization of copper(II) fluoride as the efficacious silane activator and catalyst reoxidant (Scheme 287b).<sup>451</sup>

### Scheme 287. Vinylic C–H Arylation of Enamides with Arylsilanes




During the course of their investigations, Loh and co-workers also achieved a novel Pd(II)-catalyzed C(sp<sup>2</sup>)-C(sp<sup>2</sup>) dehydrogenative cross-coupling reaction for the  $\beta$ -C–H arylation of enamides, by employing simple arenes in the reaction (Scheme 288a).<sup>452</sup> This process delivered the corresponding enamides with excellent *Z*-selectivity. Notably, this protocol accommodates a wide diversity of functional groups, such as F, Cl, OMe, CF<sub>3</sub>, COMe and CO<sub>2</sub>Me. Electron-rich arenes (such as toluene and diphenyl ether) and halobenzenes produced the corresponding *para*-, *meta*- and *ortho*- regioisomers. Highly electron-deficient arenes such as methyl benzoate and benzotrifluoride predominantly afforded *meta*-isomers, while the *ortho*-isomers were not obtained in the reaction. In addition, a double arylation done in sequence via the monoarylated product was also investigated. When employing trifluoroacetic acid in replacement of acetic acid, the expected tetrasubstituted enamides were obtained in decent yields. Later in 2016, Piersanti and co-workers expanded Loh's strategy, and reported an efficient Pd(II)-catalyzed cross-dehydrogenative coupling between methyl *N*-phthaloyl dehydroalanine esters and diverse simple arenes (Scheme 288b).<sup>453</sup>

### Scheme 288. Pd(II)-Catalyzed Alkenyl C–H Arylation of Enamides with Simple Arenes



Diaryliodonium salts were identified as practical aryl sources in the copper(II)-catalyzed olefinic C–H arylation of cyclic enamides (Scheme 289a).<sup>454</sup> This process exhibited excellent functional group tolerance. Various sterically hindered diaryliodonium salts, however, gave lower reactivity, and the expected products were not obtained. One year later, Kesavan's group also reported an efficient Cu(I)-catalyzed C–H arylation of cyclic enamides with diaryliodonium salts. Of note, this reaction was conducted at room temperature without any base or additive, featuring a wide substrate scope with good to excellent yields (Scheme 289b).<sup>455</sup>



### Scheme 289. Copper-Catalyzed Vinylic C–H Arylation of Enamides with Diaryliodonium Salts



In 2017, Zhang and co-workers elaborated a general approach for the synthesis of multisubstituted oxazoles by the tandem oxidative cyclization of acyclic enamides with aryl iodides (Scheme 290).<sup>456</sup> The reaction occurred via the amide-directed vinylic C–H arylation followed by silver-

mediated oxidative cyclization sequence, thus allowing the rapid assembly of 2,4,5-trisubstituted oxazoles from readily available starting materials. As a particular highlight, the authors showcased the potential of this protocol by the straightforward construction of the core skeleton of a non-steroidal anti-inflammatory drug aristoxazole.


### Scheme 290. Synthesis of Substituted Oxazoles via Pd-Catalyzed Cascade Oxidative Cyclization of Enamides




Apart from  $\beta$ -selective alkenyl C(sp<sup>2</sup>)-H arylation of enamides, Park's group in 2011 successfully developed a palladium(II)-catalyzed  $\alpha$ -C-H arylation of *N*-substituted enamides with arylboronic acids under oxidative Heck cross-coupling conditions, enabling the stereoselective synthesis of  $\beta$ -substituted  $\beta$ -amidoacrylate derivatives in moderate to high yields (Scheme 291a).<sup>457</sup> Subsequently, Qin and co-workers expanded this strategy to report an efficient  $\alpha$ -C(sp<sup>2</sup>)-H arylation of *N*-substituted enamides with aryl iodides under a PdCl<sub>2</sub>(COD)/Ag<sub>3</sub>PO<sub>4</sub> catalytic system. The scope of both enamides and aryl iodides were found to be broad, and not sensitive to electronic and steric factors (Scheme 291b).<sup>458</sup>

### Scheme 291. Pd(II)-Catalyzed Vinylic $\alpha$ -C-H Arylation of Enamides

a) Park et al. 2011




b) Qin et al. 2013





Moreover, Loh's group continued their alkenyl C-H activation strategy onto enamides, and described the intramolecular palladium(II)-catalyzed C-H arylation of *N*-protected enamides via a 6-*endo* Heck cyclization process, providing a robust access to substituted 1,2-dihydroisoquinoline derivatives (Scheme 292).<sup>459</sup> By the combination of Pd(OAc)<sub>2</sub> catalyst, PPh<sub>3</sub> ligand and Cs<sub>2</sub>CO<sub>3</sub> base in DMF at 120 °C, a scope of 17 examples of this reaction was documented in reasonable yields of 35–84%. Notably, various nitrogen-containing products such as 1,1'-disubstituted ethylenes, 2-azabicyclo[3,3,0]octadienes as well as 5/5/6-membered pyrroloisoindolone derivatives can be obtained through a rare  $\beta$ -N-Pd elimination process by fine-tuning the phosphine ligands and bases.

### Scheme 292. Palladium(II)-Catalyzed Intramolecular Vinylic C-H Arylation of Enamides



Afterward, the group of Tong and Wang elaborated an efficient and straightforward synthesis of 2,3-dihydro-1*H*-benzo[*d*]azepine, a seven-membered heterocyclic core embedded in 7,8-dihydro-5*H*-benzo[4,5]-azepino[2,1-*a*]isoindol-5-one, via the Pd(II)-catalyzed intramolecular cross-dehydrogenative coupling of both aryl and vinyl C-H bonds (Scheme 293).<sup>460</sup> The authors showcased the practicality of this protocol by the total synthesis of aporhoadane alkaloids including palmanine, chilenammine, and lennoxamine by using this strategy as the key step.


### Scheme 293. Synthesis of *N*-Heterocyclic Scaffolds via Pd(II)-Catalyzed Dehydrogenative Cross-Coupling



**5.1.2. Alkylation.** Kobayashi's group in 2004 discovered that enamides could be employed as nucleophiles to couple with a series of electrophiles in the presence of strong Lewis or Brønsted acids, and the cross-coupling products of enamides were eventually obtained.<sup>461-463</sup> Inspired by this work, Tsogoeva's group investigated a chiral Brønsted acid-catalyzed stereoselective reaction of enamides to form alkylated enamides bearing a chiral quaternary carbon center (Scheme 294a).<sup>464</sup> Later in 2013, Guan and co-workers reported the Fe(III)-catalyzed self-condensation of various enamides, putting forward a novel and practical approach for the assembly of nitrogen-containing quaternary carbon centers (Scheme 294b).<sup>465</sup> Of note, the reaction was typically performed at room temperature, and a scope of 17 examples of this transformation was presented with acceptable yields.

### Scheme 294. Self-Condensation of Enamides for the Synthesis of Enamido-Substituted Nitrogen-Containing Quaternary Carbon Centers

a) Tsogoeva et al., 2008




b) Guan et al., 2013



Transition metal-catalyzed alkenyl C(sp<sup>2</sup>)-H alkylation through an enamide-directing group is an effective strategy for producing the alkylated enamides. In 2017, Zhang's research group reported a Cp\*Co(III)-catalyzed olefinic C(sp<sup>2</sup>)-H alkylation of enamides with maleimides. This reaction could be accomplished with the allylated Z-



enamides formed exclusively (Scheme 295).<sup>466</sup> Notably, allyl acetates were also viable coupling partners under the conditions to afford the allylation products.

### Scheme 295. Cobalt(III)-Catalyzed Cross-Coupling of Enamides with Maleimides/Allyl Acetates





Recently, tremendous efforts have been directed towards the radical-type C-H alkylation of enamides. For example, Loh's group in 2016 investigated the alkenyl C-H alkylation of enamides with  $\alpha$ -bromo carbonyl compounds through a radical pathway. Under the optimal reaction conditions, a handful of enamides coupled with  $\alpha$ -bromo carbonyls to deliver the expected alkylation products in high yields (Scheme 296).<sup>467</sup>

### Scheme 296. Cross-Coupling of Enamides with Sterically Hindered $\alpha$ -Bromocarbonyls



1,3-Dicarbonyl compounds could also be used as alkyl radicals to couple with enamides. In 2004, an efficient Mn(III)-mediated oxidative CDC reaction between acyclic enamides and 1,3-dicarbonyl compounds was reported by Li's group. In the presence of Mn(OAc)<sub>2</sub>·2H<sub>2</sub>O, a broad scope of 31 examples was presented in reasonable yields of 30-88%. (Scheme 297).<sup>468</sup>


### Scheme 297. Direct Oxidative Coupling of Enamides and 1,3-Dicarbonyl Compounds




Moreover, ether compounds have also been exploited as alkyl radical precursors in the presence of excess amounts of oxidants. Specifically, Ding et al. in 2019 accomplished a Cu-catalyzed oxidative cross-dehydrogenative coupling (CDC) reaction of cyclic enamides with ethers, yielding the desired alkylated enamides in good yields (Scheme 298a).<sup>469</sup> Quite recently, Zhao and co-workers also achieved an analogous transformation under photoredox catalysis. In this report, non-expensive and non-poisonous Eosin Y was employed as the photocatalyst. A broad range of ethers, thioethers or even simple alkanes were all competent coupling partners in this metal-free protocol (Scheme 298b).<sup>470</sup>

### Scheme 298. Alkenyl C–H Alkylation of Enamides with Ethers or Alkanes

a) Ding et al., 2019





b) Zhao et al., 2022





It is established that readily available alkyl carboxylic acids are extensively exploited as C(sp<sup>3</sup>)-radical precursors via a single-electron transfer (SET) and decarboxylation pathway. In continuation with their investigations of β-C(sp<sup>2</sup>)-H functionalization of enamides, Loh and co-workers successfully reported a direct Ag(I)-catalyzed decarboxylative β-C(sp<sup>2</sup>)-H alkylation of enamides with commercially available alkyl carboxylic acids as alkylating reagents (Scheme 299).<sup>471</sup> By the combination of Ag<sub>2</sub>CO<sub>3</sub> catalyst, K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> oxidant and NaHCO<sub>3</sub> base in MeCN/H<sub>2</sub>O at 50 °C, a broad array of enamides bearing various synthetically useful functional groups alkylated in reasonable yields of 30–78%.

### Scheme 299. Ag(I)-Catalyzed Decarboxylative Cross-Coupling of Enamides with Unactivated Aliphatic Carboxylic Acids




Moreover, Loh's group also developed a visible-light-promoted, *fac*-Ir(ppy)<sub>3</sub>-catalyzed decarboxylative C–H alkylation of enamides with activated alkyl carboxylic acids, alkyl *N*-hydroxyphthalimide esters (NHPI-esters). Under the optimized conditions, a broad diversity of enamides could be smoothly alkylated in this process, giving rise to the functionalized enamides in 53–91% yields. Notably, *N*-hydroxyphthalimide esters bearing primary, secondary, and even tertiary alkyl groups were all found to be viable substrates in this case (Scheme 300).<sup>472</sup> Quite recently, Chen and co-workers expanded to realize a similar visible-light-promoted decarboxylative alkylation under simple transition metal- and photocatalyst-free conditions.<sup>473</sup>

### Scheme 300. Photoredox-Catalyzed C–H Alkylation of Enamides with *N*-Hydroxyphthalimide Esters




In 2019, Xiao's research group expanded their visible-light photoredox catalysis,<sup>474</sup> and reported an example of Ir(4-Fppy)<sub>2</sub>(bpy)PF<sub>6</sub>-catalyzed deaminative approach for the alkenyl C–H alkylation of enamide by making use of easily prepared Katritzky salts as the alkylating reagent, affording an alkylated enamide in 40% yield (Scheme 301).<sup>475</sup>

### Scheme 301. Photoredox-Catalyzed Deaminative C–H Alkylation of Enamides with Katritzky salts




The use of rare and expensive iridium photocatalyst undoubtedly raises remarkable concerns about the sustainability and economics of this protocol. To this end, Fu's group in 2021 expanded this visible-light-induced deaminative alkylation of enamides with a large variety of Katritzky's *N*-alkylpyridinium salts of amino acids in the presence of 10 mol% LiI under purple LED irradiation conditions (Scheme 302).<sup>476</sup> Gratifyingly, this transition metal-free strategy is also compatible with redox-active NHPI-esters to generate the decarboxylative alkylation products by using a combination of NaI (10 mol%) and PPh<sub>3</sub> (10 mol%) in acetone. Both primary, secondary, and even tertiary unactivated alkyl carboxylic acids could participate in this stereoselective alkylation with modest to high yields (38–82%).

### Scheme 302. Visible-Light-Induced Iodine-Anion-Catalyzed Decarboxylative/Deaminative Vinylic C–H Alkylation of Enamides




In 2020, Gillaizeau's group successfully applied xanthate-based radical chemistry to describe an efficient olefinic  $\beta$ -C–H alkylation of nonaromatic enamides with readily biodegradable lauroyl peroxide (DLP) as the radical initiator (Scheme 303).<sup>477</sup> This approach features a broad substrate scope and excellent functional group tolerance, thus leading to the expedient synthesis of a broad array of synthetically appealing  $\gamma$ -amino- $\beta,\gamma$ -unsaturated acyl scaffolds in satisfactory yields.

### Scheme 303. Vinylic C(sp<sup>2</sup>)-H Alkylation of Enamides Using Xanthate Chemistry




Around the same time, Hu and co-workers employed 1,2-oxazetidines as the formalimine precursors to participate in a novel Ru(II)-catalyzed regioselective alkenyl C–H aminomethylation of enamides through a ring-opening strategy (Scheme 304).<sup>478</sup> Notably, the corresponding aminomethylated enamides could be readily converted into value-added nitrogen-containing scaffolds, which remarkably renders this protocol more synthetically useful for the synthesis of highly functionalized enamides.

### Scheme 304. Ru-Catalyzed Aminomethylation of Enamides with 1,2-Oxazetidines



Apart from the above-mentioned examples using a pre-installed directing group for the vinylic C–H bond functionalizations, the use of transient directing groups allows for a transient ligand to be used, potentially in catalytic quantities, thus without the need for pre-installation or subsequent removal steps.<sup>479,480</sup> In this regard, Dong's group in 2017 elegantly devised a novel *in situ* installed enamide-directing strategy for the synthesis of  $\alpha$ -alkylated ketones enabled by a cationic iridium catalyst (Scheme 305),<sup>481</sup> affording the branched-selective products which are extremely valuable building blocks but are nontrivial to obtain by conventional methods.<sup>482</sup>


### Scheme 305. Branched-Selective Ketone $\alpha$ -Alkylation via an Enamide Directing Strategy



**5.1.3. Fluoroalkylation.** Fluoroalkyl compounds have received elevated attention in the last ten years because of their peculiar biological activities. In this context, the alkenyl C–H fluoroalkylation of enamides has been extensively investigated as an appealing strategy to obtain structurally diverse fluoroalkylated enamides.<sup>483</sup>


In particular, Togni's reagent has been widely utilized as the trifluoromethylation reagent in numerous C–H bond functionalization reactions.<sup>484</sup> In 2012, Loh's group disclosed an efficient and straightforward  $\beta$ -C(sp<sup>2</sup>)-H trifluoromethylation of enamides catalyzed by a Cu(I) species (Scheme 306).<sup>485</sup> A large number of enamides with different substitution patterns were subjected to the optimized conditions, yielding the trifluoromethylated enamides in modest to good yields.

### Scheme 306. Copper-Catalyzed Olefinic C–H Trifluoromethylation of Enamides




Contrastingly, radical oxidative cross-coupling could also offer a powerful strategy for the olefinic  $\beta$ -C–H trifluoromethylation of enamides. In this regard, Gillaizeau and co-workers in 2015 elaborated an efficient Fe(II)-catalyzed direct C<sub>3</sub>-trifluoromethylation of enamides with Togni's reagent through a radical pathway. The desired trifluoromethylated products were obtained in 31–95% yields (Scheme 307).<sup>486</sup> By decreasing the reaction time to 2 hours, it was established that the oxotrifluoromethylated byproduct was detected in 55% yield. However, the deprotonation product was not formed when using methanol as the solvent instead.

### Scheme 307. Fe-Catalyzed C–H Trifluoromethylation of Enamides




In 2017, Selander's group utilized inexpensive and air-stable Langlois' reagent (CF<sub>3</sub>SO<sub>2</sub>Na) as the radical fluoroalkyl source for the redox-economical synthesis of  $\beta$ -trifluoromethylated enamides via a Cu(II)-catalyzed reaction with oxime acetates (Scheme 308).<sup>487</sup>

### Scheme 308. Synthesis of Trifluoromethylated Enamides with Langlois' Reagent



Very recently, the group of Chen and Yang illustrated a convenient visible-light-promoted approach for the vinylic  $\beta$ -C–H trifluoromethylation of enamides with Langlois' reagent as the CF<sub>3</sub> radical source (Scheme 309).<sup>488</sup> An array of trifluoromethylated enamides were produced in modest to high yields with good *E*-selectivity under typically mild conditions.

### Scheme 309. Visible-Light-Promoted Olefinic C–H Trifluoromethylation of Enamides with Langlois' Reagent




The direct C–H difluoromethylation of enamides could also be realized under photoredox catalysis. In 2019, Loh and co-workers disclosed a direct Ir(ppy)<sub>3</sub>-catalyzed  $\beta$ -C(sp<sup>2</sup>)-H difluoromethylation of both cyclic and acyclic enamides by making use of a difluoromethylating reagent [PPh<sub>3</sub>CF<sub>2</sub>H]Br promoted by visible-light (Scheme 310a).<sup>489</sup>


A large variety of  $\beta$ -difluoromethylated enamides bearing various synthetically useful functional groups were obtained in moderate to excellent yields (35–91%). Following this, Wu's group later employed easily accessible hypervalent iodine(III) [bis(difluoroacetoxy)iodo]benzene reagents to engage this photoredox-catalyzed process to generate a broad range of (*E*)- $\beta$ -difluoromethylated enamides (Scheme 310b).<sup>490</sup>

### Scheme 310. Difluoromethylation of Enamides via Photoredox Catalysis

a) Loh et al., 2019




b) Wu et al., 2021




Loh and co-workers further achieved a photoredox-catalyzed  $\beta$ -C(sp<sup>2</sup>)-H trifluoroethylation of enamides with commercially available 2,2,2-trifluoroethyl iodide, thus allowing a robust and efficacious synthetic route to various pharmaceutically pivotal  $\beta$ -trifluoroethylated enamides with excellent regio- and stereoselectivities (Scheme 311a).<sup>491</sup> Subsequently, the same group further described a metal-free visible-light-induced stereoselective alkenyl C(sp<sup>2</sup>)-H perfluoroalkylation of structurally diverse enamides with perfluoroalkyl iodides in the presence of organic dye eosin Y as the photocatalyst, giving rise to a sustainable and environmentally benign approach to the synthesis of perfluoroalkyl-containing enamides (Scheme 311b).<sup>492</sup>

### Scheme 311. Visible-Light-Induced Vinylic C-H Trifluoroethylation and Perfluoroalkylation of Enamides

a) Loh et al., 2020




b) Loh et al., 2021




Commercially available fluoroalkyl halides have been extensively employed as fluoroalkyl radical sources under transition metal-catalyzed reaction conditions. The copper-based catalytic system has been established as an effective approach for the  $\beta$ -C(sp<sup>2</sup>)-H functionalization of enamides. In the very first report of its kind, Gillaizeau's group disclosed a Cu(I)-catalyzed alkenyl C-H fluoroalkylation of enamides using  $\text{BrCF}_2\text{CO}_2\text{Et}$  (Scheme 312).<sup>493</sup> On the basis of mechanistic studies, it was established that the fluorinated enamide was obtained in a comparable yield when one equivalent of TEMPO was added. More recently, the same group further achieved a similar (fluoro)alkylation under iron catalysis.<sup>494</sup>

### Scheme 312. Copper-Catalyzed Olefinic C-H Difluoroacetylation of Enamides




In 2017, an easily accessible Cu(I)-catalyzed  $\beta$ -C(sp<sup>2</sup>)-H fluoroalkylation of enamides with fluoroalkyl halides via a radical addition/oxidation/deprotonation sequence was reported by Wang's group (Scheme 313).<sup>495</sup>

### Scheme 313. Copper-Catalyzed Vinylic C-H Difluoroalkylations and Perfluoroalkylations of Alkenes




The palladium(II)-catalyzed Heck-type fluoroalkylation of enamides and simple alkenes with fluoroalkyl halides was investigated by Zhang's group (Scheme 314).<sup>496,497</sup> In majority of the examples, the  $\beta$ -fluoroalkylated derivatives were obtained in excellent yields.

### Scheme 314. Palladium-Catalyzed Heck-Type C-H Difluoroalkylation of Alkenes




Moreover,  $\beta$ -C(sp<sup>2</sup>)-H fluoroalkylation of enamides promoted by visible-light via photocatalysis was also established. In 2012, Yu's group reported a radical C-H alkylation of enamides promoted by visible-light by taking advantage of [Ir(ppy)<sub>2</sub>(dtbbpy)PF<sub>6</sub>] as a photocatalyst. In this approach, electron-withdrawing groups bearing halides were utilized as the alkylating radical agents to generate the desired products in decent yields (Scheme 315).<sup>498</sup> Remarkably, CF<sub>3</sub>SO<sub>2</sub>Cl was also a competent trifluoromethyl radical source, affording the desired trifluoromethylated enamide in 81% yield.

### Scheme 315. Direct Vinylic C-H Functionalization of Enamides Using Visible-Light Photoredox Catalysis



In 2016, Yu's group extended to investigate the vinylic  $\beta$ -C-H trifluoromethylation of enamides promoted by visible-light in the absence of any photocatalyst. By making use of Umemoto's reagent as the trifluoromethylating reagent, the desired trifluoromethylated enamides were produced in modest to good yields (Scheme 316).<sup>499</sup>


### Scheme 316. Visible-Light-Promoted and Photocatalyst-Free Trifluoromethylation of Enamides



**5.1.4. Olefination.** The direct alkenyl  $\beta$ -C(sp<sup>2</sup>)-H olefination of enamides can be readily achieved through a vinylic C-H activation strategy. In 2011, Loh's group reported the first example of Pd(II)-catalyzed oxidative


cross-coupling reaction of structurally diverse enamides with acylates by using molecular oxygen as the terminal oxidant. The X-ray structure of the cyclic vinylpalladium(II) complex was obtained, suggesting that C-H activation was most probably involved in the mechanism (Scheme 317).<sup>500</sup>

### Scheme 317. Synthesis and Characterization of a Cyclic Vinylpalladium(II) Complex




Later, Gillaizeau's group devised a mild and effective strategy for the  $\beta$ -C-H olefination of non-aromatic enamides via a Pd(II)-catalyzed alkenyl C-H activation process (Scheme 318).<sup>501</sup> This work exhibited a wide substrate scope of both non-aromatic cyclic enamides and electronically activated alkenes.

### Scheme 318. Pd(II)-Catalyzed Direct C-H Olefination of Nonaromatic Enamides





Rhodium complex was also proved to be an efficient catalyst for the  $\beta$ -C-H alkenylation of enamides. Glorius and co-workers used [RhCp\*Cl<sub>2</sub>]<sub>2</sub> and AgSbF<sub>6</sub> along with Cu(OAc)<sub>2</sub> as the oxidant. They demonstrated a Cp\*Rh(III)-catalytic system for the alkenylation of enamides with terminal olefins (Scheme 319).<sup>502</sup> However, simple enamides were found to be inappropriate for this process. Ester-substituted enamides presented an appropriate electronic disposition, which underwent a smooth transformation with olefins to afford the Z-isomers as major products.

### Scheme 319. Cp\*Rh(III)-Catalyzed Oxidative Vinylic C-H Olefination of Enamides





In 2018, Dong's group presented a  $\text{Cp}^*\text{Rh(III)}$ -catalyzed amide directing group-assisted C–H alkenylation between oxindoles and activated olefins, thus allowing an atom-economical and straightforward strategy to construct valuable and multipurpose functionalized *N*-(2*E*,4*Z*)-butadiene substituted oxindoles (Scheme 320).<sup>503</sup> Oxindoles functioned well as a directing-group was utilized extensively in this protocol.

### Scheme 320. $\text{Cp}^*\text{Rh(III)}$ -Catalyzed Alkenylation of Oxindoles and Olefins to Synthesize *N*-(2*E*,4*Z*)-Butadiene Substituted Oxindoles





**5.1.5. Alkynylation.** In a similar fashion, nitrogen-containing conjugated enynes could be readily acquired by the  $\beta\text{-C}(\text{sp}^2)\text{-H}$  alkylation of enamides. Since the early 2010s, Loh's group reported an effective  $\text{Cp}^*\text{Rh(III)}$ -catalyzed olefinic  $\beta\text{-C-H}$  alkylation of enamides by using an alkynylidonium(III) reagent (TIPS-EBS) for the stereospecific configuration of synthetically practical *Z*-enynes under extremely benign conditions (Scheme 321).<sup>504</sup> This catalytic system portrayed an excellent tolerance of various functional groups. The process was well tolerated with aryl-substituted enamides bearing electron-rich groups (Me, OMe, Ph, *N*-piperidyl), electron-deficient groups ( $\text{CF}_3$ , CN,  $\text{SO}_2\text{Me}$ , COMe) as well as halogen groups (F, Cl, Br). In addition, both naphthyl-substituted and heterocyclic enamides reacted smoothly in this transformation. Remarkably, non-aromatic enamides also gave the alkynylated products in excellent yields, and ester-substituted enamides reacted smoothly in this process.

### Scheme 321. Rhodium(III)-Catalyzed Olefinic C–H Alkynylation of Enamides





On the basis of the six-membered palladacycle complex, Loh and co-workers further established an efficient  $\text{Pd(II)}$ -catalyzed C–H alkylation of cyclic enamides on the  $\text{C}_3$ -position with readily available alkynyl bromides (Scheme 322).<sup>505</sup> Of onte, DMSO was necessarily added as a ligand to stabilize the transition state in this process.

### Scheme 322. Palladium-Catalyzed C–H Alkynylation of Enamides with Alkynyl Bromides



A couple of years later, Beng and co-workers described an atom-economical  $\text{Cp}^*\text{Ir(III)}$ -catalyzed dehydrogenative vinylic  $\beta\text{-C}(\text{sp}^2)\text{-H}$  alkylation of cyclic non-aromatic eneformamides with diverse terminal alkynes (Scheme 323).<sup>506</sup> The versatility of the corresponding  $\alpha$ -alkynyl eneformamides renders this protocol synthetically appealing for the synthesis of azapolycyclic architectures and relevant saturated cyclic amines.




### Scheme 323. Iridium-Catalyzed $\alpha\text{-C-H}$ Alkynylation of Cyclic Nonaromatic Eneformamides



**5.1.6. Carbonylation.** Carbonyl compounds are crucial intermediates in synthetic organic chemistry and a large number of robust strategies for the incorporation of this functional group into synthetically useful organic molecules have been reported over the decades.<sup>507</sup> Lately, numerous investigations have been focused on the alkenyl C–H carbonylation of enamides catalyzed by transition metals, and a handful of acylating agents are utilized for the incorporation of carbonyl groups.



Direct carbonylation reactions with CO have been and will continue to be a hot research topic in the fertile field of C–H functionalization reactions. In this regard, Guan and co-workers in 2013 reported a Pd(II)-catalyzed vinylic C–H carbonylation of acyclic enamides with atmospheric pressure of CO in the presence of KI and Ac<sub>2</sub>O as additives (Scheme 324a).<sup>508</sup> Afterward, Lei's group also achieved this oxidative alkenyl C–H carbonylation by combination of photoredox and palladium catalysis in conjunction with molecular oxygen as the terminal oxidant, which remarkably obviated the use of stoichiometric amounts of Cu(OAc)<sub>2</sub>, thus providing an environmentally friendly approach for the rapid synthesis of 1,3-oxazin-6-ones in 44–83% yields (Scheme 324b).<sup>509</sup>

**Scheme 324. Oxidative C–H Carbonylation of Enamides with Carbon Monoxide**





In 2011, Bergman, Ellman, and co-workers investigated a vinylic C–H activation strategy through a Cp\*Rh(III)-catalyzed amidation of enamides with isocyanates. Under the standard conditions, the expected enaminone products were obtained in satisfactory yields (Scheme 325).<sup>510</sup>

**Scheme 325. Rh(III)-Catalyzed Vinyl C–H Amidation of Enamides with Isocyanates**





Later, Loh and colleagues were able to establish a practical and environmentally friendly approach for the straightforward construction of pyrimidin-4-ones through an efficient Ru(II)-catalyzed heteroannulation between enamides and isocyanates (Scheme 326).<sup>511</sup> This strategy proceeded with a diverse substrate scope and excellent functional group compatibility, only releasing water as an innocuous waste. Of note, the authors demonstrated the feasibility of this strategy by late-stage modification of biologically active natural products.

**Scheme 326. Synthesis of Pyrimidin-4-ones through a Ruthenium-Catalyzed Heteroannulation between Enamides and Isocyanates**





The group of Guo and Duan illustrated a Pd(II)-catalyzed alkenyl C–H functionalization approach via a decarboxylative acylation of cyclic enamides with various  $\alpha$ -oxocarboxylic acids as the efficacious acylating agents under typically mild conditions, allowing a practical and efficient method for the preparation of acylated enamides from readily available starting materials. (Scheme 327).<sup>512</sup>

### Scheme 327. Palladium-Catalyzed Decarboxylative Acylation of Enamides with $\alpha$ -Oxocarboxylic Acids





Liang and co-workers achieved a palladium(II)-catalyzed chelation-assisted  $\beta$ -C(sp<sup>2</sup>)-H carboxamidation of acyclic enamides through the incorporation of isocyanides. This strategy portrayed excellent functional group compatibility, yielding the corresponding enamino products in modest yields. (Scheme 328).<sup>513</sup>

### Scheme 328. Synthesis of *N*-Acyl Enamine Amides via Pd(II)-Catalyzed C–H Carboxamidation of Enamides with Isocyanides





Moreover, the sustainable iron-catalyzed alkenyl  $\beta$ -C–H carbonylation of enamides was also investigated through a carbonyl radical pathway. In 2014, Loh's group reported the C–H alkoxycarbonylation of acyclic enamides with carbazates enabled by an iron(II)phthalocyanine [ $\text{Fe}(\text{Pc})$ ] catalyst in conjunction with *tert*-butyl hydroperoxide (TBHP) as an oxidant, generating the enamino products in modest yields (Scheme 329).<sup>514</sup>

### Scheme 329. Iron-Catalyzed C–H Alkoxycarbonylation of Enamides with Carbazates



Enlightened by this work, Loh and co-workers continued to investigate the Fe(II)-catalyzed C–H carbonylation of enamides, and reported the dehydrogenative cross-coupling reactions between acyclic enamides and formamides (Scheme 330).<sup>515</sup> This protocol gave rise to various carbamoylated enamides with complete *Z*-selectivities.


### Scheme 330. Iron-Catalyzed C–H Carbamoylation of Enamides with Formamides



Subsequently, the same group further reported an efficient dehydrogenative carbonylation of enamides with simple aldehydes enabled by an iron catalyst to deliver a


diverse series of synthetically valuable  $\beta$ -ketoenamides with excellent stereoselectivity and good functional group tolerance. Notably, this approach was applicable to late-stage diversification of biologically active natural products (Scheme 331a).<sup>516</sup> Meanwhile, Loh and co-workers also achieved this transformation through a sustainable and environmentally friendly transition-metal-free photoredox catalysis, affording the  $\beta$ -ketoenamides in 35–91% yields. (Scheme 331b).<sup>517</sup>

### Scheme 331. Synthesis of $\beta$ -ketoenamides via Olefinic C–H Carbonylation of Enamides with Aldehydes




**5.1.7. Sulfonylation.**  $\beta$ -Amido sulfones serve as crucial scaffolds widespread in a variety of pharmaceutically relevant compounds and bioactive natural products. The development of efficient strategies for the synthesis of  $\beta$ -amidovinyl sulfones is highly sought-after. Among the established methods, the direct alkenyl C–H sulfonylation of enamides offers a rapid route to acquire  $\beta$ -amidovinyl sulfones. In this regard, Loh's group in 2013 reported a Pd(II)-catalyzed alkenyl C–H sulfonylation of cyclic enamides with commercially available sulfonyl chlorides (Scheme 332).<sup>441</sup> A handful of structurally diverse arylsulfonyl chlorides were studied under the optimized conditions, and the expected coupling products were formed in modest yields.

### Scheme 332. Palladium-Catalyzed Alkenyl C–H Bond Sulfonylation of Enamides



By making use of  $\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$  as a photoredox catalyst, Yu's group in 2013 described a similar C–H sulfonylation of acyclic enamides promoted by visible-light through a radical mechanism. The process features a broad substrate scope of both enamides and sulfonyl chlorides, yielding a series of  $\beta$ -amidovinyl sulfones in 65–99% yields (Scheme 333).<sup>518</sup>


### Scheme 333. Photoredox-Catalyzed Alkenyl C–H Bond Sulfonylation of Enamides




In 2018, the oxidative cross-coupling synthesis of  $\beta$ -amino sulfones from sodium sulfinates and enamides through a metal-free, visible-light promoted approach was reported by Zhang's group (Scheme 334a).<sup>519</sup> This process utilized a cost-effective, readily available organic dye rose bengal (RB) as the photoredox catalyst in combination with nitrobenzene or atmospheric oxygen as the oxidant. The scope of enamides was found to be broad and not sensitive to electronic and steric factors. The desired sulfonylated enamides were obtained in high yields (68–92%). Subsequently, Manolikakes' group also realized the C–H sulfonylation of acyclic enamides with comparable efficiency through a manganese(III) acetate-mediated radical process (Scheme 334b).<sup>520</sup> More recently, Zeng and co-workers further achieved a sustainable electrochemical sulfonylation of acyclic enamides under metal- and oxidant-free conditions (Scheme 334c).<sup>521</sup>

### Scheme 334. Direct Alkenyl C–H Sulfonylation of Enamides with Sodium Sulfinates


a) Zhang et al., 2018



b) Manolikakes et al., 2019




c) Zeng et al., 2021




The direct incorporation of sulfur dioxide ( $\text{SO}_2$ ) has been used as an appealing approach for the construction of sulfonyl derivatives.<sup>522</sup> In 2019, Loh and colleagues elaborated a copper-mediated three-component coupling reaction between acyclic enamides, aryldiazonium tetrafluoroborates, and 1,4-diazabicyclo[2.2.2]octane bis(sulfur dioxide) (DABSO) for the synthesis of  $\beta$ -arylsulfonylated enamides (Scheme 335a).<sup>523</sup> In the same year, they also investigated this vinylic C–H arylsulfonylation of enamides under  $\text{Ir}(\text{ppy})_3$  catalysis in the absence of blue LEDs irradiation (Scheme 335b).<sup>524</sup> This process occurred under benign conditions, giving rise to the corresponding  $\beta$ -amidovinyl sulfones in up to 91% yield. More recently, Zhang and Lian further established a transition-metal- and base-free strategy for the arylsulfonylation of enamides by using a cheap and bench-stable  $\text{SO}_2$  surrogate (tetrabromothiophene  $S,S$ -dioxide, SOGen),<sup>525</sup> which could *ex situ* generate  $\text{SO}_2$  gas in a controlled and predictable manner (Scheme 335c).<sup>526</sup>

### Scheme 335. Alkenyl C–H Sulfonylation of Enamides with DABSO and SOGen


a) Loh et al., 2019



b) Loh et al., 2019




c) Zhang, Lian et al., 2021




**5.1.8. Phosphorylation.** Phosphorus is one of the most abundant elements in the earth's crust which plays an indispensable role in the growth of living organisms.<sup>527</sup> Accordingly, it is crucial to establish robust strategies for the synthesis of structurally diverse phosphorus-containing compounds (especially  $\beta$ -phosphorylated enamides) because of their widespread applications in synthetic chemistry. The groups of Zhang and Zou independently described the Mn(III)-mediated oxidative CDC reaction of acyclic enamides with *H*-phosphonates or *H*-phosphine oxides. Specifically, Zhang's group demonstrated the synthesis by means of  $\text{Mn}(\text{acac})_3$  as the oxidant in conjunction with toluene as the solvent without the use of any base in the reaction. The protocol exhibited a diverse scope of substrate with excellent *Z*-selectivity (Scheme 336a).<sup>528</sup> On the other hand, Zou's group discovered that  $\text{Mn}(\text{OAc})_3$  could provide comparable efficiency in the presence of  $\text{K}_2\text{CO}_3$  in methanol. The reaction was typically finished within 0.5 h at room temperature, affording  $\beta$ -phosphorylated enamides as *E/Z*-configurations mixtures with the *E*-isomers formed predominantly (Scheme 336b).<sup>529</sup> Quite recently, Xu et al. accomplished an analogous Mn(III)-promoted C–H phosphorylation of tertiary acyclic enamides with comparable efficiency (Scheme 336c).<sup>530</sup>

### Scheme 336. Manganese(III)-Mediated Olefinic C–H Phosphorylation of Enamides


a) Zhang, Xiong et al., 2018



b) Zou, Zeng et al., 2019




c) Xu et al., 2022




Apart from these radical-based processes, the group of Zhang and Ma in 2018 established an efficient Pd(II)-catalyzed C(sp<sup>2</sup>)-H functionalization protocol for the direct phosphorylation of enamides with high chemo- and stereoselectivity (Scheme 337).<sup>531</sup> An extraordinarily broad scope of 60 examples was documented in yield up to 96%. Both cyclic and acyclic enamides with different substitution patterns were well tolerated. Notably, this protocol was applicable to late-stage modification of bioactive natural products such as pregnenolone and prasterone derivatives.

### Scheme 337. Palladium(II)-Catalyzed Vinylic C-H Phosphorylation of Enamides




Later, Gillaizeau's group devised an alternative approach for the direct C-H phosphorylation of enamides by using diphenyliodonium salt combined with Et<sub>3</sub>N as the efficacious radical initiator under transition-metal-free conditions (Scheme 338).<sup>532</sup> Based on their mechanistic studies, the authors proposed a tentative catalytic mechanism involving a single-electron-transfer process initiated by the electron donor-acceptor (EDA) complex between Et<sub>3</sub>N and diphenyliodonium salts.

### Scheme 338. Diphenyliodonium Ion/Et<sub>3</sub>N Promoted C-H Phosphorylation of Enamides




**5.1.9. Annulation Reaction.** The outstanding step-economy of annulation reactions makes it highly sought-after for the assembly of diverse complex polycyclics in organic synthesis. Diazabicyclic olefins are versatile synthons for the synthesis of highly functionalized cyclopentenes with extensive structural diversity. In 2017, John and Radhakrishnan reported the Pd(II)-catalyzed oxidative annulation of acyclic enamides with strained diazabicyclics via tandem ring-opening and ring-closing sequences (Scheme 339),<sup>533</sup> thereafter enabling the stereoselective synthesis of multisubstituted cyclopentene fused 2-pyrrolines in moderate to high yields. By making use of [RhCp\*Cl<sub>2</sub>]<sub>2</sub> as the catalyst, a broad array of *trans*-disubstituted cyclopentenes could be obtained through a ring-opening desymmetrization process.

### Scheme 339. Synthesis of Cyclopentene Fused 2-Pyrrolines via Pd-Catalyzed Oxidative Annulation of Enamides with Diazabicyclics




More recently, the formal oxidative annulation between enamides and maleimides was established by Luo's group through an efficient manganese(II) acetate-promoted dehydrative dehydroaromatizing [4 + 2] cycloaddition, providing access a broad series of highly valuable pyrrolo[3,4-c]-pyridine derivatives (Scheme 340).<sup>534</sup> Various substituted maleimides were suitable coupling partners for this transformation. NH-free maleimide, however, was proven to be much less efficient (17%). Of note, alkylenamides failed to deliver the expected product in this case. Interestingly, the target product exhibited typical aggregation-induced emission (AIE) characteristics, which remarkably expand the potential application of these pyrrolo[3,4-c]-pyridine derivatives.

**Scheme 340. Mn(OAc)<sub>2</sub>-Promoted Formal [4 + 2] Cycloaddition of Enamides with Maleimides**



Pioneering work of Fagnou, Stuart, and their co-workers revealed an efficient pyrrole synthesis utilizing enamides with diverse internal alkynes under cationic Cp<sup>\*</sup>Rh(III) catalysis. In 2010, they expanded their rhodium(III) catalysis strategy to disclose the vinylic C–H bond functionalization of enamides for the expedient synthesis of multisubstituted pyrroles (Scheme 341).<sup>535</sup> Easily prepared [Cp<sup>\*</sup>Rh(MeCN)<sub>3</sub>][SbF<sub>6</sub>]<sub>2</sub> was readily employed as the efficacious catalyst to achieve this transformation. The reaction occurred smoothly in the presence of a catalytic amount of Cu(OAc)<sub>2</sub>·H<sub>2</sub>O (20 mol%) in conjunction with molecular oxygen as the terminal oxidant. Notably, both symmetrical and unsymmetrical internal alkynes were tolerated to deliver the corresponding pyrroles with excellent regioselectivities.


**Scheme 341. Rhodium(III)-Catalyzed Vinylic C–H Functionalization of Enamides with Alkynes**



Later in 2013, the groups of Wang<sup>536</sup> and Ackermann<sup>537</sup> independently disclosed the inexpensive ruthenium(II)-catalyzed oxidative annulation reaction of enamides with internal alkynes to construct substituted pyrroles via C–H/N–H bond functionalizations (Scheme 342). In Ackermann's report, they employed a catalytic amount of Cu(OAc)<sub>2</sub>·H<sub>2</sub>O (30 mol%) under an ambient atmosphere of air as the terminal oxidant. Interestingly, *N*-unsubstituted


pyrroles could be produced directly with the addition of AgSbF<sub>6</sub> and MeOH as illustrated in Wang's work.

**Scheme 342. Ruthenium-Catalyzed Pyrrole Synthesis via Oxidative Annulation of Enamides with Alkynes**



In 2014, Guan and colleagues elaborated the synthesis of *N*-substituted pyrroles enable by a palladium(II)-catalyzed annulative reaction of enamides with a diverse array of internal alkynes (Scheme 343a).<sup>538</sup> Around the same time, Loh's group also achieved an analogous pyrrole synthesis in synthetically useful yields using molecular oxygen as an oxidant for the palladium catalyst regeneration (Scheme 343b).<sup>539</sup> However, poor regioselectivities were observed with unsymmetrical internal alkynes as the annulation partners in this protocol.


**Scheme 343. Synthesis of Substituted Pyrroles via Pd(II)-Catalyzed Oxidative Annulation of Enamides with Internal Alkynes**



However, when employing electron-deficient internal alkynes as coupling partners, the annulation reaction of enamides could be carried out under copper-promoted


conditions, which undoubtedly providing a cost-effective strategy for the expeditious synthesis of NH-free pyrroles (Scheme 344).<sup>540</sup>

### Scheme 344. Synthesis of Substituted Pyrroles via Copper-Promoted Oxidative Annulation of Enamides and Alkynes



Pawar's group later illustrated an efficient earth-abundant, environmentally benign Cp\*Co(III)-catalyzed pyrrole synthesis from enamides and alkynes (Scheme 345a).<sup>541</sup> The reaction occurred smoothly at room temperature, resulting in the formation of *N*-acetyl pyrroles in high yields (60–88%). In this protocol, a broad range of *N*-unsubstituted pyrroles can also be formed at elevated temperature. In the same year, Zhang and co-workers also achieved this cobalt-catalyzed formal oxidative [3 + 2] annulation in the presence of CuO as an oxidant (Scheme 345b).<sup>542</sup>


### Scheme 345. Synthesis of Substituted Pyrroles via Cp\*Co(III)-Catalyzed [3 + 2] Annulation of Enamides and Alkynes



Apart from the formation of five-membered pyrrole derivatives,<sup>540</sup> Guan and co-workers demonstrated that the copper-catalyzed oxidative annulation of enamides with electron-deficient internal alkynes could also deliver


substituted pyridines (Scheme 346).<sup>543</sup> More interestingly, this reliable strategy can extend to couple with aryne precursors for the facile assembly of multisubstituted isoquinoline derivatives under metal-free reaction conditions.

### Scheme 346. Formal Dehydrative [4 + 2] Cycloaddition of Enamides with Electron-Deficient Internal Alkynes



In 2015, Wang's group devised an economic approach for the convenient and regioselective assembly of multisubstituted pyridines through the formal dehydrative [4 + 2] cycloaddition of acyclic enamides with a broad array of internal alkynes under ruthenium catalysis (Scheme 347).<sup>544</sup> Unsymmetric internal alkynes participated well in this cyclization reaction, exclusively generating a single regioisomer. When employing diynes as coupling partners, multisubstituted bipyridines could also be produced by this methodology under slightly modified conditions. Combined mechanistic studies and DFT calculations clearly demonstrated that this dehydrative cycloaddition reaction probably occurred through a concerted metalation deprotonation (CMD) process.


### Scheme 347. Ru-Catalyzed Formal Dehydrative [4 + 2] Cycloaddition of Enamides and Alkynes



**5.1.10. Other Useful Reactions.** In 2014, Sun's group devised a rapid route for the synthesis of  $\beta$ -sulfinyl


enamides via a direct alkenyl C–H sulfonylation of enamides with the sulfonylative agent (PhSOCl), which is conveniently generated *in situ* from PhSO<sub>2</sub>Na and MeSiCl<sub>3</sub> (Scheme 348).<sup>545</sup> The presence of DMAc served as a Lewis base in this process.

### Scheme 348. Direct Synthesis of *N*-Protected- $\beta$ -Sulfinylenamines via C-Sulfonylation of Enamides and Enecarbamates



The Ag(I)-mediated oxidative C–H sulfonylation of enamides with disulfides for the synthesis of  $\beta$ -sulfonylated enamides was illustrated by Yang, Deng, and their co-workers in 2014 (Scheme 349).<sup>546</sup> Gratifyingly, this approach was applicable to the C–H selenation of enamides with diaryl diselenide, albeit in a 28% yield.


### Scheme 349. Silver-Mediated C–H Sulfonylation of Enamides with Disulfides




Zhang's group investigated a Cp\*Rh(III) and Ag(I) co-catalyzed alkenyl C–H acetoxylation of enamides by utilizing Cu(OAc)<sub>2</sub>·H<sub>2</sub>O as both acetoxyating reagent and oxidant. The process was highly stereoselective, exclusively giving rise to *Z*-configuration products in modest yields (Scheme 350a).<sup>547</sup> Later, Xu and colleagues also achieved an efficient alkenyl C–H acyloxylation of enamides with simple carboxylic acids enabled by a cationic Rh(III) catalyst (Scheme 350b).<sup>548</sup>

### Scheme 350. Cp\*Rh(III)-Catalyzed Vinylic C–H Acetoxylation of Enamides

a) Zhang et al., 2014




b) Xu, Li et al., 2018





In addition, Prakash's group illustrated a direct olefinic C–H fluorination of enamides with commercially available Selectfluor as the fluorinating reagent under Brønsted acidic conditions, affording a series of hitherto unknown fluoro-containing olefins in excellent yields (Scheme 351).<sup>549</sup>

### Scheme 351. Direct Olefinic C(sp<sup>2</sup>)-H Fluorination of Enamides




Xu and co-workers were able to synthesize highly useful vinylsilanes through an efficient and sustainable transition-metal-free dehydrogenative silylation of various enamides with simple silanes (Scheme 352).<sup>550</sup> This reaction proceeded smoothly in the presence of *tert*-butyl peroxybenzoate (TBPB) to trigger the formation of silyl radical. The authors highlighted the potential of this protocol by the derivatization of the corresponding vinylsilanes into a variety of value-added complex molecules.

### Scheme 352. Synthesis of Vinylsilanes via Metal-Free Dehydrogenative Silylation of Enamides



Very recently, Zeng's group devised a robust approach for the metal-free C–H thiocyanation of enamides with easily accessible  $\text{NH}_4\text{SCN}$  to assemble a diverse array of (*E*)- $\beta$ -thiocyanoenamides with high regio- and stereoselectivities (Scheme 353).<sup>551</sup>

### Scheme 353. Regio- and Stereo-Selective Olefinic C–H Thiocyanation of Enamides with $\text{NH}_4\text{SCN}$





## 5.2. Enaminones

$\beta$ -Keto enamines, or generally known as enaminones, contain unique electronic effects in its core structure – an electron-donating amino group and an electron-withdrawing carbonyl on either end of the C=C double bond. In addition, enaminones possess interesting properties and are versatile building blocks in organic synthesis. The application of enaminones bearing different substitution patterns on the nitrogen of the amino group were extensively investigated. Despite the extensive studies on this topic, *N,N*-disubstituted enaminones such as *N,N*-dimethyl enaminones, were not similarly scrutinized. In contrast to other conventional enaminones, *N,N*-dimethyl enaminones present a highly polarized C=C double bond due to the neighboring electron-donating amino group and electron-withdrawing carbonyl, which consequently result in a higher electron density at the  $\alpha$ -carbon as compared to the amino group. In addition, the *N,N*-dimethyl moiety functions as a better leaving group in contrast to other enaminones, thus allowing a number of transformations to be carried out with ease. As such, there have been an ever-increasing interest dedicated towards the study of *N,N*-dimethyl enaminones in recent years.

In 2018, Loh and co-workers reported a  $\text{Cp}^*\text{Rh(III)}$ -catalyzed regioselective [4 + 2] cycloaddition of

enaminones with  $\alpha$ -diazo- $\beta$ -ketoesters, allowing the construction of a series of highly functionalized salicylaldehydes with good functional group compatibility (Scheme 354).<sup>552</sup> In this case, alkenyl C–H bond functionalization/cyclization cascade reaction and subsequent re-aromatization gave rise to the corresponding products in a single step under benign conditions. The scope of the reaction was broad, both with regard to the enaminones and the  $\alpha$ -diazo- $\beta$ -ketoesters.


### Scheme 354. Synthesis of Salicylaldehydes through Regioselective [4 + 2] Cycloadditions of Enaminones with Diazocarbonyls




In the same year, the groups of Gao<sup>553</sup> and Zhu<sup>554</sup> independently published the efficient and straightforward assembly of multisubstituted salicylaldehyde derivatives through an enaminone-directed regioselective C–H [4 + 2] annulation of vinyl enaminones with diverse alkynes under  $\text{Cp}^*\text{Rh(III)}$  catalysis (Scheme 355).

### Scheme 355. Synthesis of Salicylaldehydes through Regioselective [4 + 2] Annulations of Enaminones with Alkynes

#### a) Gao et al., 2018




#### b) Zhu et al., 2018



A  $\text{Cp}^*\text{Rh(III)}$ -catalyzed C–H [5 + 1] annulation strategy for the construction of polyaromatic rings from enaminones was developed by Loh's group (Scheme 356).<sup>555</sup> The reaction featured a wide functional group tolerance and offered a novel route to synthesize polycyclic aromatic compounds bearing amino and formyl substituents. On the basis of mechanistic studies, the



authors proposed that the active Rh(III) catalyst activated the *ortho*-position of the phenyl ring through a weak keto-coordination to yield an intermediate. Subsequent coordination and migratory insertion of the olefin gave a rhodacycle intermediate which then underwent a  $\beta$ -hydride elimination to give another intermediate. Upon rotation of the C–C bond, another C–H activation step promoted by the pivalate group formed another Rh(III) complex. Reductive elimination of this complex afforded a six-membered cyclic intermediate. The amino group and rhodium complex was subsequently removed and upon condensation of the carbonyl formed an iminium ion intermediate. Lastly, hydrolysis and aromatization of the iminium ion generated the desired product.

### Scheme 356. Cp\*Rh(III)-Catalyzed [5 + 1] Annulation of Enaminones with Vinyl Esters





Subsequently, Loh and co-workers continued to present the Rh<sup>III</sup>-catalyzed formal oxidative [3 + 3] annulation of easily prepared enaminones with electron-deficient acrylates through alkenyl C–H activation and subsequent intramolecular aminolysis cyclization (Scheme 357),<sup>556</sup> allowing an efficient and practical access to highly functionalized *N*-substituted 2-pyridones in up to 91% yield with good functional group compatibility.

### Scheme 357. Synthesis of 2-Pyridones through Rh<sup>III</sup>-Catalyzed Formal Oxidative [3 + 3] Annulation of Enaminones with Acrylates



More recently, Guo and colleagues were able to assemble multisubstituted pyridines through an efficient [4 + 2] annulation of enaminones with diverse  $\alpha,\beta$ -unsaturated ketoxime acetates (Scheme 358).<sup>557</sup> In the presence of  $\text{FeCl}_2$  (10 mol%), the annulation reaction proceeded uneventfully, and a broad scope of 49 examples of the desired products was documented in appreciable yields of up to 87%. Of note, the reaction could be performed in the *avapourtec* flow reactor which definitely allowed for a substantial shortening of the reaction time, furnishing the annulation products in comparable yields. Preliminary mechanistic investigations were carried out which revealed that  $\text{FeCl}_2$  may play as a Lewis acid to activate the  $\alpha,\beta$ -unsaturated ketoxime acetates for the subsequent nucleophilic addition in this case.

### Scheme 358. Synthesis of Pyridines through Fe(II)-Catalyzed [4 + 2] Annulation of Enaminones with $\alpha,\beta$ -Unsaturated Ketoxime Acetates




The reaction of enaminones with diverse  $\alpha$ -diazoketones could produce *NH*-free pyrroles. In 2013, Reddy and co-workers described the synthesis of trisubstituted pyrroles through the cross-coupling reaction of  $\alpha$ -diazoketones with  $\beta$ -enaminoketones by using  $\text{Cu}(\text{OTf})_2$  (10 mol%) as the catalyst (Scheme 359a).<sup>558</sup> Gratifyingly, Park's group


demonstrated that the copper-catalyzed [3 + 2] cycloaddition of *N,N*-dimethyl enaminones with diazoketones exclusively afforded 2-amino-2,3-dihydrofurans which could be readily converted into highly functionalized furans in the presence of *p*-toluenesulfonic acid (Scheme 359b).<sup>559</sup>

### Scheme 359. Synthesis of Multisubstituted Furans and Pyrroles with Enaminones with $\alpha$ -Diazoketones

a) Reddy et al., 2013





b) Park et al., 2012





The group of Liu and Wan recently illustrated a Pd(II)-catalyzed annulation between enaminones and activated olefins for the construction of *NH*-free pyrroles (Scheme 360).<sup>560</sup> The three C(sp<sup>3</sup>)-H bonds were activated in the reaction to yield the corresponding pyrroles with good functional group tolerance. Evolution of H<sub>2</sub> gas from the reaction was also detected. Upon generation of the active Pd(II) catalyst, C-H activation of enaminones with the catalyst gave a Pd(II) complex. Subsequently, a 1,2-migratory insertion of the olefin to the Pd(II) complex formed a palladacycle which readily underwent  $\beta$ -H elimination to yield a dienamine intermediate. Dehydrogenative C-,N-palladation followed by a reductive elimination afforded the corresponding pyrroles. More recently, the same group further reported the construction of pyrazoles via a metal-free [2 + 2 + 1] annulation reaction of enaminones with hydrazines (Scheme 361).<sup>561</sup>

### Scheme 360. Pd-Catalyzed Triple-Fold C-H Activation with Enaminones and Alkenes




### Scheme 361. Synthesis of Pyrazoles via Metal-Free [2 + 2 + 1] Annulation of Enaminones with Hydrazines




Apart from the above-mentioned heterocycles, Cacchi and colleagues accomplished the rapid synthesis of multisubstituted indoles by the copper-catalyzed intramolecular oxidative cyclization of *N*-aryl enaminones (Scheme 362a).<sup>562</sup> Later on, the group of Chang and Yu also published an analogous transformation by means of a CuI/I<sub>2</sub> system (Scheme 362b).<sup>563</sup> Remarkably, the protocol could afford a diverse range of imidazo[1,2-*a*]pyridines in decent yields.

### Scheme 362. Cu-Catalyzed Intramolecular Oxidative Cyclization of Enaminones

a) Cacchi et al., 2009




b) Chang, Yu et al., 2016




The direct alkenyl C-H sulfenylation of enaminones represents a straightforward approach to the synthesis of polyfunctionalized aminothioalkenes. In 2016, Wan et al. reported a metal-free aerobic coupling of enaminones with thiophenols in the presence of a catalytic amount of KIO<sub>3</sub> as an efficacious hypervalent iodate catalyst (Scheme 363a).<sup>564</sup> The sustainable bio-based ethyl lactate (EL) was employed as a green solvent in this protocol to furnish the aminothioalkenes in excellent yields (75–85%). Almost at the same time, Loh and co-workers also performed a similar transformation by a Pd(OAc)<sub>2</sub>/dppe catalytic system (Scheme 363b).<sup>565</sup>

### Scheme 363. Synthesis of Polyfunctionalized Alkenes through KIO<sub>3</sub>-Catalyzed Aerobic Cross-Coupling of Enaminones with Thiophenols

a) Wang et al., 2016




b) Loh, Jiang et al., 2016



The incorporation of a  $\text{SCF}_3$  group into organic molecules will substantially enhance their biological activities. By using readily accessible  $\text{AgSCF}_3$  and trichloroisocyanuric acid (TCCA) to *in situ* generate the electrophilic trifluoromethylthio radical, Yang and co-workers were able to assemble a diverse range of  $\text{SCF}_3$ -containing chromones from *o*-hydroxyarylenaminones. A scope of 23 examples of this transformation was presented with yields of 52–95% (Scheme 364).<sup>566</sup> More recently, a similar trifluoromethylselenolation ( $\text{CF}_3\text{Se}$ ) of *ortho*-hydroxyarylenaminones was achieved under photoredox catalysis.<sup>567</sup>


#### Scheme 364. Synthesis of 3-((Trifluoromethyl)thio)-4*H*-chromen-4-one from *o*-Hydroxyarylenaminones and $\text{AgSCF}_3$




Additionally, Wan et al. described the metal-free C–H thiocyanation of enaminones with  $\text{NH}_4\text{SCN}$  as the thiocyanate source under the visible-light irradiation of a 14 W compact fluorescent lamp (CFL) with Rose Bengal as the non-metal photocatalyst (Scheme 365a).<sup>568</sup> In the same year, the groups of Duan<sup>569</sup> and Zhou<sup>570</sup> independently elaborated the efficient synthesis of SCN-containing enaminones under mild *N*-bromosuccinimide (NBS)- and iodobenzene diacetate (PIDA)-mediated conditions (Scheme 365b and c). Meanwhile, the olefinic C–H thiocyanation of *o*-hydroxyarylenaminones was also achieved, affording a large variety of biologically important 3-thiocyanato-4*H*-chromen-4-ones<sup>568,570,571</sup>

#### Scheme 365. Synthesis of SCN-Containing Enaminones via Olefinic C–H Thiocyanation


a) Wan et al., 2019



b) Duan et al., 2019




c) Zhou et al., 2021



In 2018, Yan and co-workers developed an efficient iron-based approach to prepare diverse polysubstituted thiazoles from readily available enaminones and elemental sulfur. In this case, the C–S bond is formed with sulfur powder through the direct functionalization of  $\text{C}(\text{sp}^2)/\text{C}(\text{sp}^3)$ –H bonds (Scheme 366).<sup>572</sup> Both electron-rich and electron-deficient functional groups were compatible with the reaction, affording the products in excellent yields.


#### Scheme 366. Synthesis of Thiazoles from Enaminones and Elemental Sulfur through the Vinylic C–H Functionalization/C–S Bond Formation



Quite recently, Fu and co-workers described the use of similar starting materials to perform an unprecedented TBPB-promoted oxidative cyclization for the synthesis of structurally diverse five-membered thiazole-2-thiones (Scheme 367).<sup>573</sup> This strategy undergoes  $\text{C}(\text{sp}^2)$ –H/ $\text{C}(\text{sp}^3)$ –H bond sulfuration of enaminones between the alkenyl and *N*-alkyl moieties along with a subsequent  $\text{C}(\text{sp}^3)$ –H bond thiocarbonylation, leading to the highly appealing


synthesis of thiazole-2-thiones in modest to excellent yields with a broad substrate scope.

### Scheme 367. TBPB-Promoted Oxidative Cascade Cyclization of Enaminones with Elemental Sulfur




The Maddani group demonstrated a metal-free alkenyl C-H halogenation of *N*-aryl enaminones by using a readily available and inexpensive DMSO-halo acid combination strategy under typically mild conditions (Scheme 368).<sup>574</sup> The scope of enaminones was found to be broad and not sensitive to electronic and steric factors. Interestingly, this protocol was applicable to *o*-hydroxyarylenaminone substrates, furnishing the halogenated chromenones in excellent yields.

### Scheme 368. Transition Metal-Free Oxidative Olefinic C-H Halogenation of *N*-Aryl Enaminones




In a subsequent report, Liu and colleagues described the facile synthesis of 3-halochromones with simple KX (X = Br, I) salts as halogen source in the presence of PhI(OAc)<sub>2</sub> oxidant (Scheme 369).<sup>575</sup> Instead of a radical-based pathway, control experiments definitely indicated that this protocol proceeded through an electrophilic halogenation involving the formation of a halogenium intermediate. Shortly thereafter, the same group extended to establish the electrochemical method for this transformation with sodium halide (NaX, X = Cl, Br, I) as the halogen source (Scheme 369).<sup>576</sup>

### Scheme 369. Synthesis of 3-Halochromones from 2-Hydroxyphenyl Enaminones and Metal Halides




In an effort to forge C-O bond, Loh and co-workers in 2018 disclosed an efficient alkenyl C-H acyloxylation of enaminones with PhI(OAc)<sub>2</sub> under transition-metal-free conditions (Scheme 370).<sup>577</sup> This transformation occurred smoothly at room temperature to deliver a diverse array of (*E*)-vinyl acetates in yields ranging from 45% to 91%.

### Scheme 370. Site-Selective C-H Acyloxylation of Enaminones with Phenyliodine(III) Diacetate



Meanwhile, Wan's group employed aryl peroxide as *O*-centered free-radical precursor, and accomplished the catalyst-free synthesis of both 3-acyloxy chromones and  $\alpha$ -acyloxy enaminones through a radical-involved pathway under extremely mild conditions (Scheme 371).<sup>578</sup>


### Scheme 371. Synthesis of Acyloxy Enaminones and Chromones via Free-Radical C-H Acyloxylation




The group of Yang and Chen in 2017 elaborated a direct synthesis of 3-CF<sub>2</sub>/CF<sub>3</sub>-containing chromone derivatives by the reaction of *o*-hydroxyphenylenaminones with BrCF<sub>2</sub>COOEt or Ph<sub>2</sub>SCF<sub>3</sub>OTf under visible-light photoredox catalysis (Scheme 372a).<sup>579</sup> The reaction was conducted under irradiation of white LEDs (18 W) at room temperature to give the corresponding products in 51–88% and 32–75% yield, respectively. Around the same time, Zhang and co-workers performed a similar transformation for the synthesis of 3-CF<sub>2</sub>-containing chromones using inexpensive Ru(bpy)<sub>3</sub>Cl<sub>2</sub> as the photocatalyst (Scheme 372b).<sup>580</sup> Afterward, Wan's laboratory utilized Langlois' reagent (CF<sub>3</sub>SO<sub>2</sub>Na) as the trifluoromethyl radical source, and accomplished the metal-free synthesis of 3-CF<sub>3</sub>-containing chromones in the presence of K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> as the oxidant. (Scheme 372c).<sup>581</sup>

### Scheme 372. Direct Synthesis of 3-CF<sub>2</sub>/CF<sub>3</sub>-Containing Chromones from 2-Hydroxyphenyl Enaminones


a) Yang, Chen et al., 2017



b) Zhang et al., 2017




c) Wan et al., 2020




Moreover, the Patil group explored the reaction of *o*-hydroxyphenylenaminones with TIPS-EBX reagent in the presence of AuCl as a catalyst, leading to a diverse array of 3-alkynyl chromones (Scheme 373a).<sup>582</sup> Afterward, they extended to investigate the reaction between *o*-hydroxyarylenaminones and diazo compounds under gold catalysis, affording a series of 3-alkylated chromones in up to 89% yield (Scheme 373b).<sup>583</sup> Detailed experimental and computational studies revealed a plausible pathway involving the OH-group assisted C–H bond alkylation of enaminones.

### Scheme 373. Synthesis of 3-Substituted Chromones via Gold-Catalyzed *o*-Hydroxyarylenaminones with TIPS-EBX Reagent and Diazo Compounds

a) Patil et al., 2016






b) Baik, Patil et al., 2019



In 2020, the group of Iaroshenko and Mkrtchyan was able to synthesize isoflavones through a consecutive cascade arylation between *o*-hydroxyarylenaminones and *in situ* generated aryl radicals under photoredox catalysis (Scheme 374).<sup>584</sup> In this report, the authors employed diaryliodonium and diazonium salts as the aryl radical precursors. The photo-Meerwein arylation reaction of aryldiazonium tetrafluoroborates proceeded with Eosin Y as a photocatalyst under irradiation of green LEDs, while the arylation process by diaryliodonium triflates underwent smoothly at room temperature in the presence of Ru(bpy)<sub>3</sub>Cl<sub>2</sub>·6H<sub>2</sub>O under irradiation of blue LEDs.


### Scheme 374. Visible-Light-Mediated Arylation of *ortho*-Hydroxyarylenaminones



The same year, Liu and co-workers illustrated a highly regioselective β-C(sp<sup>2</sup>)-H activation of enaminones with various nitrogen heterocycles for the construction of 2-nitrogenated chromones (Scheme 375).<sup>585</sup> This strategy unveiled a novel approach to access useful heterocyclic structures. The reaction was promoted by molecular iodine, which gave a 3-iodochromone intermediate and HI. In the presence of the oxidant PhI(OAc)<sub>2</sub>, molecular iodine was

regenerated from HI. The 3-iodochromone intermediate was subsequently activated by the Cu(II) species to form a complex which was subjected to nucleophilic attack by the nitrogen heterocycles to form a Michael intermediate. The elimination of HI with the assistance of a base afforded the 2-nitrogenated chromones.


### Scheme 375. Copper-Catalyzed $\beta$ -C(sp<sup>2</sup>)-H *N*-Heteroarylation of Enaminones




Cyclic six-membered enaminones, also known as 2,3-dihydropyrid-4(1*H*)-ones, are versatile and synthetically useful intermediates that demonstrate unique biological properties, and could undergo a series of transformations to afford various biologically active piperidine-containing molecules. As such, the vinylic C–H functionalization of nonaromatic, cyclic enaminones has been extensively investigated over the past years. Specifically, Georg's group in 2008 presented the first example of Pd(II)-catalyzed lakenyl C–H arylation of 2,3-dihydropyrid-4(1*H*)-ones with aryltrifluoroborates, giving rise to an efficient approach for the assembly of 3-arylpiperidines which are a privileged structure and prevalent scaffold in numerous natural products (Scheme 376a).<sup>586</sup> Although an elegant reaction has been developed, the protocol suffers from a relatively high catalyst loading (30 mol%). To circumvent this issue, Georg et al. expanded to establish a new strategy that accommodates readily available boronic acids as the coupling partner in the presence of 10 mol% Pd(OAc)<sub>2</sub> (Scheme 376b).<sup>587</sup> The reaction was typically finished within 1 h at 60 °C in DMF. The authors claimed this significant improvement can be attributed to the addition of stoichiometric copper(II) salts, which can greatly facilitate the delivery of the aryl group to the Pd(II) catalyst center through the facile formation of arylcopper species, thereafter significantly increasing the efficiency of the transmetalation process. Later in 2014, the same group further accomplished a highly regioselective C–H arylation at the C6 position of cyclic enaminones enabled by a Pd(II)/bpy catalytic system with molecular oxygen as the sole oxidant (Scheme 376c).<sup>588</sup>

### Scheme 376. Palladium(II)-Catalyzed C–H Arylation of Cyclic Enaminones with Organoboron Reagents


a) Georg et al., 2008



b) Georg et al., 2012





c) Georg et al., 2014





Meanwhile, Georg and co-workers also investigated the alkenyl C–H Hiyama cross-coupling of enaminones with organosilicon reagents (Scheme 377).<sup>589</sup> By taking advantage of copper(II) fluoride (CuF<sub>2</sub>) as both silane activator and reoxidant, the coupling reaction between 2,3-dihydropyrid-4(1*H*)-ones and aryltriethoxysilanes occurred smoothly to afford C5-arylated enaminones. Gratifyingly, Denmark's reagent such as dimethylphenylsilanol can also participate without the need to use extra fluoride activator. Soon after, the authors further reported the C–H arylation of cyclic enaminones with aryl iodides (Scheme 378).<sup>590</sup>

### Scheme 377. Palladium(II)-Catalyzed C–H Arylation of Cyclic Enaminones with Organosilicon Reagents




### Scheme 378. Palladium(II)-Catalyzed C–H Arylation of Cyclic Enaminones with Aryl Iodides




Moreover, Georg and colleagues also investigated the palladium(II)-catalyzed C–H olefination of cyclic enaminones with both primary and secondary electron-deficient alkenes (Scheme 379a).<sup>591</sup> By using a combination of Cu(OAc)<sub>2</sub> (2.0 equiv) and KOTFA (1.0 equiv) in DMF at 80 °C, the reaction occurred smoothly, and excellent conversion was observed for most substrates. Later in 2014, the authors expanded this alkenylation to establish the biomimetic strategy of aerobic C–H olefination of enaminones at room temperature (Scheme 379b).<sup>592</sup>

### Scheme 379. Pd(II)-Catalyzed C–H Olefination of Cyclic Enaminones

a) Georg et al., 2011



b) Georg et al., 2014




## 5.3. Enamines

Arguably, enamines have long been used as extremely versatile building blocks for the synthesis of a variety of nitrogen-containing molecules,<sup>593,594</sup> and also play a prominent role in the field of organocatalysis.<sup>595</sup> Considering the unique property of enamines, the alkenyl C(sp<sup>2</sup>)-H functionalization of readily available enamines has drawn considerable attention from the synthetic community, and a myriad of efficient and straightforward methodologies have been established over the decades for the purpose of streamlining synthetic routes to fabricate highly functionalized enamines. Specifically, the group of



Zhao and Du in 2012 reported an efficient transition-metal-free C–H acyloxylation of enamines with carboxylic acids by means of iodosobenzene (PhIO) as the oxidant, exclusively furnishing a diverse array of β-acyloxy enamines that can be readily converted into oxazole derivatives (Scheme 380).<sup>596</sup> The protocol was proposed to take place through the in situ generation of PhI(OCOR)<sub>2</sub> followed by the facile β-acyloxylation of enamines.

### Scheme 380. PhIO-Mediated β-C–H Acyloxylation of Enamines with Carboxylic Acids




Subsequently, the same group further developed a coupling reaction between structurally diverse enamines and electron-deficient amines by using a TBAI/TBHP oxidative system, enabling an efficient assembly of synthetically appealing diaminoalkenes in decent yields (37–95%) (Scheme 381).<sup>597</sup> This oxidative C(sp<sup>2</sup>)-N bond formation reaction may involve the formation of tetra-*n*-butylammonium iodite from TBAI and TBHP, which then undergoes an electrophilic addition process by enamines to generate the adduct species. Subsequent nucleophilical substitution reaction with amines affords the corresponding imines which further isomerized into the thermodynamically more stable diaminoalkene derivatives.

### Scheme 381. TBAI/TBHP-Mediated Oxidative Coupling of Enamines with Electron-Deficient Amines





In the following year, the group of Wang and Fan disclosed an efficient metal-free oxidative cross-coupling of enamines with *N*-protected propargylamines by taking advantage of a PhIO/Bu<sub>4</sub>NI-mediated system (Scheme 382).<sup>598</sup> A scope of 22 examples of this transformation was presented with yield up to 87%. In this report, tetrabutylammonium iodide (Bu<sub>4</sub>NI) serves as an activator of PhIO. Interestingly, the resulting oxidative coupling products can undergo a copper(II)-mediated electrophilic cyclization process to afford structurally diverse multisubstituted 3-pyrrolines, which are recognized as a key substructure in many pharmaceutically relevant molecules.

**Scheme 382. Iodine(III)-Mediated Cross-Coupling of Enamines with Propargylamines and Subsequent Cyclization to Multisubstituted 3-Pyrrolines**




The group of Sun and Shi in 2014 published the Rh-catalyzed hydrogen-bonding assisted alkenyl β-C-H functionalization of both five- and six-membered cyclic enamines with carboxylic acids by means of 3-methoxy-2-pyridinyl amino group as the directing group (Scheme 383a).<sup>599</sup> A broad range of aryl, alkenyl carboxylic acids as well as aliphatic acids are all competent coupling partners for this decarbonylative reaction. Of note, the directing group can be easily removed by the treatment with TsOH in aqueous media, offering an alternative approach to highly functionalized cyclic 1,2-diketones. Almost at the same time, Dong and colleagues made use of a similar strategy to the synthesis of β-alkenylated α-enamino-ketones via enamine-directed vinyl C-H/alkyne couplings (Scheme 383b).<sup>600</sup>

**Scheme 383. Rh-Catalyzed Cross-Coupling of Cyclic Enamines with Carboxylic Acids or Alkynes**



Several years later, Song and colleagues successfully established a general denitrogenative C-H arylation of enamines with 3-aminoindazoles as the arylating agents via the C-N bond cleavage, providing a series of arylated enamines in good yields (Scheme 384).<sup>601</sup> Mechanistically, the authors proposed that a benzonitrile radical may be probably involved in this C-H arylation process.


**Scheme 384. Copper-Catalyzed Alkenyl C-H Arylation of Enamines with 3-Aminoindazoles**



The introduction of a trifluoromethylthio group (CF<sub>3</sub>S-) into enamines through vinylic C-H cleavage has been also


explored. In 2013, Shibata's group elaborately designed a novel and shelf-stable electrophilic trifluoromethylthiation reagent, that is trifluoromethanesulfonyl hypervalent iodonium ylide. A variety of trifluoromethylthiolated enamines were nicely obtained in excellent yields (74–97%) in the presence of a catalytic amount of CuCl (20 mol%) (Scheme 385).<sup>602</sup> The reaction was typically finished within 5 min at room temperature in 1,4-dioxane. The scope was impressive as indoles and  $\beta$ -keto esters were also suitable substrates in this case, furnishing the trifluoromethylthiolated products in high yields. From a mechanistic point of view, the authors suggested that the in situ reduction of trifluoromethanesulfonyl group by an intramolecular rearrangement process gives rise to the CF<sub>3</sub>S group, which is identified as the key step in this protocol.

### Scheme 385. Copper-Catalyzed Vinylic C–H Bond Trifluoromethylthiation of Enamines




Later in 2017, Jiang and Wu established a general protocol for the cross-coupling of enamines with Langlois' reagent (Scheme 386).<sup>603</sup> A series of enamines can smoothly couple with CF<sub>3</sub>SO<sub>2</sub>Na by using TBHP as both initiator and oxidant under transition-metal-free conditions, giving rise to a facile synthesis of  $\beta$ -trifluoromethylated enamines in moderate to excellent yields (42–92%) with exclusive *E*-configuration.

### Scheme 386. Synthesis of $\beta$ -Trifluoromethylated Enamines with Langlois' Reagent



Vinyl sulfides are privileged and extremely important scaffolds that are widespread in a series of biologically active natural products and pharmaceutically relevant molecules. In 2016, the Du group found that the reaction of enamines with diaryl disulfides could afford a broad array of  $\alpha$ -thioenamines in appreciable yields (45–91%) by using a catalytic amount of tetrabutylammonium iodide (10 mol%) in combination with 3.0 equiv. of *tert*-butyl hydroperoxide (TBHP) (Scheme 387).<sup>604</sup> This metal-free TBAI/TBHP oxidative system also enables the formation of C–Se bonds for the synthesis of  $\alpha$ -selenylated enamines, albeit with relatively low yields (28–42%).


### Scheme 387. TBAI/TBHP-Mediated Oxidative Coupling of Enamines with Disulfides



Afterward, Lei and co-workers illustrated a rare example of electrochemical oxidative C–H/S–H coupling to forge the C–S bond, providing a sustainable, atom-economical, and environmentally friendly approach for the synthesis of vinyl sulfides with H<sub>2</sub> as the only byproduct (Scheme 388).<sup>605</sup> A broad range of enamines and thiophenols could be readily converted under electrochemical oxidative conditions, significantly obviating the use of external


oxidants. A plausible mechanism involving the generation of a thiyl radical was tentatively proposed to elucidate this C–S formation process. More recently, Li and Xie followed this work in an attempt to expand the substrate scopes of this electrochemical protocol by employing a modified conditions with a carbon felt anode and a nickel plate cathode as the electrode materials.<sup>606</sup>

### Scheme 388. Electrochemical Oxidative Cross-Coupling of Enamines with Thiophenols




Subsequently, Wu's group presented a general and highly efficient strategy for the vinylic  $\beta$ -C–H phosphorylation of enamines enabled by cost-effective cobaloxime catalysis under visible-light irradiation with hydrogen evolution (Scheme 389).<sup>607</sup> The reaction occurred under especially mild conditions and exhibited excellent functional group tolerance for both enamines and H-phosphine oxides, providing a concise route to a series of  $\beta$ -phosphinoyl products in 52–99% yield with modest *Z*-selectivity. On the basis of experimental and spectroscopic investigations, a reductive quenching pathway of cobaloxime catalyst to generate phosphinoyl radical was proposed to elucidate the mechanism.

### Scheme 389. Direct Alkenyl C–H Phosphorylation of Enamines Enabled by Cobaloxime Catalysis




In a recent report, Hashmi, Tian, and their co-workers reported the use of enamines as substrates to couple with hypervalent iodine(III) reagents under gold catalysis, in which a series of tetrasubstituted conjugated 1,3-enynes were obtained in a regioselective manner (Scheme 390).<sup>608</sup> Mechanistically, the Au(I) species **I** was initially produced under the typical conditions, which then undergoes oxidative addition with hypervalent iodine reagent to afford the alkynyl Au(III) intermediate **II**. Subsequent alkenyl C–H functionalization of enamines with alkynyl Au(III) species **III** followed by a facile reductive elimination process would release the corresponding 1,3-enynes and simultaneously regenerate the Au(I) catalyst.

### Scheme 390. Synthesis of Tetrasubstituted 1,3-Enynes by Gold-Catalyzed C–H Alkynylation of Enamines





#### Proposed mechanism



A diverse array of highly functionalized naphthalene derivatives were readily prepared by the group of Wei from readily available enamines and alkynes via a general, practical, and efficient iodine(III)-mediated oxidative benzannulation strategy (Scheme 391).<sup>609</sup> This metal-free protocol features a wide substrate scope and especially mild conditions. In this case, the nucleophilic attack of alkyne on the iodine(III) species generated from the reaction of enamine with PhIO was crucial for subsequent intramolecular electrophilic benzannulation process.


#### Scheme 391. Synthesis of Polysubstituted Naphthalene Derivatives via Iodine(III)-Mediated Benzannulation of Enamines with Alkynes




Glorius and co-workers in 2008 took the lead in realizing a direct and broadly applicable approach for the efficient synthesis of 2,3-disubstituted NH-free indoles through the palladium-catalyzed intramolecular oxidative cyclization of *N*-aryl enamines, which can be readily prepared from commercially available aniline derivatives. (Scheme 392a).<sup>610</sup> Initial mechanistic studies unambiguously favored a  $\sigma$ -bond-metathesis or deprotonation pathway for this transformation. Later in 2011, they continued to uncover the details of this dehydrogenative cross-coupling reaction.<sup>611</sup> In the aftermath of this work, the Rueping group achieved a similar transformation enabled by the synergistic palladium and photoredox catalysis while obviating the use of stoichiometric Cu(OAc)<sub>2</sub> as the oxidant (Scheme 392b).<sup>612</sup> Around the same time, Guan's group carried out the intramolecular oxidative cyclization of tertiary enamines in an effort to prepare various multisubstituted pyrroles and indoles in decent yields with a stoichiometric amount of TFA as an additive (Scheme 392c).<sup>613</sup>

#### Scheme 392. Synthesis of Indoles and Pyrroles via Pd-Catalyzed Intramolecular Oxidative Cyclization of *N*-Aryl Enamines


a) Glorius et al., 2008



b) Rueping et al., 2014




c) Guan et al., 2014



Moreover, Kurth and co-workers in 2013 described the intramolecular coupling of *o*-bromoaniline-based *N*-aryl  $\beta$ -nitroenamines by the combination of Pd(PPh<sub>3</sub>)<sub>4</sub> catalyst and Et<sub>3</sub>N additive, offering an efficient route to access 3-nitroindoles in decent yields with excellent regioselectivity (Scheme 393).<sup>614</sup> Noteworthy is that the reaction occurs under microwave irradiation conditions. Moreover, a variety of 3-carboalkoxy- and 3-cyanoindoles can be obtained by this microwave-assisted protocol.


#### Scheme 393. Microwave-Assisted Pd-Catalyzed Indole Synthesis from *N*-Aryl Enamines




By taking advantage of environmentally benign, sustainable, and inexpensive iron catalyst combined with a highly active Cu(OAc)<sub>2</sub>•CuCl<sub>2</sub> as the oxidant, Liang and colleagues were able to prepare a series of multisubstituted indoles in synthetically satisfactory yields (Scheme 394a).<sup>615</sup> The generated iron and copper bimetallic chelate species by the coordination of the *N*-atom and the C=C bond was assumed to greatly enhance the activity of the enoate intermediate in this protocol. Later in 2015, Taylor and co-workers performed analogous investigations by their Cu(II)-mediated protocol at relatively high temperature (Scheme 394b).<sup>616</sup>

### Scheme 394. Sustainable Synthesis of Indoles via Oxidative Cyclization of *N*-Aryl Enamines

a) Liang et al., 2010





b) Taylor et al., 2015



Subsequently, Zhang et al. achieved an efficient one-pot synthesis of 2,3-disubstituted indoles through a facile oxidative cyclization reaction of *N*-aryl enamines by a combination of CuCl, *N*-bromosuccinimide (NBS), and NaOAc, leading to the corresponding polysubstituted indoles in up to 99% yield (Scheme 395).<sup>617</sup>


### Scheme 395. CuCl/NBS Mediated Oxidative Cyclization of *N*-Aryl Enamines




However, it should be noted that the intramolecular cyclization of *N*-aryl enamines for indole synthesis can be performed under transition-metal-free conditions. For instance, Zhao and co-workers in 2009 expanded their PIDA-mediated strategy, and established a novel metal-free approach for indoles synthesis from *N*-aryl enamines in the presence of phenyliodine(III) diacetate (PIDA) (Scheme 396a).<sup>618</sup> The mechanism of this process was not investigated in detail, but the authors suggested an intramolecular S<sub>N</sub>2'-type cyclization pathway to elucidate the formation of indoles. Shortly afterwards, Li and co-workers found that *N*-aryl enamines could also undergo an intramolecular oxidative cyclization process to produce a series of functionalized indoles using a I<sub>2</sub>/NBS catalytic system. A scope of 13 examples of this transformation was presented with yields of 36–95% (Scheme 396b).<sup>619</sup>

### Scheme 396. Metal-free Indole Synthesis via Oxidative Cyclization of *N*-Aryl Enamines

a) Zhao et al., 2009



b) Li et al., 2011




The merging of photoredox catalysis with Co-catalysis opened new avenues on the route toward the synthesis of indoles from enamines. In 2016, Wu and colleagues accomplished an elegant synergistic iridium/cobalt-catalyzed strategy for indole synthesis by the intramolecular dehydrogenative annulation of *N*-aryl enamines under visible-light irradiation (Scheme 397a).<sup>620</sup> Using Ir(ppy)<sub>3</sub> as the photosensitizer and Co(dmgh)<sub>2</sub>(4-


CO<sub>2</sub>Mepy)Cl as the cobaloxime catalyst, a series of *N*-aryl enamines can be readily converted into the expected indoles under ambient conditions with H<sub>2</sub> as the only byproduct, with 23 examples documented in yields of 55–95%. Combined spectroscopic and electrochemical investigations indicated that this formal dehydrogenative annulation protocol proceeds through a visible-light-catalyzed oxidation of *N*-aryl enamines, followed by an intramolecular radical addition process. One year later, Lei's group also achieved the same transformation in the presence of KI under electrochemical conditions (Scheme 397b).<sup>621</sup>

### Scheme 397. Synthesis of Indoles by Intramolecular Dehydrogenative Cyclization of *N*-Aryl Enamines

a) Wu et al., 2016




b) Lei et al., 2017




Moreover, substituted pyrrole synthesis has been also realized from enamine starting materials. In 2012, Li and colleagues elaborated a general approach to access tetrasubstituted pyrroles by the dehydrogenative cross-coupling between enamines and acetone with 1.0 equiv of silver carbonate as an oxidant (Scheme 398a).<sup>622</sup> Later, the group of Zhang and Cui discovered that enamines reacted smoothly with propargyl acetates through a copper-catalyzed cascade propargylation/alkyne azacyclization/isomerization sequence to produce fully substituted pyrroles in up to 93% yield (Scheme 398b).<sup>623</sup> The reaction was carried out in toluene at 150 °C under microwave irradiation. Soon after, Luo, Deng, and their colleagues obtained a similar transformation with allenes as the coupling partner by the combination of KI, TBHP, and K<sub>2</sub>CO<sub>3</sub> in DCE solvent (Scheme 398c).<sup>624</sup> A plausible mechanism was discussed, and the reaction may proceed via a tandem Michael addition/oxidative annulation pathway. More recently, Liu et al. established the copper-mediated annulation of enamines with internal dialkyl ethylenedicarboxylates at room temperature, giving rise to diverse 2,3-dicarboxylate-functionalized pyrroles in high yields (61–80%). Interestingly, a broad array of 2-vinyl pyrroles could be synthesized when employing terminal propiolates as coupling partners under modified conditions (Scheme 398d).<sup>625</sup>

### Scheme 398. Direct Synthesis of Multisubstituted Pyrroles from Enamines


a) Li et al., 2012




b) Zhang, Cui et al., 2016



c) Luo, Deng et al., 2016



d) Liu et al., 2022




Meanwhile, efficient and practical synthetic strategies for the homo- and heterocoupling of enamines have been also established for the synthesis of structurally diverse pyrroles in recent years. Specifically, Guan and co-workers in 2016 elaborated a robust approach for the synthesis of a large variety of multisubstituted NH-pyrroles by a K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>-mediated oxidative homocyclization of enamines (Scheme 399a).<sup>626</sup> The same group further expanded to describe a novel Cu-catalyzed tandem oxidative cyclization/1,2-amino migration of enamines, affording aminomethylsubstituted pyrroles in good yields (Scheme 399b).<sup>627</sup> An optimized set of conditions consisted of substrate solutions in MeCN in the presence of Cu(TFA)<sub>2</sub> catalyst, AgOAc additive and O<sub>2</sub> oxidant. One year later, Chen and colleagues elaborated the synthesis of unsymmetrically multisubstituted pyrroles via copper-catalyzed aerobic dimerization (Scheme 399c).<sup>628</sup> Subsequently, they continued to report the synthesis of symmetric fully substituted pyrrole derivatives from enamines by an electrochemical-oxidation-induced annulation process (Scheme 399d).<sup>629</sup> Meanwhile, Roy, Sarkar, and their co-workers described a similar approach for multisubstituted NH-pyrrole synthesis under electrochemical conditions. Notably, with trifluoroethanol (TFE) as a unique additive, this electrochemical strategy also enables the heterocoupling of aryl- and


alkylsubstituted enamines to afford a series of tetrasubstituted NH pyrroles in comparable yields. Combined experimental and theoretical studies were also performed to elucidate the “magic effect” of TFE additive in promoting this heterocoupling reactions (Scheme 399e).<sup>630</sup>

### Scheme 399. Synthesis of Multisubstituted Pyrroles via Oxidative Cyclization of Enamines


a) Guan et al., 2016




b) Guan et al., 2018




c) Chen et al., 2019



d) Chen et al., 2021





e) Roy, Sarkar et al., 2021




Apart from the above-described examples for pyrrole synthesis, the group of Li and Gao in 2015 developed a novel method for the convenient synthesis of diverse multisubstituted *trans*-2,3-dihydropyrroles from enamines and chalcones via an iodine-promoted, one-pot cascade Michael/cyclization sequence, affording the products in modest to excellent yields (Scheme 400).<sup>631</sup>

### Scheme 400. Rapid Synthesis of Multisubstituted 2,3-Dihydropyrroles from $\beta$ -Enamine Esters




In the course of their investigations on the intramolecular cyclization of enamines for indole synthesis,<sup>610</sup> Glorius and co-workers found that the use of acetonitrile as the solvent instead of DMF exclusively resulted in the formation of tetrasubstituted pyrazoles (Scheme 401).<sup>632</sup> The reaction proceeded involving an oxidative N–N bond formation in the presence of copper acetate as both a Lewis acid activator and as an oxidant. Mechanistically, the nitrile substrate is nicely activated by the copper(II) Lewis acid, greatly facilitating the addition of the nucleophilic enamines. Subsequent loss of HOAc results in the formation a 1,3-bisimine. After rotation around the former *Z* double bond followed by elimination of another HOAc, a Cu<sup>II</sup>-chelate species was produced. Finally, reductive elimination furnishes the desired pyrazoles and releases copper(0) simultaneously.

### Scheme 401. Synthesis of Tetrasubstituted Pyrazoles from Enamines and Nitriles



Ding's group in 2015 published the efficient palladium-catalyzed intermolecular cyclization reaction of various *N*-aryl enamines with isocyanides as versatile C<sub>1</sub> building blocks through both aromatic and olefinic C(sp<sup>2</sup>)–H bonds cleavage (Scheme 402).<sup>633</sup> By using a combination of Pd(OAc)<sub>2</sub> catalyst and 1,10-phen ligand, a series of highly valuable 4-aminoquinolines were produced in decent yields (30–80%). This cascade oxidative cyclization protocol was proposed to undergo sp<sup>2</sup> C–H bonds activation of enamines, isocyanides migratory insertion, and reductive elimination sequences.


### Scheme 402. Pd-Catalyzed Intermolecular Oxidative Cyclization of Enamines with Isocyanides




Quinoxalines, also known as benzo[*a*]pyrazines, are an extremely important class of nitrogen-containing heterocyclic scaffolds widespread in numerous natural products and drug candidates with remarkable biological activities.<sup>634–636</sup> As such, tremendous efforts have been made for the preparation of quinoxalines from *N*-aryl enamines. Specifically, Zhang and colleagues in 2014 reported that *N*-aryl enamines could react with nitromethane through an oxidative tandem nitrosation/cyclization sequence enabled by the KI/TBHP catalytic system, furnishing the metal-free synthesis of 3-trifluoromethylquinoxalines in a single step with high efficiency (Scheme 403a).<sup>637</sup> In a related synthesis, Mo, Pan, Su, and their co-workers utilized readily available *iso*-butyl nitrite (IBN) as a novel nitrogen reagent to construct C–N bonds, and reported on a reaction between *N*-aryl enamines and IBN in the presence of  $\text{Ac}_2\text{O}$  (Scheme 403b).<sup>638</sup> This process presumably involved the generation of the key  $\beta$ -imino oxime ester intermediate and subsequent  $6\pi e$ -azacyclization.

### Scheme 403. Synthesis of Quinoxalines through Alkenyl C–H Functionalization of *N*-Aryl Enamines

a) Zhang et al., 2014



b) Mo, Pan, Su et al., 2018




In addition to the examples discussed above, in which enamines were used directly as the starting materials for the reaction, there are a few processes employing the combination of simple ketones or aldehydes with suitable directing groups to *in situ* generate enamines. Such carbonyl-involved synergistic transition-metal/enamine

catalysis would unambiguously convert the ketone  $\text{sp}^3$   $\alpha$ -C–H bonds to  $\text{sp}^2$  bonds, thus significantly enhancing the reactivity toward oxidative addition by a low-valent transition metal. Though the overall reaction of this strategy is the  $\alpha$ -C–H transformation of ketones, the inherent catalytic mechanism involves the enamine vinylic C–H functionalization step.<sup>639–644</sup>




In demonstrating the proof of concept,<sup>645</sup> Dong and co-workers pioneered to explore the feasibility of such a dual activation strategy, and successfully accomplished an unprecedented Rh-catalyzed  $\alpha$ -C–H alkylation of cyclic 1,2-diketones by using terminal olefins as the alkylating agents (Scheme 404).<sup>646</sup> In this ketone  $\alpha$ -alkylation protocol, a stoichiometric amount of 2-aminopyridine was employed as the efficacious *in situ*-installed directing group. A wide range of simple olefins including aliphatic, aromatic olefins, and even ethylene gas were compatible with the conditions, exclusively affording the linear alkylation products.

### Scheme 404. Rh(I)-Catalyzed C–H Bond Alkylation of Enamines Generated from 1,2-Diketones





In 2014, Dong's group elegantly disclosed a distinctive and regioselective  $\alpha$ -alkylation of cyclopentanones with simple olefins (Scheme 405).<sup>647</sup> In this report, readily available 7-azaindoline was identified as the optimal bifunctional aminocatalyst. Using the combination of  $[\text{Rh}(\text{coe})_2\text{Cl}]_2$  catalyst (2.5 mol%), *N*-heterocyclic carbene (NHC) ligand (IMes, 5 mol%), 7-azaindoline (25 mol%), and  $\text{TsOH}\cdot\text{H}_2\text{O}$  additive (10 mol%) in toluene, a range of 3-substituted cyclopentanones coupled regioselectively with ethylene at the less-hindered position under both pH and redox-neutral conditions, delivering the products in decent yields with moderate diastereoselectivity. Notably, other cyclic, acyclic ketones as well as terminal  $\alpha$ -olefins were also compatible substrates under slightly modified conditions, albeit with modest efficiency.

### Scheme 405. Regioselective $\alpha$ -Alkylation of Ketones with Olefins via Rh-Enamine Cooperative Catalysis



Inspired by Córdova's seminal work of palladium and enamine cooperative catalysis to obtain  $\alpha$ -allylic alkylation of ketones with allylic acetates,<sup>648,649</sup> Lei, Luo, and colleagues in 2014 elaborated an efficient oxidative  $\alpha$ -C-H alkylation of regular ketones with allylarenes enabled by synergistic nucleophilic enamine activation and Pd-catalyzed allylic C-H functionalization (Scheme 406).<sup>650</sup> However, this Pd/proline co-catalyzed oxidative coupling protocol was limited to allylarenes and, furthermore, the asymmetric version of this process did not furnish the enantioselective products.


#### Scheme 406. Oxidative Coupling of Allylarenes with Unactivated Ketones via Synergistic Pd/Enamine Catalysis



Later in 2015, Dong and Lim continued to describe a complementary protocol for the intramolecular Conia-ene-type ketone  $\alpha$ -C-H alkylation with olefins by taking advantage of a similar dual-activation strategy (Scheme 407).<sup>651</sup> Likewise, a stoichiometric amount of 2-aminopyridine was identified as the bifunctional



aminocatalyst. The authors found that rhodium catalyst system was recognized to be more efficient for the generation of five-membered rings, whereas a ruthenium-based catalyst was discovered to furnish the six-membered ring products.

#### Scheme 407. Rh- and Ru-Catalyzed Intramolecular Ketone $\alpha$ -Alkylation via Dual Activation





Almost at the same time, Dong and colleagues also devised an amino/Rh cooperative catalysis protocol for the  $\alpha$ -C-H alkenylation of regular ketones with various internal alkynes by the combination of Wilkinson's catalyst and bifunctional 7-azaindoline aminocatalyst, enabling the controllable synthesis of thermodynamically stable  $\alpha,\beta$ -enones and kinetically favored  $\beta,\gamma$ -enones with decent yields (Scheme 408).<sup>652</sup> Gratifyingly, a broad range of functionalities were compatible, and excellent *E*-selectivity was observed. Moreover, a series of control, kinetic monitoring, and deuterium labeling experiments were performed to investigate the catalytic mechanism of this process. Mechanistically, the reaction may involve a sequence of the insertion of the rhodium(I) species into the in situ formed enamine vinyl C-H bond, subsequent alkyne coordination, Rh-H migratory insertion, and reductive elimination.

#### Scheme 408. Rhodium-Catalyzed $\alpha$ -Alkenylation of Ketones with Internal Alkynes



Furthermore, Dong and co-workers also elaborated a palladium and enamine cooperative catalysis approach for the direct α-C–H monoarylation of cyclopentanones with aryl bromides (Scheme 409).<sup>653</sup> Both aminocatalyst and Pd(OAc)<sub>2</sub> proved to be indispensable for achieving high efficiency. Of note, no coordinating moiety on the aminocatalyst was required in this reaction, which is quite different from that of synergistic Rh/enamine catalysis.



#### Scheme 409. Direct α-Arylation of Cyclopentanones by Palladium/Enamine Cooperative Catalysis



A couple of months later, a general procedure for the enantioselective α-arylation desymmetrization reaction of cyclohexanones by the combination of Pd(OAc)<sub>2</sub>/PPh<sub>3</sub> and (L)-proline catalysis was presented by Jia and colleagues, unexpectedly giving rise to a diverse array of chiral bridged

morphan derivatives bearing α-carbonyl tertiary stereocenters (Scheme 410).<sup>654</sup> Stoichiometric amount of K<sub>3</sub>PO<sub>4</sub> base and HOAc additive were required for the reaction. A broad scope of 22 examples of this reaction was documented with yields of 39–96% and excellent enantiomeric excesses (up to 99%).



#### Scheme 410. Synergistic Pd/(L)-Proline Catalyzed Enantioselective α-Arylation Desymmetrization of Cyclohexanones



#### 5.4. Enolates



**5.4.1. Enol Acetates.** Early in 2004, the Iishi group first investigated the Pd(OAc)<sub>2</sub>/HPMoV catalyzed aerobic oxidative cross-coupling of vinyl carboxylates with acrylates using O<sub>2</sub> as the sole oxidant in acetic acid, affording the conjugated 1,3-dienes in 45–76% yields (Scheme 411).<sup>655</sup>

#### Scheme 411. Pd(OAc)<sub>2</sub>/HPMoV-Catalyzed Oxidative Cross-Coupling of Vinyl Carboxylates with Acrylates





Later in 2015, Loh and co-workers explored the intermolecular reaction of vinylacetates with various aryl iodides in the presence of a superstoichiometric amount of carboxylic acids (Scheme 412).<sup>656</sup> Contrast to the direct olefinic C–H arylation, this protocol underwent a three-component oxyarylation of vinylacetates with the retention of C=C bonds, leading to the difunctionalized vinyl esters in modest to excellent yields.

#### Scheme 412. Palladium(II)-Catalyzed Intermolecular Oxyarylation of Vinylacetates



**5.4.2. Enol Carbamates.** Enol carbamates have been long well-recognized as key structural motifs of numerous biologically relevant pharmaceuticals, and also served as versatile synthons for a series of transformations.<sup>657</sup> As a consequence, the vinylic C–H functionalization of enol carbamates has been extensively investigated in recent years. Specifically, Fu's group in 2014 reported the Cp<sup>\*</sup>Rh(III)-catalyzed C–H activation of enol carbamates with allenes to afford highly unsaturated conjugated alkenes with excellent regioselectivity (Scheme 413).<sup>658</sup> The produced olefins could be easily converted into synthetically important cyclic skeletons through an intramolecular transesterification or Diels-Alder reaction, which definitely renders this protocol highly appealing in organic synthesis.



**Scheme 413. Synthesis of Conjugated Olefins through Rh(III)-Catalyzed C–H Activation of Enol Carbamates with Allenes**



The same year, Glorius and co-workers described a vinylic C–H alkenylation of enol carbamates with a series of activated olefins under Cp<sup>\*</sup>Rh(III) catalysis (Scheme 414).<sup>659</sup> Notably, both cyclic and acyclic enol-carbamates were compatible with the conditions. More interestingly, bis-protected estrone derivative only afforded the vinylic C–H olefination product, thus providing an efficient strategy for the highly regioselective functionalization of ketone-containing molecules.


**Scheme 414. Cp<sup>\*</sup>Rh(III)-Catalyzed C–H Olefination of**

**Enol Carbamates with Activated Alkenes**




Successfully, Zhang, Zhong, and co-workers presented an inexpensive ruthenium(II)-catalyzed alkenyl C–H olefination of enol carbamates, affording a diverse array of linear (*Z,E*)-configured butadiene skeletons in modest to high yields (Scheme 415).<sup>660</sup> In this report, the valuable branched (*Z,Z*)-butadienes could be also obtained by the regio- and stereoselective atom-economic *syn*-hydrovinylation of enol-carbamates with internal alkynes under oxidant-free conditions.

**Scheme 415. Ru-Catalyzed Olefinic C–H Alkenylation of Enol Carbamates**




Moreover, Kim's group presented a Cp<sup>\*</sup>Rh(III)-catalyzed regioselective vinylic C–H allylation of enol carbamates with 4-vinyl-1,3-dioxolan-2-ones and allylic carbonates to furnish a diverse array of allylic alcohols and terminal allylated products, respectively (Scheme 416).<sup>661</sup> The authors demonstrated that the efficacious assistance of a carbamoyl directing group enables a highly efficient synthesis of synthetically useful allylated enol carbamates. In addition, the same group also described the alkenyl C–H alkylation of enol carbamates with maleimides under rhodium(III) catalysis (Scheme 417).<sup>662</sup>

**Scheme 416. Rh(III)-Catalyzed Vinylic C–H Allylation of Enol Carbamates with Allylic Carbonates**





**Scheme 417. Rh(III)-Catalyzed Vinylic C–H Alkylation of Enol Carbamates with Maleimides**




More recently, Niu's group reported the regioselective thiocarbamate-directed vinylic C–H amidation of alkenes with dioxazolones enabled by earth-abundant and cost-effective Cp\*Co(III) catalysis under especially mild conditions, exclusively giving rise to a broad array of (*Z*)-selective enamines in up to 99% yield with excellent functional group compatibility (Scheme 418).<sup>663</sup>

**Scheme 418. Thiocarbamate-Directed Cp\*Co(III)-Catalyzed Olefinic C–H Amidation**




**5.4.3. Enol Phosphates.** Enol phosphates often serve as a versatile alternative to the corresponding halides and triflates to engage in the transition-metal-catalyzed coupling reactions because of their high stability and ready availability. Despite the remarkable achievements in phosphorus-containing groups assisted C–H activation reactions,<sup>664</sup> the direct alkenyl C(sp<sup>2</sup>)–H functionalization of enol phosphates is extremely difficult to achieve. To address this issue, Loh and co-workers expanded their olefinic C–H functionalization strategy, and reported in 2015 the unprecedented example of Cp\*Rh(III)-catalyzed coupling reactions of enol phosphates with a broad array of activated alkenes, alkynes as well as allenes, leading to the synthesis of a series of synthetically valuable alkenylated and hydroalkenylated enol phosphates with good regio- and stereoselectivity (Scheme 419).<sup>665</sup>

**Scheme 419. Cp\*Rh(III)-Catalyzed C–H Alkenylation and Hydroalkenylation of Enol Phosphates**




**5.4.4. N-Enoxyphthalimides.** Cyclopropanes are reasonably widespread in numerous biologically related pharmaceuticals and natural products,<sup>666,667</sup> but their stereoselective synthesis remains challenging, especially through the C–H functionalization strategy. In 2014, Rovis and co-workers accomplished a novel Rh(III)-catalyzed cyclopropanation reaction of *N*-enoxyphthalimides with electron-deficient alkenes, producing a diverse array of *trans*-1,2-disubstituted cyclopropanes with modest stereocontrol (Scheme 420).<sup>668</sup> In this report, the authors elegantly designed a new monosubstituted isopropylcyclopentadienyl ligand (Cp<sup>*i*-Pr</sup>) on the Rh(III) catalyst which significantly improves the catalytic efficiency and diastereoselectivity of this transformation. The protocol was proposed to occur via the irreversible activation of vinylic C–H bond followed by two migratory insertion steps including across the electron-deficient olefin to generate the  $\sigma$ -alkylrhodium(III) species and subsequent intramolecular carboration through a 3-exo-trig cyclization process.

**Scheme 420. Rh(III)-Catalyzed Diastereoselective [2 + 1] Annulation of *N*-Enoxyphthalimides and Alkenes**






The development of stereodivergent transformation is definitely of great interest as it enables funneling of a given reaction pathway toward one stereochemical product or another by slightly modifying the reaction parameters. By judicious modulation of the stereoelectronic properties of Rh(III) catalyst and alkene substrates, Rovis and co-workers were able to switch the diastereoselectivity of this Cp<sup>X</sup>Rh(III)-catalyzed cyclopropanation methodology from their *trans*-diastereomers to the corresponding *cis*-congeners (Scheme 421).<sup>669</sup> The authors systematically screened 4,5-dichloro-substituted phthalimides together with a cyclohexyl-Cp-ligated Rh(III) catalyst as the best combination to synthesize a variety of *cis*-cyclopropane scaffolds in decent yields (32–88%) and high diastereoselectivities (up to >20:1 dr).

**Scheme 421. Rh(III)-Catalyzed *cis*-Cyclopropanation of *N*-Enoxyphthalimides and Alkenes**




The cyclopropanation of allylic alcohols is especially difficult to achieve by the well-established Simmons-Smith and diazo decomposition reactions. The Rovis group continued their rhodium(III)-catalyzed cyclopropanation of *N*-enoxyphthalimides with alkenes, and expanded to report a diastereoselective [2 + 1] annulation of allylic alcohols with *N*-enoxyphthalimides. This transformation was greatly promoted by the electron-deficient trifluoromethyl-*tetra*-methyl-cyclopentadienyl (Cp<sup>CF<sub>3</sub></sup>) ligand, yielding the cyclopropanation products which are not accessible by other routes (Scheme 422).<sup>670</sup> Of note, both *trans*- and *cis*-1,2-disubstituted primary allylic alcohols were compatible with this stereospecific protocol. Mechanistic studies demonstrated that the pendant hydroxyl group is indispensable for both reactivity and diastereocontrol. The diastereoselectivity of this transformation probably arises from an intermediate generated from the ring-opening acylation of allylic alcohol. Meanwhile, Rovis and co-worker also achieved a similar cyclopropanation protocol for *N*-enoxyphthalimides and unbiased olefins under modified conditions.<sup>671</sup>

**Scheme 422. Rh(III)-Catalyzed Cyclopropanation of *N*-Enoxyphthalimides with Allylic Alcohols**



The enantioselective and diastereoselective Rovis-cyclopropanation of electron-deficient alkenes with *N*-enoxysuccinimides was accomplished by Cramer's group through a chiral cyclopentadienyl Rh(III)-catalyzed olefinic C–H bond functionalization (Scheme 423).<sup>672</sup> The reaction occurred uneventfully under typically mild and open-flask conditions, yielding diverse chiral cyclopropane products in 49–90% yields with decent stereocontrol. The authors highlighted the potential of this strategy by using this protocol as a key step for the synthesis of oxylipin family of natural products and inhibitor UPC-648.


#### Scheme 423. Chiral Cyclopentadienyl Rh<sup>III</sup>-Catalyzed Enantioselective Cyclopropanation of Electron-Deficient Olefins



More recently, Cramer et al. made further efforts to report the cyclopropanation of *N*-enoxysuccinimides with acrolein enabled by a tailored  $\text{Cp}^*\text{Rh}^{\text{III}}$ -catalyzed vinylic C–H functionalization to obtain diverse disubstituted *cis*-cyclopropanes in good enantio- and diastereoselectivity (Scheme 424).<sup>673</sup> In this report, the bulky ethylcyclohexyl group as C<sub>4</sub> substituent of the second generation  $\text{Cp}^*$  ligand greatly enforced the *cis*-selectivity of the cyclopropanation event. More interestingly, upon treatment with primary amines under  $\text{Cp}^*\text{Ir}^{\text{III}}$  catalysis, the resulting dicarbonyl *cis*-cyclopropanes could undergo an iterative aminative transfer hydrogen process to afford an exquisite set of rigid saturated 3-



azabicyclo[3.1.0]hexanes which are extremely important motifs widespread in biologically active compounds.

#### Scheme 424. $\text{Cp}^*\text{Rh}^{\text{III}}$ -Catalyzed *cis*-Cyclopropanation of Enoxysuccinimides with Acrolein




Apart from the formation of cyclopropane adducts, the reaction of *N*-enoxypthalimides with alkenes could also undergo *syn*-carboamination to generate the alkene difunctionalization products. In this regard, Rovis and co-workers elaborated an unprecedented Rh(III)-catalyzed intermolecular *syn*-carboamination of 1,2-disubstituted alkenes initiated by an alkenyl C–H activation event that employs *N*-enoxypthalimides as the bifunctional source of both carbon- and nitrogen-based functionalities (Scheme 425),<sup>674</sup> thereafter allowing the stereospecific formation of one C–C bond and one C–N bond across an alkene in one step. Of note, the use of sterically hindered *tert*-butyl-tetramethyl-cyclopentadienyl ligand ( $\text{Cp}^{*\text{tBu}}$ ) dramatically alters the inherent chemoselectivity, and enables the formation of the *syn*-carboamination with excellent diastereoselectivity. Key to the success of this protocol is the judicious choice of methanol as a solvent which could significantly promote the phthalimide group to undergo an in situ ring-opening process.

#### Scheme 425. Rh(III)-Catalyzed *syn*-Carboamination of Internal Alkenes



Moreover, Cramer and co-workers continued their elegant enantioselective  $\text{Cp}^x\text{Rh}^{\text{III}}$ -catalyzed alkenyl C–H activation strategy, and further illustrated the intermolecular asymmetric carboamination of electron-deficient acrylates with *N*-enoxysuccinimides at ambient temperature (Scheme 426).<sup>675</sup> A fine-tailored sterically bulky chiral  $\text{Cp}^x$  ligand was proven to be essential for ensuring high reactivity and enabling the reaction towards the intermolecular carboamination process, yielding a diverse array of enantioenriched non-natural  $\alpha$ -amino esters in 49–88% yields and excellent enantiomeric ratios of >99.5:0.5.


**Scheme 426. Enantioselective  $\text{Cp}^x\text{Rh}^{\text{III}}$ -Catalyzed Carboaminations of Acrylates**




**5.5. Ketene Dithioacetals**

Ketene dithioacetals are highly valuable intermediates and versatile synthons which have found widespread applications in the synthesis of diverse functional organic molecules.<sup>676</sup> The direct  $\alpha$ -functionalization of ketene dithioacetals through alkenyl  $\text{C}(\text{sp}^2)$ –H bond functionalization provides a robust tool for the synthesis of highly functionalized ketene dithioacetal derivatives. In 2010, the Yu group first reported a highly regioselective coupling reaction between terminal alkenes and diverse  $\alpha$ -oxoketene dithioacetals to synthesize a wide range of functionalized 1,3-butadienes in presence of  $\text{Pd}(\text{OAc})_2$  (20 mol%) catalyst (Scheme 427).<sup>677</sup> Further condensation of the corresponding 1,3-butadienes by diamines gave rise to biologically-active bicyclic pyridones.

**Scheme 427. Palladium(II)-Catalyzed Oxidative Cross-Coupling of  $\alpha$ -Oxoketene Dithioacetals with Alkenes and Its Proposed Mechanism**




**Proposed mechanism**




The same group further expanded this  $\text{Pd}(\text{II})$ -catalyzed alkenyl  $\text{C}(\text{sp}^2)$ –H olefination to  $\alpha$ -cyanoketene dithioacetal substrates (Scheme 428).<sup>678</sup> A diverse range of electron-deficient alkenes reacted well to yield the highly functionalized linear 1,3-butadienes with air as the terminal oxidant, while styrene derivatives underwent the oxidative cross-coupling to give both linear and branched 1,3-butadienes using  $\text{AgOAc}$  as the oxidant. Interestingly, various unbiased cyclic and internal linear alkene substrates both coupled smoothly in the presence of a catalytic amount of benzoquinone (BQ) in air, generating the skipped 1,4-butadienes.


**Scheme 428. Pd-Catalyzed Oxidative Cross-Coupling of  $\alpha$ -Cyanoketene Dithioacetals with Olefins**




selected out of 17 examples



selected out of 8 examples




selected out of 10 examples




Later, Li's group also achieved the vinylic  $\alpha$ -olefination of ketene dithioacetals with a broad array of styrenes under visible-light photoredox catalysis and external oxidant-free conditions with  $H_2$  gas as the only byproduct (Scheme 429).<sup>679</sup> This atom-economical protocol employed 5 mol% of Mes-Acr<sup>+</sup>ClO<sub>4</sub><sup>-</sup> as the photocatalyst in conjunction with 15 mol% of Co(dmgH)<sub>2</sub>PyCl as a proton-reduction catalyst, yielding a diverse variety of multisubstituted 1,3-dienes in modest to excellent yields.

#### Scheme 429. Visible-Light Photoredox-Catalyzed C–H Olefination of Ketene Dithioacetal with Styrenes





selected out of 35 examples



Moreover, Yu's group demonstrated a palladium(II)-catalyzed oxidative C–H bond allylation of  $\alpha$ -oxoketene dithioacetals with a series of allyl carbonates (Scheme 430).<sup>680</sup> This methodology allows a concise route to diverse highly functionalized skipped 1,4-butadienes in modest to good yields (35–75%).


#### Scheme 430. Oxidative C–H Allylation of Ketene Dithioacetals with Allyl Carbonates



In 2013, Liu and co-workers described a direct C(sp<sup>2</sup>)–H trifluoromethylation of ketene dithioacetals by using PhI<sup>+</sup>CF<sub>3</sub> as the trifluoromethyl source, which is readily generated *in situ* by simply mixing PhI(OAc)<sub>2</sub>, TMSCF<sub>3</sub> and KF (Scheme 431a).<sup>681</sup> The reaction could proceed smoothly at room temperature under metal-free conditions, delivering the trifluoromethylated ketene dithioacetals in moderate to good yields. Following this, the Yu group also introduced an efficient copper-catalyzed alkenyl C(sp<sup>2</sup>)–H trifluoromethylation of  $\alpha$ -oxoketene dithioacetals by using Cu(OH)<sub>2</sub> as the catalyst in combination with TMSCF<sub>3</sub> as the trifluoromethylating agent (Scheme 431b).<sup>682</sup> Both cyclic and acyclic dithioalkyl  $\alpha$ -oxoketene acetals were viable substrates in this strategy, and various synthetically useful substituents were well tolerated. The authors performed detailed mechanistic studies, which clearly demonstrated that the olefinic C–H bond cleavage was not

involved in the rate-determining step, and a radical-involved catalytic mechanism is tentatively proposed on the basis of a TEMPO-quenching experiment of the trifluoromethylation reaction.


### Scheme 431. $\alpha$ -Trifluoromethylation of Ketene Dithioacetals with TMSCF<sub>3</sub>




Besides, the sustainable C–H trifluoromethylation of  $\alpha$ -oxoketene dithioacetals was later realized by employing Ru(bpy)<sub>3</sub>Cl<sub>2</sub> as the photocatalyst and Togni's reagent as the trifluoromethylating reagent under visible-light photoredox catalysis (Scheme 432a).<sup>683</sup> Very recently, Zeng et al. achieved a similar transformation with cheap and bench-stable CF<sub>3</sub>SO<sub>2</sub>Na (Langlois' reagent) as the trifluoromethyl radical source, affording an array of trifluoromethylated  $\alpha$ -oxoketene dithioacetals in modest yields (Scheme 432b).<sup>684</sup>

### Scheme 432. Photoredox Catalysis and Electrochemical Oxidative Trifluoromethylation of $\alpha$ -Oxoketene Ketene Dithioacetals

a) Yang, Xia et al., 2017




b) Zeng et al., 2022




The palladium(II)-catalyzed C–H difluoroalkylation of ketene dithioacetals with bromodifluoroacetates was reported by Wang, Zhu, and co-workers (Scheme 433),<sup>685</sup> which enables the rapid synthesis of a class of CF<sub>2</sub>-containing *tetra*-substituted ketene dithioacetals in decent yields with high functional group compatibility and wide substrate scope. Due to the unique structural properties of  $\alpha$ -oxoketene dithioacetals, the authors proposed an alternative Pd(o)/Pd(I) catalytic cycle involving two single electron transfer (SET) process in this transformation.

### Scheme 433. Pd-Catalyzed C–H Difluoroalkylation of Ketene Dithioacetals with Bromodifluoroacetates




The formation of C–O bonds is one of the fundamental transformations in synthetic organic chemistry. As a consequence, a myriad of synthetically useful methods have been extensively established to forge diverse C–O bonds.<sup>686</sup> The formation of C–O bonds through the activation of inert C–H bonds has particularly aroused much interests in the last few years.<sup>687,688</sup> In 2014, Wang, Liu, and co-workers reported a highly regioselective palladium-catalyzed C(sp<sup>2</sup>)-H acyloxylation between various ketene dithioacetals and carboxylic acids by utilizing PhI(OAc)<sub>2</sub> as an oxidant, delivering a large variety of functionalized vinyl esters with good efficiency and excellent functional group tolerance (Scheme 434).<sup>689</sup>

### Scheme 434. Palladium-Catalyzed Vinylic C–H Bond Acyloxylation of Ketene Dithioacetals



Additionally, the Yu group also demonstrated a Cu(II)-promoted C–H alkoxylation of  $\alpha$ -oxoketene dithioacetals with alcohols by means of a combination of  $\text{PhI}(\text{OAc})_2$  and benzoquinone as the oxidants (Scheme 435).<sup>690</sup> A broad range of alkoxyated ketene dithioacetals were produced with good yields (53–81%). Primary mechanism investigations were carried out. Based on a series of control and kinetic isotope effect (KIE) experiments, a plausible reaction mechanism involving a single-electron-transfer (SET) process was tentatively proposed.


### Scheme 435. Copper-Promoted C–H Alkoxylation of Ketene Dithioacetals and Its Proposed Mechanism




The C–S bond formation reactions of ketene dithioacetals through the direct alkenyl  $\text{C}(\text{sp}^2)\text{--H}$  bond functionalization were also realized by Singh and co-workers in 2015 (Scheme 436a).<sup>691</sup> This operationally simple one-pot reaction proceeded smoothly in the presence of iodine and a copper(I) salt with readily available dimethyl sulfoxide (DMSO) as both the thiomethyl source and solvent, leading to the synthesis of polythiolated alkenes in high yields (75–95%). Later, Liu and co-workers further reported a practical alkenyl C–H sulfenylation/alkyl thiolation reaction of ketene dithioacetals under transition-metal-free conditions (Scheme 436b).<sup>692</sup>

### Scheme 436. Vinyl C–H Sulfenylation/Alkyl Thiolation of Ketene Dithioacetals

a) Singh et al., 2015





b) Liu et al., 2018



Organic thiocyanates are versatile synthetic intermediates in the synthesis of many important pharmaceuticals and other sulfur-containing organic compounds.<sup>693</sup> In 2017, Wang and Xu presented the first transition-metal-free C–H thiocyanation reaction of ketene dithioacetals by using *in situ* generated *N*-thiocyanatosuccinimide (NTS) as the thiocyanation reagent in the presence of *N*-chlorosuccinimide (NCS) and  $\text{NH}_4\text{SCN}$  under ambient conditions (Scheme 437).<sup>694</sup> The methodology accommodated a wide range of olefin substrates to afford SCN-containing ketene dithioacetals with excellent yields. More importantly, the SCN group in the resulting products could be readily converted into diverse sulfur-containing groups, such as  $-\text{SCF}_3$  and thiotetrazole, thus rendering the strategy potentially useful for the discovery of new drug candidates.


### Scheme 437. Transition-Metal-Free Alkenyl C–H Bond Thiocyanation of Ketene Dithioacetals



Apart from the oxidative C–O, C–S cross-couplings, the direct alkenyl C–H phosphorylation of ketene dithioacetals for the construction of C–P bonds was also efficiently achieved under metal-free or  $\text{AgNO}_3$ -mediated conditions (Scheme 438).<sup>695</sup> Gratifyingly, both *H*-phosphonates and *H*-phosphine oxides were compatible in this reaction with yields ranging from 38% to 97% with good regioselectivity. Further elaboration of the resulting phosphorylated products with hydrazine hydrate could provide an




alternative method for the synthesis of phosphorylated pyrazole derivative. The authors performed kinetic isotope effect (KIE) study to gain more insights, and the results clearly suggested that the rupture of the vinylic C–H bond should be not involved in the turnover-limiting step. Radical trapping experiments indicated that a radical pathway might be probably involved in this process.

### Scheme 438. Transition-Metal-Free Alkenyl C–H Bond Phosphorylation of Ketene Dithioacetals




Successfully, Yu and co-workers developed an iron-catalyzed C–H alkylation of  $\alpha$ -oxoketene dithioacetals by using styrenes as the alkylating reagents, providing a variety of functionalized ketene dithioacetal derivatives in modest to high yields (Scheme 439a).<sup>696</sup> Later, Yu and Lei independently described the direct oxidative C–H alkylation of ketene dithioacetals with simple ethers and toluene derivatives using di-*tert*-butyl peroxide (DTBP) as the terminal oxidant (Scheme 439b and c).<sup>697,698</sup> However, a catalytic amount of FeCl<sub>3</sub> (10 mol%) and DABCO·6H<sub>2</sub>O (10 mol%) were both indispensable in Yu's strategy in order to obtain high yields (55–88%).

### Scheme 439. Direct C–H Bond Alkylation of Ketene Dithioacetals




Subsequently, Yu's group further disclosed another example of iron-catalyzed olefinic C–H alkylation of  $\alpha$ -oxoketene dithioacetals with various cyclobutanone oxime esters via a ring-opening radical C–C bond cleavage strategy under redox-neutral conditions (Scheme 440).<sup>699</sup> Interestingly, the C–H alkoxylation of  $\alpha$ -oxoketene dithioacetals could be also achieved when conducting the reaction under an O<sub>2</sub> atmosphere in the presence of CuCl<sub>2</sub> as the catalyst.

### Scheme 440. Direct C–H Alkylation and Alkoxylation of Ketene Dithioacetals




Moreover, Yu and co-workers extended their investigations to illustrate an efficient visible-light-induced alkenyl C(sp<sup>2</sup>)-H arylation of *S,S*-functionalized internal alkenes that is,  $\alpha$ -oxoketene dithioacetals, using aryldiazonium salts (ArN<sub>2</sub>BF<sub>4</sub>) as the coupling partners in conjunction with Ru(bpy)<sub>3</sub>Cl<sub>2</sub>·6H<sub>2</sub>O as the photosensitizer at ambient temperature (Scheme 441),<sup>700</sup> offering an environmental-friendly strategy to synthesize all-carbon tetrasubstituted alkenes including anticancer drug tamoxifen.

### Scheme 441. Photoredox-Catalyzed C–H Arylation of Ketene Dithioacetals




One year later, Li and co-workers disclosed an efficient chemodivergent cross-coupling reaction of  $\alpha$ -oxo ketene dithioacetals with diazo compounds by the catalyst-controlled alkenyl C–H activation/functionalization (Scheme 442).<sup>701</sup> The ruthenium(II)-catalyzed C–H activation exclusively occurred at the  $\alpha$ -position of ketene dithioacetals, and 1:2 coupling with  $\alpha$ -diazoketoesters where dimerized carbene was identified as a key intermediate, leading to the formation of C–H furfurylation products. In contrast, the  $\text{Cp}^*\text{Rh}(\text{III})$ -catalyzed C–H activation of  $\alpha$ -benzoyl ketene dithioacetals occurred via both aryl and alkenyl C–H bond cleavage to produce the [4 + 2] annulation naphthalenone derivatives. In a subsequent manuscript, the same group further described the example of silver-catalyzed vinylic C–H bond functionalization of  $\alpha$ -oxo ketene dithioacetals with tertiary propargylic carbonates through a cascade nucleophilic addition and intramolecular Friedel–Crafts alkylation process (Scheme 443).<sup>702</sup> This novel protocol features a broad substrate scope, operational simplicity and simple reaction conditions, providing a powerful method to the synthesis of various functionalized indenes with moderate to good yields (30–85%).

### Scheme 442. Ruthenium- and Rhodium-Catalyzed Chemodivergent Couplings of Ketene Dithioacetals and $\alpha$ -Diazo Ketones





### Scheme 443. Ag(I)-Catalyzed Tandem Nucleophilic Addition and Friedel–Crafts Alkylation between Ketene Dithioacetals and Propargyl Carbonates



More recently, Xiao and colleagues demonstrated that the annulative coupling reaction of  $\alpha$ -aroyl ketene dithioacetals with diazo compounds under  $\text{Cp}^*\text{Rh}(\text{III})$  catalysis via aryl and alkenyl C–H bond functionalizations exclusively furnish the highly functionalized indanone derivatives bearing an all-carbon quaternary stereocenter at the  $\beta$ -position (Scheme 444),<sup>703</sup> which is quite different from that of [4 + 2] annulation to afford naphthalenone derivatives as demonstrated in Li's work.<sup>702</sup>


### Scheme 444. Synthesis of $\beta$ -Quaternary Indanones via Alkenyl C–H Activation



## 6. MISCELLANEOUS ALKENYL C–H BOND FUNCTIONALIZATIONS



In 2011, Liu and co-workers achieved a Pd(II)-catalyzed cross-coupling between glycals and activated olefins under benign conditions (Scheme 445).<sup>704</sup> This strategy provides a convenient access to C<sub>2</sub>-functionalized glycals, which are conventional intermediates in natural products and biologically important structures. The substrate scope features a variety of sugars, protecting groups and substituents on the olefins. The desired products were obtained in decent yields with excellent *E*-selectivity.

### Scheme 445. Palladium(II)-Catalyzed Cross-Coupling of Glycals with Activated Olefins




Around the same time, Toste, Bergman, and co-workers developed a straightforward one-pot Co(I)-catalyzed C–H functionalization and Michael addition of alkenes to synthesize the corresponding cyclic tetraalkyl-substituted,  $\gamma,\delta$ -unsaturated compounds in good to excellent yields (Scheme 446).<sup>705</sup> The postulated intermediates in the proposed mechanism were characterized. The catalytic turnover for the base and active catalyst was also documented in this investigation.

### Scheme 446. Cobalt-Dinitrosyl Mediated Vinylic C–H Functionalization Reaction





Later, Dong's research group illustrated an efficacious Cp\*Rh(III)-catalyzed alkenyl C(sp<sup>2</sup>)-H of 7-azaindoles with internal alkynes to stereoselectively construct dienes and polycyclic heteroaromatic ring products (Scheme 447).<sup>706</sup> This groundbreaking method was significantly influenced by the reaction conditions and the substituents on the vinylic moiety.

### Scheme 447. Cp\*Rh(III)-Catalyzed Vinylic C(sp<sup>2</sup>)-H of 7-Azaindoles with Alkynes




Almost at the same time, Cook's group elaborated a direct synthesis of 2-aminopyridines by a Ru(II)-catalyzed alkenyl C–H functionalization followed by an amide alcoholysis (Scheme 448).<sup>707</sup> This strategy featured a highly chemo- and regioselective synthesis of the pyridine ring due to a site-specific C–H activation of the substrates. An array of multisubstituted 2-aminopyridines were obtained in decent yields. Mechanistic investigations shed light on the key oxidative C–H activation/cyclization processes.

### Scheme 448. Ru(II)-Catalyzed C–H Functionalization for the Synthesis of 2-Aminopyridines





In the following year, Zhu and colleagues illustrated a  $\text{Cp}^*\text{Rh(III)}$ -catalyzed olefinic C–H functionalization for the cross-coupling of oxadiazoles with alkynes for the construction of 2-acylamino and 2-amino pyridines derivatives which are crucial heterocyclic core structures for many natural products and pharmaceuticals containing a reactive functional group. By alternating the solvent, selective protection/deprotection of amino groups can be achieved. Notably, the reaction features exceptional functional group tolerance, excellent yields and regioselectivity (Scheme 449).<sup>708</sup>

**Scheme 449. Rhodium(III)-Catalyzed Cross-Coupling of Oxadiazoles with Alkynes**




Meanwhile, the same group also disclosed a fascinating associative covalent relay process through a  $\text{Cp}^*\text{Rh(III)}$ -catalyzed oxadiazolone-directed olefinic C–H cross-coupling with alkynes, providing an efficient access to structurally diverse 2-aminepyridines (Scheme 450).<sup>709</sup> Although the mechanism was justified by previous works, the proposed relay formalism in this report provided a mechanistic conceptual framework for investigations.

**Scheme 450. Rhodium(III)-Catalyzed Oxadiazolone-Directed Olefinic C–H Cross-Coupling with Alkynes**





Yang's group in 2019 elaborated a palladium(II)-catalyzed Mizoroki-Heck type reaction to construct a diverse of mono- or diaryl-substituted phosphine ligands with excellent yields and diastereoselectivity (Scheme 451).<sup>710</sup> Mechanistic investigations suggested that this P=O directed protocol goes through a cationic Heck reaction promoted by the Ag salt.

**Scheme 451. Pd(II)-Catalyzed Mizoroki-Heck Type Reaction to Synthesize Phosphine Ligands**



Recently, an innovative approach for producing cycloalkeno[*c*]-fused pyridines has been established by Jiang, Zhang, and their co-workers (Scheme 452).<sup>711</sup> By coupling *N*-acetyl hydrazones bearing medium to large cycloalkenes with vinyl azides, a variety of regioselective fused pyridines can be obtained. A broad scope of 41 examples was presented in up to 81% yield.


**Scheme 452. Pd-Catalyzed Alkenyl C–H Activation and Cyclization of *N*-Acetyl Hydrazones of Acylcycloalkenes with Vinyl Azides**




Heterocyclic alkenes are an important class of feedstocks and synthetic valuable synthons for the synthesis of biologically important heterocyclic scaffolds. Remarkable advances on the direct vinylic C-H functionalization of heterocyclic alkenes have been made over the decades.<sup>712</sup> For selected examples, Georg's group used uracils as alkene substrates to couple with acrylates or similarly electron-deficient olefins (Scheme 453a).<sup>713</sup> Pd(OAc)<sub>2</sub>/AgOAc was the best catalyst/oxidant combination for this reaction. Of note, the additive was essential to improve the efficiency of reaction, and PivOH was found to be the best choice. Moreover, Hong et al. reported the coupling of coumarin with acrylates and styrenes, where Pd(OPiv)<sub>2</sub> was used as the catalyst precursor with 1 atm of O<sub>2</sub> as the oxidant. The reaction was regioselective, and occurred at the C<sub>3</sub>-position of coumarins (Scheme 453b).<sup>714</sup>

### Scheme 453. Selected Examples of Vinylic C-H Bond Olefination of Heterocyclic Alkenes

#### a) Georg et al., 2013




#### b) Hong et al., 2013



The group of Castarlenas and Oro disclosed the first example of cross-coupling between *N*-vinylpyrazoles and alkynes via directed vinylic C-H activation, resulting in the formation of Markovnikov-selective butadienyl pyrazoles in decent yields with moderate to good regioselectivities (Scheme 454).<sup>715</sup> The reaction occurred smoothly in the presence of rhodium catalyst bearing a carbene and *cis*-cyclooctene ligand which is crucial for this hydrovinylation of the electron-rich nitrogenated alkenes. Notably, this

protocol allows the rapid synthesis of diverse conjugated acyclic trienes through a cascade alkyne dimerization and hydrovinylation sequence in a one-pot fashion.


### Scheme 454. Rh-Catalyzed Cross-Coupling between *N*-Vinylpyrazoles and Alkynes




Later, Ackermann's group illustrated a Cp\*Co(III)-catalyzed reaction of *N*-vinylpyrazoles with isocyanates, affording the vinylic C-H aminocarbonylation products in moderate yields (Scheme 455a).<sup>716</sup> Subsequently, Ellman and co-workers demonstrated a highly efficient, stereoselective C(sp<sup>2</sup>)-H bond addition across activated alkenes and polarized π-bonds (Scheme 455b).<sup>717</sup> Both aromatic and alkenyl C-H bonds can undergo this three-component addition cascade in this case. Notably, enones, aldehydes, as well as *N*-tertbutanesulfinyl imines are all compatible with this protocol.

### Scheme 455. Cp\*Co(III)-Catalyzed Alkenyl C-H Bond Functionalization of *N*-Vinylpyrazoles

#### a) Ackermann et al., 2015




#### b) Ellman et al., 2016




Recently, Iwasawa et al. achieved the Rh(III)-catalyzed alkenyl C-H carboxylation of alkenylpyrazoles (Scheme 456).<sup>718</sup> By the combination of RhCl<sub>3</sub>·3H<sub>2</sub>O catalyst, trimesitylphosphine ligand, and AlMe<sub>2</sub>(OMe) additive in DMA at 80 °C, the alkenyl C-H bonds of both cyclic and acyclic alkenylpyrazoles was smoothly carboxylated in decent yields under CO<sub>2</sub> atmosphere. Notably, the carboxylation reaction proceeded smoothly on a preparative scale even with a decreased catalyst loading (1.0 mol%) at lower reaction temperature.

### Scheme 456. Rhodium(III)-Catalyzed Alkenyl C–H Carboxylation of *N*-Vinylpyrazoles



Moreover, Ackermann and colleagues expanded to employ imidazole as the directing group, and established a general protocol for the Ru(II)-catalyzed electrochemical dehydrogenative annulation reaction of alkenyl imidazoles with internal alkynes, leading to the formation of a variety of bridgehead *N*-fused [5,6]-bicyclic heteroarenes (Scheme 457).<sup>719</sup> In this report, the authors identified the unprecedented azaruthenabicyclo[3.2.0]heptadienes as key intermediates which were unambiguously characterized by X-ray analysis. It is noteworthy that this ruthenaelectro-catalyzed dehydrogenative C–H/N–H annulation protocol was not restricted to alkenyl imidazoles, a broad array of 2-arylimidazoles could be smoothly annulated with both symmetrical and unsymmetrical internal alkynes under modified conditions. Detailed mechanistic investigations are suggestive of an oxidation-induced reductive elimination pathway within a ruthenium(II/III) manifold.


### Scheme 457. Ru(II)-Catalyzed Electrochemical Dehydrogenative Annulation of Alkenyl Imidazoles with Internal Alkynes



### 7. ALKENYL C–F BOND FUNCTIONALIZATIONS OF GEM-DIFLUOROALKENES

Fluoroalkenes represent an extremely important class of privileged structural motifs, which have found widespread applications in pharmaceutical and material sciences (Scheme 458).<sup>720–727</sup> For example, tetrafluorinated ethylene (TFE) has been used as the monomer for the synthesis of Teflon. The ability to access various  $\alpha$ -fluoroalkenes will certainly facilitate the development of novel drugs and functional materials. Accordingly, there have been much effort directed towards the development of novel synthetic methods to obtain this class of compounds including the reactions using fluorinating reagents with vinylic halides or metals. However, most of the reported protocols suffer from the need to use pre-activation substrates or non-readily available starting materials, low regio- or stereo-selectivity as well as poor functional group compatibility due to the use of sensitive reagents.


### Scheme 458. Representative Examples of Mono- and Difluorovinyl-Containing Bioactive Molecules



Indeed, *gem*-Difluoroalkenes are synthetically valuable fluorinated scaffolds frequently encountered in numerous biologically active pharmaceuticals and agrochemicals. The judicious use of readily prepared *gem*-difluoroalkenes as starting materials to generate alkenyl fluorinated metal complexes through alkenyl C–F functionalization or as electrophiles offers a highly efficient and operationally simple approach for the incorporation of  $\alpha$ -fluoroalkenyl motifs onto structurally diverse organic molecules. As a consequence, the cleavage, activation, and further functionalization of the alkenyl C(sp<sup>2</sup>)-F bond of *gem*-difluoroalkenes have received considerable attention in recent years.<sup>18–26</sup> In this section, we systematically

summarized the recent advances on the alkenyl  $sp^2$  C–F bond functionalization methods mainly focusing on the use of readily available *gem*-difluoroalkenes as starting materials. Moreover, polyfluoroalkenes such as tetrafluoroethylene (TFE) and its analogues will also be covered. In contrast to alkenyl  $sp^2$  C–H bond functionalization, we will organize this section according to the cross-coupling partners (Scheme 459).


### Scheme 459. General Scheme of Alkenyl C–F Bond Functionalizations of *gem*-Difluoroalkenes




#### 7.1. C–H Bond Formation

Monofluoroalkenes are versatile fluorinated building blocks with widespread applications in medicinal chemistry and material sciences, which can also be used as an important fluorinated synthon for further synthetic transformations.<sup>728</sup> Hydrodefluorination (HDF) generally provides the simplest synthetic transformation of C–F bonds,<sup>729</sup> and the hydrodefluorination of *gem*-difluoroalkenes allows the straightforward synthesis of monofluoroalkenes. Early in 1979, Burton's group first reported the reduction of *gem*-difluoroalkenes in the presence of commercially available sodium bis(2-methoxyethoxy)aluminumhydride (Red-Al<sup>®</sup>) as the efficacious reductant, yielding the  $\beta$ -fluoroalkenes in high yields with up to 93:7 stereoselectivity (Scheme 460a).<sup>730</sup> Following this, Wu and Cao in 2015 also reported a general and efficient method for the selective hydrodefluorination of *gem*-difluoroalkenes by utilizing Red-Al<sup>®</sup> as the reducing agent without the use of any base and catalyst under mild conditions (Scheme 460b).<sup>731</sup> A large variety of monofluoroalkenes were synthesized in modest to excellent yields with decent *E*-selectivity.

### Scheme 460. Synthesis of Monofluoroalkenes through Hydrodefluorination of *gem*-Difluoroalkenes




b) Wu, Cao et al., 2015





By treatment of 3,3-difluoro allylic alcohols with methyllithium followed by reduction with LiAlH<sub>4</sub>, Tellier and co-workers were able to synthesize a series of monofluoroalkenyl alcohols in a stereoselective manner (Scheme 461a).<sup>732,733</sup> Later, Paquin's group successfully achieved a synthetic route to *trans*- $\beta$ -fluorostyrenes from (*Z*)-1-aryl-2-fluoro-1-(trimethylsilyl)ethenes by employing LiEt<sub>3</sub>H as the reductant followed by a stereospecific removal of the silyl group in the presence of water and TBAF (Scheme 461b).<sup>734</sup>

### Scheme 461. Synthesis of Monofluoroalkenes through Selective Reduction of *gem*-Difluoroalkenes

a) Tellier et al., 1995





b) Paquin et al., 2011



By means of water as the proton source, *gem*-difluoroalkenes can undergo a stereoselective and regioselective copper-catalyzed hydrodefluorination


reaction through vinylic C–F activation to form a diverse array of di- and tri-substituted *Z*-fluoroalkenes (Scheme 462).<sup>735</sup> A wide range of substrates tolerated in this strategy, including aliphatic, aromatic,  $\alpha,\beta$ -unsaturated, and substituted *gem*-difluoroalkenes. More importantly, a series of *gem*-difluoroalkenes derivatived from natural products were also able to react smoothly under the mild conditions to generate the corresponding late-stage diversification products in excellent yields.

### Scheme 462. Synthesis of *Z*-Monofluoroalkenes via Hydrodefluorination of *gem*-Difluoroalkenes





Around the same time, Ito and co-workers achieved a stereodivergent hydrodefluorination reaction of *gem*-difluoroalkenes (Scheme 463).<sup>736</sup> This method stereoselectively afforded both (*Z*)- or (*E*)-terminal monofluoroalkenes from the same starting material. In the presence of (Bpin)<sub>2</sub>, *Z*-product was formed after deborylation, due to the specific conformation of key intermediate that avoids steric repulsion between bulky Bpin and aryl group. In contrast, the use of copper(I)/hydrosilane catalytic system exclusively afforded an array of *E*-monofluoroalkenes due to the electronic repulsion between F and aryl group in the elimination transition state of copper(I)-hydride adducts.

### Scheme 463. Copper-Catalyzed Stereodivergent Hydrodefluorination of *gem*-Difluoroalkenes



Several years later, Tsui's group described a highly stereoselective palladium-catalyzed hydrodefluorination reaction of tetrasubstituted *gem*-difluoroalkenes using dimethylphenylsilane (Me<sub>2</sub>PhSiH) as the "H source" (Scheme 464).<sup>737</sup> A series of trisubstituted terminal (*E*)-monofluoroalkenes can be obtained in 52–99% yields with excellent *E/Z* selectivity (>99:1). The stereocontrol of this reaction was proposed to be influenced by the ester-directing C–F bond oxidative addition step in the catalytic mechanism.

### Scheme 464. Pd-Catalyzed Hydrodefluorination of Tetrasubstituted *gem*-Difluoroalkenes




## 7.2. C–C Bond Formation

### 7.2.1. C–H/C–F Functionalization Reaction.

As mentioned above, fluoroalkenes are important core structures which are widely found in medicinal molecules. Nevertheless, the synthetic pathway to this class of compounds is very sparse, previous strategies often suffer from poor atom-economy or harsh reaction conditions. By using appropriate directing groups, *gem*-difluoroalkenes can readily engage in transition metal-catalysed C–F/C–H functionalization reactions, thus providing an efficient route to monofluoroarylalkenes. Various transition-metal catalysts have been used to realize this C–C bond coupling reaction via alkenyl C–F activation in recent years.


Seminal example of such process was first disclosed by Loh's group in 2015. They pioneered an impressive Cp<sup>\*</sup>Rh(III)-catalyzed C–H/C–F activation for the synthesis of (hetero)arylated monofluoroalkenes (Scheme 465).<sup>738</sup> Exploiting the abundant nature of *gem*-difluoroalkenes as electrophiles, this strategy presented a highly effective and convenient way to introduce  $\alpha$ -fluoroalkenyl moieties onto (hetero)aryls without the presence of an oxidant. In addition, the use of an alcoholic solvent and the generation of HF in the reaction was established to be advantageous towards the outcome of the reaction, suggesting the likelihood of a hydrogen-bonding interaction during the cleavage of the C–F bond.

### Scheme 465. Synthesis of Monofluoroalkenes through Cp<sup>\*</sup>Rh(III)-Catalyzed C–H/C–F Activation




Following this, Loh's group extended to report the [4 + 1] annulation of acrylamides and *gem*-difluoroacrylates to produce 5-methylene-1*H*-pyrrol-2(5*H*)-one derivatives through a Cp<sup>\*</sup>Rh(III)-catalyzed alkenyl C–H activation combined with twofold C–F bond cleavage (Scheme 466).<sup>739</sup> This stereospecific strategy was similar to the formal dehydrogenative alkylidene carbene insertion reaction. To identify the key intermediate of this process, the authors performed control experiments. By reacting acrylamide with *gem*-difluoroacrylate in the presence of [RhCp<sup>\*</sup>(MeCN)<sub>3</sub>](SbF<sub>6</sub>)<sub>2</sub> catalyst and NaOAc, the expected annulation product was produced along with an acyclic monofluoroacrylate. Elevating the reaction temperature to 80 °C, the acyclic monofluoroacrylate was readily converted into the annulation product, which clearly confirmed that this protocol should undergo a stepwise quasinucleophilic displacement route. In the same year, Wang and co-workers achieved a similar heteroannulation reaction by using a combination of Rh(III) and Ag(I) in a relay catalysis process.<sup>740</sup> The OTs group was used instead of the CO<sub>2</sub>R group, which allowed further synthetic transformations via traditional cross-coupling.

### Scheme 466. Cp<sup>\*</sup>Rh(III)-Catalyzed [4 + 1] Annulation of Acrylamides with *gem*-Difluoroacrylates





The direct synthesis of fluorinated heterocycles was achieved by Wang's group with 2,2-difluorovinyl tosylate as starting material through a Cp<sup>\*</sup>Rh(III)-catalyzed C(sp<sup>2</sup>)–H functionalization of arenes and alkenes (Scheme 467).<sup>741</sup> In this process, *N*-substituted benzamides can be coupled with 2,2-difluorovinyl tosylate to produce a range of fluorinated quinoline-1(2*H*)-ones. Interestingly, reactions between an alcohol and *N*-OPiv benzamides gave *gem*-difluorinated dihydroisoquinolin-1(2*H*)-ones, while reactions with *N*-OMe benzamides afforded acyclic monofluorinated alkenes which could be further cyclized in the presence of a Brønsted acid (H<sub>2</sub>SO<sub>4</sub>) to produce 4-fluoroisoquinolin-1(2*H*)-ones. Both reactions involved a seven-membered rhodacycle species as the key intermediate.

### Scheme 467. Synthesis of Fluorinated Heterocycles via Rhodium-Catalyzed Coupling of Benzamides with 2,2-Difluorovinyl Tosylate





Subsequently, Loh and colleagues further demonstrated that *N*-pyrimidylindoles undergoing C–H alkylation with  $\alpha,\alpha$ -difluorovinyl tosylate in the presence of Cp<sup>\*</sup>Rh(III) catalyst produced a broad range of C<sub>2</sub> alkylated indoles (Scheme 468).<sup>742</sup> In this case, the tosyl difluorinated enol ether acted as an unconventional substitute for an acetate building block.

### Scheme 468. Rh(III)-Catalyzed Alkylation of Indoles and $\alpha,\alpha$ -Difluorovinyl Tosylate



Moreover, Loh and co-workers investigated the  $\text{Cp}^*\text{Rh(III)}$ -catalyzed defluorinative C–H vinylation of acrylamides with *gem*-difluoroolefins in order to construct a variety of 2-fluoro-1,3-dienes (Scheme 469).<sup>743</sup> An initial alkenyl  $\text{C}(\text{sp}^2)\text{--H}$  bond activation accompanied by nucleophilic addition and  $\beta$ -fluoride elimination gave the coupling products in decent yields with exceptional stereocontrol. Mechanistic studies illustrated that due to the peculiar effects of the fluorine substituent, the reactivity of this transformation was not observed for analogues containing heavier halide atoms.



### Scheme 469. $\text{Cp}^*\text{Rh(III)}$ -Catalyzed Defluorinative C–H Vinylation of Acrylamides with *gem*-Difluoroalkenes



However, *N*-nitrosoanilines and 2,2-difluorovinyl tosylates undergo regio- and stereoselective  $\alpha$ -fluoroalkenylation under  $\text{Cp}^*\text{Rh(III)}$  catalysis to give *ortho*-substituted products in basic conditions (Scheme 470).<sup>744</sup> An array of *Z*-monofluoroalkenes were obtained in modest to excellent yields via chelation-assisted C–H bond



activation, migratory insertion, and  $\beta$ -F elimination. In this report, the nitrosoaniline directing group was also able to give *ortho*-substituted products in high regioselectivity. Many of such reactions involve  $\text{Ca}^{2+}$  additives but the potential role of this cation was not mentioned. However, it is possible that the high affinity of  $\text{Ca}^{2+}$  for  $\text{F}^-$  makes it easier to remove the fluoride ion from the reaction, facilitate  $\beta$ -F elimination and regenerate the active  $\text{Cp}^*\text{Rh(III)}$  catalyst.

### Scheme 470. Rh(III)-Catalyzed $\alpha$ -Fluoroalkenylation of *N*-Nitrosoanilines with 2,2-Difluorovinyl Tosylates



Generally, the use of native directing group sometimes substantially increases the overall efficiency of the reaction as it remarkably avoids the pre-installation and subsequent removal of the auxiliary. Recently, Peng and colleagues made use of the quinazolinone ring as the inherent coordinating group to produce a broad array of 2-(*o*-monofluoroalkenyl)quinazolinones by means of 2,2-difluorovinyl tosylate as the monofluoroolefin synthon (Scheme 471).<sup>745</sup>


### Scheme 471. Rh-Catalyzed C–H $\alpha$ -Fluoroalkenylation of 2-Arylquinazolinones with 2,2-Difluorovinyl Tosylate



Fu and Xiao et al. reported the  $\text{Cp}^*\text{Rh(III)}$ -catalyzed C–H activation of 1-arylpyrazoles and further cross-coupling with methyl trifluoroacrylate for the straightforward


synthesis of *E*-monofluoroalkenes (Scheme 472).<sup>746</sup> Notably, complex heterocycles can be obtained by this strategy as the reaction between  $\beta$ -carboxymethyl  $\alpha,\alpha$ -difluoroalkene and 2-phenylpyridines afforded a broad range of benzoindolizines by removing three F-atoms in a single step. This protocol can be applied to complex bioactive molecules such as purine and estrone derivatives to give the corresponding products in 51% and 80% yield, respectively.

### Scheme 472. Cp\**Rh*(III)-Catalyzed C–H Activation of 1-Arylpyrazoles with Methyl Trifluoroacrylate





In addition, Li's group successfully developed the Cp\**Rh*(III)-catalyzed benzylic C(sp<sup>3</sup>)-H fluoroalkenylation of 8-methylquinoline derivatives with *gem*-difluoroalkenes under redox-neutral conditions (Scheme 473).<sup>747</sup> The mechanism is similar to the other reactions in this section, and likewise gave *Z*-monofluoroalkenes in high regio- and stereoselectivity.

### Scheme 473. *Rh*(III)-Catalyzed Fluoroalkenylation of 8-Methylquinolines with *gem*-Difluoroalkenes



Ru catalysts can be also used to couple *gem*-difluoroalkenes with indoles at the C2 position assisted by a 2-pyrimidinyl directing group (Scheme 474).<sup>748</sup> The reaction is redox-neutral and uses both C–H bond activation and C–F bond cleavage to produce a variety of 1,2-diarylstubstituted monofluoroalkenes with high stereo- and regio-selectivities and *Z*-monofluoroalkenes as the major product. In this  $\alpha$ -fluoroalkenylation reaction, the 2-pyridyl and 2-pyrimidinyl groups were essential to achieve excellent selectivity. Mechanistically, the Ru(II) catalyst undergoes cyclometallation with *N*-pyrimidinylindole to give a five-membered metalacyclic intermediate I. *gem*-Difluoroalkene then coordinates to the intermediate to form a Ru–C(alkyl) complex II, which then regioselectively undergoes migratory insertion to afford a seven-membered Ru complex III. The electronic structure of the *gem*-difluoroalkene controls the migratory insertion reaction, by having the nucleophilic aryl group add to the electrophilic fluorinated position. The Ru is added to the  $\beta$ -position relative to F as a result of this reaction. This intermediate then selectively undergoes  $\beta$ -F elimination through a *syn*-coplanar transition state to give the product and Ru(II) fluoride. Finally, anion exchange can be performed with a chloride to regenerate the Ru(II) catalyst.


### Scheme 474. Ru(II)-Catalyzed $\alpha$ -Fluoroalkenylation of Arenes via C–H Activation and C–F Cleavage



Ji's group illustrated a general and efficient example of  $\alpha$ -fluoroalkenylation of oxime ethers with various *gem*-difluorostyrenes through a Ru(II)-catalyzed C–H/C–F bond cleavage strategy (Scheme 475).<sup>749</sup> Remarkably, the


alkenyl moiety in the resulting products existed exclusively in the *Z*-configuration. This method worked with a variety of substrates along with good tolerance of functional groups. The proposed mechanism was ascertained through the identification of a cycloruthenated intermediate. The *O*-methyl oximyl-directing group can be removed with ease to acquire  $\alpha$ -fluoroalkenylated acetophenones.

#### Scheme 475. Ru(II)-Catalyzed $\alpha$ -Fluoroalkenylation of Oxime Ethers with *gem*-Difluorostyrenes



Coupling reactions between *gem*-difluoroalkenes and substituted indoles via the sustainable cobalt-catalyzed chelation-assisted C–H/C–F functionalizations are also accessible. Li and co-workers in 2016 described the first example of  $\text{Cp}^*\text{Co}(\text{III})$ -catalyzed  $\alpha$ -fluoroalkenylation of diverse (hetero)arenes with *gem*-difluorostyrenes under redox-neutral conditions (Scheme 476).<sup>750</sup> The reaction proceeded smoothly to furnish a series of 1,2-diaryl-substituted monofluoroalkenes in moderate to good yields with excellent *Z*-selectivity.


#### Scheme 476. Co(III)-Catalyzed $\alpha$ -Fluoroalkenylation of Heteroarenes with *gem*-Difluorostyrenes



Subsequently, Ackermann's group elaborate an efficient room-temperature C–F/C–H functionalization reactions


enabled by user-friendly  $\text{Cp}^*\text{Co}(\text{CO})\text{I}_2$  catalyst, affording an array of monofluorinated heteroarenes in excellent yields (84–99%) with high diastereoselectivity (Scheme 477).<sup>751</sup> This alkenylative reaction occurred under exceedingly mild conditions via *syn* Co–F elimination to produce the corresponding *Z*-fluoroalkenes. Extensive mechanistic studies suggested a fast cleavage of the C–F bond involved in this process.

#### Scheme 477. Room-Temperature Co(III)-Catalyzed C–H/C–F Bond Functionalizations of Indoles with *gem*-Difluoroalkenes



The  $\text{Cp}^*\text{Co}(\text{III})$ -catalyzed alkenyl C–H functionalization of biologically active 6-arylpurine derivatives accompanied by selective olefinic C–F bond cleavage of *gem*-difluoroalkenes was reported by Yoshino and Matsunaga (Scheme 478).<sup>752</sup> Such reactions employed the purine moiety as a native directing group for the C–H activation process to generate a range of monofluoroalkenes with satisfactory *Z*-selectivity.


#### Scheme 478. Co(III)-Catalyzed $\alpha$ -Fluoroalkenylation of 6-Arylpurines with *gem*-Difluoroalkenes



Loh and colleagues explored the use of less toxic Mn catalysts to perform C–H bond activation/C–F bond


cleavage reactions (Scheme 479).<sup>753</sup> However, unlike previous reactions, this protocol produces the thermodynamically unfavourable *E*-monofluoroalkene as the major product, without the need of any additional oxidants. This is probably due to the selective  $\beta$ -defluorination from the key intermediate. Despite this, reactions involving 1-phenyl-pyrazole delivered *Z*-monofluoroalkenes, and sterically hindered *gem*-difluoroalkenes similarly selectively gave the *Z*-products.

### Scheme 479. Synthesis of Monofluoroalkenes via Mn-Catalyzed C–H Activation and C–F Cleavage




Meanwhile, the C–H/C–F functionalization of *gem*-difluoroalkenes and perfluoroalkenes were also performed by using a commercially available  $\text{Mn}(\text{CO})_5\text{Br}$  catalyst to produce monofluorinated and polyfluorinated alkenes respectively (Scheme 480).<sup>754</sup> These reactions occurred with high chemo-, positional- and diastereo-selectivity. In this case, the 2-pyridyl auxiliary could be easily removed by the treatment with  $\text{NaSPh}$  and  $\text{MeOTf}$ , affording free (*NH*)-indoles.

### Scheme 480. Mn(I)-Catalyzed C–H/C–F Activation of (Hetero)arenes with *gem*-Difluoroalkenes and Perfluoroalkenes





In 2015, Cao et al. reported a straightforward metal-free C–H  $\alpha$ -fluorovinylation of azole heterocycles, including 2-phenyl-1,3,4-oxadiazole, benzothiazole and benzoxazole with diverse *gem*-difluoroalkenes through a room-temperature nucleophilic vinylic substitution ( $\text{S}_{\text{N}}\text{V}$ ) with the assistance of  $\text{KHMDS}$  or  $\text{NaH}$  (Scheme 481).<sup>755</sup>

### Scheme 481. Base-Mediated $\alpha$ -Fluoroalkenylation of Azole Heterocycles with *gem*-Difluoroalkenes






In a recent report, Zhou and Yi et al. illustrated an effective and metal-free [3 + 2] annulation reaction of *N*-phenoxy amides with *gem*-difluoroalkenes through a  $\text{Cs}_2\text{CO}_3$ -promoted tandem [3,3]-sigmatropic rearrangement, providing a straightforward approach to assemble a variety of 2-aminobenzofurans in a one-pot reaction (Scheme 482).<sup>756</sup> In this case, the reaction proceeded through the cleavage of multiple bonds involving C–H, O–N and two C–F bonds. The utility of this strategy was highlighted by the application of the on-DNA synthesis as well as the products' applicability as potential anticancer agents.

### Scheme 482. $\text{Cs}_2\text{CO}_3$ -Mediated [3 + 2] Annulation of *N*-Phenoxy Amides with *gem*-Difluoroalkenes





**7.2.2. Alkylation.** Sterically hindered secondary and tertiary alkyl halides engage in the nickel-catalyzed C(sp<sup>2</sup>)-C(sp<sup>3</sup>) defluorinative reductive cross-coupling with *gem*-difluoroalkenes to produce monofluoroalkenylated products via selective C-F bond cleavage (Scheme 483).<sup>757</sup> A range of functional groups tolerated in this reaction to generate the expected products with excellent *Z*-stereoselectivity. Mechanistically, Ni(COD)<sub>2</sub> is first transformed into the active Ni(I)-L<sub>n</sub> complex I, which then undergoes borylation to form L<sub>n</sub>Ni(I)-Bpin species II. With a single electron transfer (SET) to an alkyl halide, a caged alkyl radical/Ni(II) intermediate III was produced. Then, radical addition of the intermediate onto *gem*-difluoroalkene occurs regioselectively at the difluorinated position to give a caged complex IV which then undergoes another SET to form a key Ni(III) species V, which subsequently undergoes β-F elimination to give *Z*-monofluoroalkene and a Ni(III) compound VI. Further reduction regenerated the active Ni(I) catalyst. This reaction displays excellent *Z*-stereoselectivity because the steric crush is minimized during the β-F elimination step.

**Scheme 483. Nickel-Catalyzed Defluorinative Reductive Cross-Coupling of Secondary and Tertiary Alkyl Halides with *gem*-Difluoroalkenes**





A variety of monofluoroalkenes can be readily formed at room temperature by Zn(o)-mediated decarboxylative defluoroalkylation of *gem*-difluoroalkenes with *N*-hydroxyphthalimide (NHP) esters as the alkyl radical precursors (Scheme 484).<sup>758</sup> The reactions could be scaled up and gave the products with great *Z*-stereoselectivity. Mechanistically, Zn(o) first undergoes SET to the NHP ester to produce a radical anion, which then fragments into a stabilized NHP anion and *O*-based carboxyl radical. The radical further undergoes decarboxylation to afford the thermodynamically preferred alkyl radical (R•), which subsequently undergoes attack to the *gem*-difluoroalkene to furnish a difluorinated radical intermediate. Another SET reduction by Zn<sup>+</sup> occurs again to generate a β,β-difluorinated anion, then β-F elimination occurs using the conformationally favorable anti-coplanar pathway to give the *Z*-monofluoroalkenes. Very recently, a similar defluorinative alkylation was also achieved under photoredox catalysis.<sup>759</sup>

**Scheme 484. Zinc-Mediated Decarboxylative Defluoroalkylation of *gem*-Difluoroalkenes**



Interestingly, Fe catalyst can also facilitate the cross-coupling reaction of *gem*-difluoroalkenes with several unactivated and heteroatom substituted alkenes via hydrogen atom transfer (HAT) reaction. This allows a small group of structurally diverse alkylated monofluoroalkenes to be synthesized with great Z-selectivity (Scheme 485).<sup>760</sup>



### Scheme 485. Iron-Catalyzed Defluorinative Cross-Coupling of Alkenes with *gem*-Difluoroalkenes




A distinctive and efficient method of reductive coupling between *gem*-difluoroalkenes and unactivated alkyl bromides using Ni catalyst is able to regioselectively and stereoselectively furnish a series of  $\alpha$ -benzylfluoroalkenes (Scheme 486).<sup>761</sup> A large variety of functionalized monofluoroalkenes could be synthesized via a characteristic catalytic mechanism involving alkyl-Ni chain migration and defluorinative cross-coupling. This reaction occurs via oxidative addition of C-Br bond to Ni(0) to form a Ni(II) species II, which then undergoes single-electron reduction by Mn(0) and subsequent chain migration to generate Ni(I) V with a Ni-C bond at the stabilised benzylic position. *gem*-Difluoroalkene coordinates to the complex to form a Ni(I) intermediate VI. The intermediate is then subjected to regioselective migratory insertion, producing the alkyl-Ni(I) adduct VII. Finally,  $\beta$ -F elimination occurs to produce the coupling

monofluoroalkenes accompanied by the Ni(I)-F species VIII which undergoes further reduction by Mn(0) to regenerate the active Ni(0) catalyst.

### Scheme 486. Ni(II)-Catalyzed Reductive Coupling of Alkyl Bromides with *gem*-Difluoroalkenes and Its Proposed Mechanism







#### Proposed mechanism






Synthetically useful chiral 3,3-disubstituted oxindoles bearing a monofluoroalkenyl moiety can be obtained via an efficient Ni-catalyzed enantioselective reductive monofluoroalkenylation between aryl bromides and *gem*-difluoroalkenes (Scheme 487).<sup>762</sup> The reaction occurred smoothly at room temperature, affording the products in decent yields with 85-95% enantiomeric excess. The chelating Pfaltz-type ligand was used to regulate the enantioselectivity of this transformation. Of note, this protocol was applicable to the manipulation of biologically active molecules.

### Scheme 487. Nickel(II)-Catalyzed Enantioselective Reductive Monofluoroalkenylation of Aryl Bromides with *gem*-Difluoroalkenes



In 2019, Terada's research group reported a novel organocatalytic nucleophilic substitution reaction between *gem*-difluoroalkenes and ketene silyl acetals (Scheme 488).<sup>763</sup> The reaction occurred under especially benign conditions by the use of phosphazene P4-*t*Bu as an organobase catalyst to afford a series of monofluoroalkenes bearing an alkoxy carbonylmethyl group in excellent yields and *Z*-selectivities.



### Scheme 488. Phosphazene P4-*t*Bu-Catalyzed Nucleophilic Substitution of *gem*-Difluoroalkenes with Ketene Silyl Acetals



In the same year, Li's group accomplished a highly regio- and stereoselective method for the synthesis of fluorinated 1,4-enynes bearing an all-carbon quaternary center at the


propargylic position (Scheme 489).<sup>764</sup> The strategy proceeded through an alkynyl enolate intermediate followed by a nucleophilic addition and subsequent β-F elimination. Notably, this protocol accommodated broad substrate scope and functional groups under mild reaction conditions. The resulting products can be further converted into other synthetic useful intermediates. Moreover, this strategy can be applied towards the rapid synthesis of various α-alkenyl allenates by exploiting 3,3-disubstituted allenates.

### Scheme 489. Regio- and Stereoselective Synthesis of Fluorinated 1,4-Enynes




Transition metal-catalyzed alkenyl C–F bond alkylation of *gem*-difluoroalkenes with organometallic reagents such as Grignard reagents, organozinc reagents, and organoaluminum reagents have been well documented in recent years. In 2014, Cao et al. reported a novel ligand-free Kumada–Tamao–Corriu cross-coupling of *gem*-di- and monofluoroalkenes with alkyl/aryl magnesium halides enabled by palladium- and nickel-based catalytic systems, giving rise to di-cross- or mono-cross-coupled products in high yields. (Scheme 490).<sup>765</sup> Later, the same group further expanded to report the Ni(II)-catalyzed Kumada-type double cross-coupling between *gem*-difluoroalkenes and 1,4- or 1,5-Grignard reagents (Scheme 491).<sup>766</sup> The protocol occurred smoothly under especially ambient conditions to furnish a diverse array of synthetically appealing exocyclic trisubstituted alkenes which are frequently encountered in numerous pharmaceutically relevant compounds and important organic materials.<sup>767</sup>

### Scheme 490. Kumada–Tamao–Corriu Cross-Coupling Reaction of *gem*-Di- and Monofluoroalkenes with Grignard Reagents




**Scheme 491. Ni(II)-Catalyzed Kumada-Type Double Cross-Coupling of *gem*-Difluoroalkenes with Di-Grignard Reagents**





Subsequently, Cao and colleagues continued their research on a powerful Cu(I)-catalyzed or metal-free cross-coupling of *gem*-difluoroolefins with Grignard reagents (Scheme 492).<sup>768</sup> Interestingly, the tertiary and secondary C–F bond alkylation of *gem*-difluoroalkenes progressed with ease in the presence of CuCN catalyst or under catalyst-free conditions, yielding the alkylated fluoroalkenes in satisfactory yields with excellent *Z*-selectivity.

**Scheme 492. Cu(I)-Catalyzed/Transition Metal-Free Cross-Coupling of *gem*-Difluoroalkenes with Grignard Reagents**





Beside these trisubstituted *gem*-difluoroalkenes for the stereoselective synthesis of monofluorostyrenes by steric control, very recently, Tsui and colleagues established a chelation-controlled alkenyl C–F bond functionalization of tetrasubstituted *gem*-difluoroalkenes with Grignard reagents under transition metal-free catalysis, giving rise to stereodefined tetrasubstituted (*E*)-monofluoroalkenes in moderate yields (Scheme 493).<sup>769</sup>

**Scheme 493. Selective C–F Bond Functionalization of Tetrasubstituted of *gem*-Difluoroalkenes with Grignard Reagents**





Moreover, Cao's group was able to prepare alkylated monofluoroalkenes by reacting organozinc reagents with *gem*-difluoroalkenes through a nickel-catalyzed cross-coupling reaction (Scheme 494).<sup>770</sup> In this report, LiCl is used to facilitate the reaction, and mild reaction conditions are sufficient to selectively synthesize diverse *Z*-monofluoroalkenes in appreciable yields.

**Scheme 494. Nickel-Catalyzed Cross-Coupling of Organozinc Reagents with *gem*-Difluoroalkenes**



In a recent report by the group of Liu and Ma, an efficient defluorinative alkylation of aryl and vinyl C(sp<sup>2</sup>)-F bond has been accomplished under nickel catalysis with trialkylaluminum reagents as the alkyl source (Scheme 495).<sup>771</sup> Of note, a wide range of vinyl monofluorides and vinyl *gem*-difluorides were successfully alkylated in the presence Ni(COD)<sub>2</sub> and dppbz.


#### Scheme 495. Ni-Catalyzed Defluorinative Alkylation of with Trialkylaluminium Reagents



Sparse examples have been reported involving the formation of monofluoroalkenes bearing a chiral carbon center. More recently, Shi et al. presented the first example of asymmetric reductive defluorinative cross-coupling between *gem*-difluoroalkenes and racemic benzyl electrophiles by means of B<sub>2</sub>pin<sub>2</sub> as a stoichiometric nonmetallic reductant under nickel catalysis (Scheme 496).<sup>772</sup> In this strategy, a diverse range of monofluoroalkenes bearing a stereogenic allylic center were obtained in 54–96% yield. Mechanistic studies demonstrated that a radical-involved pathway may be operative in this stereo- and enantioselective protocol, and the ester group in *gem*-difluoroalkenes may greatly


facilitate the C–F bond cleavage through oxidative addition to a chiral nickel complex.

#### Scheme 496. Enantioselective Nickel-Catalyzed Reductive Defluorinative Cross-Coupling of *gem*-Difluoroalkenes and Carbon Electrophiles



In another instance, Liao and colleagues realized a highly enantioselective nucleophilic addition of fluorinated enolates to *gem*-difluoroalkenes under copper catalysis with their sulfoxide phosphine (SOP) as a ligand, leading to fabricate a wide range of structurally diverse chiral vicinal difluorides bearing a tertiary fluorinated carbon center in moderate to good yields and high *Z/E*-selectivity together with excellent enantioselectivity (Scheme 497).<sup>773</sup> Different types of acetates and acetamides were tolerated with the mild conditions. It should be pointed out, however, that a slight detrimental effect on stereocontrol was found in some amide substrates.


#### Scheme 497. Synthesis of Chiral Vicinal Difluorides through Enantioselective Cu-Catalyzed Nucleophilic Addition of Fluorinated Reagents




**7.2.3. Alkenylation and Allylation.** Hydroalkenylation of *gem*-difluoroalkenes and alkynes can also be catalysed by Ni to produce 2-fluoro-1,3-dienes (Scheme 498).<sup>774</sup> By using a combination of Ni(COD)<sub>2</sub>, PCy<sub>3</sub>, BEt<sub>3</sub>, LiO<sup>t</sup>Pr, and ZrF<sub>4</sub>, the products can be obtained in up to 89% yield. The combination of BEt<sub>3</sub> and LiO<sup>t</sup>Pr acted as a hydride source, while ZrF<sub>4</sub> helped to increase yields as a co-catalyst. Mechanistically, the oxidative cyclization of *gem*-difluoroalkene and alkyne in the presence of Ni(0) results in the regioselective formation of the β,β-difluorinated nickelacyclopentene intermediate I. The new metal-carbon bond is placed β with respect to F, similar to the previous migratory insertion reactions. Subsequent β-F elimination occurs to produce vinylnickel fluoride II containing a *Z*-fluoroalkene. The vinylnickel fluoride II

then reacts with  $\text{BEt}_3$  and  $\text{LiO}^i\text{Pr}$  to form vinylnickel hydride **III** which experiences reductive elimination to release the 2-fluoro-1,3-dienes and regenerate the active  $\text{Ni}(0)$  catalyst.

### Scheme 498. Nickel-Catalyzed Hydroalkenylation of *gem*-Difluoroalkenes and Alkynes




#### Proposed mechanism




Quite recently, Tsui's group illustrated a highly diastereoselective  $\text{Pd}(0)$ -catalyzed twofold Miyaura borylation/Suzuki-Miyaura cross-coupling of C-F bonds for the direct preparation of stereodefined difluorinated 1,3-dienes (Scheme 499).<sup>775</sup> In this report, a diverse array of symmetrical 1,3-dienes bearing a vicinal difluoro fragment were produced exclusively as a single diastereomer from the corresponding tetrasubstituted *gem*-difluoroalkenes. Interestingly, a series of unsymmetrical difluorinated 1,3-dienes can also be obtained through the  $\text{Pd}(0)$ -catalyzed cross-coupling of *gem*-difluoroalkenes with borylated monofluoroalkenes.

### Scheme 499. Synthesis of Difluorinated 1,3-Dienes via $\text{Pd}$ -Catalyzed C-F Activation of Tetrasubstituted *gem*-Difluoroalkenes



Moverover, the same group also expanded to report a  $\text{Pd}(0)$ -catalyzed stereoselective C-F bond vinylation and allylation of tetrasubstituted *gem*-difluoroalkenes for the synthesis of synthetically useful monofluorinated 1,3- and skipped 1,4-dienes in high yields and excellent diastereoselectivities (Scheme 500).<sup>776</sup> Unlike the widely used  $\text{Pd}(\text{PPh}_3)_4$ , the catalytic system incorporated a  $\text{Pd}(0)$  and *dpe* as the ligand was illustrated for Stille-type cross-coupling between *gem*-difluoroolefins and vinyl- or allyltin reagents.

### Scheme 500. $\text{Pd}(0)$ -Catalyzed Stereoselective C-F Bond Vinylation and Allylation of Tetrasubstituted *gem*-Difluoroalkenes




**7.2.4. Arylation.** Early in 1991, Heitz and Knebelkamp pioneered to report the first example of  $\text{Pd}$ -catalyzed defluorinative cross-coupling between 1,1-difluoroethylene and aryl iodides (Scheme 501a).<sup>777</sup> In this work, the migratory insertion of 1,1-difluoroethylene into C-Pd



bonds of arylpalladium(II) iodides generated by initial oxidative addition proceeds smoothly to afford the  $\beta,\beta$ -difluorinated phenethylpalladium(II) complex, which subsequently undergoes a facile  $\beta$ -F elimination to furnish  $\alpha$ -fluorostyrenes. A couple of years later, the group of Ichikawa and Minami further discovered that the treatment of 1,1-difluoro-1-ethylenes with a low-valent zirconocene reagent ( $\text{ZrCp}_2$ ) generated the thermostable 1-fluorovinylzirconocenes via a vinylic C-F bond cleavage, which engaged in palladium-catalyzed coupling reaction with aryl iodides in a one-pot operation to afford arylated fluoroethylenes in acceptable yields (Scheme 501b).<sup>778</sup>

### Scheme 501. Synthesis of Fluorostyrenes via Vinylic C-F Activation of Fluoroolefins with Aryl Iodides

a) Heitz et al., 1991





b) Ichikawa, Minami et al., 1999





The Suzuki-Miyaura coupling is one of the most dominant strategies for the construction of C-C bonds in modern synthetic chemistry. The defluorinative Suzuki-Miyaura type cross-coupling of *gem*-difluoroalkenes with boronic acids through alkenyl C-F activation gives rise to monofluoro-substituted olefins. In 2015, Cao's group first reported the stereoselective Ni-catalyzed Suzuki-type cross-coupling between *gem*-difluoroalkenes and a series of arylboronic acids (Scheme 502).<sup>779</sup> In the presence of  $\text{NiCl}_2(\text{PCy}_3)_2$  (5 mol%) and  $\text{K}_3\text{PO}_4$ , a number of *Z*-monofluorostilbenes were produced in good yields with exceptional regioselectivity. The authors tentatively proposed a plausible mechanism that proceeds through the oxidative addition of the alkenyl C-F bond accompanied by cross-coupling with arylboronic acids.

### Scheme 502. Nickel-Catalyzed Suzuki-Miyaura Type Cross-Coupling of *gem*-Difluoroalkenes with Arylboronic Acids



Another opportunity for the defluorinative Suzuki-Miyaura cross-coupling of *gem*-difluoroalkenes with arylboronic acids has been established by Toste and co-workers (Scheme 503).<sup>780</sup> Overall, this redox-neutral protocol features a broad functional group scope, which conveniently proceeds via a  $\beta$ -F elimination process to yield the products with excellent diastereoselectivity ( $\geq 50:1$ ). The utility of this strategy was highlighted in the rapid synthesis of a Gleevec<sup>®</sup> analogue by utilizing a monofluorostilbene to function as an amide isostere.


### Scheme 503. Pd(II)-Catalyzed Defluorinative Suzuki-Miyaura Cross-Coupling of *gem*-Difluoroalkenes with Arylboronic Acids



Recently, Tsui's group established a conceptually similar approach to prepare a series of tetrasubstituted (*E*)-monofluoroalkenes (Scheme 504).<sup>781</sup> Detailed DFT calculations suggested that the reaction proceeds through a formal [4 + 1] cycloaddition pathway accompanied by a 1,5-sigmatropic fluoride migration which is promoted through the chelation of the ester-substituent group to the Pd-catalyst. The mechanism also justified the low C-F bond strength for its successive cleavage (slow step) and the stereoselectivity control. Mechanistically, the C-F activation readily generates a vinylpalladium(II) fluoride intermediate which subsequently participates in a



transmetalation effortlessly with arylboronic acids without the need for an external base.

### Scheme 504. Pd-Catalyzed Suzuki-Miyaura Coupling of *gem*-Difluoroalkenes with Arylboronic Acids



Shen and colleagues found that the combination of lithium organoborates with zinc bromide ( $\text{ZnBr}_2$ ) readily *in situ* generated a highly reactive lithium aryl zincate species which significantly facilitates transmetalating the aryl group from boron to nickel catalyst.<sup>782</sup> Continuing this approach, they applied readily available “ate-type” aryl pinacol boronates for the ready assembly of *Z*-selective monofluoroalkenes from *gem*-difluoroalkenes in the presence of a catalytic amount of  $\text{ZnBr}_2$  (Scheme 505).<sup>783</sup>


### Scheme 505. Ni-Catalyzed Defluorinative Coupling of *gem*-Difluoroalkenes with Lithium Organoborates



Readily available tetrafluoroethylene (TFE) is an economical bulk organofluorine feedstock for the mass production of commonly used poly(tetrafluoroethylene) and copolymers. In 2013, Ogoshi and co-workers achieved a defluorinative Suzuki-Miyaura type coupling of tetrafluoroethylene with arylboronates under Pd-catalyzed base-free conditions (Scheme 506a).<sup>784</sup> The reaction took place smoothly through the formation of the key


palladium(II) fluoride intermediate without the use of a Lewis acid additive to facilitate the oxidative addition of a vinylic C–F bond to Pd(0) catalyst. Later, they successfully expanded to elaborate the efficient synthesis of trifluorostyrenes via a facile  $\beta$ -F elimination of 2-aryl-1,1,2,2-tetrafluoroethylcopper intermediate which is *in situ* generated from arylboronates, copper *tert*-butoxide, and tetrafluoroethylene (TFE) with 1,10-phenanthroline as a ligand (Scheme 506b).<sup>785</sup>

### Scheme 506. Synthesis of Trifluorostyrene Derivatives from Tetrafluoroethylene and Arylboron Reagents



In 2011, Ogoshi and colleagues also reported the palladium-catalyzed C–F bond monoarylation of TFE with *in situ*-prepared arylzinc agents, affording a variety of  $\alpha,\beta,\beta$ -trifluorostyrenes in synthetically satisfactory yields (Scheme 507).<sup>786</sup> This strategy employs lithium iodide as a Lewis acid to facilitate the oxidative addition of a vinylic C–F bond to Pd(0) even at room temperature, thereafter generating a well-defined trifluorovinylpalladium(II) iodide species which was readily isolated and unambiguously determined by X-ray diffraction.


### Scheme 507. Pd-Catalyzed C–F Bond Arylation of Tetrafluoroethylene (TFE) with Arylzinc Reagents



Moreover, Ishihara's group reported the examples of C–F bond arylation of readily available benzyl 2,3,3-trifluoroacrylate with various arylmagnesium bromides


(Scheme 508).<sup>787</sup> The reaction occurred smoothly at  $-78\text{ }^{\circ}\text{C}$  in the presence of CuBr (13 mol%) as the catalyst, delivering a variety of  $\alpha,\beta$ -difluoroacrylates in up to 90% yield with high *Z*-selectivity. Both alkyl- and alkenylmagnesium halides were compatible with this protocol to afford the corresponding defluorinative products.

### Scheme 508. Copper-Catalyzed C–F Bond Arylation of *gem*-Difluoroalkenes with Grignard Reagents



**7.2.5. Alkynylation.** The fluorinated conjugated enynes represent an important class of synthetically useful building blocks in organic synthesis. In 2014, Cao et al. first disclosed a benign and practical strategy for the stereoselective synthesis of fluorinated conjugated enynes by reacting readily available *gem*-difluoroalkenes with terminal alkynes in the presence of  $n\text{BuLi}$  and  $\text{K}_3\text{PO}_4$  (Scheme 509).<sup>788</sup> This Sonogashira-type defluorinative cross-coupling afforded a series of conjugated  $\beta$ -fluoroenynes in up to 93% yield with high *E*-selectivity.


### Scheme 509. Synthesis of $\beta$ -Fluoroenynes by Reaction of *gem*-Difluoroalkenes with Terminal Alkynes



In a related report, Tsui and co-workers illustrated an impressive stereoselective Pd-catalyzed C–F alkylation of tetrasubstituted *gem*-difluoroalkenes with diverse terminal alkynes (Scheme 510).<sup>789</sup> This defluorinative Sonogashira-type coupling enables the efficient synthesis of a broad series of conjugated tetrasubstituted monofluoroenynes with a defined stereochemistry. The high stereoselectivity of this protocol could be rationalized by the chelation-assisted oxidative addition of the Pd-

catalyst to the alkenyl C–F bond. The key intermediate, a monofluorovinyl Pd(II) species, was readily isolated alongside with an X-ray crystal structure as proof for the proposed mechanism.



### Scheme 510. Pd(0)-Catalyzed C–F Alkylation of Tetrasubstituted *gem*-Difluoroalkenes with Alkynes



**7.2.6. Carboxylation.**  $\alpha$ -Fluoroacrylic acid derivatives are versatile motifs widespread in many pharmaceutically relevant molecules.<sup>790</sup> Consequently, remarkable attention has been devoted towards the development of efficient and practical methods for the rapid synthesis of  $\alpha$ -fluoroacrylic acids. In this regard, Yu's group in 2019 synthesized a diverse array of  $\alpha$ -fluoroacrylic acids through the selective alkenyl C–F carboxylation of *gem*-difluoroalkenes with an atmospheric pressure of  $\text{CO}_2$  by a copper(I)/diboron catalytic system (Scheme 511a).<sup>791</sup> This formal *ipso* monocarboxylation of C–F bonds can be performed via telescoping synthesis to stereoselectively afford the (*Z*)-product. Intriguingly, the challenging *gem*-difluorodienes were also compatible with this protocol. Later, detailed theoretical calculations were carried out by Shi's group,<sup>792</sup> and the results clearly revealed that the reaction proceeds successively via the migratory insertion of difluoroalkene into the Cu(I)-B intermediate, *syn*  $\beta$ -F elimination, transmetalation, and rate-determining carboxylation steps. The regioselectivity of this reaction is rationalized by the migration insertion step. Almost simultaneously, the group of Zhou and Wu also established a silimar approach for the stereoselective alkenyl C–F bond carboxylation (Scheme 511b).<sup>793</sup> Due to the use of stoichiometric bis(pinacolatodiboron) as a nonmetallic reductant, a large number of synthetically useful functionalities were compatible under the conditions. More challenging *gem*-difluorodienes engaged in this C–F carboxylation strategy to furnish conjugated  $\alpha$ -fluorocarboxylic acids in decent yields with high *Z*-selectivity. Impressively, the less reactive (*E*)-fluorostyrenes delivered the corresponding cinnamic acids with the retention of stereochemistry (*E/Z* > 20/1), while no reaction occurred when the corresponding



(*Z*)-isomers were employed. This reveals that a *trans*- $\beta$ -F elimination may be operative in this process.

### Scheme 511. Stereoselective Defluorinative C–F Bond Carboxylation of *gem*-Difluoroalkenes with CO<sub>2</sub>



The group of Wu and Zhou extended to disclose a green and sustainable strategy for the straightforward construction of  $\alpha$ -fluoroacrylic acids through the electrochemical defluorinative C–F carboxylation of *gem*-difluoroalkenes with CO<sub>2</sub> (Scheme 512).<sup>794</sup> Utilizing a Pt plate as the cathode and an inexpensive Ni plate as the anode in an easy-to-use undivided cell, the reaction progressed with ease under typically ambient conditions in the absence of any exorbitant transition-metal catalysts, ligands, external base or reductant. A series of  $\alpha$ -fluoroacrylic acids were obtained in 35–83% yield with up to 20:1 *Z/E* ratio. The authors performed a cyclic voltammetry experiment and tentatively proposed an unorthodox process involving single-electron reduction, reaction with CO<sub>2</sub>, single-electron reduction,  $\beta$ -F elimination.



### Scheme 512. Electrochemical Defluorinative C–F Bond Carboxylation of *gem*-Difluoroalkenes with CO<sub>2</sub>



**7.2.7. Other Useful Reactions.** In 2016, Cao's group disclosed a stereoselective C–F bond cyanation of *gem*-



difluoroolefins with readily available benzyl nitriles as the environmentally benign cyanating agent (Scheme 513).<sup>795</sup> The reaction operated under a copper-free system and ambient conditions. A series of synthetically versatile monofluorinated alkenyl nitriles were prepared in yields ranging from 50% to 85% with excellent stereoselectivity. A plausible catalytic mechanism was proposed involving the oxidation of the C–H bond, cleavage of the C–CN bond and subsequent nucleophilic vinylic substitution (S<sub>N</sub>V) to afford the expected products.

### Scheme 513. Stereoselective C–F Bond Cyanation of *gem*-Difluoroolefins with Benzyl Nitriles




Later, He's group also demonstrated the synthesis of  $\alpha$ -fluoroacrylonitriles through a transition metal-free organocatalytic C–F bond cyanation of *gem*-difluoroolefins with TMSCN. In the presence of a catalytic amount of DBU, the *gem*-difluoroalkenes participated in a nucleophilic addition- $\beta$ -elimination reaction to generate the  $\alpha$ -fluoroacrylonitriles in satisfactory yields with exceptional *Z/E* stereoselectivity (Scheme 514).<sup>796</sup>

### Scheme 514. Organocatalytic C–F Bond Cyanation of *gem*-Difluoroolefins with TMSCN





Around the same time, Deng et al. reported the feasible strategy for the nucleophilic substitution of *gem*-difluoroalkenes with trimethylsilyl nucleophiles (Scheme 515).<sup>797</sup> In this case, a catalytic amount of Cs<sub>2</sub>CO<sub>3</sub> (10 mol%) was used as a promoter to maintain a high concentration of the nucleophilic anions which was proposed to kickstart the catalytic cycle. This strategy enables the construction of diverse highly functionalized monofluoroalkenes bearing a cyano or trifluoromethyl group with modest to excellent stereoselectivities.

#### Scheme 515. Nucleophilic Substitution of *gem*-Difluoroalkenes with TMSNu



More recently, Wu and colleagues continued their carbonylation strategy,<sup>798</sup> and presented an unprecedented Pd/Cu synergistic system for the formal defluorinative carbonylative coupling reaction between *gem*-difluoroalkenes and aryl iodides, providing an efficient access to a variety of  $\alpha$ -fluoroaldehydes under benign conditions in appreciable yields with excellent *Z*-selectivity (Scheme 516).<sup>799</sup> Preliminary mechanistic experiments suggested that the transmetalation of the alkyl-Pd(II)I complex and CuBpin species was a crucial step for the catalytic cycle.



#### Scheme 516. Pd/Cu-Catalyzed Formal Defluorinative Carbonylative Coupling of *gem*-Difluoroalkenes with Aryl Iodides



### 7.3. C-Het Bond Formation



**7.3.1. Borylation and Silation.** Organoboron- and organosilicon-based reagents are widely used in organic synthesis and drug discovery. Accordingly, there have been considerable efforts towards the development of broadly applicable methods to prepare this extremely important class of compounds. The direct alkenyl C–F bond functionalization of *gem*-difluoroalkenes undoubtedly offers a practical and unique tool for the synthesis of fluorinated alkenyl borons and silanes, which could also act as versatile building blocks for further elaborations to synthesize value-added complex molecules.<sup>25</sup> In 2017, Cao and colleagues continued their C–F bond functionalization strategy, and described the first example of copper-catalyzed stereoselective C–F bond borylation of *gem*-difluoroalkenes with bis(pinacolato)diboron (B<sub>2</sub>pin<sub>2</sub>) in the presence of NaO<sup>t</sup>Bu base and Xantphos ligand at room temperature, enabling a convenient synthesis of structurally diverse (*Z*)-fluorinated alkenylboronic pinacol esters in 51–81% yields (Scheme 517).<sup>800</sup>

#### Scheme 517. Cu-Catalyzed Stereoselective Borylation of *gem*-Difluoroalkenes with Bis(pinacolato)diboron



Meanwhile, the research group of Niwa, Ogoshi, and Hosoya also established a straightforward approach to fabricate a series of borylated fluoroalkenes via the copper-catalyzed regioselective monodefluoroborylation of diverse polyfluoroalkene substrates, including (difluorovinyl)arenes, (trifluorovinyl)arenes, tetrafluoroethylene (TFE), and trifluoromethylated monofluoroalkenes (Scheme 518).<sup>801</sup> The type of boron reagent used in the reaction depends on the electronic character of *gem*-difluoroalkenes. Substrates bearing an electron-deficient aryl group made use of (Bpin)<sub>2</sub> as coupling partners, while those with electron-rich aryl group employed (Bnep)<sub>2</sub>. The resulting (fluoroalkenyl)-boronic esters could be readily converted into the corresponding potassium trifluoroborate salts by the treatment with KHF<sub>2</sub> in MeOH.



### Scheme 518. Copper-Catalyzed Regioselective Monodefluoroborylation of Polyfluoroalkenes



Subsequently, Ito's group employed a Cu(I)/Xantphos catalytic system for the defluoroborylation of aliphatic *gem*-difluoroalkenes by means of (Bpin)<sub>2</sub> as the borylating reagent in conjunction with KO<sup>t</sup>Bu as the base (Scheme 519).<sup>802</sup>  $\alpha$ -Boryl- $\alpha$ -fluoroalkenes were synthesized with high *Z*-stereoselectivity which can be isolated after


converting into corresponding trifluoroborates. On the basis of preliminary DFT calculations, the authors proposed a plausible mechanism for this process. Initially, Cu(I) alkoxide is formed from CuCl, the ligand, and KO<sup>t</sup>Bu, which then reacts with (Bpin)<sub>2</sub> to produce a boryl-Cu(I) intermediate. An alkyl-Cu(I) species is then produced via a four-centred reaction between the key boryl-Cu(I) intermediate and *gem*-difluoroalkenes, which subsequently undergoes  $\beta$ -F elimination to give the (*Z*)-defluoroborylated products accompanied by the release of copper(I) fluoride. Finally, CuF reacts with diboron and KO<sup>t</sup>Bu to reform the transmetalated active boryl-Cu(I) intermediate.

### Scheme 519. Cu(I)-Catalyzed Defluoroborylation of Aliphatic *gem*-Difluoroalkenes





The chemoselective poly-borylation reaction of *gem*-difluoroalkenes with (Bpin)<sub>2</sub> is greatly facilitated by Cu(I) catalyst via dual alkenyl C–F bond activation (Scheme 520).<sup>803</sup> By slightly adjusting the type of Cu salts, additives and solvents, together with the reaction temperature, a large variety of multi-borylated compounds including 1,2-alkyldiboronates, 1,1,2-alkyltriboronates, and 1,1,1,2-alkyltetraboronates can be selectively produced. Notably, Xantphos was kept as the ligand in all three C–F borylation reactions. The borylated products could subsequently be further functionalized and serve as helpful starting materials.

### Scheme 520. Highly Tunable Copper-Catalyzed Multi-Borylation of *gem*-Difluoroalkenes



Cu catalysts with different oxidation states, ligands, and additives can also be used to catalyze the defluorinative borylation and silylation of *gem*-difluoroalkenes (Scheme 521).<sup>804</sup> Using CuCl<sub>2</sub> together with DPEphos will promote C–F bond borylation, while employing the CuCl/PCy<sub>3</sub> catalytic system will promote C–F bond silylation. In this report, aliphatic *gem*-difluoroalkenes was not compatible with this protocol to undergo defluorinative silylation. In contrast, the defluorinative borylation of aliphatic *gem*-difluoroalkenes occurred smoothly to afford monofluoroalkenyl boronic esters in 50% yield.



### Scheme 521. Cu(I)/Cu(II)-Catalyzed Defluorinative Borylation and Silylation of *gem*-Difluoroalkenes



Silylated fluoroolefins could be served as synthetically important building blocks for the synthesis of bioactive natural products and pharmaceuticals. Using the (IPr)CuF catalyst, Ogoshi's group was able to perform the C–F bond defluorosilylation of diverse substrates including *gem*-difluoroalkenes, tetrafluoroethylene (TFE), and other polyfluoroalkenes with silylborane reagent, affording a variety of fluorinated vinylsilanes (Scheme 522).<sup>805</sup> The obtained (Trifluorovinyl)phenyldimethylsilanes can react with iodobenzene in a copper-mediated cross-coupling reaction to produce  $\alpha,\beta,\beta$ -trifluorostyrene. Mechanistic studies illustrated that this defluorosilylation process


proceeded through the 1,2-addition of a silylcopper species to the polyfluoroalkene followed by a facile selective  $\beta$ -F elimination which is greatly facilitated by the *in situ* generated Lewis acidic F–Bpin.

### Scheme 522. Synthesis of Fluorinated Vinylsilanes via Copper-Catalyzed C–F Bond Defluorosilylation of Polyfluoroalkenes with Silylborane Reagent



Generally, it is difficult to selectively defluorosilylate and convert a highly stable fluoroolefin into the corresponding silylated fluoroalkenes. In 2019, Shi's group developed a convenient and economical approach for the defluorosilylation of fluoroalkenes with silylboronates to assemble a series of silylated fluoroolefins under transition metal- and ligand-free conditions (Scheme 523).<sup>806</sup> The reaction occurred in the presence of NaOMe and tolerated a broad range of substrate scope. DFT calculations were performed and illustrated that transient silyl anion complex participated in an S<sub>N</sub>2' or S<sub>N</sub>Y' substitution type process, which accounts for this alkoxy base-promoted defluorosilylation.


### Scheme 523. Metal-Free Defluorosilylation of Fluoroalkenes with Silylboronates



In 2021, the group of Zhu and Feng uncovered the example of Fe(II)-catalyzed defluoroborylation of diverse unactivated *gem*-difluoroalkenes, *gem*-dichloroalkenes, and *gem*-dibromoalkenes, affording the corresponding 1,2-bis(boryl)-alkanes in modest to good yields (Scheme 524).<sup>807</sup> This strategy featured high regioselectivity, a wide



substrate scope, and excellent functional group compatibility. Initial mechanistic experiments suggested a double  $\beta$ -F elimination involved in the mechanism, and 1,1-diborylated olefins could have been possible intermediates in this transformation.

### Scheme 524. Synthesis of 1,2-Bis(boryl)alkanes via Defluoroborylation of *gem*-Dihaloalkenes




The same group continued to describe the regio- and stereoselective defluorosilylation of unbiased aliphatic *gem*-difluoroalkenes under via iron-catalyzed controllable C-F bond cleavage (Scheme 525).<sup>808</sup> By slightly adjusting the conditions, a diverse array of *gem*-disilylated and (*E*)-silylated alkenes were readily prepared in high yields. Combined experimental and DFT calculation studies demonstrated that the chemoselectivity of disilylation might undergo a thermodynamically favored process through an insertion/elimination/addition mechanism.

### Scheme 525. Regio- and Stereoselective Iron-Catalyzed Controllable Defluorosilylation of Unactivated *gem*-Difluoroalkenes





**7.3.2. Miscellaneous.** In 2014, Cao's group illustrated a versatile transition-metal-free approach for the formation of C-N bonds through a vinylic nucleophilic substitution reaction (S<sub>N</sub>V) of (2,2-difluorovinyl)arenes with a large variety of nitrogen-based heterocycles under basic and relatively benign conditions (Scheme 526).<sup>809</sup> This Ullmann-type transformation occurred uneventfully in the presence of K<sub>3</sub>PO<sub>4</sub> under ambient conditions, yielding the (*E*)-*N*- $\alpha$ -fluorovinyl derivatives of azoles in decent yields with high stereocontrol.

### Scheme 526. Synthesis of *N*-( $\alpha$ -Fluorovinyl)azoles from *gem*-Difluoroalkenes and Azoles





Subsequently, the same group continued to report a unique three-component reaction in an effort to synthesize fluorovinyl aryl ethers from *gem*-difluoroalkenes, arylboronic acids and oxygen under catalyst-free conditions (Scheme 527).<sup>810</sup>

### Scheme 527. Synthesis of Fluorovinyl Aryl Ethers from *gem*-Difluoroalkenes, Arylboronic Acids and Oxygen





In 2016, Cao's group further described a practical and benign strategy for the efficient synthesis of fluorovinyl pyrazolyl ethers and thioethers via the coupling reaction between aryl substituted *gem*-difluoroalkenes and pyrazolin-5-ones or pyrazolin-5-thiones in the presence of an alkoxy base (Scheme 528).<sup>81</sup>

### Scheme 528. Efficient Synthesis of Fluorovinyl Pyrazolyl(thio)ethers from *gem*-Difluoroalkenes with Pyrazolin-5-ones(thiones)




Successfully, the group of Du and He disclosed the *N*-heterocyclic carbene-catalyzed stereoselective synthesis of alkenyl C–S bond formation reaction. Under a catalytic amount of the stable carbene, a variety of thiols participated in this base-free nucleophilic substitution reaction with *gem*-difluoroalkenes to afford the corresponding  $\alpha$ -fluorovinyl thioethers in up to 99% yield with excellent *Z*-selectivity (Scheme 529).<sup>812</sup>

### Scheme 529. NHC-Catalyzed Stereoselective Synthesis of $\alpha$ -Fluorovinyl Thioethers




Moreover, Ji and co-workers elaborated a defluorinative reductive cross-coupling reaction between *gem*-difluoroolefins and thiosulfonates/selenosulfonates under nickel catalysis, enabling the synthesis of thiolated or selenylated monofluoroalkenes through the regioselective cleavage of the vinylic C–F bond. Of note, this protocol features highly accessible substrates, benign reaction conditions, and excellent *E*-selectivity (Scheme 530).<sup>813</sup>

### Scheme 530. Ni-Catalyzed Defluorinative Reductive Cross-Coupling Reaction of *gem*-Difluoroalkenes and Thiosulfonate/Selenosulfonate



An efficient approach for the preparation of diverse synthetically valuable  $\alpha$ -fluoro- $\beta$ -arylalkenyl sulfones from *gem*-difluoroalkenes and sodium sulfinates in the absence of any transition metal or base has been reported by Shi's group (Scheme 531).<sup>814</sup> The protocol demonstrated a wide functional group tolerance with excellent stereoselectivity. Remarkably, this strategy can also be used for late-stage transformation of value-added complex compounds such as drugs and natural products.


### Scheme 531. Direct Synthesis of $\alpha$ -Fluoro- $\beta$ -arylalkenyl Sulfones from *gem*-Difluoroalkenes with Sodium Sulfinates



#### 7.4. Photoredox Catalysis

Along with the ever-increasing efforts in photoredox chemistry over the last decade, a large number of novel and straightforward strategies have emerged which greatly enriched the alkenyl C–F bond functionalizations of *gem*-difluoroalkenes under visible-light-induced catalysis or visible-light redox/transition-metal synergistic catalysis to incorporate various fluorine-containing moieties into organic molecules.<sup>815</sup> Under photoredox conditions, *gem*-difluoroalkenes can be easily irradiated to undergo C–F bond cleavage, thus generating a monofluoroalkenyl radical through a SET process. The reaction can be quenched by another radical to furnish the coupling products. This radical can be formed by the oxidized photoredox catalyst or by a transition metal-based co-catalyst. Alternatively, the metalated complex can be readily transformed into many synthetically appealing moieties by further coupling with other organic electrophiles (Scheme 532).


#### Scheme 532. Visible-Light-Promoted C–F Bond Activation of *gem*-Difluoroalkenes



In 2016, Hashmi's group first initiated the C–F bond functionalization of *gem*-difluoroalkenes by photoredox catalysis (Scheme 533).<sup>816</sup> They elaborately realized an unprecedented radical–radical cross-coupling between  $\alpha$ -aminoalkyl radicals and monofluoroalkenyl radicals

generated from *gem*-difluoroalkenes, producing a large variety of tetrasubstituted monofluoroalkenes in good yields. The elegant late-stage diversification of top-selling drugs (rosiglitazone, citalopram and venlafaxine etc.) and bioactive natural products (oleanic acid, androsterone, and dihydroartemisinin) significantly illustrated the synthetic potential of this radical-involved strategy. More recently, a similar stereoselective C–F bond functionalization of *gem*-difluoroalkenes was also achieved by Li's group through a chelation-assisted, nickel- and photoredox-catalyzed strategy.<sup>817</sup>


#### Scheme 533. Visible-Light-Promoted Defluorinative C(sp<sup>3</sup>)–H Monofluoroalkenylation of Tertiary Amines with *gem*-Difluoroalkenes



Similarly, Zhou and co-workers presented the direct synthesis of fluorinated tetrahydropyridines and bridged azabicyclo[3. m. 1] frameworks which are less accessible by other methods via an unprecedented photocatalytic dehydrogenative/defluorinative cross-coupling approach from both acyclic and cyclic tertiary amines with  $\alpha$ -trifluoromethyl alkenes, followed by an intramolecular defluorinative C–H functionalization (Scheme 534a).<sup>818</sup> Subsequently, the same group further accomplished the construction of biologically important fluorinated benzo[*a*]quinolizidines and dihydrobenzoxepines from dihydroisoquinoline acetic acids (Scheme 534b)<sup>819</sup> and *o*-hydroxyphenylacetic acids (Scheme 534c)<sup>820</sup> through a radical-based decarboxylative/defluorinative cross-coupling under visible light photocatalysis. It must be noted that the combination of bifunctional nucleophiles with  $\alpha$ -trifluoromethyl alkenes involving the consecutive two C–F bond cleavage strategy was successively reported


by the groups of Ichikawa<sup>821-823</sup> and Zhang<sup>824,825</sup> under base- or transition-metal-mediated conditions.

### Scheme 534. Visible Light-Mediated Double C–F Bond Activation of $\alpha$ -Trifluoromethyl Alkenes




Loh and co-workers in 2018 established a practical photoredox-catalyzed defluorinative trifluoromethylation of  $\alpha$ -trifluoromethyl alkenes and *gem*-difluoroalkenes (Scheme 535).<sup>826</sup> The reactions progressed effectively through the addition of air-stable Langlois' reagent as the trifluoromethyl source, followed by a  $\beta$ -fluoride elimination to give the corresponding multifluorinated olefins in modest yields with excellent stereoselectivity.

### Scheme 535. Photoredox-Catalyzed Defluorinative Trifluoromethylation of $\alpha$ -Trifluoromethyl Alkenes and *gem*-Difluoroalkenes






The visible-light-promoted vicinal trifluoromethylation and monofluoroalkenylation of alkenes through a radical-radical cross-coupling process has been developed by Wang and collaborators. This redox-neutral protocol employed *gem*-difluoroalkenes and Langlois' reagent as the radical sources, enabling a mild and step-economical route to incorporate two privileged fluorinated moieties into simple alkenes in one step. The utility of this strategy is highlighted by late-stage diversification of pharmaceutically relevant compounds (Scheme 536).<sup>827</sup>

### Scheme 536. Visible-Light-Induced Defluorinative Trifluoromethylation and Monofluoroalkenylation of Simple Alkenes





Visible light can also be used to carry out photocatalytic decarboxylation of *N*-protected  $\alpha$ -amino acids with *gem*-difluoroalkenes at room temperature, producing a diverse array of  $\alpha$ -amino monofluoroalkenes which should find widespread applications in pharmacology and material chemistry (Scheme 537).<sup>828</sup> In this case, this reaction begins when the photocatalyst Ir(III) is irradiated to produce an excited  $^*Ir(III)$ . As this is a strong oxidizing agent, the excited  $^*Ir(III)$  readily removes an electron from the deprotonated *N*-Boc proline, followed by decarboxylation to generate a transient  $\alpha$ -aminoalkyl radical **I** accompanied by the formation of Ir(II). Ir(II) then causes *gem*-difluoroalkene to undergo reduction and form a radical anion, which subsequently undergoes decomposition by losing F<sup>-</sup> to produce a monofluoroalkenyl radical **II** while regenerating the Ir(III) photocatalyst via SET. Finally, radicals **II** and **I** undergo cross-coupling with each other to form the decarboxylative monofluoroalkenylation product.

### Scheme 537. Photocatalytic Decarboxylation of *N*-Protected $\alpha$ -Amino Acids with *gem*-Difluoroalkenes





Inspired by this work, An and Li et al. reported a general and practical monofluoroalkenylation of unactivated carboxylic acids through an efficient photoredox catalytic decarboxylation in the presence of 2,4,5,6-tetra(9*H*-carbazol-9-yl)isophthalonitrile (4CzIPN) as the organic photocatalyst (Scheme 538).<sup>829</sup> This strategy circumvents the exploitation of presynthesized redox-active esters and high-powered light sources, allowing a benign method for the straightforward decarboxylative cross-coupling of unactivated carboxylic acids. In addition, this strategy caters for the exceptional and gruelling late-stage monofluoroalkenylation of complex molecules such as bioactive ramipril, gemfibrozil, dehydroabiatic acid, enoxolone as well as the sugar-derivative cyclic  $\alpha$ -oxy acid.

### Scheme 538. Photoredox Catalytic Decarboxylative Monofluoroalkenylation of *gem*-Difluoroalkenes with Carboxylic Acids



In the same year, Feng's group reported the efficient C–F bond carboxylation of *gem*-difluoroalkenes through photoredox/palladium dual catalysis (Scheme 539).<sup>830</sup> In this metallaphotoredox C–F activation reactions, electronically diverse substituted *gem*-difluoroalkenes can efficiently undergo C–F carboxylation with an inert CO<sub>2</sub> electrophile to form a large variety of value-added  $\alpha$ -fluoroacrylic acids, which then undergoes further esterification to produce the respective methyl esters.


### Scheme 539. Selective Defluorinative Carboxylation of *gem*-Difluoroalkenes by Photoredox/Palladium Dual Catalysis




Defluorinative monofluoroalkenylation can also be achieved via cooperative photoredox and hydrogen-atom-

transfer (HAT) catalysis (Scheme 540).<sup>831</sup> This reaction allows the cross-coupling of diverse amines, ethers, and thioethers to regioselectively form monofluoroalkenes with different  $\alpha$ -heteroatom substituents. Mechanistically, the photocatalyst is initially irradiated with blue LED to produce the excited redox active Ir(III)\*, which then oxidizes quinuclidine to form the Ir(II) complex and the radical cation I. The Ir(II) catalyst then provides one electron to the *gem*-difluoroalkene, then loses F<sup>-</sup> to produce the radical II and regenerate the photocatalyst. At the same time, the oxidized quinuclidine radical cation I undergoes HAT with amine to form the  $\alpha$ -aminoalkyl radical III. Finally, the two radicals II and III heterodimerize to form the defluorinative monofluoroalkenylation product. By the synergetic merger of photoredox and bromine-based hydrogen atom transfer catalysis, quite recently, Deng and colleagues achieved a similar transformation under batch and continuous-flow conditions with a broad range of C-H patterns, including ethers, amides and aliphatic aldehydes.<sup>832</sup>

### Scheme 540. Defluorinative Monofluoroalkenylation via Cooperative Visible-Light Photoredox and Hydrogen-Atom-Transfer (HAT) Catalysis




#### Proposed mechanism



The group of Sun and Zhou presented the C-F bond alkylation of *gem*-difluoroalkenes under visible-light-promoted conditions with readily prepared 4-alkyl-1,4-dihydropyridines (DHPs, also known as Hantzsch esters)



as the alkyl radical precursors (Scheme 541).<sup>833</sup> A broad range of primary, secondary, and even tertiary 4-alkyl substituted DHPs were all compatible with this photoredox strategy to afford alkyl substituted monofluoroalkenes in 26–95% yield.

### Scheme 541. Visible-Light-Promoted Defluorinative Alkylation of *gem*-Difluoroalkenes with DHPs





Quite recently, Wang et al. elaborated a practical strategy for the defluorinative monofluoroalkenylation of alkyl bromides through a silyl radical-mediated halogen abstraction process under visible-light photoredox catalysis (Scheme 542).<sup>834</sup> Mechanistic studies suggested that the products were selectively formed through the coupling of aliphatic radicals and fluoroalkenyl radicals. Halogen abstraction promoted by the presence of silyl radicals allowed the strategy to be applied for the monofluoroalkenylation of a wide diversity of alkyl and heteroaryl halides. Of note, the ability to scale up the reaction and be used for cholesterol demonstrated its applicability for late-stage monofluoroalkenylation reactions.

### Scheme 542. Silyl Radical-Mediated Dehalogenative Coupling of *gem*-Difluoroalkenes with Alkyl Halides




A more recent report by Crudden et al. demonstrated that redox-active alkylsulfones could also be employed as feasible coupling partners in the photocatalytic desulfonylative alkylation of *gem*-difluoroalkenes, allowing an efficient access to diverse (*E*)-fluoroalkenes bearing quaternary centers in decent yields (30–85%). Both the Ir catalyst and  $\text{NEt}_3$  were essential to generate the products. Of note, this protocol was applicable to a broad range of *gem*-difluoroalkenes with different substitution patterns. Cyclic tertiary sulfones were also compatible in this case, albeit with a decreased efficiency. However, desulfonylative coupling reactions with electron-neutral or electron-rich *gem*-difluorostyrenes preferentially afforded the *Z*-isomers (Scheme 543).<sup>835</sup>

### Scheme 543. Visible-Light-Promoted Defluorinative Coupling of Alkylsulfones with *gem*-Difluoroalkenes





Apart from the above-mentioned C–C bond formation reactions, visible-light-mediated defluorinative cross-coupling reactions of *gem*-difluoroalkenes to forge various C–X bonds were also extensively investigated in recent years. Specifically, the group of Yang and Xia in 2019 explored a visible-light-induced monofluoroalkenylation via the defluorination of *gem*-difluoroalkenes, followed by the cross-coupling with a series of aryl, benzyl, and alkyl thiols (Scheme 544).<sup>836</sup> This protocol presents as a practical way to construct multisubstituted fluorinated vinylsulfides under benign conditions with good tolerance of functional groups. As a particular highlight, the potential of this strategy was showcased by late-stage modification of dehydrocholic acid and diosgenin derivatives.

### Scheme 544. Visible-Light-Mediated Defluorinative Cross-Coupling of *gem*-Difluoroalkenes with Thiols





An efficient regioselective synthesis of diverse  $\alpha$ -fluoro- $\beta$ -arylalkenyl sulfides with *gem*-difluoroalkenes and sodium sulfonates as starting materials was realized by Shi and co-workers under green LED irradiation conditions (Scheme 545).<sup>837</sup> In this work, the authors employed  $\text{Na}_2$  (Eosin Y) as a photosensitizer in conjunction with  $\text{PPh}_3$  as a reductant to undergo a phosphine-associated deoxygenative functionalization of S–O bonds. Control experiments were carried out to ascertain the thiol radicals and anions as key intermediates in the proposed mechanism. Remarkably, regioselective isomerization of the C=C bond (from *Z* to *E*) was observed using green light without the presence of a photoinitiator.

### Scheme 545. Highly *E*-Selective Synthesis of $\alpha$ -Fluoro- $\beta$ -arylalkenyl Sulfides via Visible-Light-Induced Deoxygenation/Isomerization





R = H, 94% (*E/Z* = 92.5/7.5)  
 R = Me, 81% (*E/Z* = 92/8)  
 R = OMe, 58% (*E/Z* = 76/24)  
 R = OCF<sub>3</sub>, 65% (*E/Z* = 91/9)  
 R = CF<sub>3</sub>, 50% (*E/Z* = 92/8)  
 R = F, 82% (*E/Z* = 93/7)  
 R = Cl, 87% (*E/Z* = 92/8)  
 R = Br, 76% (*E/Z* = 93/7)  
 R = I, 50% (*E/Z* = 92.5/7.5)  
 R = CO<sub>2</sub>Me, 76% (*E/Z* = 89/11)



Fluorinated organoboron compounds are synthetically usefully building blocks in organic chemistry. In 2020, Wu's group successfully developed a photocatalyzed defluoroborylation via a straightforward B-H activation of an *N*-heterocyclic carbene (NHC) borane with *gem*-difluoroalkenes, affording a series of mono-fluoroalkenylboranes in 32–88% yield with moderate *Z*-selectivity (Scheme 546).<sup>83,8</sup> This strategy is highly atom-economic and regioselective, simple to operate and remarkably allows for the defluoroborylation of a wide substrate scope of multi-fluorinated compounds such as polyfluoroarenes, *gem*-difluoroalkenes, and trifluoromethylalkenes.



### Scheme 546. Photocatalyzed Defluoroborylation B-H Activation of an *N*-Heterocyclic Carbene (NHC) Borane



### 7.5. Annulation Reactions

With respect to the synthesis of heterocycles, *gem*-difluoroalkenes can also be used as starting materials together with transition-metal catalysts. In such instances, one or more F atoms usually undergo substitution to form new C–C or C–heteroatom bonds. Specifically, Ichikawa and collaborators in 2005 expanded their base-promoted nucleophilic 5-*endo-trig* cyclization strategy of 1,1-difluoro-1-alkenes with an *N*, *O* or a *C*-nucleophile to assemble diverse fluorinated heterocycles,<sup>839–842</sup> and reported an intramolecular Pd-catalyzed Heck-type 5-*endo-trig* cyclization reaction of 3,3-difluoroallylketone *O*-pentafluorobenzoyloximes via aminopalladium species, affording an efficient access to 5-fluoro-3*H*-pyrroles. This protocol can also be applied to the construction of fused tricyclic system, albeit in a moderate yield (Scheme 547).<sup>843</sup>


### Scheme 547. Synthesis of 5-Fluoro-3*H*-pyrroles via Intramolecular Heck-Type 5-*Endo-Trig* Cyclization



Later, the same group further discovered a cationic palladium-promoted Friedel–Crafts-type cyclization of 1,1-difluoro-1-alkenes with cationic [Pd(MeCN)<sub>4</sub>](BF<sub>4</sub>)<sub>2</sub> as a catalyst (Scheme 548).<sup>844</sup> Through the formation of metal-alkene complexes, the electrophilic activation of electron-deficient alkene by the cationic palladium(II) catalyst enables the intermediary  $\alpha,\alpha$ -difluoroalkyl palladium species to be readily generated via a Friedel–Crafts-type




cyclization. In the presence of  $\text{BF}_3 \cdot \text{Et}_2\text{O}$ , subsequent  $\beta$ -fluorine elimination was greatly accelerated to form 4-fluoro-1,2-dihydrophenanthrenes, which finally underwent hydrolysis to afford the corresponding cyclic ketones.

### Scheme 548. Palladium(II)-Catalyzed Friedel-Crafts Cyclization of 4,4-(Difluorohomoallyl)arenes





Cao's group expanded their C–F bond functionalization strategy, and illustrated the divergent synthesis of a broad range of heterocycles. For example, they in 2015 reported an unprecedented Cu(I)-catalyzed domino reaction for the construction of multisubstituted furans through the cyclization of *gem*-difluoroalkenes with a series of activated methylene carbonyls such as 1,3-dicarbonyl compounds, acetoacetonitrile, and phenylsulfonylacetone (Scheme 549a).<sup>845</sup> Subsequently, they further established a practical strategy for the rapid synthesis of highly functionalized aminothiophenes through the cyclization reaction of *gem*-difluoroalkenes with a diverse array of  $\beta$ -keto tertiary thioamides (Scheme 549b).<sup>846</sup> The reaction took place smoothly in the presence of  $\text{K}_2\text{CO}_3$  without the need for transition-metal catalysts, giving synthetically valuable *N,N*-disubstituted 2-aminothiophenes in 51–91% yield. The same group also discovered that the reaction of *gem*-difluoroalkenes with acyl hydrazides in DMSO under heating afforded a variety of 1,3,4-oxadiazoles in modest to excellent yields (Scheme 549c).<sup>847</sup>

### Scheme 549. Divergent Synthesis of Structurally Diverse Heterocycles from *gem*-Difluoroalkenes



An intramolecular electrophilic 5-*endo-trig* cyclization of  $\beta,\beta$ -difluoro-*o*-sulfonamidostyrenes to the direct synthesis of 2-fluoroindoles via  $\beta$ -F elimination was reported by Ichikawa's group using  $\text{AgSbF}_6$  as the catalyst and *N,O*-bis(trimethylsilyl)acetamide (BSA) as a fluoride captor (Scheme 550a).<sup>848</sup> In a subsequent report, Hao and Yang demonstrated that primary arylamines can couple with 1-halo-2-(2,2-difluorovinyl)benzene to produce a range of fluorine-containing indole derivatives (Scheme 550b).<sup>849</sup> First, Buchwald-Hartwig cross-coupling occurred in the presence of Pd catalyst to generate 2-amino-(2,2-difluorovinyl)benzene. Then, base-promoted 5-*endo-trig* intramolecular nucleophilic attack by *N* on *gem*-difluoroalkene occurs, followed by  $\beta$ -fluoride elimination to afford a wide variety of 2-fluoroindoles.


### Scheme 550. Synthesis of 2-Fluoroindoles from *gem*-Difluoroalkenes via Alkenyl C–F Bond Activation



The 1,2,4,5-tetrazine heterocyclic ring is one of the most versatile aromatic heterocycles, and its rapid synthesis is of particular interest across both organic and inorganic community.<sup>850</sup> In 2017, Hu's group reported a practical strategy for the straightforward construction of synthetically appealing aromatic tetrazines from the


corresponding *gem*-difluoroalkenes, a large variety of both symmetrical and asymmetrical 3,6-disubstituted 1,2,4,5-tetrazines can be readily prepared in 61–91% yield under ambient conditions (Scheme 551).<sup>851</sup>

### Scheme 551. Synthesis of 1,2,4,5-Tetrazines from *gem*-Difluoroalkenes




Langer's group in 2018 disclosed a base-promoted synthesis of dibenzo[*b,d*]oxepines via a vinylic C–F activation of 1-bromo-2-(2,2-difluorovinyl)benzenes with structurally diverse phenols, followed by a Pd(II)-catalyzed intramolecular C–H arylation, thereby giving rise to a crucial class of pharmaceutically relevant oxepine derivatives in modest to excellent yields (Scheme 552).<sup>852</sup> Apart from reactions with phenols, 4-hydroxypyridine was also a competent nucleophile to participate in such transformations. Reacting 4-hydroxypyridine with 1-bromo-2-(2,2-difluorovinyl)benzene in the presence of K<sub>2</sub>CO<sub>3</sub> and Pd(OAc)<sub>2</sub> catalyst resulted in the formation of a cyclic product with *N*-vinylic-4-pyridone as a likely key intermediate.

### Scheme 552. Base-Promoted One-Pot Synthesis of Dibenzo[*b,d*]oxepines



In 2020, Lu's group was able to prepare substituted indolizines via a Cu(I)-catalyzed cyclization reaction between 2-(pyridin-2-yl)acetates and *gem*-difluoroalkenes (Scheme 553).<sup>853</sup> This strategy exploited the cleavage and further functionalization of vinylic C–F bonds, thereafter allowing an effective approach to synthesize a series of indolizine derivatives in up to 87% yield along with excellent functional group tolerance.

### Scheme 553. Cu(I)-Catalyzed Cyclization of 2-(Pyridin-2-yl)acetates and *gem*-Difluoroalkenes




By taking advantage of 2,2-difluorovinyl tosylate as the fluorinated synthon, the group of Wu and Wang in 2004 elaborated the direct synthesis of 2-fluoroindolizines from *in situ* generated *N*-ylides of pyridiniums (Scheme 554a).<sup>854</sup> Inspired by this precedential work, Ren's group recently illustrated a base-promoted oxidative annulation reaction of *gem*-difluoroalkenes and pyridinium ylides with ambient air as the sole oxidant under benign conditions


(Scheme 554b).<sup>855</sup> The protocol underwent a formal [3 + 2] annulation process of the generated pyridinium ylide with *gem*-difluoroalkenes followed by facile  $\beta$ -F elimination to afford a series of multisubstituted 2-fluoroindolizines in good yields. Almost simultaneously, Peng's group also presented an analogous cycloaddition procedure by means of 1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU) as the base (Scheme 554c).<sup>856</sup>

### Scheme 554. Synthesis of 2-Fluoroindolizines through Oxidative [3 + 2] Annulation of Pyridinium Salts and *gem*-Difluoroalkenes


a) Wu, Wang et al., 2004



b) Ren et al., 2021




c) Zhang, Peng et al., 2021



More recently, the group of Li and Wang disclosed an efficient one-pot procedure for the rapid preparation of fluorinated polyfused heterocycles such as benzofuro[3,2-*b*]pyridines, 5*H*-indeno-[1,2-*b*]pyridines, and 5,6-dihydrobenzo[*h*]quinolines via the TfOH-promoted domino cyclization of azadienes with readily available difluoroenoxy silanes (Scheme 555).<sup>857</sup> This formal [4 + 2] cyclization strategy undergoes 1,4-difluoroalkylation, desulfonylation, intramolecular cyclization, and dehydrated and dehydrofluorinated aromatization sequences.


### Scheme 555. Cu(I)-Catalyzed Cyclization of 2-(Pyridin-2-yl)acetates and *gem*-Difluoroalkenes




Phenacenes are an important and versatile subclass of polycyclic aromatic hydrocarbons (PAHs) which can serve as organic semiconducting materials.<sup>858</sup> In the presence of a cationic Pd(II) catalyst, *gem*-difluoroalkenes such as 1,1-difluoro-1-alkenes,<sup>859,860</sup> or 1,1,2-trifluoro-1-alkenes<sup>861</sup> can undergo electrophilic Friedel-Crafts-type cyclizations to synthesize a broad array of highly appealing pinpoint-fluorinated phenacenes (Scheme 556a) or phenacenes (Scheme 556b). Mechanistically, the Pd catalyst first coordinated to the trifluoroalkene to form a  $\pi$ -complex. Electrophilic Friedel-Crafts-type carbopalladation and rearomatization occurred to produce a cyclic Pd(II)-alkyl intermediate, which then experienced  $\beta$ -F elimination assisted by  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  to afford the fluorinated phenacenes and regenerate the active cationic Pd(II) catalyst.

### Scheme 556. Synthesis of Fluorinated Polycyclic Aromatic Hydrocarbons via Palladium-Catalyzed Electrophilic Friedel-Crafts-Type Cyclization

a) Ichikawa et al., 2015




b) Ichikawa et al., 2017



In contrast to the above-mentioned  $\beta$ -F elimination process, Chatani and colleagues in 2010 reported the elegant work of nickel-catalyzed cyclization of 1,1-difluoro-1,6-enynes with organozinc reagents involving  $\alpha$ -F elimination.<sup>862,863</sup> Mechanistically, the process is initiated by an oxidative cyclization of both alkene and alkyne moieties with the  $\text{Ni}^0$  complex, resulting in the formation of intermediary  $\alpha, \alpha$ -difluoronickelacyclopentenes. The authors claimed that organozinc reagents serves as a bifunctional activator, a Lewis base with respect to  $\text{Ni}^{\text{II}}$  and a Lewis acid with respect to the C-F bond, which greatly facilitated subsequent  $\alpha$ -F elimination to generate bicyclo[3.2.0]heptene derivatives (Scheme 557).<sup>864</sup>


### Scheme 557. Nickel-Catalyzed Cyclization of 1,1-Difluoro-1,6-enynes with Organozinc Reagents




Inspired by this work, Ichikawa's group in 2015 realized an unprecedented nickel-catalyzed [2 + 2 + 2] cyclization of 1,1-difluoroethylene with alkynes to directly assemble a series of tetrasubstituted fluoroarenes by sequential  $\alpha$ -F elimination of  $\alpha,\alpha$ -difluorinated nickelacycloheptadiene species, generated by the oxidative cyclization of 1,1-difluoroethylene and alkynes with nickel(0), and subsequent  $\beta$ -H elimination of the intermediary cyclohexadienylnickel(II) fluoride (Scheme 558a).<sup>865</sup> Afterward, the group of Ogoshi and Ohashi elaborately explored the nickel(0)-mediated defluorinative coupling of tetrafluoroethylene (TFE) with styrenes. In the presence of Ni(cod)<sub>2</sub> and PCy<sub>3</sub>, exposure of styrene to TFE resulted in the formation of a well-defined nickelacycle possessing a unique  $\eta^3$ - $\pi$ -benzyl structure in 89% isolated yield (Scheme 558b).<sup>866</sup> Treating this nickelacycle with BF<sub>3</sub>·Et<sub>2</sub>O at room temperature smoothly generated 1,2-difluoro-3-phenyl cyclobutene in 60% yield through sequential both  $\alpha$ - and  $\beta$ -F eliminations.

### Scheme 558. Defluorinative Cyclization Reaction of Fluoroolefins through $\alpha$ -Fluorine Elimination

a) Ichikawa et al., 2015



b) Ohashi, Ogoshi et al., 2017



## 8. CONCLUSIONS AND OUTLOOK

In this comprehensive review, we summarised the latest developments in alkenyl sp<sup>2</sup> C–H and C–F bond functionalization reactions. The development of practical and broadly applicable methods for alkenyl sp<sup>2</sup> C–H bond functionalization has attracted tremendous attention because they can provide efficient access to synthetically important functionalized alkenes and diene derivatives from easily accessible simple alkenes. Significant advances in developing new catalytic systems that can effectively carry out alkenyl C–H bonds of alkenes and their derivatives have been made in the past decades. Nevertheless, there are still many challenges remain. Outstanding problems that remained to be solved include the development of alkenyl sp<sup>2</sup> C–H and C–F bond functionalization that can work under greener and milder reaction conditions. Unfortunately, many of the reported methods still require the use of expensive transition metals such as palladium and rhodium catalysts. In addition, the catalytic loading of these catalysts remained high and there were very few reports that used less than 1 mol% of the catalysts. The use of cheaper main group catalysts such as Fe is still rare and not to mention the development of metal-free systems. Metal-free photoredox methods is an attractive approach and a few reports have emerged in recent years. In many cases, these reactions were carried out in organic solvents under high temperature. Moreover, the development of new alkenyl sp<sup>2</sup> C–H and C–F bond functionalization that can work under metal-free system in a biocompatible reaction condition will be an interesting area to explore in the coming years.

For these reactions to be fully adopted for the synthesis of pharmaceuticals and complex molecules, it is extremely important that the scope and limitations of these methods be well studied. As shown in this review, many types of alkenes and coupling partners have been shown to work effectively. **There are only sporadic examples using simple alkenes, alkyl halides/alcohols or alkanes as coupling partners.** Additional work to widen the scope of alkenes and the coupling partners will further help to expand the synthetic utility of these methods for the synthesis of complex molecules.

Another problem in this area of research is the lack of stereochemical studies (regio, *E/Z*, enantioselective, etc) in this exciting area. Although the use of various methods such as the use of steric effect, directing groups and 1,*n*-metal shift strategies have been successfully reported, the use of catalyst to control the regio- and *E/Z* selectivities is still lacking. More research in the asymmetric areas such as kinetic resolution, enantioselective control, etc. will certainly make this method an extremely attractive strategy for the stereoselective synthesis of alkenes.

Another area that needs to be expanded is the development of new intramolecular versions and annulation methods for the construction of cyclic alkenes. Although sporadic reports on the intramolecular coupling for the synthesis of macrocycles have reported by Loh and other researchers, extensive work to expand to different

macrocycles will greatly expand the scope of this method. Reports on the annulation to construct smaller rings such as 5, 6, etc. are still rare.

In summary, this comprehensive review covers most of the published work in the fertile area of alkenyl sp<sup>2</sup> C–H and C–F bond functionalization methods. Due to their high efficiency, atom-economy nature as well as easy accessibility of the starting materials, these methods have emerged rapidly as useful strategies for the construction of pharmaceuticals, functional materials and bioactive natural products. Meanwhile, they may serve as alternative or complimentary strategies to classical cross-coupling reactions, Wittig or olefin metathesis for stereoselective synthesis of highly functionalized alkenes. We anticipate that this review will contribute to inspire broad research interest and stimulate new breakthroughs in the development of innovative and broadly applicable strategies to expand the research of alkenyl sp<sup>2</sup> C–H and C–F bond functionalization reactions.

## AUTHOR INFORMATION

### Corresponding Author

**Teck-Peng Loh** – College of Advanced Interdisciplinary Science and Technology, Henan University of Technology, Zhengzhou, 450001, China; School of School of Chemistry, Chemical Engineering and Biotechnology, Nanyang Technological University, 637371, Singapore; orcid.org/0000-0002-2936-337X; Email: teckpeng@ntu.edu.sg

### Authors

**Ming-Zhu Lu** – College of Advanced Interdisciplinary Science and Technology, Henan University of Technology, Zhengzhou, 450001, China; School of School of Chemistry, Chemical Engineering and Biotechnology, Nanyang Technological University, 637371, Singapore; orcid.org/0000-0002-6333-0803

**Jeffrey Goh** – School of School of Chemistry, Chemical Engineering and Biotechnology, Nanyang Technological University, 637371, Singapore

**Manikantha Maraswami** – School of School of Chemistry, Chemical Engineering and Biotechnology, Nanyang Technological University, 637371, Singapore

**Zhenhua Jia** – Institute of Advanced Synthesis, School of Chemistry and Molecular Engineering Nanjing Tech University, 30 South Puzhu Road, Nanjing, 211816, China; orcid.org/0000-0002-5154-4839

**Jie-Sheng Tian** – School of Chemistry and Chemical Engineering, Northwestern Polytechnical University (NPU), Xi'an 710072, China

### Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

### Notes

The authors declare no competing financial interest.

### Biographies

Ming-Zhu Lu was born in Huaian, Jiangsu province of P. R. China. He obtained his Ph.D. degree on organic chemistry from University of Science and Technology of China (USTC) in 2015, under the supervision of Prof. Teck-Peng Loh. He then carried out postdoctoral research with Prof. Jin-Quan Yu (2015-2017) at the State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry (SIOC). Now, he is working as a senior research fellow in the research group of Prof. Teck-Peng Loh at Nanyang Technological University, Singapore. He joined Henan University of Technology as a full professor in 2022. His current research interests mainly focus on transition-metal-catalyzed C–H functionalization reactions.

Jeffrey Goh was born in Singapore in 1992. He earned a Diploma in Chemical Process Technology (Industrial Chemistry) from Singapore Polytechnic in 2012 and subsequently went on to Nanyang Technological University to complete his B.Sc. (Highest Distinction) in Chemistry and Biological Chemistry in 2018. He is currently working on his Ph.D. under the supervision of Prof. Teck-Peng Loh, focusing on the development of water-based C–X bond forming reactions. His research interests include green chemistry, C–H activation and asymmetric catalysis.

Manikantha Maraswami received his B.S. and M.S. degree from Karnatak University, Dharwad and in 2018, completed his Ph.D. degree in chemistry from Nanyang Technological University, Singapore with Associate Professor Gang Chen. In early 2018, he began to research with Professor Teck-Peng Loh at NTU, Singapore as a research fellow. Since 2021, he has been a postdoc associate at the University of Massachusetts Medical School, MA, USA. He is now working on the design and synthesis of protease inhibitors at UMASS.

Zhenhua Jia is a full professor of Institute of Advanced Synthesis, School of Chemistry and Molecular Engineering in Nanjing Tech University. He received his B.S. in 2004 from Tianjin University and Ph.D. in 2013 from Sun Yat-sen University jointed with McGill University under the supervision of Prof. Albert S. C. Chan and Prof. Chao-Jun Li. Then he continued to pursue his postdoctoral research in Marquette University, Chinese University of Hong Kong and University of Alberta with Prof. Chae S. Yi, Prof. Henry N.C. Wong and Prof. Frederick G. West respectively. At the beginning of 2018, he joined Prof. Teck-Peng Loh's team and hold his current position. His research interest includes green and sustainable synthesis, organic transformations under bio-compatible conditions.

Jie-Sheng Tian is currently a full Professor of School of Chemistry and Chemical Engineering in Northwestern Polytechnical University (China). He received his Ph.D. in organic chemistry from Nanyang Technological University in 2012 under the supervision of Prof. Teck-Peng Loh. He then did his postdoctoral studies at Nanyang Technological University with Prof. Teck-Peng Loh. After working as a Research Associate Professor in University of Science and Technology of China for around one year, he moved to Nanjing Tech University as a full Professor to start his independent academic career in 2016. His research interests focus on the development of green synthetic method for drug and polymer synthesis.

Teck-Peng Loh is a distinguished university professor of chemistry at Nanyang Technological University, Singapore. Under the tutelage of Professor E. J. Corey, he obtained his PhD (1994) from Harvard University. He has published 426 international refereed papers in reputable chemistry journals bearing high impact factor and 10 patents. He has been invited to give more than 100 lectures in many institutions in the world and plenary and keynote talks in conferences such as the ICOS, OMCOS, Asian European Symposium, etc. He has also served as chairman of many of these conferences. He has been awarded outstanding researcher awards from both National University of Singapore and Nanyang Technological University. In 2017 he received the Yoshida Prize (Japan) and the prestigious President's Science Award (individual) Singapore. He has been elected Fellow, Academia of Sciences, Singapore (2018) and Fellow of Academia of Sciences, Malaysia since 2010. His research work mainly focuses on the development of new synthetic methodology, green chemistry, and synthesis of natural and unnatural products.

## ACKNOWLEDGMENTS

We gratefully acknowledge the financial support from Distinguished University Professor grant (Nanyang Technological University), AcRF Tier 1 grants from the Ministry of Education of Singapore (RG107/19, RG11/20 and RT14/20), and the Agency for Science, Technology and Research (A\*STAR) under its MTC Individual Research Grants (IRG). The authors also gratefully acknowledge the financial support from the National Natural Science Foundation of China (No. 22101095).

## REFERENCES

- Maryanoff, B. E.; Reitz, A. B. The Wittig Olefination Reaction and Modifications Involving Phosphoryl-Stabilized Carbanions. Stereochemistry, Mechanism, and Selected Synthetic Aspects. *Chem. Rev.* **1989**, *89*, 863–927.
- Byrne, P. A.; Gilheany, D. G. The Modern Interpretation of the Wittig Reaction Mechanism. *Chem. Soc. Rev.* **2013**, *42*, 6670–6696.
- Rocha, D. H. A.; Pinto, D. C. G. A.; Silva, A. M. S. Applications of the Wittig Reaction on the Synthesis of Natural and Natural-Analogue Heterocyclic Compounds. *Eur. J. Org. Chem.* **2018**, *2018*, 2243–2257.
- Fürstner, A. Olefin Metathesis and Beyond. *Angew. Chem., Int. Ed.* **2000**, *39*, 3012–3043.
- Vougioukalakis, G. C.; Grubbs, R. H. Ruthenium-Based Heterocyclic Carbene-Coordinated Olefin Metathesis Catalysts. *Chem. Rev.* **2010**, *110*, 1746–1787.
- Montgomery, T. P.; Ahmed, T. S.; Grubbs, R. H. Stereoretentive Olefin Metathesis: An Avenue to Kinetic Selectivity. *Angew. Chem., Int. Ed.* **2017**, *56*, 11024–11036.
- Ogba, O. M.; Warner, N. C.; O'Leary, D. J.; Grubbs, R. H. Recent Advances in Ruthenium-Based Olefin Metathesis. *Chem. Soc. Rev.* **2018**, *47*, 4510–4544.
- Albright, H.; Davis, A. J.; Gomez-Lopez, J. L.; Vonesh, H. L.; Quach, P. K.; Lambert, T. H.; Schindler, C. S. Carbonyl-Olefin Metathesis. *Chem. Rev.* **2021**, *121*, 9359–9406.
- Stereoselective Alkene Synthesis; Jianbo Wang, Springer, 2012.
- Sigman, M. S.; Werner, E. W. Imparting Catalyst Control upon Classical Palladium-Catalyzed Alkenyl C–H Bond Functionalization Reactions. *Acc. Chem. Res.* **2012**, *45*, 6, 874–884.

- Tang, S.; Liu, K.; Liu, C.; Lei, A. Olefinic C–H Functionalization Through Radical Alkenylation. *Chem. Soc. Rev.* **2015**, *44*, 1070–1082.
- Blanksby, S. J.; Ellison, G. B. Bond Dissociation Energies of Organic Molecules. *Acc. Chem. Res.* **2003**, *36*, 255–263.
- Shang, X.; Liu, Z.-Q. Transition Metal-Catalyzed C<sub>vinyl</sub>–C<sub>vinyl</sub> Bond Formation via Double C<sub>vinyl</sub>–H Bond Activation. *Chem. Soc. Rev.* **2013**, *42*, 3253–3260.
- Wang, K.; Hu, F.; Zhang, Y.; Wang, J. Directing Group-Assisted Transition-Metal-Catalyzed Vinylic C–H Bond Functionalization. *Sci. China: Chem.* **2015**, *58*, 1252–1265.
- Maraswami, M.; Loh, T.-P. Transition-Metal-Catalyzed Alkenyl sp<sup>2</sup> C–H Activation: A Short Account. *Synthesis* **2019**, *51*, 1049–1062.
- Liu, B.; Yang, L.; Li, P.; Wang, F.; Li, X. Recent Advances in Transition Metal-Catalyzed Olefinic C–H Functionalization. *Org. Chem. Front.* **2021**, *8*, 1085–1101.
- Zhang, J.; Lu, X.; Shen, C.; Xu, L.; Ding, L.; Zhong, G. Recent Advances in Chelation-Assisted Site- and Stereoselective Alkenyl C–H Functionalization. *Chem. Soc. Rev.* **2021**, *50*, 3263–3314.
- Zhang, X.; Cao, S.; Recent Advances in the Synthesis and C–F Functionalization of *gem*-Difluoroalkenes. *Tetrahedron Lett.* **2017**, *58*, 375–392.
- Fujita, T.; Fuchibe, K.; Ichikawa, J. Transition-Metal-Mediated and -Catalyzed C–F Bond Activation by Fluorine Elimination. *Angew. Chem., Int. Ed.* **2019**, *58*, 390–402.
- Liu, C.; Zeng, H.; Zhu, C.; Jiang, H. Recent Advances in Three-Component Difunctionalization of *gem*-Difluoroalkenes. *Chem. Commun.* **2020**, *56*, 10442–10452.
- Koley, S.; Altman, R. A. Recent Advances in Transition Metal-Catalyzed Functionalization of *gem*-Difluoroalkenes. *Isr. J. Chem.* **2020**, *60*, 313–339.
- Wang, J.; Gao, H.; Shi, C.; Chen, G.; Tan, X.; Chen, X.; Xu, L.; Cai, X.; Huang, B.; Li, H. Recent Advances in Radical-Based C–F Bond Activation of Polyfluoroarenes and *gem*-Difluoroalkenes. *Chem. Commun.* **2021**, *57*, 12203–12217.
- Xu, W.; Zhang, Q.; Shao, Q.; Xia, C.; Wu, M. Photocatalytic C–F Bond Activation of Fluoroarenes, *gem*-Difluoroalkenes and Trifluoromethylarenes. *Asian J. Org. Chem.* **2021**, *10*, 2454–2472.
- Sorrentino, J. P.; Altman, R. A. Fluorine-Retentive Strategies for the Functionalization of *gem*-Difluoroalkenes. *Synthesis* **2021**, *53*, 3935–3950.
- Zhang, J.; Geng, S.; Feng, Z. Advances in Silylation and Borylation of Fluoroarenes and *gem*-Difluoroalkenes via C–F Bond Cleavage. *Chem. Commun.* **2021**, *57*, 11922–11934.
- Simur, T. T.; Ye, T.; Yu, Y.-J.; Zhang, F.-L.; Wang, Y.-F. C–F Bond Functionalizations of Trifluoromethyl Groups via Radical Intermediates. *Chin. Chem. Lett.* **2022**, *33*, 1193–1198.
- Ribas, X. *C–H and C–X Bond Functionalization: Transition Metal Mediation*; RSC, Cambridge, U.K., 2013.
- Joseph, P. A. J.; Priyadarshini, S. Copper-Mediated C–X Functionalization of Aryl Halides. *Org. Process Res. Dev.* **2017**, *21*, 1889–1924.
- Huang, C.-Y.; Li, J.; Li, C.-J. Photocatalytic C(sp<sup>3</sup>) Radical Generation via C–H, C–C, and C–X Bond Cleavage. *Chem. Sci.* **2022**, *13*, 5465–5504.
- Yang, X.; Wu, C.; Su, W.; Yu, J. Mechanochemical C–X/C–H Functionalization: An Alternative Strategic Access to Pharmaceuticals. *Eur. J. Org. Chem.* **2022**, *2022*, e202101440.
- da Silva, M. J.; Gusevskaya, E. V. Palladium-Catalyzed Oxidation of Monoterpenes: Novel Tandem Oxidative Coupling–Oxidation of Camphene by Dioxxygen. *J. Mol. Catal. A: Chem.* **2001**, *176*, 23–27.
- da Silva, M. J.; Goncalves, J. A.; Alves, R. B.; Howarth, O. W.; Gusevskaya, E. V. Palladium Catalyzed Transformations of Monoterpenes: Stereoselective Deuteriation and Oxidative

- Dimerization of Camphene. *J. Organomet. Chem.* **2004**, *689*, 302–308.
- (33) Hubert, P.; Seibel, E.; Beemelmans, C.; Campagne, J.-M.; Figueiredo, R. M. Stereoselective Construction of (*E,Z*)-1,3-Dienes and Its Application in Natural Product Synthesis. *Adv. Synth. Catal.* **2020**, *362*, 5532–5575.
- (34) Soengas, R. G.; Rodríguez-Solla, H. Modern Synthetic Methods for the Stereoselective Construction of 1,3-Dienes. *Molecules* **2021**, *26*, 249–289.
- (35) Xu, Y.-H.; Lu, J.; Loh, T.-P. Direct Cross-Coupling Reaction of Simple Alkenes with Acrylates Catalyzed by Palladium Catalyst. *J. Am. Chem. Soc.* **2009**, *131*, 1372–1373.
- (36) Xu, Y.-H.; Wang, W. -J.; Wen, Z.-K.; Hartley, J. J.; Loh, T.-P. Palladium-Catalyzed Direct Cross-Coupling Reaction between Indenes and Electron-Deficient Alkenes. *Tetrahedron Lett.* **2010**, *51*, 3504–3507.
- (37) Wen, Z.-K.; Xu, Y.-H.; Loh, T.-P. Palladium-Catalyzed Cross-Coupling of Unactivated Alkenes with Acrylates: Application to the Synthesis of the C<sub>13</sub>-C<sub>21</sub> Fragment of Palmerolide A. *Chem. - Eur. J.* **2012**, *18*, 13284–13287.
- (38) Wen, Z.-K.; Xu, Y.-H.; Loh, T.-P. Palladium(II)-Catalyzed Cross-Coupling of Simple Alkenes with Acrylates: A Direct Approach to 1,3-Dienes through C–H Activation. *Chem. Sci.* **2013**, *4*, 4520–4524.
- (39) Zhong, X.-M.; Cheng, G.-J.; Chen, P.; Zhang, X.; Wu, Y.-D. Mechanistic Study on Pd/Mono-*N*-protected Amino Acid Catalyzed Vinyl–Vinyl Coupling Reactions: Reactivity and *E/Z* Selectivity. *Org. Lett.* **2016**, *18*, 5240–5243.
- (40) Zhang, X.; Wang, M.; Zhang, M.-X.; Xu, Y.-H.; Loh, T.-P. Synthesis of Dienyl Ketones via Palladium(II)-Catalyzed Direct Cross-Coupling Reactions between Simple Alkenes and Vinyl Ketones: Application to the Synthesis of Vitamin A<sub>1</sub> and Bornelone. *Org. Lett.* **2013**, *15*, 5531–5533.
- (41) Zhang, Y.; Cui, Z.; Li, Z.; Liu, Z.-Q. Pd(II)-Catalyzed Dehydrogenative Olefination of Vinylic C–H Bonds with Allylic Esters: General and Selective Access to Linear 1,3-Butadienes. *Org. Lett.* **2012**, *14*, 1838–1841.
- (42) Gigant, N.; Bäckvall, J. E. Synthesis of Conjugated Dienes Via a Biomimetic Aerobic Oxidative Coupling of Two C<sub>Vinyl</sub>-H Bonds. *Chem. - Eur. J.* **2013**, *19*, 10799–10803.
- (43) Bras, J. L.; Muzart, J. Intermolecular Dehydrogenative Heck Reactions. *Chem. Rev.* **2011**, *111*, 1170–1214.
- (44) Delcamp, J. H.; Brucks, A. P.; White, M. C. A General and Highly Selective Chelate-Controlled Intermolecular Oxidative Heck Reaction. *J. Am. Chem. Soc.* **2008**, *130*, 11270–11271.
- (45) Werner, E. W.; Sigaman, M. S. A Highly Selective and General Palladium Catalyst for the Oxidative Heck Reaction of Electronically Nonbiased Olefins. *J. Am. Chem. Soc.* **2010**, *132*, 13981–13983.
- (46) Werner, E. W.; Sigaman, M. S. Operationally Simple and Highly (*E*)-Styrenyl-Selective Heck Reactions of Electronically Nonbiased Olefins. *J. Am. Chem. Soc.* **2011**, *133*, 9692–9695.
- (47) Prediger, P.; Barbosa, L. F.; Génisson, Y.; Correia, C. R. D. Substrate-Directable Heck Reactions with Arenediazonium Salts. The Regio- and Stereoselective Arylation of Allylamine Derivatives and Applications in the Synthesis of Naftifine and Abamines. *J. Org. Chem.* **2011**, *76*, 7737–7749.
- (48) Maraswami, M.; Diggins, T.; Goh, J.; Tio, R.; Ong, W. Q. R.; Hirao, H.; Loh, T.-P. Intramolecular Alkene–Alkene Coupling via Rh(III)-Catalyzed Alkenyl sp<sup>2</sup> C–H Functionalization: Divergent Pathways to Indene or  $\alpha$ -Naphthol Derivatives. *ACS Catal.* **2021**, *11*, 11494–11500.
- (49) Finkbeiner, P.; Kloekner, U.; Nachtsheim, B. J. OH-Directed Alkynylation of 2-Vinylphenols with Ethynyl Benzenes: A Fast Access to Terminal 1,3-Enynes. *Angew. Chem., Int. Ed.* **2015**, *54*, 4949–4952.
- (50) Zhang, Y.; Shang, T.; Li, L.; He, Y.; Wen, T.; Tu, Y.; Wang, S.; Lin, D. Rh(III)-Catalyzed C–H Activation-Desymmetrization of Diazabicycles Using Enol as Directing Group: A Straightforward Approach to Difunctionalized Cyclopentenes. *Tetrahedron Lett.* **2018**, *59*, 1394–1397.
- (51) He, H.; Liu, W.-B.; Dai, L.-X.; You, S.-L. Ir-Catalyzed Cross-Coupling of Styrene Derivatives with Allylic Carbonates: Free Amine Assisted Vinyl C–H Bond Activation. *J. Am. Chem. Soc.* **2009**, *131*, 8346–8347.
- (52) Caspers, L. D.; Finkbeiner, P.; Nachtsheim, B. J. Direct Electrophilic C–H Alkynylation of Unprotected 2-Vinylanilines. *Chem. - Eur. J.* **2017**, *23*, 2748–2752.
- (53) Boelke, A.; Caspers, L. D.; Nachtsheim, B. J. NH<sub>2</sub>-Directed C–H Alkynylation of 2-Vinylanilines with Vinylbenziodoxolones. *Org. Lett.* **2017**, *19*, 5344–5347.
- (54) Jin, L.; Zhang, P.; Li, Y.; Yu, X.; Shi, B.-F. Atroposelective Synthesis of Conjugated Diene-Based Axially Chiral Styrenes via Pd(II)-Catalyzed Thioether-Directed Alkenyl C–H Olefination. *J. Am. Chem. Soc.* **2021**, *143*, 12335–12344.
- (55) Chen, G.; Gui, J.; Li, L.; Liao, J. Chiral Sulfoxide-Olefin Ligands: Completely Switchable Stereoselectivity in Rhodium-Catalyzed Asymmetric Conjugate Additions. *Angew. Chem., Int. Ed.* **2011**, *50*, 7681–7685.
- (56) Qi, W.-Y.; Zhu, T.-S.; Xu, M.-H. Design of Chiral Sulfoxide Olefins as a New Class of Sulfur-Based Olefin Ligands for Asymmetric Catalysis. *Org. Lett.* **2011**, *13*, 3410–3413.
- (57) Feng, X.; Nie, Y.; Yang, J.; Du, H. Rh(I)-Catalyzed Asymmetric 1,2-Addition to  $\alpha$ -Diketones with Chiral Sulfur–Alkene Hybrid Ligands. *Org. Lett.* **2012**, *14*, 624–627.
- (58) Li, Y.; Xu, M.-H. Simple Sulfur–Olefins as New Promising Chiral Ligands for Asymmetric Catalysis. *Chem. Commun.* **2014**, *50*, 3771–3782.
- (59) Dai, D.-T.; Yang, M.-W.; Chen, Z.-Y.; Wang, Z.-L.; Xu, Y.-H. Chelation-Controlled Stereospecific Cross-Coupling Reaction between Alkenes for Atroposelective Synthesis of Axially Chiral Conjugated Dienes. *Org. Lett.* **2022**, *24*, 1979–1984.
- (60) Zhang, F.-L.; Hong, K.; Li, T.-J.; Park, H.; Yu, J.-Q. Functionalization of C(sp<sup>3</sup>)-H Bonds Using a Transient Directing Group. *Science* **2016**, *351*, 252–256.
- (61) Zhao, Q.; Poisson, T.; Pannecoucke, X.; Besset, T. The Transient Directing Group Strategy: A New Trend in Transition-Metal-Catalyzed C–H Bond Functionalization. *Synthesis* **2017**, *49*, 4808–4826.
- (62) Gandeepan, P.; Ackermann, L. Transient Directing Groups for Transformative C–H Activation by Synergistic Metal Catalysis. *Chem* **2018**, *4*, 199–222.
- (63) Higham, J. I.; Bull, J. A. Transient Imine Directing Groups for the C–H Functionalisation of Aldehydes, Ketones and Amines: An Update 2018–2020. *Org. Biomol. Chem.* **2020**, *18*, 7291–7315.
- (64) Ha, H.; Lee, J.; Park, M. H.; Jung, B.; Kim, M. Transient Directing Group-Assisted C–H Bond Functionalization of Aliphatic Amines: Strategies for Efficiency and Site-Selectivity. *Bull. Korean Chem. Soc.* **2020**, *41*, 582–587.
- (65) Zhang, M.; Zhong, Z.; Liao, L.; Zhang, A.-Q. Application of a Transient Directing Strategy in Cyclization Reactions via C–H Activation. *Org. Chem. Front.* **2022**, *9*, 3882–3896.
- (66) Lapuh, M. I.; Mazeh, S.; Besset, T. Chiral Transient Directing Groups in Transition-Metal-Catalyzed Enantioselective C–H Bond Functionalization. *ACS Catal.* **2020**, *10*, 12898–12919.
- (67) Liao, G.; Zhang, T.; Lin, Z.-K.; Shi, B.-F. Transition Metal-Catalyzed Enantioselective C–H Functionalization via Chiral Transient Directing Group Strategies. *Angew. Chem., Int. Ed.* **2020**, *59*, 19773–19786.
- (68) Shen, C.; Zhu, Y.; Shen, W.; Jin, S.; Zhong, G.; Luo, S.; Xu, L.; Zhong, L.; Zhang, J. Access to Axially Chiral Aryl 1,3-Dienes by Transient Group Directed Asymmetric C–H Alkylations. *Org. Chem. Front.* **2022**, *9*, 2109–2115.

- (69) Shen, C.; Zhu, Y.; Shen, W.; Jin, S.; Zhong, L.; Luo, S.; Xu, L.; Zhong, G.; Zhang, J. Construction of Axial Chirality by Asymmetric  $\alpha$ -C–H Alkenylation of Aryl Alkenes. *Org. Chem. Front.* **2022**, *9*, 10.1039/d2qo00803c.
- (70) Liu, M.; Sun, J.; Erbay, T. G.; Ni, H.-Q.; Martin-Montero, R. Liu, P.; Engle, K. M. Pd<sup>II</sup>-Catalyzed C(alkenyl)–H Activation Facilitated by a Transient Directing Group. *Angew. Chem., Int. Ed.* **2022**, *61*, e202203624.
- (71) Shen, C.; Zhu, Y.; Jin, S.; Xu, K.; Luo, S.; Xu, L.; Zhong, G.; Zhong, L.; Zhang, J. Regio- and Stereo-Selective Olefinic C–H Functionalization of Aryl Alkenes in Ethanol. *Org. Chem. Front.* **2022**, *9*, 989–994.
- (72) Tian, Q.; Larock, R. C. Synthesis of 9-Alkylidene-9H-fluorenes by a Novel Palladium-Catalyzed Rearrangement. *Org. Lett.* **2020**, *2*, 3329–3332.
- (73) Hu, T.-J.; Zhang, G.; Chen, Y.-H.; Feng, C.-G.; Lin, G.-Q. Borylation of Olefin C–H Bond via Aryl to Vinyl Palladium 1,4-Migration. *J. Am. Chem. Soc.* **2016**, *138*, 2897–2900.
- (74) Hu, T.-J.; Li, M.-Y.; Zhao, Q.; Feng, C.-G.; Lin, G.-Q. Highly Stereoselective Synthesis of 1,3-Dienes through an Aryl to Vinyl 1,4-Palladium Migration/Heck Sequence. *Angew. Chem., Int. Ed.* **2018**, *57*, 5871–5875.
- (75) Xue, Z.-J.; Li, M.-Y.; Zhu, B.-B.; He, Z.-T.; Feng, C.-G.; Lin, G.-Q. A 1,4-Palladium Migration/Heck Sequence with Unactivated Alkenes: Stereoselective Synthesis of Trisubstituted 1,3-Dienes. *Adv. Synth. Catal.* **2021**, *363*, 2089–2092.
- (76) Xue, Z.-J.; Li, M.-Y.; Zhu, B.-B.; He, Z.-T.; Feng, C.-G.; Lin, G.-Q. Stereoselective Synthesis of Conjugated Trienes via 1,4-Palladium Migration/Heck Sequence. *Chem. Commun.* **2020**, *56*, 14420–14422.
- (77) BouzBouz, S.; Cossy, J. Tetrafibricin: Synthesis of the C1–C13, C15–C25, and C27–C40 Fragments. *Org. Lett.* **2004**, *6*, 3469–3472.
- (78) Molander, G. A.; Dehmel, F. Formal Total Synthesis of Oximidine II via a Suzuki-Type Cross-Coupling Macrocyclization Employing Potassium Organotrifluoroborates. *J. Am. Chem. Soc.* **2004**, *126*, 10313–10318.
- (79) Shimizu, M.; Tatsumi, H.; Mochida, K.; Shimono, K.; Hiyama, T. Synthesis, Crystal Structure, and Photophysical Properties of (1*E*,3*E*,5*E*)-1,3,4,6-Tetraarylhexa-1,3,5-trienes: A New Class of Fluorophores Exhibiting Aggregation-Induced Emission. *Chem. - Asian J.* **2009**, *4*, 1289–1297.
- (80) Li, M.-Y.; Han, P.; Hu, T.-J.; Wei, D.; Zhang, G.; Qin, A.; Feng, C.-G.; Tang, B. Z.; Lin, G.-Q. Suzuki-Miyaura Coupling Enabled by Aryl to Vinyl 1,4-Palladium Migration. *iScience* **2020**, *23*, 100966.
- (81) Wang, Q.; Chen, R.; Lou, J.; Zhang, D. H.; Zhou, Y.-G.; Yu, Z. Highly Regioselective C–H Alkylation of Alkenes Through an Aryl to Vinyl 1,4-Palladium Migration/C–C Cleavage Cascade. *ACS Catal.* **2019**, *9*, 11669–11675.
- (82) Regioselective Tandem C–H Alkylation/Coupling Reaction of *ortho*-Iodophenylethylenes via C,C-Pallada(II)cycles. Zhu, B.-B.; Ye, W.-B.; He, Z.-T.; Zhang, S.-S.; Feng, C.-G.; Lin, G.-Q. *ACS Catal.* **2021**, *11*, 12123–12132.
- (83) Lu, X.; Huang, Y. Stereospecific Cyanation of the Olefinic C–H Bond Enabled by 1,4-Rhodium Migration. *Org. Chem. Front.* **2021**, *8*, 3008–3013.
- (84) Sheykhani, M.; Shafiee-Pour, M.; Abbasnia, C–H Activation under the Guise of Diels-Alder Reaction: Annulation toward the Synthesis of Benzo[*e*]isoindole-1,3-diones. *Org. Lett.* **2017**, *19*, 1270–1273.
- (85) Wei, D.; Hu, T.-J.; Feng, C.-G.; Lin, G.-Q. Synthesis of Substituted Naphthalenes by 1,4-Palladium Migration Involved Annulation with Internal Alkynes. *Chin. J. Chem.* **2018**, *36*, 743–748.
- (86) Wei, D.; Li, M.-Y.; Zhu, B.-B.; Yang, X.-D.; Zhang, F.; Feng, C.-G.; Lin, G.-Q. Sequential Cross-Coupling/Annulation of *ortho*-Vinyl Bromobenzenes with Aromatic Bromides for the Synthesis of Polycyclic Aromatic Compounds. *Angew. Chem., Int. Ed.* **2019**, *58*, 16543–16547.
- (87) Rocaboy, R.; Baudoin, O. 1,4-Palladium Shift/C(sp<sup>3</sup>)–H Activation Strategy for the Remote Construction of Five-Membered Rings. *Org. Lett.* **2019**, *21*, 1434–1437.
- (88) Quideau, S.; Deffieux, D.; Douat-Casassus, C.; Pouységu, L. Plant Polyphenols: Chemical Properties, Biological Activities, and Synthesis. *Angew. Chem., Int. Ed.* **2011**, *50*, 586–621.
- (89) Huang, Z.; Lumb, J.-P. Phenol-Directed C–H Functionalization. *ACS Catal.* **2019**, *9*, 521–555.
- (90) Kennedy, O.; Zhorenes, R. *Coumarins: Biology, Applications and Mode of Action*; John Wiley and Sons: Chichester, 1997.
- (91) Ferguson, J.; Zeng, F.; Alper, H. Synthesis of Coumarins via Pd-Catalyzed Oxidative Cyclocarbonylation of 2-Vinylphenols. *Org. Lett.* **2012**, *14*, 5602–5605.
- (92) Seoane, A.; Casanova, N.; Quiñones, N.; Mascareñas, J. L.; Gulías, M. Straightforward Assembly of Benzoxepines by Means of a Rhodium(III)-Catalyzed C–H Functionalization of *o*-Vinylphenols. *J. Am. Chem. Soc.* **2014**, *136*, 834–837.
- (93) Gulías, M.; Marcos-Atanes, D.; Mascareñas, J. L.; Font, M. Practical, Large-Scale Preparation of Benzoxepines and Coumarins through Rhodium(III)-Catalyzed C–H Activation/Annulation Reactions. *Org. Process Res. Dev.* **2019**, *23*, 1669–1673.
- (94) Liu, X.-G.; Zhang, S.-S.; Jiang, C.-Y.; Wu, J.-Q.; Li, Q.; Wang, H. Cp\*Co(III)-Catalyzed Annulations of 2-Alkenylphenols with CO: Mild Access to Coumarin Derivatives. *Org. Lett.* **2015**, *17*, 5404–5407.
- (95) Manjolinho, F.; Arndt, M.; Gooßen, K.; Gooßen, L. J. Catalytic C–H Carboxylation of Terminal Alkynes with Carbon Dioxide. *ACS Catal.* **2012**, *2*, 2014–2021.
- (96) Fenner, S.; Ackermann, L. C–H Carboxylation of Heteroarenes with Ambient CO<sub>2</sub>. *Green Chem.* **2016**, *18*, 3804–3807.
- (97) Luo, J.; Larrosa, I. C–H Carboxylation of Aromatic Compounds through CO<sub>2</sub> Fixation. *ChemSusChem* **2017**, *10*, 3317–3332.
- (98) Hong, J.; Li, M.; Zhang, J.; Sun, B.; Mo, F. C–H Bond Carboxylation with Carbon Dioxide. *ChemSusChem* **2019**, *12*, 6–39.
- (99) Song, L.; Jiang, Y.-X.; Zhang, Z.; Gui, Y.-Y.; Zhou, X.-Y.; Yu, D.-G. CO<sub>2</sub> = CO + [O]: Recent Advance in Carbonylation of C–H Bonds with CO<sub>2</sub>. *Chem. Commun.* **2020**, *56*, 8355–8367.
- (100) Yang, Z.; Yu, Y.; Lai, L.; Zhou, L.; Ye, K.; Chen, F.-E. Carbon dioxide cycle via electrocatalysis: Electrochemical Carboxylation of CO<sub>2</sub> and Decarboxylative Functionalization of Carboxylic Acids. *Green Synth. Catal.* **2021**, *2*, 19–26.
- (101) Sasano, K.; Takaya, J.; Iwasawa, N. Pd(II)-Catalyzed Direct Carboxylation of Alkenyl C–H Bonds with CO<sub>2</sub>. *J. Am. Chem. Soc.* **2013**, *135*, 10954–10957.
- (102) Zhang, Z.; Ju, T.; Miao, M.; Han, J.-L.; Zhang, Y.-H.; Zhu, X.-Y.; Ye, J.-H.; Yu, D.-G.; Zhi, Y.-G. Transition-Metal-Free Lactonization of sp<sup>2</sup> C–H Bonds with CO<sub>2</sub>. *Org. Lett.* **2017**, *19*, 396–399.
- (103) Tan, L.-P.; Liang, D.; Cheng, Y.; Xiao, W.-J.; Chen, J.-R. Visible-Light-Induced Tandem Radical Addition/Cyclization of 2-Alkenylphenols and CBr<sub>4</sub> for the Synthesis of 4-Arylcoumarins. *Org. Chem. Front.* **2021**, *8*, 5052–5057.
- (104) Seoane, A.; Casanova, N.; Quiñones, N.; Mascareñas, J. L.; Gulías, M. Rhodium(III)-Catalyzed Dearomatizing (3 + 2) Annulation of 2-Alkenylphenols and Alkynes. *J. Am. Chem. Soc.* **2014**, *136*, 7607–7610.
- (105) Kujawa, S.; Best, D.; Burns, D. J.; Lam, H. W. Synthesis of Spirocyclic Enones by Rhodium-Catalyzed Dearomatizing Oxidative Annulation of 2-Alkenylphenols with Alkynes and Enynes. *Chem. - Eur. J.* **2014**, *20*, 8599–8602.
- (106) Liu, T.; Han, L.; Zhang, J.; Lu, G. Multiple Reaction Pathways of Eight-Membered Rhodacycles in Rh-Catalyzed Annulations of 2-Alkenyl Phenols/Anilides with Alkynes. *J. Org.*

*Chem.* **2021**, *86*, 10484–10491.

- (107) Duarah, G.; Kaishap, P. P.; Sarma, B.; Gogoi, S. Ruthenium(II)-Catalyzed Dearomatized C–H Activation and Annulation Reaction of Vinyl naphthols with Alkynes: Access to Spiro-Pentacyclic Naphthalenones. *Chem. - Eur. J.* **2018**, *24*, 10196–10200.
- (108) Wang, Y.; Oliveira, J. C. A.; Lin, Z.; Ackermann, L. Electrooxidative Rhodium-Catalyzed [5 + 2] Annulations via C–H/O–H Activations. *Angew. Chem., Int. Ed.* **2021**, *60*, 6419–6424.
- (109) DeKorver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. Ynamides: A Modern Functional Group for the New Millennium. *Chem. Rev.* **2010**, *110*, 5064–5106.
- (110) Evano, G.; Coste, A.; Jouvin, K. Ynamides: Versatile Tools in Organic Synthesis. *Angew. Chem., Int. Ed.* **2010**, *49*, 2840–2859.
- (111) Hu, L.; Zhao, J. Ynamide: A New Coupling Reagent for Amide and Peptide Synthesis. *Synlett*, **2017**, *28*, 1663–1670.
- (112) Han, X.-L.; Liu, X.-G.; Lin, E.; Chen, Y.; Chen, Z.; Wang, H.; Li, Q. Cp\*Co(III)-Catalyzed Oxidative [5 + 2] Annulation: Regioselective Synthesis of 2-Aminobenzoxepines via C–H/O–H Functionalization of 2-Vinylphenols with Ynamides. *Chem. Commun.* **2018**, *54*, 11562–11565.
- (113) Lin, P.-P.; Han, X.-L.; Ye, G.-H.; Li, J.-L.; Li, Q.; Wang, H. Cp\*Co(III)-Catalyzed Dearomative [3 + 2] Spiroannulation of 2-Alkenylphenols with Ynamides via C–H Activation. *J. Org. Chem.* **2019**, *84*, 12966–12974.
- (114) Casanova, N.; Seoane, A.; Mascareñas, J. L.; Gulías, M. Rhodium-Catalyzed (5+1) Annulations Between 2-Alkenylphenols and Allenes: A Practical Entry to 2,2-Disubstituted 2*H*-Chromenes. *Angew. Chem., Int. Ed.* **2015**, *54*, 2374–2377.
- (115) Kuppasamy, R.; Muralirajan, K.; Cheng, C.-H. Cobalt(III)-Catalyzed [5 + 1] Annulation for 2*H*-Chromenes Synthesis via Vinylic C–H Activation and Intramolecular Nucleophilic Addition. *ACS Catal.* **2016**, *6*, 3909–3913.
- (116) Kumar, A.; Ramaiah, K.; Prabhu, R. Rhodium(III)-Catalyzed [5+1] Annulation of 2-Alkenylphenols with Maleimides: Access to Highly Functionalized Spirocyclic Skeletons. *Chem. Commun.* **2021**, *57*, 8194–8197.
- (117) Casanova, N.; Del Rio, K. P.; García-Fandiño, R.; Mascareñas, J. L.; Gulías, M. Palladium(II)-Catalyzed Annulation Between *ortho*-Alkenylphenols and Allenes. Key Role of the Metal Geometry in Determining the Reaction Outcome. *ACS Catal.* **2016**, *6*, 3349–3353.
- (118) Yi, W.; Li, L.; Chen, H.; Ma, K.; Zhong, Y.; Chen, W.; Gao, H.; Zhou, Z. Rh(III)-Catalyzed Oxidative [5 + 2] Annulation Using Two Transient Assisting Groups: Stereospecific Assembly of 3-Alkenylated Benzoxepine Framework. *Org. Lett.* **2018**, *20*, 6812–6816.
- (119) Nagireddy, A.; Kotipalli, R.; Nanubolu, B. J.; Reddy, M. S. Rhodium-Catalyzed Coordination-Assisted Regioselective and Migratory Three-Point Double Annulation of *o*-Alkenyl Phenols with Tertiary Propargyl Alcohols. *Org. Lett.* **2022**, *24*, 5062–5067.
- (120) Guo, L.; Zhang, F.; Hu, W.; Li, L.; Jia, Y. Palladium-Catalyzed Synthesis of Benzofurans via C–H Activation/Oxidation Tandem Reaction and Its Application to the Synthesis of Decursivine and Serotobenine. *Chem. Commun.* **2014**, *50*, 3299–3302.
- (121) Ferguson, J.; Zeng, F.; Alwis, N.; Alper, H. Synthesis of 2(*1H*)-Quinolines via Pd-Catalyzed Oxidative Cyclocarbonylation of 2-Vinylanilines. *Org. Lett.* **2013**, *15*, 1998–2001.
- (122) Ma, J.; Chen, P.; Gong, X.; Hu, Y.; Ma, Q.; Wang, B.; Ma, Y. Metal-Free C–H [5 + 1] Carbonylation of 2-Alkenyl/Pyrrylanilines Using Dioxazolones as Carbonylating Reagents. *Org. Lett.* **2021**, *23*, 3761–3766.
- (123) Zhang, Z.; Liao, L.-L.; Yan, S.-S.; Wang, L.; He, Y.-Q.; Ye, J.-H.; Li, J.; Zhi, Y.-G.; Yu, D.-G. Lactamization of sp<sup>2</sup> C–H Bonds with CO<sub>2</sub>: Transition-Metal-Free and Redox-Neutral. *Angew. Chem., Int. Ed.* **2016**, *55*, 7068–7072.
- (124) Chen, P.; Nan, J.; Hu, Y.; Ma, Q.; Ma, Y. Ru<sup>II</sup>-Catalyzed/NH<sub>2</sub>-Assisted Selective Alkenyl C–H [5 + 1] Annulation of Alkenylanilines with Sulfoxonium Ylides to Quinolines. *Org. Lett.* **2019**, *21*, 4812–4815.
- (125) Liu, L.; Lin, J.; Pang, M.; Jin, H.; Yu, X.; Wang, S. Photo-Thermo-Mechanochemical Approach to Synthesize Quinolines via Addition/Cyclization of Sulfoxonium Ylides with 2-Vinylanilines Catalyzed by Iron(II) Phthalocyanine. *Org. Lett.* **2022**, *24*, 1146–1151.
- (126) Wang, L.; Ferguson, J.; Zeng, J. Palladium-Catalyzed Direct Coupling of 2-Vinylanilines and Isocyanides: An Efficient Synthesis of 2-Aminoquinolines. *Org. Biomol. Chem.* **2015**, *13*, 11486–11491.
- (127) Cheng, Y.; Judd, T. C.; Bartberger, M. D.; Brown, J.; Chen, K.; Fremereau, R. T.; Hickman, D., Jr.; Hitchcock, S. A.; Jordan, B.; Li, V.; Lopez, P.; Louie, S. W.; Luo, Y.; Michelsen, K.; Nixey, T.; Powers, T. S.; Rattan, C.; Sickmier, E. A.; Jean, D. J. S.; Wahl, R. C., Jr.; Wen, P. H.; Wood, S. From Fragment Screening to In Vivo Efficacy: Optimization of a Series of 2-Aminoquinolines as Potent Inhibitors of Beta-Site Amyloid Precursor Protein Cleaving Enzyme 1 (BACE1). *J. Med. Chem.* **2011**, *54*, 5836–5857.
- (128) Xu, P.; Zhu, T.-H.; Wei, T.-Q.; Wang, S.-Y.; Ji, S.-J. Co(acac)<sub>2</sub>/O<sub>2</sub>-Catalyzed Oxidative Isocyanide Insertion with 2-Vinylanilines: Efficient Synthesis of 2-Aminoquinolines. *RSC Adv.* **2016**, *6*, 32467–32470.
- (129) Xiang, Y.; Luo, P.; Hao, T.; Xiong, W.; Song, X.; Ding, Q. Copper-Mediated Formal [5 + 1] Annulation of 2-Vinylanilines and Glyoxylic Acid: A Facile Approach for the Synthesis of 4-Arylated Quinolines. *Tetrahedron*. **2021**, *79*, 131832–131838.
- (130) Cendón, B.; Casanova, N.; Comanescu, C.; García-Fandiño, R.; Seoane, A.; Gulías, M.; Mascareñas, J. L. Palladium-Catalyzed Formal (5 + 2) Annulation between *ortho*-Alkenylanilides and Allenes. *Org. Lett.* **2017**, *19*, 1674–1677.
- (131) Wu, L.; Meng, Y.; Ferguson, J.; Wang, L.; Zeng, F. Palladium-Catalyzed Oxidative Annulation of *ortho*-Alkenylanilines and Allenes: An Access to Benzo[*b*]azepines. *J. Org. Chem.* **2017**, *82*, 4121–4128.
- (132) Singh, A.; Shukla, R. K.; Volla, C. M. R. Rh(III)-Catalyzed [5 + 1] Annulation of 2-Alkenylanilides and 2-Alkenylphenols with Allenyl Acetates. *Chem. Sci.* **2022**, *13*, 2043–2049.
- (133) Font, M.; Cendón, B.; Seoane, A.; Mascareñas, J. L.; Gulías, M. Rhodium(III)-Catalyzed Annulation of 2-Alkenyl Anilides with Alkynes through C–H Activation: Direct Access to 2-Substituted Indolines. *Angew. Chem., Int. Ed.* **2018**, *57*, 8255–8259.
- (134) Seoane, A.; Comanescu, C.; Casanova, N.; García-Fandiño, R.; Diz, X.; Mascareñas, J. L.; Gulías, M. Rhodium-Catalyzed Annulation of *ortho*-Alkenyl Anilides with Alkynes: Formation of Unexpected Naphthalene Adducts. *Angew. Chem., Int. Ed.* **2019**, *58*, 1700–1704.
- (135) Li, D.; Zeng, F. Palladium-Catalyzed Domino Alkenylation/Amination/Pyridination Reactions of 2-Vinylanilines with Alkynes: Access to Cyclopentaquinolines. *Org. Lett.* **2017**, *19*, 6498–6501.
- (136) Qiao, H.; Zhang, S.; Li, K.; Cao, Z.; Zeng, F. Palladium(II)/Lewis Acid Cocatalyzed Oxidative Annulation of 2-Alkenylanilines and Propargylic Esters: An Access to Benzo[*b*]azepines. *J. Org. Chem.* **2019**, *84*, 10843–10851.
- (137) Chelucci, G. Synthesis and Metal-Catalyzed Reactions of *gem*-Dihalovinyl Systems. *Chem. Rev.* **2012**, *112*, 1344–1462.
- (138) Lin, J.; Wu, C.; Tian, X. Nickel-Catalyzed Cascade Reaction of 2-Vinylanilines with *gem*-Dichloroalkenes. *Org. Lett.* **2022**, *24*, 4855–4859.
- (139) Yu, R.; Li, D.; Zeng, F. Palladium-Catalyzed Sequential Vinylic C–H Arylation/Amination of 2-Vinylanilines with Aryl Boronic Acids: Access to 2-Arylindoles. *J. Org. Chem.* **2018**, *83*, 323–329.

- (140) Li, L.; Gao, H.; Sun, M.; Zhou, Z.; Yi, W. Experimental and Computational Studies on Cp<sup>\*C</sup>Rh(III)/KOPiv-Catalyzed Intramolecular Dehydrogenative Cross-Couplings for Building Eight-Membered Sultam/Lactam Frameworks. *Org. Lett.* **2020**, *22*, 5473–5478.
- (141) Krolski, M. E.; Renaldo, A. F.; Rudisill, D. E.; Stille, J. K. Palladium-Catalyzed Coupling of 2-Bromoanilines with Vinylstannanes. A Regiocontrolled Synthesis of Substituted Indoles. *J. Org. Chem.* **1988**, *53*, 1170–1176.
- (142) Wang, M.; Li, Y.; Wu, Q.-A.; Luo, S.; Li, Y. Iron-Promoted Construction of Indoles via Intramolecular Oxidative C–N Coupling of 2-Alkenylanilines Using Persulfate. *Synthesis* **2019**, *51*, 3085–3090.
- (143) Yang, R.; Yu, J.-T.; Sun, S.; Zheng, Q.; Cheng, J. Copper-Mediated Intramolecular aza-Wacker-Type Cyclization of 2-Alkenylanilines toward 3-Aryl indoles. *Tetrahedron Lett.* **2017**, *58*, 445–448.
- (144) Ma, A.-L.; Li, Y.-L.; Deng, J. A Concise Approach to Indoles via Oxidative C–H Amination of 2-Alkenylanilines Using Dioxxygen as the Sole Oxidant. *RSC Adv.* **2016**, *6*, 35764–35770.
- (145) Jang, Y. H.; Youn, S. W. Metal-Free C–H Amination for Indole Synthesis. *Org. Lett.* **2014**, *16*, 3720–3723.
- (146) Zhang, X.; Guo, R.; Zhao, X. Organoselenium-Catalyzed Synthesis of Indoles through Intramolecular C–H Amination. *Org. Chem. Front.* **2015**, *2*, 1334–1337.
- (147) Ortgies, S.; Breder, A. Selenium-Catalyzed Oxidative C(sp<sup>2</sup>)-H Amination of Alkenes Exemplified in the Expedient Synthesis of (Aza-)Indoles. *Org. Lett.* **2015**, *17*, 2748–2751.
- (148) Fra, L.; Millán, A.; Souto, J. A.; Muñiz, K. Indole Synthesis Based On A Modified Koser Reagent. *Angew. Chem., Int. Ed.* **2014**, *53*, 7349–7353.
- (149) Zhao, C.-Y.; Li, K.; Pang, Y.; Li, J.-Q.; Liang, C.; Su, G.-F.; Mo, D.-L. Iodine(III) Reagent-Mediated Intramolecular Amination of 2-Alkenylanilines to Prepare Indoles. *Adv. Synth. Catal.* **2018**, *360*, 1919–1925.
- (150) Andries-Ulmer, A.; Brunner, C.; Rehbein, J.; Gulder, T. Fluorine as a Traceless Directing Group for the Regiodivergent Synthesis of Indoles and Tryptophans. *J. Am. Chem. Soc.* **2018**, *140*, 13034–13041.
- (151) Li, Y.-L.; Li, J.; Huang, Y.-N.; Deng, J. Metal-Free Synthesis of Indole via NIS-Mediated Cascade C–N Bond Formation/Aromatization. *J. Org. Chem.* **2015**, *80*, 3841–3851.
- (152) Maity S.; Zheng, N. A Visible-Light-Mediated Oxidative C–N Bond Formation/Aromatization Cascade: Photocatalytic Preparation of N-Arylindoles. *Angew. Chem., Int. Ed.* **2012**, *51*, 9562–9566.
- (153) Hu, K.; Zhang, Y.; Zhou, Z.; Yang, Y.; Zha, Z.; Wang, Z. Iodine-Mediated Electrochemical C(sp<sup>2</sup>)-H Amination: Switchable Synthesis of Indolines and Indoles. *Org. Lett.* **2020**, *22*, 5773–5777.
- (154) Zheng, Y.-T.; Song, J.; Xu, H.-C. Electrocatalytic Dehydrogenative Cyclization of 2-Vinylanilides for the Synthesis of Indoles. *J. Org. Chem.* **2021**, *86*, 16001–16007.
- (155) Kim, Y.; Kim, D.; Chang, S. Ir(III)-Catalyzed Electrooxidative Intramolecular Dehydrogenative C–H/N–H Coupling for the Synthesis of N–H Indoles. *Chem. Commun.* **2021**, *57*, 12309–12312.
- (156) Liang, Q.-J.; Yang, C.; Meng, F.-F.; Jiang, B.; Xu, Y.-H.; Loh, T.-P. Chelation versus Non-Chelation Control in the Stereoselective Alkenyl sp<sup>2</sup> C–H Bond Functionalization Reaction. *Angew. Chem., Int. Ed.* **2017**, *56*, 5091–5095.
- (157) Meng, K.; Li, T.; Yu, C.; Shen, C.; Zhang, J.; Zhong, G. Geminal Group-Directed Olefinic C–H Functionalization via Four- to Eight-Membered exo-Metalloacycles. *Nat. Commun.* **2019**, *10*, 5109.
- (158) Wu, Z.; Fatuzzo, N.; Dong, G. Distal Alkenyl C–H Functionalization via the Palladium/Norbornene Cooperative Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 2715–2720.
- (159) Wang, J.; Dong, G. Palladium/Norbornene Cooperative Catalysis. *Chem. Rev.* **2019**, *19*, 7478–7528.
- (160) Wu, Z.; Fatuzzo, N.; Dong, G. Palladium/Norbornene-Catalyzed Distal Alkenyl C–H Arylation and Alkylation of cis-Olefins. *Tetrahedron* **2021**, *90*, 132173–132192.
- (161) Xu, S.; Hirano, K.; Miura, M. Pd-Catalyzed Regioselective C–H Alkenylation and Alkynylation of Allylic Alcohols with the Assistance of a Bidentate Phenanthroline Auxiliary. *Org. Lett.* **2020**, *22*, 9059–9064.
- (162) Zhu, Y.; Chen, F.; Cheng, D.; Chen, Y.; Zhao, X.; Wei, W.; Lu, Y.; Zhao, J. Rhodium(III)-Catalyzed Alkenyl C–H Functionalization to Dienes and Allenes. *Org. Lett.* **2020**, *22*, 8786–8790.
- (163) Dethe, D. H.; Nagabhushana, C. B. Ruthenium-Catalyzed Direct Dehydrogenative Cross-Coupling of Allyl Alcohols and Acrylates: Application to Total Synthesis of Hydroxy β-Sanshool, ZP-Amide I, and Chondrillin. *Org. Lett.* **2020**, *22*, 1618–1623.
- (164) Hu, S.; Wang, D.; Liu, J.; Li, X. Rhodium(III)-Catalyzed Oxidative Olefination of N-Allyl Sulfonamides. *Org. Biomol. Chem.* **2013**, *11*, 2761–2765.
- (165) Parella, R.; Babu, S. A. Pd(II)-Catalyzed, Picolinamide-Assisted, Z-Selective γ-Arylation of Allylamines To Construct Z-Cinnamylamines. *J. Org. Chem.* **2017**, *82*, 6550–6567.
- (166) Luo, Y.-C.; Yang, C.; Qiu, S.-Q.; Liang, Q.-J.; Xu, Y.-H.; Loh, T.-P. Palladium(II)-Catalyzed Stereospecific Alkenyl C–H Bond Alkylation of Allylamines with Alkyl Iodides. *ACS Catal.* **2019**, *9*, 4271–4276.
- (167) Yang, Y.; Yu, B. Recent Advances in the Chemical Synthesis of C-Glycosides. *Chem. Rev.* **2017**, *117*, 12281–12356.
- (168) Kitamura, K.; Ando, Y.; Matsumoto, T.; Suzuki, K. Total Synthesis of Aryl C-Glycoside Natural Products: Strategies and Tactics. *Chem. Rev.* **2018**, *118*, 1495–1598.
- (169) Liao, H.; Ma, J.; Yao, H.; Liu, X.-W. Recent Progress of C-Glycosylation Methods in the Total Synthesis of Natural Products and Pharmaceuticals. *Org. Biomol. Chem.* **2018**, *16*, 1791–1806.
- (170) Leng, W.-L.; Liao, H.; Yao, H.; Ang, Z.-E.; Xiang, S.; Liu, X.-W. Palladium-Catalyzed Decarboxylative Allylation/Wittig Reaction: Substrate-Controlled Synthesis of C-Vinyl Glycosides. *Org. Lett.* **2017**, *19*, 416–419.
- (171) Huang, N.; Liao, H.; Yao, H.; Xie, T.; Zhang, S.; Zou, K.; Liu, X.-W. Diastereoselective Synthesis of C-Vinyl Glycosides via Gold(I)-Catalyzed Tandem 1,3-Acyloxy Migration/Ferrier Rearrangement. *Org. Lett.* **2018**, *20*, 16–19.
- (172) Liu, J.; Gong, H. Stereoselective Preparation of α-C-Vinyl/Aryl Glycosides via Nickel-Catalyzed Reductive Coupling of Glycosyl Halides with Vinyl and Aryl Halides. *Org. Lett.* **2018**, *20*, 7991–7995.
- (173) Li, M.; Qiu, Y.-F.; Wang, C.-T.; Li, X.-S.; Wei, W.-X.; Zhao, Y.-Z.; Bao, Q.-F.; Ding, Y.-N.; Shi, W.-Y.; Liang, Y.-M. Visible-Light-Induced Pd-Catalyzed Radical Strategy for Constructing C-Vinyl Glycosides. *Org. Lett.* **2020**, *22*, 6288–6293.
- (174) Sun, Q.; Zhang, H.; Wang, Q.; Qiao, T.; He, G.; Chen, G. Stereoselective Synthesis of C-Vinyl Glycosides via Palladium-Catalyzed C–H Glycosylation of Alkenes. *Angew. Chem., Int. Ed.* **2021**, *60*, 19620–19625.
- (175) Landge, V. G.; Maxwell, J. M.; Chand-Thakuri, P.; Kapoor, M.; Diemler, E. T.; Young, M. C. Palladium-Catalyzed Regioselective Arylation of Unprotected Allylamines. *JACS Au* **2021**, *1*, 13–22.
- (176) Landge, V. G.; Grant, A. J.; Fu, Y.; Rabon, A. M.; Payton, J. L.; Young, M. C. Palladium-Catalyzed γ,γ'-Diarylation of Free Alkenyl Amines. *J. Am. Chem. Soc.* **2021**, *143*, 10352–10360.
- (177) González, J. M.; Cendón, B.; Mascareñas, J. L.; Gulías, M. Kinetic Resolution of Allyltriflamides through a Pd-Catalyzed C–H Functionalization with Allenes: Asymmetric Assembly of Tetrahydropyridines. *J. Am. Chem. Soc.* **2021**, *143*, 3747–3752.

- (178) Wang, C.; Zhang, L.; Chen, C.; Han, J.; Yao, Y.; Zhao, Y. Oxalyl Amide Assisted Palladium-Catalyzed Synthesis of Pyrrolidones via Carbonylation of  $\gamma$ -C(sp<sup>3</sup>)-H Bonds of Aliphatic Amine Substrates. *Chem. Sci.* **2015**, *6*, 4610–4614.
- (179) Chen, C.; Guan, M.; Zhang, J.; Wen, Z.; Zhao, Y. Palladium-Catalyzed Oxalyl Amide Directed Silylation and Germanylation of Amine Derivatives. *Org. Lett.* **2015**, *17*, 3646–3649.
- (180) Guan, M.; Chen, C.; Zhang, J.; Zeng, S.; Zhao, Y. Palladium-Catalyzed Oxalyl Amide Assisted Direct *ortho*-Alkynylation of Arylalkylamines Derivatives at  $\delta$  and  $\epsilon$  Positions. *Chem. Commun.* **2015**, *51*, 12103–12106.
- (181) Viart, H. M.-F.; Bachmann, A.; Kayitare, W.; Sarpong, R.  $\beta$ -Carboline Amides as Intrinsic Directing Groups for C(sp<sup>2</sup>)-H Functionalization. *J. Am. Chem. Soc.* **2017**, *139*, 1325–1329.
- (182) Zang, Z.-L.; Zhao, S.; Karnakanti, S.; Liu, C.-L.; Shao, P.-L.; He, Y. Catalytic Multisite-Selective Acetoxylation Reactions at sp<sup>2</sup> vs sp<sup>3</sup> C-H Bonds in Cyclic Olefins. *Org. Lett.* **2016**, *18*, 5014–5017.
- (183) Miao, C.-L.; Zhao, S.; Zang, Z.-L.; Xiao, L.; Zhou, C.-H.; He, Y.; Cai, G.-X. Pd-Catalyzed Remote Site-Selective and Stereoselective C(Alkenyl)-H Alkenylation of Unactivated Cycloalkenes. *J. Org. Chem.* **2020**, *85*, 774–787.
- (184) Jiang, B.; Meng, F.-F.; Liang, Q.-J.; Xu, Y.-H.; Loh, T.-P. Palladium-Catalyzed Direct Intramolecular C-N Bond Formation: Access to Multisubstituted Dihydropyrroles. *Org. Lett.* **2017**, *19*, 914–917.
- (185) Guo, L.-Y.; Li, Q.; Liu, Y.-T.; Li, L.; Ni, Y.-Q.; Li, Y.; Pan, F. Palladium-Catalyzed Alkynylation of Alkenes via C-H Activation for the Preparation of Conjugated 1,3-Enynes. *Adv. Synth. Catal.* **2022**, *364*, 1109–1116.
- (186) Sharma, R.; Thakur, K.; Kumar, R.; Kumar, I.; Sharma, U. Distant C-H Activation/Functionalization: A New Horizon of Selectivity Beyond Proximity. *Catal. Rev.* **2015**, *57*, 345–405.
- (187) Liu, M.; Yang, P.; Karunananda, M. K.; Wang, Y.; Liu, P.; Engle, K. M. C(alkenyl)-H Activation via Six-Membered Palladacycles: Catalytic 1,3-Diene Synthesis. *J. Am. Chem. Soc.* **2018**, *140*, 5805–5813.
- (188) Shen, C.; Lu, X.; Zhang, J.; Ding, L.; Sun, Y.; Zhong, G. Bidentate auxiliary-directed alkenyl C-H allylation via exo-palladacycles: synthesis of branched 1,4-dienes. *Chem. Commun.* **2019**, *55*, 13582–13585.
- (189) Kotora, M.; Matsumura, H.; Gao, G.; Takahashi, T. Palladium-Catalyzed Coupling of Two Alkynes and an Alkenyl Iodide: Formation of Pentasubstituted Fulvenes. *Org. Lett.* **2001**, *3*, 3467–3470.
- (190) Mousseau, J. J.; Fortier, A.; Charette, A. B. Synthesis of 2-Substituted Pyrazolo[1,5-*a*]pyridines through Cascade Direct Alkenylation/Cyclization Reactions. *Org. Lett.* **2010**, *12*, 516–519.
- (191) Lu, X.-L.; Shannon, M.; Peng, X.-S.; Wong, H. V. C. Stereospecific Iron-Catalyzed Carbon(sp<sup>2</sup>)-Carbon(sp<sup>3</sup>) Cross-Coupling with Alkylolithium and Alkenyl Iodides. *Org. Lett.* **2019**, *21*, 2546–2549.
- (192) Desaintjean, A.; Belrhomari, S.; Rousseau, L.; Lefèvre, G. Iron-Catalyzed Cross-Coupling of Functionalized Benzylmanganese Halides with Alkenyl Iodides, Bromides, and Triflates. *Org. Lett.* **2019**, *21*, 8684–8688.
- (193) Schreiber, B. S.; Carreira, E. M. Palladium-Catalyzed Regioselective C-H Iodination of Unactivated Alkenes. *J. Am. Chem. Soc.* **2019**, *141*, 8758–8763.
- (194) Schreiber, B. S.; Fadel, M.; Carreira, E. M. Palladium-Catalyzed C-H Alkynylation of Unactivated Alkenes. *Angew. Chem., Int. Ed.* **2020**, *59*, 7818–7822.
- (195) Tang, C.; Zhang, R.; Zhu, B.; Fu, J.; Deng, Y.; Tian, L.; Guan, W. Bi, X. Directed Copper-Catalyzed Intermolecular Heck-Type Reaction of Unactivated Olefins and Alkyl Halides. *J. Am. Chem. Soc.* **2018**, *140*, 16929–16935.
- (196) White, C. J.; Yudin, A. K. Contemporary Strategies for Peptide Macrocyclization. *Nat. Chem.* **2011**, *3*, 509–524.
- (197) Han, B.; Li, B.; Qi, L.; Yang, P.; He, G.; Chen, G. Construction of Cyclophane-Braced Peptide Macrocycles via Palladium-Catalyzed Picolinamide-Directed Intramolecular C(sp<sup>2</sup>)-H Arylation. *Org. Lett.* **2020**, *22*, 6879–6883.
- (198) Giri, R.; Yu, J.-Q. Synthesis of 1,2- and 1,3-Dicarboxylic Acids via Pd(II)-Catalyzed Carboxylation of Aryl and Vinyl C-H Bonds. *J. Am. Chem. Soc.* **2008**, *130*, 14082–14083.
- (199) Mochida, S.; Hirano, K.; Satoh, T.; Miura, M. Synthesis of Functionalized  $\alpha$ -Pyrone and Butenolide Derivatives by Rhodium-Catalyzed Oxidative Coupling of Substituted Acrylic Acids with Alkynes and Alkenes. *J. Org. Chem.* **2009**, *74*, 6295–6298.
- (200) Itoh, M.; Shimizu, M.; Hirano, K.; Satoh, T.; Miura, M. Rhodium-Catalyzed Decarboxylative and Dehydrogenative Coupling of Maleic Acids with Alkynes and Alkenes. *J. Org. Chem.* **2013**, *78*, 11427–11432.
- (201) Yu, Y.; Huang, L.; Wu, W.; Jiang, H. Palladium-Catalyzed Oxidative Annulation of Acrylic Acid and Amide with Alkynes: A Practical Route to Synthesize  $\alpha$ -Pyrone and Pyridones. *Org. Lett.* **2014**, *16*, 2146–2149.
- (202) Kudo, E.; Shibata, Y.; Yamazaki, M.; Masutomi, K.; Miyauchi, Y.; Fukui, M.; Sugiyama, H.; Uekusa, H.; Satoh, T.; Miura, M.; Tanaka, K. Oxidative Annulation of Arenecarboxylic and Acrylic Acids with Alkynes under Ambient Conditions Catalyzed by an Electron-Deficient Rhodium-(III) Complex. *Chem. - Eur. J.* **2016**, *22*, 14190–14194.
- (203) Li, Y.-T.; Zhu, Y.; Tu, G.-L.; Zhang, J.-Y.; Zhao, Y.-S. Rhodium(III)-Catalyzed Oxidative Annulation of Acrylic Acid with Alkynes: An Easy Approach to the Synthesis of  $\alpha$ -Pyrone. *Chem. - Asian J.* **2018**, *13*, 3281–3284.
- (204) Prakash, R.; Shekarrao, K.; Gogoi, S. Ruthenium(II)-Catalyzed Alkene C-H Bond Functionalization on Cinnamic Acids: A Facile Synthesis of Versatile  $\alpha$ -Pyrone. *Org. Lett.* **2015**, *17*, 5264–5267.
- (205) Yang, Q.-L.; Xing, Y.-K.; Wang, X.-Y.; Ma, H.-X.; Weng, X.-J.; Yang, X.; Guo, H.-M.; Mei, T.-S. Electrochemistry-Enabled Ir-Catalyzed Vinylic C-H Functionalization. *J. Am. Chem. Soc.* **2019**, *141*, 18970–18976.
- (206) Li, H.; Jiang, Q.; Jie, X.; Shang, Y.; Zhang, Y.; Goossen, L. J.; Su, W. Rh/Cu-Catalyzed Ketone  $\beta$ -Functionalization by Merging Ketone Dehydrogenation and Carboxyl-Directed C-H Alkylation. *ACS Catal.* **2018**, *8*, 4777–4782.
- (207) Zhu, Y.-Q.; Liu, Y.; Wang, H. N.; Liu, W. B.; Li, C.-J. Reaction of Alkenecarboxylic Acids with Isocyanates via Rhodium(III)-Catalyzed C-H Activation: A Versatile Route to Cyclic Imides. *Org. Chem. Front.* **2016**, *3*, 971–974.
- (208) Yu, C.; Zhang, J.; Zhong, G. One Step Synthesis of  $\gamma$ -Alkylidenebutenolides from Simple Vinyl Carboxylic Acids and Alkenes. *Chem. Commun.* **2017**, *53*, 9902–9905.
- (209) Zhu, Y.-Q.; Han, T.-F.; He, J.-L.; Li, M.; Li, J.-X.; Zhu, K. Route to Substituted Furan-2(5*H*)-ones from cyclo-Alkenecarboxylic Acids and Acrylates via C-H Activation. *J. Org. Chem.* **2017**, *82*, 8598–8603.
- (210) Qiu, Y.; Kong, W.-J.; Struwe, J.; Sauermann, N.; Rogge, T.; Scheremetjev, A.; Ackermann, L. Electrooxidative Rhodium-Catalyzed C-H/C-H Activation: Electricity as Oxidant for Cross-Dehydrogenative Alkenylation. *Angew. Chem., Int. Ed.* **2018**, *57*, 5828–5832.
- (211) Sun, Y.; Zhang, G. Palladium-Catalyzed Formal [4+2] Cycloaddition of Benzoic and Acrylic Acids with 1,3-Dienes via C-H Bond Activation: Efficient Access to 3,4-Dihydroisocoumarin and 5,6-Dihydrocoumalins. *Chin. J. Chem.* **2018**, *36*, 708–711.
- (212) Jeganmohan, M.; Ramesh, B. Ru(II)- or Rh(III)-Catalyzed Annulation of Aromatic/Vinylic Acids with

Alkylidenecyclopropanes via C–H Activation. *J. Org. Chem.* **2022**, *87*, 5668–5681.

(213) Deb, A.; Maiti, D. Emergence of Unactivated Olefins for the Synthesis of Olefinated Arenes. *Eur. J. Org. Chem.* **2017**, *2017*, 1239–1252.

(214) Jambu, S.; Jeganmohan, M. Rhodium(III)-Catalyzed C–H Olefination of Aromatic/Vinyl Acids with Unactivated Olefins at Room Temperature. *Org. Lett.* **2020**, *22*, 5057–5062.

(215) Ma, W.; Kaplaneris, N.; Fang, X.; Gu, L.; Mei, R.; Ackermann, L. Chelation-Assisted Transition Metal-Catalyzed C–H Chalcogenylations. *Org. Chem. Front.* **2020**, *7*, 1022–1060.

(216) Liu, C.; Fang, Y.; Wang, S.-Y.; Ji, S.-J. RhCl<sub>3</sub>·3H<sub>2</sub>O-Catalyzed Ligand-Enabled Highly Regioselective Thiolation of Acrylic Acids. *ACS Catal.* **2019**, *9*, 8910–8915.

(217) Tsai, H.-C.; Huang, Y.-H.; Chou, C.-M. Rapid Access to *Ortho*-Alkylated Vinylarenes from Aromatic Acids by Dearomatization and Tandem Decarboxylative C–H Olefination/Rearomatization. *Org. Lett.* **2018**, *20*, 1328–1332.

(218) Wang, Y.-C.; Huang, Y.-H.; Tsai, H.-C.; Basha, R. S.; Chou, C.-M. Palladium-Catalyzed Proaromatic C(Alkenyl)–H Olefination: Synthesis of Densely Functionalized 1,3-Dienes. *Org. Lett.* **2020**, *22*, 6765–6770.

(219) Belitz, F.; Seitz, A.-K.; Goebel, J. F.; Hu, Z.; Gooßen, L. J. Ru-Catalyzed C–H Arylation of Acrylic Acids with Aryl Bromides. *Org. Lett.* **2022**, *24*, 3466–3470.

(220) Gao, Y.; Nie, J.; Li, Y.; Li, X.; Chen, Q.; Huo, Y.; Hu, X.-Q. Rh-Catalyzed C–H Amination/Annulation of Acrylic Acids and Anthranils by Using –COOH as a Deciduous Directing Group: An Access to Diverse Quinolines. *Org. Lett.* **2020**, *22*, 2600–2605.

(221) Jiang, Y.; Li, P.; Wang, J.; Zhao, J.; Li, Y.; Zhang, Y.; Chang, J.; Liu, B.; Li, X. Rh(III)-Catalyzed Coupling of Acrylic Acids and Ynenones via Olefinic C–H Activation and Michael Addition. *Org. Lett.* **2020**, *22*, 438–442.

(222) Yu, S.; Hong, C.; Liu, Z.; Zhang, Y. Synthesis of Cyclopentenones through Rhodium-Catalyzed C–H Annulation of Acrylic Acids with Formaldehyde and Malonates. *Org. Lett.* **2021**, *23*, 5054–5059.

(223) Yu, S.; Hong, C.; Liu, Z.; Zhang, Y. Cobalt-Catalyzed Vinylic C–H Addition to Formaldehyde: Synthesis of Butenolides from Acrylic Acids and HCHO. *Org. Lett.* **2021**, *23*, 8359–8365.

(224) Hong, C.; Yu, S.; Liu, Z.; Zhang, Y. Rh-Catalyzed Coupling of Acrylic/Benzoic Acids with  $\alpha$ -Diazocarbonyl Compounds: An Alternative Route for  $\alpha$ -Pyrones and Isocoumarins. *Org. Lett.* **2022**, *24*, 815–820.

(225) Yu, S.; Hong, C.; Liu, Z.; Zhang, Y. Synthesis of Pyranones: Ru-Catalyzed Cascade Reaction via Vinylic C–H Addition to Glyoxylate. *Org. Lett.* **2022**, *24*, 4871–4875.

(226) Jiang, Y.; Li, P.; Zhao, J.; Liu, B.; Li, X. Iodonium Ylides as Carbene Precursors in Rh(III)-Catalyzed C–H Activation. *Org. Lett.* **2020**, *22*, 7475–7479.

(227) Fernández, D. F.; Casanova, N.; Mascareñas, J. L.; Gullías, M. Rhodium(III)-Catalyzed Intramolecular Annulations of Acrylic and Benzoic Acids to Alkynes. *ACS Omega.* **2019**, *4*, 6257–6263.

(228) Gandeepan, P.; Rajamalli, P.; Cheng, C.-H. Rhodium(III)-Catalyzed [4+1] Annulation of Aromatic and Vinylic Carboxylic Acids with Allenes: An Efficient Method Towards Vinyl-Substituted Phthalides and 2-Furanones. *Chem. - Eur. J.* **2015**, *21*, 9198–9203.

(229) Ye, B.; Cramer, N. Chiral Cyclopentadienyls: Enabling Ligands for Asymmetric Rh(III)-Catalyzed C–H Functionalizations. *Acc. Chem. Res.* **2015**, *48*, 1308–1318.

(230) Wang, S.-G.; Cramer, N. Asymmetric Cp<sup>\*</sup>Rh(III)-Catalyzed Acrylic Acid C–H Functionalization with Allenes Provides Chiral  $\gamma$ -Lactones. *ACS Catal.* **2020**, *10*, 8231–8236.

(231) Trost, B. M.; Imi, K.; Davies, I. W. Elaboration of Conjugated Alkenes Initiated by Insertion into a Vinylic C–H Bond. *J. Am. Chem. Soc.* **1995**, *117*, 5371–5372.

(232) Besset, T.; Kuhl, N.; Patureau, F.W.; Glorius, F. Rh<sup>III</sup>-Catalyzed Oxidative Olefination of Vinylic C–H Bonds: Efficient and Selective Access to Di-unsaturated  $\alpha$ -Amino Acid Derivatives and Other Linear 1,3-Butadienes. *Chem. - Eur. J.* **2011**, *17*, 7167–7171.

(233) Sasaki, I.; Doi, H.; Hashimoto, T.; Kikuchi, T.; Ito, H.; Ishiyama, T. Iridium(I)-Catalyzed Vinylic C–H Borylation of 1-Cycloalkenecarboxylates with Bis(pinacolato)diboron. *Chem. Commun.* **2013**, *49*, 7546–7548.

(234) Hu, X.-H.; Zhang, J.; Yang, X.-F.; Xu, Y.-H.; Loh, T.-P. Stereo- and Chemoselective Cross-Coupling between Two Electron-Deficient Acrylates: An Efficient Route to (*Z,E*)-Muconate Derivatives. *J. Am. Chem. Soc.* **2015**, *137*, 3169–3172.

(235) Feng, R.; Yu, W.; Wang, K.; Liu, Z.; Zhang, Y. Ester-Directed Selective Olefination of Acrylates by Rhodium Catalysis. *Adv. Synth. Catal.* **2014**, *356*, 1501–1508.

(236) Tan, E.; Quinonero, O.; Elena de Orbe, M.; Echavarren, A. M. Broad-Scope Rh-Catalyzed Inverse-Sonogashira Reaction Directed by Weakly Coordinating Groups. *ACS Catal.* **2018**, *8*, 2166–2172.

(237) Das, R.; Kumar, G. S.; Kapur, M. Amides as Weak Coordinating Groups in Proximal C–H Bond Activation. *Eur. J. Org. Chem.* **2017**, *2017*, 5439–5459.

(238) Wencel-Delord, J.; Nimphius, C.; Patureau, F. W.; Glorius, F. Undirected Arene and Chelate-Assisted Olefin C–H Bond Activation: [Rh<sup>III</sup>Cp<sup>\*</sup>]-Catalyzed Dehydrogenative Alkene-Arene Coupling as a New Pathway for the Selective Synthesis of Highly Substituted *Z* Olefins. *Chem. - Asian J.* **2012**, *7*, 1208–1212.

(239) Parella, R.; Babu, S. A. Pd(OAc)<sub>2</sub>-Catalyzed, AgOAc-Promoted *Z* Selective Directed  $\beta$ -Arylation of Acrylamide Systems and Stereoselective Construction of *Z*-Cinnamamide Scaffolds. *J. Org. Chem.* **2015**, *80*, 12379–12396.

(240) Cheng, X.; Chen, Z.; Gao, Y.; Xue, F.; Jiang, C. Aminoquinoline-Assisted Vinylic C–H Arylation of Unsubstituted Acrylamide for the Selective Synthesis of *Z* Olefins. *Org. Biomol. Chem.* **2016**, *14*, 3298–3306.

(241) de Robichon, M.; Bordessa, A.; Malinowski, M.; Uziel, J.; Lubin-Germain, N.; Ferry, A. Access to C-Aryl/Alkenyl Glycosides by Directed Pd-Catalyzed C–H Functionalization of the Anomeric Position in Glycal-Type Substrate. *Chem. Commun.* **2019**, *55*, 11806–11808.

(242) Gu, Q.; Al Mamari, H. H.; Graczyk, K.; Diers, E.; Ackermann, L. Iron-Catalyzed C(sp<sup>2</sup>)–H and C(sp<sup>3</sup>)–H Arylation by Triazole Assistance. *Angew. Chem., Int. Ed.* **2014**, *53*, 3868–3871.

(243) Hu, L.; Gui, Q.; Chen, X.; Tan, Z.; Zhu, G. Cobalt-Promoted Selective Arylation of Benzamides and Acrylamides with Arylboronic Acids. *Org. Biomol. Chem.* **2016**, *14*, 11070–11075.

(244) Wang, H.-W.; Qiao, Y.-H.; Wu, J.-X.; Wang, Q.-P.; Tian, M.-X.; Li, Y.-F.; Yao, Q.-X.; Li, D.-C.; Dou, J.-M.; Lu, Y. Rh<sup>III</sup>-Catalyzed C–H (Het)arylation/Vinylation of *N*-2,6-Difluoroaryl Acrylamides. *Org. Lett.* **2021**, *23*, 656–662.

(245) Aihara, Y.; Chatani, N. Nickel-Catalyzed Direct Alkylation of C–H Bonds in Benzamides and Acrylamides with Functionalized Alkyl Halides via Bidentate-Chelation Assistance. *J. Am. Chem. Soc.* **2013**, *135*, 5308–5311.

(246) Ilies, L.; Matsubara, T.; Ichikawa, S.; Asako, S.; Nakamura, E. Iron-Catalyzed Directed Alkylation of Aromatic and Olefinic Carboxamides with Primary and Secondary Alkyl Tosylates, Mesylates, and Halides. *J. Am. Chem. Soc.* **2014**, *136*, 13126–13129.

(247) Monks, B. M.; Fruchey, E. R.; Cook, S. P. Iron-Catalyzed C(sp<sup>2</sup>)–H Alkylation of Carboxamides with Primary Electrophiles. *Angew. Chem., Int. Ed.* **2014**, *53*, 11065–11069.

(248) Ilies, L.; Ichikawa, S.; Asako, S.; Matsubara, T.; Nakamura, E. Iron-Catalyzed Directed Alkylation of Alkenes and Arenes with Alkylzinc Halides. *Adv. Synth. Catal.* **2015**, *357*, 2175–2179.

(249) Song, S.; Lu, P.; Liu, H.; Cai, S.-H.; Feng, C.; Loh, T.-P. Switchable C–H Functionalization of *N*-Tosyl Acrylamides with Acryloylsilanes. *Org. Lett.* **2017**, *19*, 2869–2872.

- (250) Sharma, S.; Han, S. H.; Oh, Y.; Mishra, N. K.; Lee, S. H.; Oh, J. S.; Kim, I. S. Cross-Coupling of Acrylamides and Maleimides under Rhodium Catalysis: Controlled Olefin Migration. *Org. Lett.* **2016**, *18*, 2568–2571.
- (251) Keerthana, M. S.; Manoharan, R.; Jegamohan, M. Cobalt(III)-Catalyzed Redox-Neutral Coupling of Acrylamides with Activated Alkenes via C–H Bond Activation. *Synthesis* **2020**, *52*, 1625–1633.
- (252) Potter, T. J.; Kamber, D. N.; Mercado, S. Q.; Ellman, J. A. Rh(III)-Catalyzed Aryl and Alkenyl C–H Bond Addition to Diverse Nitroalkenes. *ACS Catal.* **2017**, *7*, 150–153.
- (253) Shibata, T.; Kojima, M.; Onoda, S.; Ito, M. Enantioselective Cross-Coupling of Electron-Deficient Alkenes via Ir-Catalyzed Vinylic  $sp^2$  C–H Alkylation. *Org. Lett.* **2021**, *23*, 8158–8162.
- (254) Fernández, D. F.; Guliás, M.; Mascareñas, J. L.; López, F. Iridium(I)-Catalyzed Intramolecular Hydrocarbonation of Alkenes: Efficient Access to Cyclic Systems Bearing Quaternary Stereocenters. *Angew. Chem., Int. Ed.* **2017**, *56*, 9541–9545.
- (255) Zhang, J.; Loh, T.-P. Ruthenium- and Rhodium-Catalyzed Cross-Coupling Reaction of Acrylamides with Alkenes: Efficient Access to (*Z,E*)-Dienamides. *Chem. Commun.* **2012**, *48*, 11232–11234.
- (256) Shang, R.; Ilies, L.; Asako, S.; Nakamura, N.; Iron-Catalyzed  $C(sp^2)$ -H Bond Functionalization with Organoboron Compounds. *J. Am. Chem. Soc.* **2014**, *136*, 14349–14352.
- (257) Li, F.; Yu, C.; Zhang, J.; Zhong, G. Weinreb Amide Directed Cross-Coupling Reaction between Electron-Deficient Alkenes Catalyzed by Rhodium Catalyst. *Org. Biomol. Chem.* **2017**, *15*, 1236–1244.
- (258) Li, F.; Yu, C.; Zhang, J.; Zhong, G. Olefination of Electron-Deficient Alkenes with Allyl Acetate: Stereo- and Regioselective Access to (*2Z,4E*)-Dienamides. *Org. Lett.* **2016**, *18*, 4582–4585.
- (259) Yu, C.; Li, F.; Zhang, J.; Zhong, G. Direct Cross-Coupling Reaction of Electron-Deficient Alkenes Using an Oxidizing Directing Group. *Chem. Commun.* **2017**, *53*, 533–536.
- (260) Li, T.; Shen, C.; Sun, Y.; Zhang, J.; Xiang, P.; Lu, X.; Zhong, G. Cobalt-Catalyzed Olefinic C–H Alkenylation/Alkylation Switched by Carbonyl Groups. *Org. Lett.* **2019**, *21*, 7772–7777.
- (261) Meng, K.; Sun, Y.; Zhang, J.; Zhang, K.; Ji, X.; Ding, L.; Zhong, G. Iridium-Catalyzed Cross-Coupling Reactions of Alkenes by Hydrogen Transfer. *Org. Lett.* **2019**, *21*, 8219–8224.
- (262) Yoshimura, R.; Shibata, Y.; Tanaka, K. Aerobic Oxidative Cross-Coupling of Substituted Acrylamides with Alkenes Catalyzed by an Electron-Deficient  $CpRh^{III}$  Complex. *J. Org. Chem.* **2019**, *84*, 13164–13171.
- (263) Shibata, Y.; Otake, Y.; Hirano, M.; Tanaka, K. Amide-Directed Alkenylation of  $sp^2$  C–H Bonds Catalyzed by a Cationic Rh(I)/BIPHEP Complex Under Mild Conditions: Dramatic Rate Acceleration by a 1-Pyrrolidinecarbonyl Group. *Org. Lett.* **2009**, *11*, 689–692.
- (264) Meng, K.; Zhang, J.; Li, F.; Lin, Z.; Zhang, K.; Zhong, G. Amide Directed Cross-Coupling between Alkenes and Alkynes: A Regio- and Stereoselective Approach to Substituted (*2Z,4Z*)-Dienamides. *Org. Lett.* **2017**, *19*, 2498–2501.
- (265) Sun, Y.; Meng, K.; Zhang, J.; Jin, M.; Huang, N.; Zhong, G. Additive- and Ligand-Free Cross-Coupling Reactions between Alkenes and Alkynes by Iridium Catalysis. *Org. Lett.* **2019**, *21*, 4868–4872.
- (266) Fernández, D. F.; Rodrigues, C. A. B.; Calvelo, M.; Guliás, M.; Mascareñas, J. L.; López, F. Iridium(I)-Catalyzed Intramolecular Cycloisomerization of Enynes: Scope and Mechanistic Course. *ACS Catal.* **2018**, *8*, 7397–7402.
- (267) Li, J.-J.; Wang, C.-G.; Yu, J.-F.; Wang, P.; Yu, J.-Q. Cu-Catalyzed C–H Alkenylation of Benzoic Acid and Acrylic Acid Derivatives with Vinyl Boronates. *Org. Lett.* **2020**, *22*, 4692–4696.
- (268) Logeswaran, R.; Jegamohan, M. Rhodium(III)-Catalyzed Aerobic Oxidative C–H Olefination of Unsaturated Acrylamides with Unactivated Olefins. *Org. Lett.* **2021**, *23*, 767–771.
- (269) Logeswaran, R.; Jegamohan, M. Effect of Transition Metals on Chemodivergent Cross-Coupling of Acrylamides with Vinyl Acetate via C–H Activation. *Org. Lett.* **2021**, *23*, 5679–5683.
- (270) Wang, H.; Beiring, B.; Yu, D.-G.; Collins, K. D.; Glorius, F. [3]Dendralene Synthesis: Rhodium(III)-Catalyzed Alkenyl C–H Activation and Coupling Reaction with Allenyl Carbiniol Carbonate. *Angew. Chem., Int. Ed.* **2013**, *52*, 12430–12434.
- (271) Zhao, Q.; Tognetti, V.; Joubert, L.; Besset, T.; Pannecoucke, X.; Bouillon, J.-P.; Poisson, T. Palladium-Catalyzed Synthesis of 3-Trifluoromethyl-Substituted 1,3-Butadienes by Means of Directed C–H Bond Functionalization. *Org. Lett.* **2017**, *19*, 2106–2109.
- (272) Sengupta, S.; Mehta, G. Macrocyclization via C–H Functionalization: A New Paradigm in Macrocyclic Synthesis. *Org. Biomol. Chem.* **2020**, *18*, 1851–1876.
- (273) Jiang, B.; Zhao, M.; Li, S.-S.; Xu, Y.-H.; Loh, T.-P. Macrolide Synthesis through Intramolecular Oxidative Cross-Coupling of Alkenes. *Angew. Chem., Int. Ed.* **2018**, *57*, 555–559.
- (274) Maraswami, M.; Goh, J.; Loh, T.-P. Macrolactam Synthesis via Ring-Closing Alkene–Alkene Cross-Coupling Reactions. *Org. Lett.* **2020**, *22*, 9724–9728.
- (275) Misha, N. K.; Sharma, S.; Park, J.; Han, S.; Kim, I. S. Recent Advances in Catalytic  $C(sp^2)$ -H Allylation Reactions. *ACS Catal.* **2017**, *7*, 2821–2847.
- (276) Feng, C.; Feng, D.; Loh, T.-P. Rhodium(III)-Catalyzed C–H Allylation of Electron-Deficient Alkenes with Allyl Acetates. *Chem. Commun.* **2015**, *51*, 342–345.
- (277) Gensch, T.; Vásquez-Céspedes, S.; Yu, D.-G.; Glorius, F. Cobalt(III)-Catalyzed Directed C–H Allylation. *Org. Lett.* **2015**, *17*, 3714–3717.
- (278) Kim, M.; Sharma, S.; Mishra, N. K.; Han, S.; Park, J.; Kim, M.; Shin, Y.; Kwak, J. H.; Han, S. H.; Kim, I. S. Direct Allylation of Aromatic and  $\alpha,\beta$ -Unsaturated Carboxamides Under Ruthenium Catalysis. *Chem. Commun.* **2014**, *50*, 11303–11306.
- (279) Zhang, S.-S.; Wu, J.-Q.; Lao, Y.-X.; Liu, X.-G.; Liu, Y.; Lv, W.-X.; Tan, D.-H.; Zeng, Y.-F.; Wang, H. Mild Rhodium(III)-Catalyzed C–H Allylation with 4-Vinyl-1,3-dioxolan-2-ones: Direct and Stereoselective Synthesis of (*E*)-Allylic Alcohols. *Org. Lett.* **2014**, *16*, 6412–6415.
- (280) Lumbroso, A.; Cooke, M. L.; Breit, B. Catalytic Asymmetric Synthesis of Allylic Alcohols and Derivatives and their Applications in Organic Synthesis. *Angew. Chem., Int. Ed.* **2013**, *52*, 1890–1932.
- (281) Wu, J.-Q.; Qiu, Z.-P.; Zhang, S.-S.; Liu, J.-G.; Lao, Y.-X.; Gu, L.-Q.; Huang, Z.-S.; Li, J.; Wang, H. Rhodium(III)-Catalyzed C–H/C–C Activation Sequence: Vinylcyclopropanes as Versatile Synthons in Direct C–H Allylation Reactions. *Chem. Commun.* **2015**, *51*, 77–80.
- (282) Wu, X.; Ji, H. Ruthenium-Catalyzed C–H Allylation of Alkenes with Allyl Alcohols via C–H Bond Activation in Aqueous Solution. *J. Org. Chem.* **2018**, *83*, 12094–12102.
- (283) Cera, G.; Haven, T.; Ackermann, L. Expedient Iron-Catalyzed C–H Allylation/Alkylation by Triazole Assistance with Ample Scope. *Angew. Chem., Int. Ed.* **2016**, *55*, 1484–1488.
- (284) Xu, L.; Meng, K.; Zhang, J.; Sun, Y.; Lu, X.; Li, T.; Jiang, Y.; Zhong, G. Iridium-Catalyzed Alkenyl C–H Allylation Using Conjugated Dienes. *Chem. Commun.* **2019**, *55*, 9757–9760.
- (285) Huang, Y.; Xu, L.; Yu, F.; Shen, W.; Lu, X.; Ding, L.; Zhong, L.; Zhong, G.; Zhang, J. Stereoselective and Atom-Economic Alkenyl C–H Allylation/Alkenylation in Aqueous Media by Iridium Catalysis. *J. Org. Chem.* **2020**, *85*, 7225–7237.
- (286) Zhang, Y. J.; Skucas, E.; Krische, M. J. Direct Prenylation of Aromatic and  $\alpha,\beta$ -Unsaturated Carboxamides via Iridium-Catalyzed C–H Oxidative Addition–Allene Insertion. *Org. Lett.* **2009**, *11*, 4248–4250.
- (287) Wang, M.-M.; Wang, Z.-X.; Shang, M.; Dai, H.-X. Transition-Metal-Catalyzed C–H Alkynylation. *Chin. J. Org. Chem.* **2015**, *35*, 570–577.

- (288) Caspers, L. D.; Nachtsheim, B. J. Directing-Group-mediated C–H-Alkynylations. *Chem. - Asian J.* **2018**, *13*, 1231–1247.
- (289) Ano, Y.; Tobisu, M.; Chatani, N. Palladium-Catalyzed Direct *ortho*-Alkynylation of Aromatic Carboxylic Acid Derivatives. *Org. Lett.* **2012**, *14*, 354–357.
- (290) Yi, J.; Yang, L.; Xia, C.; Li, F. Nickel-Catalyzed Alkynylation of a C(sp<sup>2</sup>)–H Bond Directed by an 8-Aminoquinoline Moiety. *J. Org. Chem.* **2015**, *80*, 6213–6221.
- (291) Landge, V. G.; Shewale, C. H.; Jaiswal, G.; Sahoo, M. K.; Midya, S. P.; Balaraman, E. Nickel-Catalyzed Direct Alkynylation of C(sp<sup>2</sup>)–H Bonds of Amides: An “Inverse Sonogashira Strategy” to *ortho*-Alkynylbenzoic Acids. *Catal. Sci. Technol.* **2016**, *6*, 1946–1951.
- (292) Zhao, T.; Pu, X.; Han, W.; Gao, G. Nickel-Catalyzed 3,3-Dialkynylation of 2-Aryl Acrylamides: Direct Access to *gem*-Diethynylethenes via Double Vinylic C–H Bond Activation. *Org. Lett.* **2021**, *23*, 1199–1203.
- (293) Cera, G.; Haven, T.; Ackermann, L. Iron-Catalyzed C–H Alkynylation through Triazole Assistance: Expedient Access to Bioactive Heterocycles. *Chem. - Eur. J.* **2017**, *23*, 3577–3582.
- (294) Weak Coordinated Nitrogen Functionality Enabled Regioselective C–H Alkynylation via Pd(II)/Mono-*N*-Protected Amino Acid Catalysis. *Chem. Commun.* **2020**, *56*, 11255–11258.
- (295) Feng, C.; Feng, D.; Luo, Y.; Loh, T.-P. Rhodium(III)-Catalyzed Olefinic C–H Alkynylation of Acrylamides Using Tosyl-Imide as Directing Group. *Org. Lett.* **2014**, *16*, 5956–5959.
- (296) Collins, K. D.; Lied, F.; Glorius, F. Preparation of Conjugated 1,3-Enynes by Rh(III)-Catalyzed Alkynylation of Alkenes via C–H Activation. *Chem. Commun.* **2014**, *50*, 4459–4461.
- (297) Xie, F.; Qi, Z.; Yu, S.; Li, X. Rh(III)- and Ir(III)-Catalyzed C–H Alkynylation of Arenes under Chelation Assistance. *J. Am. Chem. Soc.* **2014**, *136*, 4780–4787.
- (298) Zhao, T.; Qin, D.; Han, W.; Yang, S.; Feng, B.; Gao, G.; You, J. Co(III)-Catalyzed *Z*-Selective Oxidative C–H/C–H Cross-Coupling of Alkenes with Triisopropylsilylacetylene. *Chem. Commun.* **2019**, *55*, 6118–6121.
- (299) Wu, G.; Ouyang, W.; Chen, Q.; Huo, Y.; Li, X. Cross-Dehydrogenative Alkynylation of Sulfonamides and Amides with Terminal Alkynes via Ir(III) Catalysis. *Org. Chem. Front.* **2019**, *6*, 284–289.
- (300) Yang, C.; Hassanpour, A.; Ghorbanpour, K.; Abdolmohammadi, S.; Vessally, E. Recent Advances in Direct Trifluoromethylation of Olefinic C–H Bonds. *RSC Adv.* **2019**, *9*, 27625–27639.
- (301) Feng, C.; Loh, T.-P. Directing-Group-Assisted Copper-Catalyzed Olefinic Trifluoromethylation of Electron-Deficient Alkenes. *Angew. Chem., Int. Ed.* **2013**, *52*, 12414–12417.
- (302) Besset, T.; Cahard, D.; Pannecoucke, X. Regio- and Diastereoselective Cu-Mediated Trifluoromethylation of Functionalized Alkenes. *J. Org. Chem.* **2014**, *79*, 413–418.
- (303) Li, L.; Guo, J.-Y.; Liu, X.-G.; Chen, S.; Wang, Y.; Tan, B.; Liu, X.-Y. Amide Groups Switch Selectivity: C–H Trifluoromethylation of  $\alpha,\beta$ -Unsaturated Amides and Subsequent Asymmetric Transformation. *Org. Lett.* **2014**, *16*, 6032–6035.
- (304) Kong, W.; Casimiro, M.; Fuentes, N.; Merino, E.; Nevado, C.; Metal-Free Aryltrifluoromethylation of Activated Alkenes. *Angew. Chem., Int. Ed.* **2013**, *52*, 13086–13090.
- (305) Zhang, B.; Mück-Lichtenfeld, C.; Daniliuc, C. G.; Studer, A. 6-Trifluoromethyl-Phenanthridines through Radical Trifluoromethylation of Isonitriles. *Angew. Chem., Int. Ed.* **2013**, *52*, 10792–10795.
- (306) Zhang, B.; Studer, A. 2-Trifluoromethylated Indoles via Radical Trifluoromethylation of Isonitriles. *Org. Lett.* **2014**, *16*, 1216–1219.
- (307) Fang, Z.; Ning, Y.; Mi, P.; Liao, P.; Bi, X. Catalytic C–H  $\alpha$ -Trifluoromethylation of  $\alpha,\beta$ -Unsaturated Carbonyl Compounds. *Org. Lett.* **2014**, *16*, 1522–1525.
- (308) Sun, S.-Z.; Xu, H.; Dai, H.-X. Copper-catalyzed  $\alpha$ -Selective C–H Trifluoromethylation of Acrylamides with TMSCF<sub>3</sub>. *Chin. Chem. Lett.* **2019**, *30*, 969–972.
- (309) Chen, N.; Xu, H.-C. Electrochemical Generation of Nitrogen-Centered Radicals for Organic Synthesis. *Green Synth. Catal.* **2021**, *2*, 165–178.
- (310) Xu, H.-H.; Song, J.; Xu, H.-C. Electrochemical Difluoromethylation of Electron-Deficient Alkenes. *ChemSusChem*, **2019**, *12*, 3060–3063.
- (311) Ruyet, L.; Lapuh, M. I.; Koshti, V. K.; Földesi, T.; Jubault, P.; Poisson, T.; Zoltán, N.; Besset, T. *Z*-Selective Pd-Catalyzed 2,2,2-Trifluoroethylation of Acrylamides at Room Temperature. *Chem. Commun.* **2021**, *57*, 6241–6244.
- (312) Lied, F.; Patra, T.; Glorius, F. Group 9 Transition Metal-Catalyzed C–H Halogenations. *Isr. J. Chem.* **2017**, *57*, 945–952.
- (313) Das, R.; Kapur, M. Transition-Metal-Catalyzed Site-Selective C–H Halogenation Reactions. *Asian J. Org. Chem.* **2018**, *7*, 1524–1541.
- (314) Mal, S.; Jana, M.; Sarkar, S. Recent Update on Transition Metal-Free C(sp<sup>2</sup>)–H Bond Halogenation in (Hetero) Arenes. *ChemistrySelect* **2021**, *6*, 11299–11330.
- (315) Paik, A.; Paul, S.; Bhowmik, S.; Das, R.; Naveen, T.; Rana, S. Recent Advances in First-Row Transition-Metal-Mediated C–H Halogenation of (Hetero)arenes and Alkanes. *Asian J. Org. Chem.* **2022**, *11*, e202200060.
- (316) Kuhl, N.; Schröder, N.; Glorius, F. Rh(III)-Catalyzed Halogenation of Vinylic C–H Bonds: Rapid and General Access to *Z*-Halo Acrylamides. *Org. Lett.* **2013**, *15*, 3860–3863.
- (317) Yu, D.-G.; Gensch, T.; de Azambuja, F.; Vásquez-Céspedes, S.; Glorius, F. Co(III)-Catalyzed C–H Activation/Formal S<sub>N</sub>-Type Reactions: Selective and Efficient Cyanation, Halogenation, and Allylation. *J. Am. Chem. Soc.* **2014**, *136*, 17722–17725.
- (318) Chen, M.-Y.; Pannecoucke, X.; Jubault, P.; Besset, T. Pd-Catalyzed Selective Chlorination of Acrylamides at Room Temperature. *Org. Lett.* **2020**, *22*, 7556–7561.
- (319) Harnedy, J.; Hareram, M. D.; Tizzard, G. J.; Coles, S. J.; Morrill, L. C. Electrochemical Oxidative *Z*-selective C(sp<sup>2</sup>)–H Chlorination of Acrylamides. *Chem. Commun.* **2021**, *57*, 12643–12646.
- (320) Pan, S.; Sarkar, S.; Ghosh, B.; Samanta, R. Transition Metal Catalysed Direct Construction of 2-Pyridone Scaffolds through C–H Bond Functionalizations. *Org. Biomol. Chem.* **2021**, *19*, 10516–10529.
- (321) Su, Y.; Zhao, M.; Han, K.; Song, G.; Li, X. Synthesis of 2-Pyridones and Iminoesters via Rh(III)-Catalyzed Oxidative Coupling between Acrylamides and Alkynes. *Org. Lett.* **2010**, *12*, 5462–5465.
- (322) Piou, T.; Rovis, T. Electronic and Steric Tuning of a Prototypical Piano Stool Complex: Rh(III) Catalysis for C–H Functionalization. *Acc. Chem. Res.* **2018**, *51*, 170–180.
- (323) Hyster, T. K.; Rovis, T. An Improved Catalyst Architecture for Rhodium(III) Catalyzed C–H Activation and Its Application to Pyridone Synthesis. *Chem. Sci.* **2011**, *2*, 1606–1610.
- (324) Ackermann, L.; Lygin, A. V.; Hofmann, N. Ruthenium-Catalyzed Oxidative Synthesis of 2-Pyridones through C–H/N–H Bond Functionalizations. *Org. Lett.* **2011**, *13*, 3278–3281.
- (325) Matsubara, T.; Ilies, L.; Nakamura, E. Oxidative C–H Activation Approach to Pyridone and Isoquinolone through an Iron-Catalyzed Coupling of Amides with Alkynes. *Chem. - Asian J.* **2016**, *11*, 380–384.
- (326) Shankar, M.; Guntreddi, T.; Ramesh, E.; Sahoo, A. K. Transformable Sulfoximine Assisted One-Pot Double Annulation

of Vinylic C–H Bonds with Unactivated Alkynes. *Org. Lett.* **2017**, *19*, 5665–5668.

(327) Mukherjee, K.; Shankar, M.; Ghosh, K.; Sahoo, A. K. An Orchestrated Unsymmetrical Annulation Episode of C(sp<sup>2</sup>)–H Bonds with Alkynes and Quinones: Access to Spiro-isoquinolones. *Org. Lett.* **2018**, *20*, 1914–1918.

(328) Xu, Y.; Li, B.; Zhang, X.; Fan, X. One-pot Synthesis of Fused N,O-Heterocycles through Rh(III)-Catalyzed Cascade Reactions of Aromatic/Vinylic N-Alkoxyamides with 4-Hydroxy-2-alkynoates. *Adv. Synth. Catal.* **2018**, *360*, 2613–2620.

(329) Garad, D. N.; Mhaske, S. B. Ru-Catalyzed Regioselective Cascade Annulation of Acrylamides with 2-Alkynoates for the Synthesis of Various 6-Oxo Nicotinic Acid Esters. *J. Org. Chem.* **2019**, *84*, 1863–1870.

(330) Grigorjeva, L.; Daugulis, O. Cobalt-Catalyzed, Aminoquinoline-Directed C(sp<sup>2</sup>)–H Bond Alkenylation by Alkynes. *Angew. Chem., Int. Ed.* **2014**, *53*, 10209–10212.

(331) Grigorjeva, L.; Daugulis, O. Cobalt-Catalyzed Direct Carbonylation of Aminoquinoline Benzamides. *Org. Lett.* **2014**, *16*, 4688–4690.

(332) Grigorjeva, L.; Daugulis, O. Cobalt-Catalyzed, Aminoquinoline-Directed Coupling of sp<sup>2</sup> C–H Bonds with Alkenes. *Org. Lett.* **2014**, *16*, 4684–4687.

(333) Tan, J.-F.; Bormann, C.T.; Severin, K.; Cramer, N. Alkynyl Triazenes as Fluoroalkyne Surrogates: Regioselective Access to 4-Fluoro-2-pyridones by a Rh(III)-Catalyzed C–H Activation–Lossen Rearrangement–Wallach Reaction. *ACS Catal.* **2020**, *10*, 3790–3796.

(334) Ma, C.; Fang, P.; Mei, T.-S. Recent Advances in C–H Functionalization Using Electrochemical Transition Metal Catalysis. *ACS Catal.* **2018**, *8*, 7179–7189.

(335) Xing, Y.-K.; Chen, X.-R.; Yang, Q.-L.; Zhang, S.-Q.; Guo, H.-M.; Hong, X.; Mei, T.-S. Divergent Rhodium-Catalyzed Electrochemical Vinylic C–H Annulation of Acrylamides with Alkynes. *Nat. Commun.* **2021**, *12*, 930.

(336) Han, Z.-J.; Zhang, Z.-X.; Li, W.-P.; Du, Z.-H.; Tao, B.-X.; Da, C.-S.; Jiao, Z.-Y.; Chen, H.; Li, Y. Ruthenium-Catalyzed Double C(sp<sup>2</sup>)–H Functionalizations of Fumaramides with Alkynes for the Divergent Synthesis of Pyridones and Naphthyridinediones. *ChemCatChem* **2020**, *12*, 2538–2547.

(337) Quiñones, N.; Seoane, A.; García-Fandiño, R.; Mascareñas, J. L.; Gulías, M. Rhodium(III)-Catalyzed Intramolecular Annulations Involving Amide-Directed C–H Activations: Synthetic Scope and Mechanistic Studies. *Chem. Sci.* **2013**, *4*, 2874–2879.

(338) Xu, X.; Liu, Y.; Park, C.-M. Rhodium(III)-Catalyzed Intramolecular Annulation through C–H Activation: Total Synthesis of (±)-Antofine, (±)-Septicine, (±)-Tylophorine, and Rosettacin. *Angew. Chem., Int. Ed.* **2012**, *51*, 9372–9376.

(339) Saiegh, T. J.; Chédotal, H.; Meyer, C.; Coossy, J. Rhodium(III)-Catalyzed C(sp<sup>2</sup>)–H Functionalization of Cyclobutenes. Access to Cyclobuta[c]pyridones and -Pyridines. *Org. Lett.* **2019**, *21*, 8364–8368.

(340) Krieger, J.-P.; Lesuisse, D.; Ricci, G.; Perrin, M.-A.; Meyer, C.; Coossy, J. Rhodium(III)-Catalyzed C–H Activation/Heterocyclization as a Macrocyclization Strategy. Synthesis of Macrocyclic Pyridones. *Org. Lett.* **2017**, *19*, 2706–2709.

(341) Lee, S.; Semakul, N.; Rovis, T. Direct Regio- and Diastereoselective Synthesis of δ-Lactams from Acrylamides and Unactivated Alkenes Initiated by Rh<sup>III</sup>-Catalyzed C–H Activation. *Angew. Chem., Int. Ed.* **2020**, *59*, 4965–4969.

(342) Zhong, R.; Xu, Y.; Sun, M.; Wang, Y. Palladium-Catalyzed Regioselective C–H Functionalization/Annulation Reaction of Amides and Allylbenzenes for the Synthesis of Isoquinolinones and Pyridinones. *J. Org. Chem.* **2021**, *86*, 5255–5264.

(343) Zhang, D.; Gao, F.; Nian, Y.; Zhou, Y.; Jiang, H.; Liu, H. Palladium-Catalyzed Picolinamide-Directed Coupling of C(sp<sup>2</sup>)–

H and C(sp<sup>2</sup>)–H: A Straightforward Approach to Quinolinone and Pyridone Scaffolds. *Chem. Commun.* **2015**, *51*, 7509–7511.

(344) Zhou, Z.; Liu, G.; Lu, X. Regiocontrolled Coupling of Aromatic and Vinylic Amides with α-Allenols To Form γ-Lactams via Rhodium(III)-Catalyzed C–H Activation. *Org. Lett.* **2016**, *18*, 5668–5671.

(345) Thrimurtulu, N.; Dey, A.; Maiti, D.; Volla, C. M. R. Cobalt-Catalyzed sp<sup>2</sup>-C–H Activation: Intermolecular Heterocyclization with Allenes at Room Temperature. *Angew. Chem., Int. Ed.* **2016**, *55*, 12361–12365.

(346) Wang, W.; Peng, X.; Qin, X.; Zhao, X.; Ma, C.; Tung, C.-H.; Xu, Z. Synthesis of Quinolinones with Palladium-Catalyzed Oxidative Annulation between Acrylamides and Alkynes. *J. Org. Chem.* **2015**, *80*, 2835–2841.

(347) Zhu, Y.-Q.; Hui, L.-W.; Niu, Y.-X.; Lv, L.-G.; Zhu, K. Reaction of Cycloalkene-1-carboxamides with Aryl Boronates via Rhodium(III)-Catalyzed C–H Activation: A Versatile Route to 3,4-Cycloalkaquinolin-2(1H)-ones. *Adv. Synth. Catal.* **2019**, *361*, 5400–5405.

(348) Giri, R.; Shi, B.-F.; Engle, K. M.; Mangel, N.; Yu, J.-Q. Transition Metal-Catalyzed C–H Activation Reactions: Diastereoselectivity and Enantioselectivity. *Chem. Soc. Rev.* **2009**, *38*, 3242–3272.

(349) Newton, C. G.; Wang, S. G.; Oliveira, C. C.; Cramer, N. Catalytic Enantioselective Transformations Involving C–H Bond Cleavage by Transition-Metal Complexes. *Chem. Rev.* **2017**, *117*, 8908–8976.

(350) Diesel, J.; Cramer, N. Generation of Heteroatom Stereocenters by Enantioselective C–H Functionalization. *ACS Catal.* **2019**, *9*, 9164–9177.

(351) Woźniak, L.; Tan, J.-F.; Nguyen, Q.-H.; du Vigné, A. M.; Smal, V.; Cao, Y.-X.; Cramer, N. Catalytic Enantioselective Functionalizations of C–H Bonds by Chiral Iridium Complexes. *Chem. Rev.* **2020**, *120*, 10516–10543.

(352) Achar, T. K.; Maiti, S.; Jana, S.; Maiti, D. Transition Metal Catalyzed Enantioselective C(sp<sup>2</sup>)–H Bond Functionalization. *ACS Catal.* **2020**, *10*, 13748–13793.

(353) Hassan, I. S.; Ta, A. N.; Danneman, M. W.; Semakul, N.; Burns, M.; Basch, C. H.; Dippon, V. N.; McNaughton, B. R.; Rovis, T. Asymmetric δ-Lactam Synthesis with a Monomeric Streptavidin Artificial Metalloenzyme. *J. Am. Chem. Soc.* **2019**, *141*, 4815–4819.

(354) Wilson, M. E.; Whitesides, G. M. Conversion of a Protein to a Homogeneous Asymmetric Hydrogenation Catalyst by Site-Specific Modification with a Diphosphinerhodium(I) Moiety. *J. Am. Chem. Soc.* **1978**, *100*, 306–307.

(355) Schwizer, F.; Okamoto, Y.; Heinisch, T.; Gu, Y.; Pellizzoni, M. M.; Lebrun, V.; Reuter, R.; Köhler, V. Lewis, J. C.; Ward, T. R. Artificial Metalloenzymes: Reaction Scope and Optimization Strategies. *Chem. Rev.* **2018**, *118*, 142–231.

(356) Wang, S.-G.; Liu, Y.; Cramer, N. Asymmetric Alkenyl C–H Functionalization by Cp<sup>\*</sup>Rh<sup>III</sup> forms 2H-Pyrrol-2-ones through [4+1]-Annulation of Acryl Amides and Allenes. *Angew. Chem., Int. Ed.* **2019**, *58*, 18136–18140.

(357) Shi, L.; Yu, K.; Wang, B. Regioselective Synthesis of Multisubstituted Isoquinolones and Pyridones via Rh(III)-Catalyzed Annulation Reactions. *Chem. Commun.* **2015**, *51*, 17277–17280.

(358) Wu, J.; Wang, D.; Wan, Y.; Ma, C. Rhodium-Catalyzed Tunable Oxidative Cyclization toward the Selective Synthesis of α-Pyrones and Furans. *Chem. Commun.* **2016**, *52*, 1661–1664.

(359) Ma, B.; Wu, P.; Wang, X.; Wang, Z.; Lin, H.-X.; Dai, H.-X. Efficient Synthesis of Spirooxindole Pyrrolones by a Rhodium(III)-Catalyzed C–H Activation/Carbene Insertion/Lossen Rearrangement Sequence. *Angew. Chem., Int. Ed.* **2019**, *58*, 13335–13339.

- (360) Kong, W.-J.; Liu, Y.-J.; Xu, H.; Chen, Y.-Q.; Dai, H.-X.; Yu, J.-Q. Pd-Catalyzed  $\alpha$ -Selective C–H Functionalization of Olefins: En Route to 4-Imino- $\beta$ -Lactams. *J. Am. Chem. Soc.* **2016**, *138*, 2146–2149.
- (361) Iwasaki, M.; Miki, N.; Tsuchiya, Y.; Nakajima, K.; Nishihara, Y. Synthesis of Benzoiselenazolone Derivatives by Nickel-Catalyzed Dehydrogenative Direct Selenation of C(sp<sup>2</sup>)–H Bonds with Elemental Selenium in Air. *Org. Lett.* **2017**, *19*, 1092–1095.
- (362) Chen, M.-Y.; Pannecoucke, X.; Jubault, P.; Besset, T. Access to Isothiazolones from Simple Acrylamides by Pd-Catalyzed C–H Bond Activation. *J. Org. Chem.* **2019**, *84*, 13194–13202.
- (363) Li, W.; Zhou, H.; He, Y.; Zeng, G.; Zheng, Y.; Hu, Y.; Chen, Z.; Ge, J.-Y.; Lv, N.; Chen, J. Synthesis of Diverse  $\gamma$ -Lactams via Rh-Catalyzed C(sp<sup>2</sup>)–H Addition to Aliphatic Nitriles. *Org. Lett.* **2022**, *24*, 5090–5094.
- (364) Lin, C.; Li, D.; Wang, B.; Yao, J.; Zhang, Y. Direct *ortho*-Thiolation of Arenes and Alkenes by Nickel Catalysis. *Org. Lett.* **2015**, *17*, 1328–1331.
- (365) Liu, C.; Fang, Y.; Wang, S.-Y.; Ji, S.-J. Highly Regioselective Rh<sup>III</sup>-Catalyzed Thiolation of *N*-Tosyl Acrylamides: General Access to (*Z*)- $\beta$ -Alkenyl Sulfides. *Org. Lett.* **2018**, *20*, 6112–6116.
- (366) Li, L.; Shen, Q.; Yin, C.; Zheng, X.; Zhong, T.; Yu, C. Regioselective Synthesis of (*Z*)-Alkenyl Thioethers via Rh(III)-Catalyzed Thiolation of *N*-2,6-Difluoroaryl Acrylamides. *Tetrahedron Lett.*, **2022**, *103*, 153981.
- (367) Singh, B. K.; Bairy, G.; Jana, R. A General Copper/Manganese Cocatalyzed C–H Selenation of Arenes, Heteroarenes, and Alkenes under Air. *ChemistrySelect* **2017**, *2*, 9227–9232.
- (368) Zhao, Q.; Poisson, T.; Pannecoucke, X.; Bouillon, J.-P.; Besset, T. Pd-Catalyzed Diastereoselective Trifluoromethylthiolation of Functionalized Acrylamides. *Org. Lett.* **2017**, *19*, 5106–5109.
- (369) Lin, C.; Chen, Z.; Liu, Z.; Zhang, Y. Direct *ortho*-Acyloxylation of Arenes and Alkenes by Cobalt Catalysis. *Adv. Synth. Catal.* **2018**, *360*, 519–532.
- (370) Shaaban, S.; Liang, S.; Liu, N.-W.; Manolikakes, G. Synthesis of Sulfones via Selective C–H-functionalization. *Org. Biomol. Chem.* **2017**, *15*, 1947–1955.
- (371) Zhu, T.; Wu, J. Directing-Group-Assisted C(sp<sup>2</sup>)–H Arylsulfonylation from Sulfur Dioxide. *Org. Lett.* **2020**, *22*, 7094–7097.
- (372) Zhu, T.; Rojsitthisak, P.; Wu, J. Generation of (*Z*)- $\beta$ -Alkenyl Alkylsulfones via a Copper-Catalyzed Decarboxylative Alkylsulfonylation. *Org. Chem. Front.* **2020**, *7*, 4050–4056.
- (373) Su, W.; Gong, T.-J.; Xiao, B.; Fu, Y. Rhodium(III)-Catalyzed Cyanation of Vinylic C–H Bonds: *N*-cyano-*N*-phenyl-*p*-toluenesulfonamide as a Cyanation Reagent. *Chem. Commun.* **2015**, *51*, 11848–11851.
- (374) Denmark, S. E.; Regens, C. S. Palladium-Catalyzed Cross-Coupling Reactions of Organosilanols and Their Salts: Practical Alternatives to Boron- and Tin-Based Methods. *Acc. Chem. Res.* **2008**, *41*, 1486–1499.
- (375) Sore, H. F.; Galloway, W. R. J. D.; Spring, D. R. Palladium-Catalyzed Cross-Coupling of Organosilicon Reagents. *Chem. Soc. Rev.* **2012**, *41*, 1845–1866.
- (376) Pan, J.-L.; Chen, C.; Ma, Z.-G.; Zhou, J.; Wang, L.-R.; Zhang, S.-Y.; Stereoselective Synthesis of *Z*-Vinylsilanes via Palladium-Catalyzed Direct Intermolecular Silylation of C(sp<sup>2</sup>)–H Bonds. *Org. Lett.* **2017**, *19*, 5216–5219.
- (377) Zhang, L.-B.; Hao, X.-Q.; Zhang, S.-K.; Liu, Z.-J.; Zheng, X.-X.; Gong, J.-F.; Niu, J.-L.; Song, M.-P. Cobalt-Catalyzed C(sp<sup>2</sup>)–H Alkoxylation of Aromatic and Olefinic Carboxamides. *Angew. Chem., Int. Ed.* **2015**, *54*, 272–275.
- (378) Sauermann, N.; Meyer, T. H.; Tian, C.; Ackermann, L. Electrochemical Cobalt-Catalyzed C–H Oxygenation at Room Temperature. *J. Am. Chem. Soc.* **2017**, *139*, 18452–18455.
- (379) Ryu, J.; Kwak, J.; Shin, K.; Lee, D.; Chang, S. Ir(III)-Catalyzed Mild C–H Amidation of Arenes and Alkenes: An Efficient Usage of Acyl Azides as the Nitrogen Source. *J. Am. Chem. Soc.* **2013**, *135*, 12861–12868.
- (380) Kim, H.; Park, G.; Park, J.; Chang, S. A Facile Access to Primary Alkylamines and Anilines via Ir(III)-Catalyzed C–H Amidation Using Azidoformates. *ACS Catal.* **2016**, *6*, 5922–5929.
- (381) Liu, Y.; Xie, F.; Jia, A.-Q.; Li, X. Cp\*Co(III)-Catalyzed Amidation of Olefinic and Aryl C–H Bonds: Highly Selective Synthesis of Enamides and Pyrimidones. *Chem. Commun.* **2018**, *54*, 4345–4348.
- (382) Kakiuchi, F.; Tanaka, Y.; Sato, T.; Chatani, N.; Murai, S. Catalytic Addition of Olefinic C–H Bonds to Olefins. *Chem. Lett.* **1995**, *24*, 679–680.
- (383) Sato, T.; Kakiuchi, F.; Chatani, N.; Murai, S. Ruthenium-Catalyzed Reactions of Acyclic  $\alpha,\beta$ -Enones with Olefins and Their Reaction Mechanisms. *Chem. Lett.* **1998**, *27*, 893–894.
- (384) Dethle, D. H.; Nirpal, A. K.; Beeralingappa, N. C.; Kumar, V.; Srivastava, A.; Mishra, K. B.; Bhat, A. A. Ruthenium-Catalyzed Stereo- and Chemoselective Oxidative Coupling of Vinyl Ketones: Efficient Access to (*E,E*)-1,6-Dioxo-2,4-dienes. *Chem. Commun.* **2022**, *58*, 3063–3066.
- (385) Luo, C.-Z.; Jayakumar, J.; Gandeepan, P.; Wu, Y.-C.; Cheng, C.-H. Rhodium(III)-Catalyzed Vinylic C–H Activation: A Direct Route toward Pyridinium Salts. *Org. Lett.* **2015**, *17*, 924–927.
- (386) Zhao, Y.; Li, S.; Zheng, X.; Tang, J.; She, Z.; Gao, G.; You, J. Rh/Cu-Catalyzed Cascade [4+2] Vinylic C–H O-Annulation and Ring Contraction of  $\alpha$ -Aryl Enones with Alkynes in Air. *Angew. Chem., Int. Ed.* **2017**, *56*, 4286–4289.
- (387) Zhao, Y.; Yu, C.; Wang, T.; She, Z.; Zheng, X.; You, J.; Gao, G. Tandem Rh-Catalyzed [4 + 2] Vinylic C–H O-Annulation of Exocyclic Enones with Alkynes and 1,5-H Shift. *Org. Lett.* **2018**, *20*, 1074–1077.
- (388) Koech, P. K.; Krische, M. J. Phosphine Catalyzed  $\alpha$ -Arylation of Enones and Enals Using Hypervalent Bismuth Reagents: Regiospecific Enolate Arylation via Nucleophilic Catalysis. *J. Am. Chem. Soc.* **2004**, *126*, 5350–5351.
- (389) Sousa E Silva, F. C.; Van, N. T.; Wengryniuk, S. E. Direct C–H  $\alpha$ -Arylation of Enones with ArI(O<sub>2</sub>C)<sub>2</sub> Reagents. *J. Am. Chem. Soc.* **2020**, *142*, 64–69.
- (390) Qin, G.; Chen, X.; Yang, L.; Huang, H. Copper-Catalyzed  $\alpha$ -Benzoylation of Enones via Radical-Triggered Oxidative Coupling of Two C–H Bonds. *ACS Catal.* **2015**, *5*, 2882–2885.
- (391) Yoo, J.; Ha, H.-J.; Kim, B.; Cho, C.-W. Synthesis of  $\alpha$ -Trifluoromethylthio- $\alpha,\beta$ -Unsaturated Carbonyl Compounds by DABCO-Mediated Electrophilic Trifluoromethylthiolation with *N*-SCF<sub>3</sub>-Dibenzenesulfonimide. *J. Org. Chem.* **2020**, *85*, 7077–7085.
- (392) Lee, S. I.; Kang, B. C.; Hwang, G.-S.; Ryu, D. H. Catalytic Carbon Insertion into the  $\beta$ -Vinyl C–H Bond of Cyclic Enones with Alkyl Diazoacetates. *Org. Lett.* **2013**, *15*, 1428–1431.
- (393) Lee, S. I.; Hwang, G.-S.; Ryu, D. H. Catalytic Enantioselective Carbon Insertion into the  $\beta$ -Vinyl C–H Bond of Cyclic Enones. *J. Am. Chem. Soc.* **2013**, *135*, 7126–7129.
- (394) Xie, Y.-Y.; Wang, Y.-C.; Qu, H.-E.; Tan, X.-C.; Wang, H.-S.; Pan, Y.-M. Regioselective Synthesis of  $\beta$ -Aryl Enaminones and 1,4,5-Trisubstituted 1,2,3-Triazoles from Chalcones and Benzyl Azides. *Adv. Synth. Catal.* **2014**, *356*, 3347–3355.
- (395) Nguyen, K. X.; Pham, P. H.; Nguyen, T. T.; Yang, C.-H.; Pham, H. T. B.; Nguyen, T. T.; Wang, H.; Phan, N. T. S. Trisulfur-Radical-Anion-Triggered C(sp<sup>2</sup>)–H Amination of Electron-Deficient Alkenes. *Org. Lett.* **2020**, *22*, 9751–9756.
- (396) Chatani, N.; Kamitani, A.; Murai, S. Ruthenium-Catalyzed Reaction of  $\alpha,\beta$ -Unsaturated Imines with Carbon Monoxide and Alkenes Leading to  $\beta,\gamma$ -Unsaturated  $\gamma$ -Butyrolactams: Involvement of Direct Carbonylation at Olefinic C–H Bonds as a Key Step. *J. Org. Chem.* **2002**, *67*, 7014–7018.
- (397) Colby, D. A.; Bergman, R. G.; Ellman, J. A. Stereoselective

- Alkylation of  $\alpha,\beta$ -Unsaturated Imines via C–H Bond Activation. *J. Am. Chem. Soc.* **2006**, *128*, 5604–5605.
- (398) Colby, D. A.; Bergman, R. G.; Ellman, J. A. Synthesis of Dihydropyridines and Pyridines from Imines and Alkynes via C–H Activation. *J. Am. Chem. Soc.* **2008**, *130*, 3645–3651.
- (399) Tsai, A. S.; Bergman, R. G.; Ellman, J. A. Asymmetric Synthesis of (–)-Incarvilleatine Employing an Intramolecular Alkylation via Rh-Catalyzed Olefinic C–H Bond Activation. *J. Am. Chem. Soc.* **2008**, *130*, 6316–6317.
- (400) Kuninobu, Y.; Nishina, Y.; Matsuki, T.; Takai, K. Synthesis of Cp–Re Complexes via Olefinic C–H Activation and Successive Formation of Cyclopentadienes. *J. Am. Chem. Soc.* **2008**, *130*, 14062–14063.
- (401) Mesganaw, T.; Ellman, J. A. Convergent Synthesis of Diverse Tetrahydropyridines via Rh(I)-Catalyzed C–H Functionalization Sequences. *Org. Process Res. Dev.* **2014**, *18*, 1097–1104.
- (402) Sun, D.; Confair, D. N.; Ellman, J. A. Rhodium-Catalyzed C–H Alkenylation/Electrocyclization Cascade Provides Dihydropyridines That Serve as Versatile Intermediates to Diverse Nitrogen Heterocycles. *Acc. Chem. Res.* **2021**, *54*, 1766–1778.
- (403) Duttwyler, S.; Lu, C.; Rheingold, A. L.; Bergman, R. G.; Ellman, J. A. Highly Diastereoselective Synthesis of Tetrahydropyridines by a C–H Activation–Cyclization–Reduction Cascade. *J. Am. Chem. Soc.* **2012**, *134*, 4064–4067.
- (404) Duttwyler, S.; Chen, S.; Takase, M. K.; Wiberg, K. B.; Bergman, R. G.; Ellman, J. A. Proton Donor Acidity Controls Selectivity in Nonaromatic Nitrogen Heterocycle Synthesis. *Science* **2013**, *339*, 678–682.
- (405) Yamakawa, T.; Yoshikai, N. Annulation of  $\alpha,\beta$ -Unsaturated Imines and Alkynes via Cobalt-Catalyzed Olefinic C–H Activation. *Org. Lett.* **2013**, *15*, 196–199.
- (406) Ischay, M. A.; Takase, M. K.; Bergman, R. G.; Ellman, J. A. Unstabilized Azomethine Ylides for the Stereoselective Synthesis of Substituted Piperidines, Tropanes, and Azabicyclo[3.1.0] Systems. *J. Am. Chem. Soc.* **2013**, *135*, 2478–2481.
- (407) Mesganaw, T.; Ellman, J. A. Preparative Synthesis of Highly Substituted Tetrahydropyridines via a Rh(I)-Catalyzed C–H Functionalization Sequence. *Org. Process Res. Dev.* **2014**, *18*, 1105–1109.
- (408) Zhang, Q.-R.; Huang, J.-R.; Zhang, W.; Dong, L. Highly Functionalized Pyridines Synthesis from *N*-Sulfonyl Ketimines and Alkynes Using the N–S Bond as an Internal Oxidant. *Org. Lett.* **2014**, *16*, 1684–1687.
- (409) Bolotin, D. S.; Bokach, N. A.; Demakova, M. Y.; Kukushkin, V. K. Metal-Involving Synthesis and Reactions of Oximes. *Chem. Rev.* **2017**, *117*, 13039–13122.
- (410) Li, J.; Hu, Y.; Zhang, D.; Liu, Q.; Dong, Y.; Liu, H. Transition Metal-Catalyzed Reactions Involving Oximes. *Adv. Synth. Catal.* **2017**, *359*, 710–771.
- (411) Parthasarathy, K.; Jeganmohan, M.; Cheng, C.-H. Rhodium-Catalyzed One-Pot Synthesis of Substituted Pyridine Derivatives from  $\alpha,\beta$ -Unsaturated Ketoximes and Alkynes. *Org. Lett.* **2008**, *10*, 325–328.
- (412) Parthasarathy, K.; Cheng, C.-H. Rhodium-Catalyzed Gram-Scale Synthesis of Highly Substituted Pyridine Derivatives. *Synthesis* **2009**, *8*, 1400–1402.
- (413) Hyster, T. K.; Rovis, T. Pyridine Synthesis from Oximes and Alkynes via Rhodium(III) Catalysis: Cp\* and Cp<sup>t</sup> Provide Complementary Selectivity. *Chem. Commun.* **2011**, *47*, 11846–11848.
- (414) Too, P. C.; Noji, T.; Lim, Y. J.; Li, X.; Chiba, S. Rhodium(III)-Catalyzed Synthesis of Pyridines from  $\alpha,\beta$ -Unsaturated Ketoximes and Internal Alkynes. *Synlett* **2011**, *19*, 2789–2794.
- (415) Martin, R. M.; Bergman, R. G.; Ellman, J. A. Synthesis of Pyridines from Ketoximes and Terminal Alkynes via C–H Bond Functionalization. *J. Org. Chem.* **2012**, *77*, 2501–2507.
- (416) Chen, S.; Bergman, R. G.; Ellman, J. A. Facile Rh(III)-Catalyzed Synthesis of Fluorinated Pyridines. *Org. Lett.* **2015**, *17*, 2567–2569.
- (417) Shi, Z.; Koester, D. C.; Bouladakis-Arapinis, M.; Glorius, F. Rh(III)-Catalyzed Synthesis of Multisubstituted Isoquinoline and Pyridine *N*-Oxides from Oximes and Diazo Compounds. *J. Am. Chem. Soc.* **2013**, *135*, 12204–12207.
- (418) Das, D.; Sahoo, G.; Biswas, A.; Samanta, R. Rh<sup>III</sup>-Catalyzed Synthesis of Highly Substituted 2-Pyridones using Fluorinated Diazomalonnate. *Chem. - Asian J.* **2020**, *15*, 360–364.
- (419) Lian, Y.; Huber, T.; Hesp, K. D.; Bergman, R. G.; Ellman, J. A. Rhodium(III)-Catalyzed Alkenyl C–H Bond Functionalization: Convergent Synthesis of Furans and Pyrroles. *Angew. Chem., Int. Ed.* **2013**, *52*, 629–633.
- (420) Hummel, J. R.; Ellman, J. A. Cobalt(III)-Catalyzed Synthesis of Indazoles and Furans by C–H Bond Functionalization/Addition/Cyclization Cascades. *J. Am. Chem. Soc.* **2015**, *137*, 490–498.
- (421) Hou, W.; Zhou, B.; Yang, Y.; Feng, H.; Li, Y. Rh(III)-Catalyzed Addition of Alkenyl C–H Bond to Isocyanates and Intramolecular Cyclization: Direct Synthesis 5-Ylidene-pyrrol-2(5H)-ones. *Org. Lett.* **2013**, *15*, 1814–1817.
- (422) Mohanty, S. R.; Pati, B. V.; Banjare, S. K.; Adhikari, G. K. D.; Ravikumar, P. C. Redox-Neutral Cobalt(III)-Catalyzed C–H Activation/Annulation of  $\alpha,\beta$ -Unsaturated Oxime Ether with Alkyne: One-Step Access to Multisubstituted Pyridine. *J. Org. Chem.* **2021**, *86*, 1074–1083.
- (423) Neely, J. M.; Rovis, T. Rh(III)-Catalyzed Regioselective Synthesis of Pyridines from Alkenes and  $\alpha,\beta$ -Unsaturated Oxime Esters. *J. Am. Chem. Soc.* **2013**, *135*, 66–69.
- (424) Yamada, T.; Hashimoto, Y.; Tanaka, K.; Morita, N.; Tamura, O. Palladium(II)-Catalyzed Substituted Pyridine Synthesis from  $\alpha,\beta$ -Unsaturated Oxime Ethers via a C–H Alkenylation/Aza-6 $\pi$ -Electrocyclization Approach. *Org. Lett.* **2021**, *23*, 1659–1663.
- (425) Neely, J. M.; Rovis, T. Rh(III)-Catalyzed Decarboxylative Coupling of Acrylic Acids with Unsaturated Oxime Esters: Carboxylic Acids Serve as Traceless Activators. *J. Am. Chem. Soc.* **2014**, *136*, 2735–2738.
- (426) Zhao, D.; Lied, F.; Glorius, F. Rh(III)-Catalyzed C–H Functionalization/Aromatization Cascade with 1,3-Dienes: A Redox-Neutral and Regioselective Access to Isoquinolines. *Chem. Sci.* **2014**, *5*, 2869–2873.
- (427) Romanov-Mikhailidis, F.; Sedillo, K. F.; Neely, J. M.; Rovis, T. Expedient Access to 2,3-Dihydropyridines from Unsaturated Oximes by Rh(III)-Catalyzed C–H Activation. *J. Am. Chem. Soc.* **2015**, *137*, 8892–8895.
- (428) Lou, S.-J.; Xu, D.-Q.; Xu, Z.-Y. Mild and Versatile Nitrate-Promoted C–H Bond Fluorination. *Angew. Chem., Int. Ed.* **2014**, *53*, 10330–10335.
- (429) Yamada, T.; Hashimoto, Y.; Tanaka, K.; Morita, N.; Tamura, O. Thioether Ligand-Enabled Cationic Palladium(II)-Catalyzed Electrophilic C–H Arylation of  $\alpha,\beta$ -Unsaturated Oxime Ethers. *J. Org. Chem.* **2020**, *85*, 12315–12328.
- (430) Wang, S.-G.; Cramer, N. An Enantioselective Cp\*Rh(III)-Catalyzed C–H Functionalization/Ring-Opening Route to Chiral Cyclopentylamines. *Angew. Chem., Int. Ed.* **2019**, *58*, 2514–2518.
- (431) Chatani, N.; Ishii, Y.; Ie, Y.; Kakiuchi, F.; Murai, S. Ru<sub>3</sub>(CO)<sub>12</sub>- and Rh<sub>4</sub>(CO)<sub>12</sub>-Catalyzed Reactions of Pyridylolefins or *N*-(2-Pyridyl)enamines with CO and Olefins. Carbonylation at Olefinic C–H Bonds. *J. Org. Chem.* **1998**, *63*, 5129–5136.
- (432) Oi, S.; Sakai, K.; Inoue, Y. Ruthenium-Catalyzed Arylation of 2-Alkenylpyridines with Aryl Bromides: Alternative *E,Z*-Selectivity to Mizoroki-Heck Reaction. *Org. Lett.* **2005**, *7*, 4009–4011.
- (433) Ackermann, L.; Born, R.; Álvarez-Bercedo, P. Ruthenium(IV) Alkylidenes as Precatalysts for Direct Arylations

- of Alkenes with Aryl Chlorides and an Application to Sequential Catalysis. *Angew. Chem., Int. Ed.* **2007**, *46*, 6364–6367.
- (434) Zell, D.; Warratz, S.; Gelman, D.; Garden, S. J.; Ackermann, L. Single-Component Phosphinous Acid Ruthenium(II) Catalysts for Versatile C–H Activation by Metal–Ligand Cooperation. *Chem. - Eur. J.* **2016**, *22*, 1248–1252.
- (435) Gramage-Doria, R.; Achelle, S.; Bruneau, C.; Robin-le Guen, F.; Dorcet, V.; Roisnel, T. Ruthenium(II)-Catalyzed C–H (Hetero)-Arylation of Alkenylic 1, n-Diazines (n = 2, 3, and 4): Scope, Mechanism, and Application in Tandem Hydrogenations. *J. Org. Chem.* **2018**, *83*, 1462–1477.
- (436) Huang, Y.; Pan, W.-J.; Wang, Z.-X. Rhodium-Catalyzed Alkenyl C–H Functionalization with Amides. *Org. Chem. Front.* **2019**, *6*, 2284–2290.
- (437) Kuninobu, Y.; Fujii, Y.; Matsuki, T.; Nishina, Y.; Takai, K. Rhodium-Catalyzed Insertion of Nonpolar and Polar Unsaturated Molecules into an Olefinic C–H Bond. *Org. Lett.* **2009**, *11*, 271–274.
- (438) Zhou, B.; Hu, Y.; Wang, C. Manganese-Catalyzed Direct Nucleophilic C(sp<sup>2</sup>)-H Addition to Aldehydes and Nitriles. *Angew. Chem., Int. Ed.* **2015**, *54*, 13659–13663.
- (439) Iliés, L.; Asako, S.; Nakamura, E. Iron-Catalyzed Stereospecific Activation of Olefinic C–H Bonds with Grignard Reagent for Synthesis of Substituted Olefins. *J. Am. Chem. Soc.* **2011**, *133*, 7672–7675.
- (440) Li, Y.; Zhang, X.-S.; Zhu, Q.-L.; Shi, Z.-J. Olefinic C–H Bond Addition to Aryl Aldehyde and Its N-Sulfonylimine via Rh Catalysis. *Org. Lett.* **2012**, *14*, 4498–4501.
- (441) Xu, Y.-H.; Wang, M.; Lu, P.; Loh, T.-P. Palladium-Catalyzed Alkenyl C–H Bond Sulfonylation Reaction Using Organosulfonyl Chlorides. *Tetrahedron* **2013**, *69*, 4403–4407.
- (442) Jiang, Q.; Guo, T.; Wu, K.; Yu, Z. Rhodium(III)-Catalyzed sp<sup>2</sup> C–H Addition to CF<sub>3</sub>-Substituted Unsaturated Ketones. *Chem. Commun.* **2016**, *52*, 2913–2915.
- (443) Yan, R.; Wang, Z.-X. Rhodium-Catalyzed Alkenyl C–H Activation and Oxidative Coupling with Allylic Alcohols. *Asian J. Org. Chem.* **2017**, *7*, 240–247.
- (444) Chaitanya, M.; Anbarasan, P. Rhodium-Catalyzed Cyanation of C(sp<sup>2</sup>)-H Bond of Alkenes. *Org. Lett.* **2015**, *17*, 3766–3769.
- (445) Miao, H.; Wang, Z.-X. Ruthenium-Catalyzed Oxidative Cross-Coupling of Alkenes with Triisopropylsilylacetylene. *Asian J. Org. Chem.* **2022**, *11*, e202200172.
- (446) Carbery, D. R. Enamides: Valuable Organic Substrates. *Org. Biomol. Chem.* **2008**, *6*, 3455–3460.
- (447) Gigant, N.; Chausset-Boissarie, L.; Gillaizeau, I. Direct Metal-Catalyzed Regioselective Functionalization of Enamides. *Chem. - Eur. J.* **2014**, *20*, 7548–7564.
- (448) Zhu, T.; Xie, S.; Rojstithsak, P.; Wu, J. Recent Advances in the Direct β-C(sp<sup>2</sup>)-H Functionalization of Enamides. *Org. Biomol. Chem.* **2020**, *18*, 1504–1521.
- (449) Zhou, H.; Chung, W.-J.; Xu, Y.-H.; Loh, T.-P. Direct Arylation of Cyclic Enamides via Pd(II)-Catalyzed C–H Activation. *Chem. Commun.* **2009**, *48*, 3472–3474.
- (450) Zhou, H.; Xu, Y.-H.; Chung, W.-J.; Loh, T.-P. Palladium-Catalyzed Direct Arylation of Cyclic Enamides with Aryl Silanes by sp<sup>2</sup> C–H Activation. *Angew. Chem., Int. Ed.* **2009**, *48*, 5355–5357.
- (451) Li, X.; Sun, K.; Shen, W.; Zhang, Y.; Lu, M.-Z.; Luo, X.; Luo, H. Rhodium(III)-Catalyzed Direct C–H Arylation of Various Acyclic Enamides with Arylsilanes. *Org. Lett.* **2021**, *23*, 31–36.
- (452) Pankajakshan, S.; Xu, Y.-H.; Cheng, J. K.; Low, M. T.; Loh, T.-P. Palladium-Catalyzed Direct C–H Arylation of Enamides with Simple Arenes. *Angew. Chem., Int. Ed.* **2012**, *51*, 5701–5705.
- (453) Bartocchini, F.; Cannas, D. M.; Fini, F.; Piersanti, G. Palladium(II)-Catalyzed Cross-Dehydrogenative Coupling (CDC) of N-Phthaloyl Dehydroalanine Esters with Simple Arenes: Stereoselective Synthesis of Z-Dehydrophenylalanine Derivatives. *Org. Lett.* **2016**, *18*, 2762–2765.
- (454) Gigant, N.; Chausset-Boissarie, L.; Belhomme, M.; Poisson, T.; Pannecoucke, X.; Gillaizeau, I. Copper-Catalyzed Direct Arylation of Cyclic Enamides Using Diaryliodonium Salts. *Org. Lett.* **2013**, *15*, 278–281.
- (455) Prakash, M.; Muthusamy, S.; Kesavan, V. Copper(I) Bromide Catalyzed Arylation of Cyclic Enamides and Naphthyl-1-acetamides Using Diaryliodonium Salts. *J. Org. Chem.* **2014**, *79*, 7836–7843.
- (456) Zhang, W.; Yu, W.; Yan, Q.; Liu, Z.; Zhang, Y. Synthesis of Substituted Oxazoles via Pd-Catalyzed Tandem Oxidative Cyclization. *Org. Chem. Front.* **2017**, *4*, 2428–2432.
- (457) Liu, Y.; Li, D.; Park, C.-M. Stereoselective Synthesis of Highly Substituted Enamides by an Oxidative Heck Reaction. *Angew. Chem., Int. Ed.* **2011**, *50*, 7333–7336.
- (458) Gou, Q.; Deng, B.; Zhang, H.; Qin, J. Arylations of Substituted Enamides by Aryl Iodides: Regio- and Stereoselective Synthesis of (Z)-β-Amido-β-Arylacrylates. *Org. Lett.* **2013**, *15*, 4604–4607.
- (459) Wang, M.; Zhang, X.; Zhuang, Y.-X.; Xu, Y.-H.; Loh, T.-P. Pd-Catalyzed Intramolecular C–N Bond Cleavage, 1,4-Migration, sp<sup>3</sup> C–H Activation, and Heck Reaction: Four Controllable Diverse Pathways Depending on the Judicious Choice of the Base and Ligand. *J. Am. Chem. Soc.* **2015**, *137*, 1341–1347.
- (460) Zhu, W.; Tong, S.; Zhu, J.; Wang, M.-X. Intramolecular Arylation of Tertiary Enamides through Pd(OAc)<sub>2</sub>-Catalyzed Dehydrogenative Cross-Coupling Reaction: Construction of Fused N-Heterocyclic Scaffolds and Synthesis of Isoindolobenzazepine Alkaloids. *J. Org. Chem.* **2019**, *84*, 2870–2878.
- (461) Matsubara, R.; Nakamura, Y.; Kobayashi, S. Highly Diastereo- and Enantioselective Reactions of Enecarbamates with Ethyl Glyoxylate to Give Optically Active *syn* and *anti* α-Alkyl-β-Hydroxy Imines and Ketones. *Angew. Chem., Int. Ed.* **2004**, *43*, 3258–3260.
- (462) Matsubara, R.; Kawai, N.; Kobayashi, S. Catalytic Enantioselective and Diastereoselective Addition of Aldehyde-Derived Enecarbamates to α-Oxo Aldehydes. *Angew. Chem., Int. Ed.* **2006**, *45*, 3814–3816.
- (463) Kobayashi, S.; Gustafsson, T.; Shimizu, Y.; Kiyohara, H.; Matsubara, R. Enecarbamates as Imine Surrogates: Nucleophilic Addition of 1,3-Dicarbonyl Compounds to Enecarbamates. *Org. Lett.* **2006**, *8*, 4923–4925.
- (464) Baudequin, C.; Zamfira, A.; Tsogoeva, S. B. Highly Enantioselective Organocatalytic Formation of a Quaternary Carbon Center via Chiral Brønsted Acid Catalyzed Self-coupling of Enamides. *Chem. Commun.* **2008**, 4637–4639.
- (465) Zhao, M.-N.; Du, W.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. FeCl<sub>3</sub>-Catalyzed Self-Condensation of Enamides for the Synthesis of Enamido-Substituted Nitrogen-Containing Quaternary Carbon Centers. *Eur. J. Org. Chem.* **2013**, *2013*, 7989–7995.
- (466) Yu, W.; Zhang, W.; Liu, Y.; Liu, Z.; Zhang, Y. Cobalt(III)-Catalyzed Cross-Coupling of Enamides with Allyl Acetates/Maleimides. *Org. Chem. Front.* **2017**, *4*, 77–80.
- (467) Ding, R.; Huang, Z.-D.; Liu, Z.-L.; Wang, T.-X.; Xu, Y.-H.; Loh, T.-P. Palladium-Catalyzed Cross-Coupling of Enamides with Sterically Hindered α-Bromocarbonyls. *Chem. Commun.* **2016**, *52*, 5617–5620.
- (468) Li, P.; Zhao, J.; Xia, C.; Li, F. Direct Oxidative Coupling of Enamides and 1,3-Dicarbonyl Compounds: A Facile and Versatile Approach to Dihydrofurans, Furans, Pyrroles, and Dicarbonyl Enamides. *Org. Lett.* **2014**, *16*, 5992–5995.
- (469) Ding, R.; Lu, W.-G.; Ci, H.; Mao, Y.-Y.; Liu, L. Copper-Catalyzed Oxidative Alkylation of Vinylic C<sub>β</sub>-H of Enamides with Cyclic Ethers. *ChemistrySelect* **2019**, *4*, 6954–6957.

- (470) Tao, J.-Y.; Wang, Y.-X.; Zhang, Q.-H.; Ni, K.; Zhu, T.-H.; Zhu, T.-H.; Zhao, K. Transition-Metal-Free Regioselective and Stereoselective C(sp<sup>2</sup>)-C(sp<sup>3</sup>) Coupling of Enamides with Ethers or Alkanes via Photoredox-catalyzed Cross-Dehydrogenative Coupling Reactions. *Green Chem.* **2022**, *24*, 4004–4011.
- (471) Guo, J.-Y.; Guan, T.; Tao, J.-Y.; Zhao, K.; Loh, T.-P. Stereoselective C(sp<sup>2</sup>)-H Alkylation of Enamides with Unactivated Aliphatic Carboxylic Acids via Decarboxylative Cross-Coupling Reactions. *Org. Lett.* **2019**, *21*, 8395–8399.
- (472) Guo, J.-Y.; Zhang, Z.-Y.; Guan, T.; Mao, L.-W.; Ban, Q.; Zhao, K.; Loh, T.-P. Photoredox-Catalyzed Stereoselective Alkylation of Enamides with *N*-Hydroxyphthalimide Esters via Decarboxylative Cross-Coupling Reactions. *Chem. Sci.* **2019**, *10*, 8792–8798.
- (473) Su, X.-D.; Zhang, B.-B.; Liu, Q.; Cheng, J.-T.; Wang, Z.-X.; Chen, X.-Y. Additive-Free, Visible-Light-Enabled Decarboxylative Alkylation of Enamides. *Org. Lett.* **2021**, *23*, 8262–8266.
- (474) Wang, P.; Zhao, Q.; Xiao, W.; Chen, J. Recent Advances in Visible-Light Photoredox-Catalyzed Nitrogen Radical Cyclization. *Green Synth. Catal.* **2020**, *1*, 42–51.
- (475) Jiang, X.; Zhang, M.-M.; Xiong, W.; Lu, L.-Q.; Xiao, W.-J. Deaminative (Carbonylative) Alkyl-Heck-Type Reactions Enabled by Photocatalytic C–N Bond Activation. *Angew. Chem., Int. Ed.* **2019**, *58*, 2402–2406.
- (476) Wang, J.-X.; Wang, Y.-T.; Zhang, H.; Fu, M.-C. Visible-Light-Induced Iodine-Anion-Catalyzed Decarboxylative/Deaminative C–H Alkylation of Enamides. *Org. Chem. Front.* **2021**, *8*, 4466–4472.
- (477) Bertho, S.; Dondasse, I.; Retailleau, P.; Nicolas, C.; Gillaizeau, I.  $\beta$ -C(sp<sup>2</sup>)-H Alkylation of Enamides Using Xanthate Chemistry. *New J. Chem.* **2020**, *44*, 7129–7141.
- (478) Li, S.; Shan, Q.-C.; Hu, L.-M.; Ma, X.-Q.; Hu, X.-H. Merging Alkenyl C–H Activation with Ring-Opening of 1,2-Oxazetidines: Ruthenium-Catalyzed Aminomethylation of Enamides. *Chem. Commun.* **2020**, *56*, 7969–7972.
- (479) Bhattacharya, T.; Pimparkar, S.; Maiti, D. Combining Transition Metals and Transient Directing Groups for C–H Functionalizations. *RSC Adv.* **2018**, *8*, 19456–19464.
- (480) Higham, J. I.; Bull, J. A. Transient Imine Directing Groups for the C–H Functionalisation of Aldehydes, Ketones and Amines: An Update 2018–2020. *Org. Biomol. Chem.* **2020**, *18*, 7291–7315.
- (481) Xing, D.; Dong, G.-B. Branched-Selective Intermolecular Ketone  $\alpha$ -Alkylation with Unactivated Alkenes via an Enamide Directing Strategy. *J. Am. Chem. Soc.* **2017**, *139*, 13664–13667.
- (482) Crisenza, G. E. M.; Bower, J. F. Branch Selective Murai-type Alkene Hydroarylation Reactions. *Chem. Lett.* **2016**, *45*, 2–9.
- (483) Yang, C.; Hassanpour, A.; Ghorbanpour, K.; Abdolmohammadi, S.; Vessally, E. Recent Advances in Direct Trifluoromethylation of Olefinic C–H Bonds. *RSC Adv.* **2019**, *9*, 27625–27639.
- (484) Charpentier, J.; Früh, N.; Togni, A. Electrophilic Trifluoromethylation by Use of Hypervalent Iodine Reagents. *Chem. Rev.* **2015**, *115*, 650–682.
- (485) Feng, C.; Loh, T.-P. Copper-Catalyzed Olefinic Trifluoromethylation of Enamides at Room Temperature. *Chem. Sci.* **2012**, *3*, 3458–3462.
- (486) Rey-Rodriguez, R.; Retailleau, P.; Bonnet, P.; Gillaizeau, I. Iron-Catalyzed Trifluoromethylation of Enamide. *Chem. - Eur. J.* **2015**, *21*, 3572–3575.
- (487) Yang, H.-B.; Selander, N. A Redox-Economical Synthesis of Trifluoromethylated Enamides with the Langlois Reagent. *Org. Biomol. Chem.* **2017**, *15*, 1771–1775.
- (488) Tang, K.; Chen, Y.; Guan, J.; Wang, Z.; Chen, K.; Xiang, H.; Yang, H. Visible-Light-Promoted Olefinic Trifluoromethylation of Enamides with CF<sub>3</sub>SO<sub>2</sub>Na. *Org. Biomol. Chem.* **2021**, *19*, 7475–7479.
- (489) Zhu, T.-H.; Zhang, Z.-Y.; Tao, J.-Y.; Zhao, K.; Loh, T.-P. Regioselective and Stereoselective Difluoromethylation of Enamides with Difluoromethyltriphenylphosphonium Bromide via Photoredox Catalysis. *Org. Lett.* **2019**, *21*, 6155–6159.
- (490) Zhao, Y.; Zhang, Y.; Liu, Y.; Zhu, T.; Wu, J. Photoredox-Catalyzed Direct C(sp<sup>2</sup>)-H Difluoromethylation of Enamides or Heterocycles with [bis(Difluoroacetoxy)iodo]benzene. *Org. Chem. Front.* **2021**, *8*, 5948–5954.
- (491) Zhao, K.; Zhang, Z.-Y.; Cui, X.-L.; Wang, Y.-X.; Wu, X.-D.; Li, W.-M.; Wu, J.-X.; Zhao, L.-L.; Guo, J.-Y.; Loh, T.-P. Visible-Light-Induced Regio- and Stereoselective C(sp<sup>2</sup>)-H Trifluoroethylation of Enamides with 2,2,2-Trifluoroethyl Iodide. *Org. Lett.* **2020**, *22*, 9029–9035.
- (492) Zhao, K.; Guo, J.-Y.; Guan, T.; Wang, Y.-X.; Tao, J.-Y.; Zhang, Y.; Zhang, Q.-H.; Ni, K.; Loh, T.-P. Photoinitiated Stereoselective Direct C(sp<sup>2</sup>)-H Perfluoroalkylation and Difluoroacetylation of Enamides. *Org. Chem. Front.* **2021**, *8*, 4086–4094.
- (493) Caillot, G.; Dufour, J.; Belhomme, M.-C.; Poisson, T.; Grimaud, L.; Pannecoucke, X.; Gillaizeau, I. Copper-Catalyzed Olefinic C–H Difluoroacetylation of Enamides. *Chem. Commun.* **2014**, *50*, 5887–5890.
- (494) Bertho, S.; Maazaoui, R.; Torun, D.; Dondasse, I.; Abderrahim, R.; Nicolas, C.; Gillaizeau, I. Iron Catalyzed  $\beta$ -C(sp<sup>2</sup>)-H Alkylation of Enamides. *New J. Chem.* **2021**, *45*, 17475–17482.
- (495) Wang, X.; Zhao, S.; Liu, J.; Zhu, D.; Guo, M.; Tang, X.; Wang, G. Copper-Catalyzed C–H Difluoroalkylations and Perfluoroalkylations of Alkenes and (Hetero)arenes. *Org. Lett.* **2017**, *19*, 4187–4190.
- (496) Feng, Z.; Min, Q.-Q.; Zhao, H.-Y.; Gu, J.-W.; Zhang, X. A General Synthesis of Fluoroalkylated Alkenes by Palladium-Catalyzed Heck-Type Reaction of Fluoroalkyl Bromides. *Angew. Chem., Int. Ed.* **2015**, *54*, 1270–1274.
- (497) Feng, Z.; Min, Q.-Q.; Zhang, X. Palladium-Catalyzed Heck-Type Difluoroalkylation of Alkenes with Functionalized Difluoromethyl Bromides. *Synthesis* **2015**, *47*, 2912–2923.
- (498) Jiang, H.; Huang, C.; Guo, J.; Zeng, C.; Zhang, Y.; Yu, S. Direct C–H Functionalization of Enamides and Enecarbamates by Using Visible-Light Photoredox Catalysis. *Chem. - Eur. J.* **2012**, *18*, 15158–15166.
- (499) Wang, H.; Cheng, Y.-Z.; Yu, S. Visible-Light-Promoted and Photocatalyst-Free Trifluoromethylation of Enamides. *Sci. China Chem.* **2016**, *59*, 195–198.
- (500) Xu, Y.-H.; Chok, Y. K.; Loh, T.-P. Synthesis and Characterization of a Cyclic Vinylpalladium(II) Complex: Vinylpalladium Species as the Possible Intermediate in the Catalytic Direct Olefination Reaction of Enamide. *Chem. Sci.* **2011**, *2*, 1822–1825.
- (501) Gigant, N.; Gillaizeau, I. Palladium(II)-Catalyzed Direct Alkenylation of Nonaromatic Enamides. *Org. Lett.* **2012**, *14*, 3304–3307.
- (502) Besset, T.; Kuhl, N.; Patureau, F. W.; Glorius, F. Rh<sup>III</sup>-Catalyzed Oxidative Olefination of Vinylic C–H Bonds: Efficient and Selective Access to Di-unsaturated  $\alpha$ -Amino Acid Derivatives and Other Linear 1,3-Butadienes. *Chem. - Eur. J.* **2011**, *17*, 7167–7171.
- (503) Li, C.; Li, W.-H.; Dong, L. Efficient Synthesis of *N*-Butadiene Substituted Oxindole Derivatives. *Org. Chem. Front.* **2018**, *5*, 3460–3463.
- (504) Feng, C.; Feng, D.; Loh, T.-P. Rhodium(III)-Catalyzed Olefinic C–H Alkynylation of Enamides at Room Temperature. *Chem. Commun.* **2014**, *50*, 9865–9868.
- (505) Xu, Y.-H.; Zhang, Q.-C.; He, T.; Meng, F.-F.; Loh, T.-P. *Adv. Synth. Catal.* **2014**, *356*, 1539–1543.
- (506) Beng, T. K.; Wanjiku, F. Iridium-Catalyzed  $\alpha$ -Alkynylation of Cyclic Nonaromatic Eneformamides: Application to the Synthesis of Azapolycyclic Architectures. *New J. Chem.* **2019**, *43*, 4664–4668.
- (507) Ning, Y.; Ohwada, T.; Chen, F.-E. Transition Metal-Catalyzed Branch-Selective Hydroformylation of Olefins in Organic Synthesis. *Green Synth. Catal.* **2021**, *2*, 247–266.

- (508) Chen, M.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. Palladium-Catalyzed Oxidative Carbonylation of the Alkenyl C-H Bonds of Enamides: Synthesis of 1,3-Oxazin-6-ones. *Angew. Chem., Int. Ed.* **2013**, *52*, 14196–14199.
- (509) Liu, K.; Zou, M.; Lei, A. Aerobic Oxidative Carbonylation of Enamides by Merging Palladium with Photoredox Catalysis. *J. Org. Chem.* **2016**, *81*, 7088–7092.
- (510) Hesp, K. D.; Bergman, R. G.; Ellman, J. A. Expedient Synthesis of *N*-Acyl Anthranilamides and  $\beta$ -Enamine Amides by the Rh(III)-Catalyzed Amidation of Aryl and Vinyl C-H Bonds with Isocyanates. *J. Am. Chem. Soc.* **2011**, *133*, 11430–11433.
- (511) Shi, P.; Li, S.; Hu, L.-M.; Wang, C.; Loh, T.-P.; Hu, X.-H. Ruthenium-Catalyzed C-H Functionalization of Enamides with Isocyanates: Easy Entry to Pyrimidin-4-ones. *Chem. Commun.* **2019**, *55*, 11115–11118.
- (512) Wang, H.; Guo, L.-N.; Duan, X.-H. Decarboxylative Acylation of Cyclic Enamides with  $\alpha$ -Oxocarboxylic Acids by Palladium-Catalyzed C-H Activation at Room Temperature. *Org. Lett.* **2012**, *14*, 4358–4361.
- (513) Xiong, Z.; Liang, D.; Luo, S. Palladium-Catalyzed  $\beta$ -Selective C(sp<sup>2</sup>)-H Carboxamidation of Enamides by Isocyanide Insertion: Synthesis of *N*-Acyl Enamine Amides. *Org. Chem. Front.* **2017**, *4*, 1103–1106.
- (514) Ding, R.; Zhang, Q.-C.; Xu, Y.-H.; Loh, T.-P. Preparation of Highly Substituted ( $\beta$ -Acylamino)Acrylates via Iron-Catalyzed Alkoxy carbonylation of *N*-Vinylacetamides with Carbazates. *Chem. Commun.* **2014**, *50*, 11661–11664.
- (515) Shen, Z.-Y.; Cheng, J.-K.; Wang, C.; Yuan, C.; Loh, T.-P.; Hu, X.-H. Iron-Catalyzed Carbamoylation of Enamides with Formamides as a Direct Approach to *N*-Acyl Enamine Amides. *ACS Catal.* **2019**, *9*, 8128–8135.
- (516) Liu, R.-H.; Shen, Z.-Y.; Wang, C.; Loh, T.-P.; Hu, X.-H. Selective Dehydrogenative Acylation of Enamides with Aldehydes Leading to Valuable  $\beta$ -Ketoenamides. *Org. Lett.* **2020**, *22*, 944–949.
- (517) Zhao, K.; Zhang, X.-C.; Tao, J.-Y.; Wu, X.-D.; Wu, J.-X.; Li, W.-M.; Zhu, T.-H.; Loh, T.-P. Regio- and Stereoselective C(sp<sup>2</sup>)-H Acylation of Enamides with Aldehydes via Transition-Metal Free Photoredox Catalysis. *Green Chem.* **2020**, *22*, 5497–5503.
- (518) Jiang, H.; Chen, X.; Zhang, Y.; Yu, S. C-H Functionalization of Enamides: Synthesis of  $\beta$ -Amidovinyl Sulfones via Visible-Light Photoredox Catalysis. *Adv. Synth. Catal.* **2013**, *355*, 809–813.
- (519) Sun, D.; Zhang, R. Transition-Metal-Free, Visible-Light-Induced Oxidative Cross-Coupling for Constructing  $\beta$ -Acetylamino Acrylosulfones from Sodium Sulfonates and Enamides. *Org. Chem. Front.* **2018**, *5*, 92–97.
- (520) Kramer, P.; Krieg, S.-C.; Kelma, H.; Manolikakes, G. Manganese(III) Acetate-Mediated direct C(sp<sup>2</sup>)-H-Sulfonylation of Enamides with Sodium and Lithium Sulfonates. *Org. Biomol. Chem.* **2019**, *17*, 5538–5544.
- (521) Gu, Q.; Wang, X.; Liu, X.; Wu, G.; Xie, Y.; Shao, Y.; Zhao, Y.; Zeng, X. Electrochemical Sulfonylation of Enamides with Sodium Sulfonates to Access  $\beta$ -Amidovinyl Sulfones. *Org. Biomol. Chem.* **2021**, *19*, 8295–8300.
- (522) Qiu, G.; Zhou, K.; Gao, L.; Wu, J. Insertion of Sulfur Dioxide via Radical Process: An Efficient Route to Sulfonyl Compounds. *Org. Chem. Front.* **2018**, *5*, 691–705.
- (523) Zhu, T.-H.; Zhang, X.-C.; Zhao, K.; Loh, T.-P. Cu(OTf)<sub>2</sub>-Mediated C(sp<sup>2</sup>)-H Arylsulfonylation of Enamides via the Insertion of Sulfur Dioxide. *Org. Chem. Front.* **2019**, *6*, 94–98.
- (524) Zhu, T.-H.; Zhang, X.-C.; Cui, X.-L.; Zhang, Z.-Y.; Jiang, H.; Sun, S.-S.; Zhao, L.-L.; Zhao, K.; Loh, T.-P. Direct C(sp<sup>2</sup>)-H Arylsulfonylation of Enamides via Iridium(III)-Catalyzed Insertion of Sulfur Dioxide with Aryldiazonium Tetrafluoroborates. *Adv. Synth. Catal.* **2019**, *361*, 3593–3598.
- (525) Jia, X.-W.; Kramer, S.; Skrydstrup, T.; Lian, Z. Design and Applications of a SO<sub>2</sub> Surrogate in Palladium-Catalyzed Direct Aminosulfonylation between Aryl Iodides and Amines. *Angew. Chem., Int. Ed.* **2021**, *60*, 7353–7359.
- (526) Chen, L.; Zhou, M.; Shen, L.; He, X.; Li, X.; Zhang, X.; Lian, Z. Metal- and Base-Free C(sp<sup>2</sup>)-H Arylsulfonylation of Enamides for Synthesis of (*E*)- $\beta$ -Amidovinyl Sulfones via the Insertion of Sulfur Dioxide. *Org. Lett.* **2021**, *23*, 4991–4996.
- (527) Ye, X.; Peng, L.; Bao, X.; Tan, C.-H.; Wang, H. Recent Developments in Highly Efficient Construction of P-Stereogenic Centers. *Green Synth. Catal.* **2021**, *2*, 6–18.
- (528) Liu, Y.; Liu, Z.; Zhang, Y.-H.; Xiong, C. Manganese(III) Acetylacetonate-Mediated Phosphorylation of Enamides at Room Temperature. *Adv. Synth. Catal.* **2018**, *360*, 3492–3496.
- (529) Zhang, D.-L.; Li, C.-K.; Zeng, R.-S.; Shoberu, A.; Zou, J.-P. Manganese(III)-Mediated Selective Phosphorylation of Enamides: Direct Synthesis of  $\beta$ -Phosphoryl Enamides. *Org. Chem. Front.* **2019**, *6*, 236–240.
- (530) Xu, X.-M.; Li, W.; Li, Q.; Chen, S.; Zhang, X.; Yang, B.; Wang, W.-L. Manganese(III)-Promoted Highly Stereoselective Phosphorylation of Acyclic Tertiary Enamides to Synthesize *E*-Selective  $\beta$ -Phosphoryl Enamides. *Org. Biomol. Chem.* **2022**, *20*, 5566–5574.
- (531) Qiao, B.; Cao, H.-Q.; Huang, Y.-J.; Zhang, Y.; Nie, J.; Zhang, F.-G.; Ma, J.-A. Pd(II)-Catalyzed Phosphorylation of Enamido C(sp<sup>2</sup>)-H Bonds: A General Route to  $\beta$ -Amido-vinylphosphonates. *Chin. J. Chem.* **2018**, *36*, 809–814.
- (532) Pal, S.; Gaumont, A.-C.; Lakhdar, S.; Gillaizeau, I. Diphenyliodonium Ion/Et<sub>3</sub>N Promoted Csp<sup>2</sup>-H Radical Phosphorylation of Enamides. *Org. Lett.* **2019**, *21*, 5621–5625.
- (533) Santhini, P. V.; Nimisha, G.; John, J.; Suresh, E.; Varma, R. L.; Radhakrishnan, K. V. Pd-Catalyzed Oxidative Annulation of Enamides with Diazabicyclic Olefins: A Rapid Access to Cyclopentene Fused 2-Pyrrolines. *Chem. Commun.* **2017**, *53*, 1848–1851.
- (534) Li, X.; Zhang, J.; Zhang, F.; Luo, X.; Luo, H. Construction of Pyridine Ring Systems by Mn(OAc)<sub>2</sub>-Promoted Formal Dehydrative Dehydroaromatizing [4 + 2] Cycloaddition of Enamides with Maleimides. *Adv. Synth. Catal.* **2022**, *364*, 1683–1688.
- (535) Stuart, D. R.; Alsabeh, P.; Kuhn, M.; Fagnou, K. Rhodium(III)-Catalyzed Arene and Alkene C-H Bond Functionalization Leading to Indoles and Pyrroles. *J. Am. Chem. Soc.* **2010**, *132*, 18326–18339.
- (536) Li, B.; Wang, N.; Liang, Y.; Xu, S.; Wang, B. Ruthenium-Catalyzed Pyrrole Synthesis via Oxidative Annulation of Enamides and Alkynes. *Org. Lett.* **2013**, *15*, 136–139.
- (537) Wang, L.; Ackermann, L. Versatile Pyrrole Synthesis through Ruthenium(II)-Catalyzed Alkene C-H Bond Functionalization on Enamines. *Org. Lett.* **2013**, *15*, 176–179.
- (537) Zhao, M.-N.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. Pd-Catalyzed Oxidative Coupling of Enamides and Alkynes for Synthesis of Substituted Pyrroles. *Org. Lett.* **2014**, *16*, 608–611.
- (539) Xu, Y.-H.; He, T.; Zhang, Q.-C.; Loh, T.-P. Synthesis of Multi-Substituted Pyrroles Using Enamides and Alkynes Catalyzed by Pd(OAc)<sub>2</sub> with Molecular Oxygen as an Oxidant. *Chem. Commun.* **2014**, *50*, 2784–2786.
- (540) Zhao, M.-N.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. Copper-Promoted Oxidative Coupling of Enamides and Alkynes for the Synthesis of Substituted Pyrroles. *Chem. - Eur. J.* **2014**, *20*, 1839–1842.
- (541) Lade, D. M.; Pawar, A. B. Cp\*Co(III)-Catalyzed Vinylic C-H Bond Activation Under Mild Conditions: Expedient Pyrrole Synthesis via (3 + 2) Annulation of Enamides and Alkynes. *Org. Chem. Front.* **2016**, *3*, 836–840.
- (542) Yu, W.; Zhang, W.; Liu, Y.; Zhou, Y.; Liu, Z.; Zhang, Y. Cobalt(III)-Catalyzed Synthesis of Pyrroles from Enamides and Alkynes. *RSC Adv.* **2016**, *6*, 24768–24772.

- (543) Zhao, M.-N.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. Coupling of Enamides with Alkynes or Arynes for Synthesis of Substituted Pyridines and Isoquinolines via Amide Activation. *Chem. Commun.* **2012**, *48*, 8105–8107.
- (544) Wu, J.; Xu, W.; Yu, Z.-X.; Wang, J. Ruthenium-Catalyzed Formal Dehydrative [4 + 2] Cycloaddition of Enamides and Alkynes for the Synthesis of Highly Substituted Pyridines: Reaction Development and Mechanistic Study. *J. Am. Chem. Soc.* **2015**, *137*, 9489–9496.
- (545) Li, Y.; Cheng, K.; Lu, X.; Sun, J. A Facile and Efficient Approach to *N*-Protected- $\beta$ -Sulfinylenamines via C-Sulfinylation of Enamides and Enecarbamates. *Adv. Synth. Catal.* **2010**, *352*, 1876–1880.
- (546) Yang, L.; Wen, Q.; Xiao, F.-H.; Deng, G.-J. Silver-Mediated Oxidative Vinylic C–H Bond Sulfonylation of Enamides with Disulfides. *Org. Biomol. Chem.* **2014**, *12*, 9519–9523.
- (547) Yu, W.; Chen, J.; Gao, K.; Liu, Z.; Zhang, Y. Amide-Assisted Acetoxylation of Vinyl C(sp<sup>2</sup>)–H Bonds by Rhodium Catalysis. *Org. Lett.* **2014**, *16*, 4870–4873.
- (548) Chen, C.; Pan, Y.; Zhao, H.; Xu, X.; Xu, J.; Zhang, Z.; Xi, S.; Xu, L.; Li, H. A Versatile Rhodium(III) Catalyst for Direct Acyloxylation of Aryl and Alkenyl C–H Bonds with Carboxylic Acids. *Org. Chem. Front.* **2018**, *5*, 415–422.
- (549) Munoz, S. B.; Krishnamurti, V.; Barrio, P.; Mathew, T.; Prakash, G. K. S. Direct Difluorination-Hydroxylation, Trifluorination, and C(sp<sup>2</sup>)–H Fluorination of Enamides. *Org. Lett.* **2010**, *20*, 1042–1045.
- (550) Chang, X.-C.; Wang, Z.-L.; Zhao, M.; Yang, C.; Li, J.-J.; Ma, W.-W.; Xu, Y.-H. Synthesis of Functionalized Vinylsilanes via Metal-Free Dehydrogenative Silylation of Enamides. *Org. Lett.* **2020**, *22*, 1326–1330.
- (551) Gu, Q.; Wang, Q.; Dai, W.; Wang, X.; Ban, Y.; Liu, T.; Zhao, Y.; Zhang, Y.; Ling, Y.; Zeng, X. K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>-Mediated Regio- and Stereo-Selective Thiocyanation of Enamides with NH<sub>4</sub>SCN. *Org. Biomol. Chem.* **2021**, *19*, 2512–2516.
- (552) Zhou, S.; Yan, B.-W.; Fan, S.-X.; Tian, J.-S.; Loh, T.-P. Regioselective Formal [4 + 2] Cycloadditions of Enaminones with Diazocarbonyls through Rh<sup>III</sup>-Catalyzed C–H Bond Functionalization. *Org. Lett.* **2018**, *20*, 3975–3979.
- (553) Zhao, Y.; Zheng, Q.; Yu, C.; Liu, Z.; Wang, D.; You, J.; Gao, G. Rh(III)-Catalyzed Regioselective C–H [4+2] C-Annulation of Vinyl Enaminones with Alkynes to Form Polysubstituted Salicylaldehydes. *Org. Chem. Front.* **2018**, *5*, 2875–2879.
- (554) Qi, B.; Guo, S.; Zhang, W.; Yu, X.; Song, C.; Zhu, J. Rh(III)-Catalyzed Enaminone-Directed Alkenyl C–H Activation for the Synthesis of Salicylaldehydes. *Org. Lett.* **2018**, *20*, 3996–3999.
- (555) Liang, G.; Rong, J.; Sun, W.; Chen, G.; Jiang, Y.; Loh, T.-P. Synthesis of Polyaromatic Rings: Rh(III)-Catalyzed [5 + 1] Annulation of Enaminones with Vinyl Esters through C–H Bond Functionalization. *Org. Lett.* **2018**, *20*, 7326–7331.
- (556) Zhou, S.; Liu, D.-Y.; Wang, S.; Tian, J.-S.; Loh, T.-P. An Efficient Method for the Synthesis of 2-Pyridones via C–H Bond Functionalization. *Chem. Commun.* **2020**, *56*, 15020–15023.
- (557) Duan, J.; Xu, G.; Rong, B.; Yan, H.; Zhang, S.; Wu, Q.; Zhu, N.; Guo, K. Iron-Catalyzed [4 + 2] Annulation of  $\alpha,\beta$ -Unsaturated Ketoxime Acetates with Enaminones toward Functionalized Pyridines. *Green Synth. Catal.* **2021**, *2*, 237–240.
- (558) Reddy, B. V. S.; Reddy, M. R.; Rao, Y. G.; Yadav, J. S.; Sridhar, B. Cu(OTf)<sub>2</sub>-Catalyzed Synthesis of 2,3-Disubstituted Indoles and 2,4,5-Trisubstituted Pyrroles from  $\alpha$ -Diazoketones. *Org. Lett.* **2013**, *15*, 464–467.
- (559) Jiang, Y.; Khong, V. Z. Y.; Lourdasamy, E.; Park, C.-M. Synthesis of 2-Aminofurans and 2-Unsubstituted Furans via Carbenoid-Mediated [3 + 2] Cycloaddition. *Chem. Commun.* **2012**, *48*, 3133–3135.
- (560) Fu, L.; Liu, Y.; Wan, J.-P. Pd-Catalyzed Triple-Fold C(sp<sup>2</sup>)–H Activation with Enaminones and Alkenes for Pyrrole Synthesis via Hydrogen Evolution. *Org. Lett.* **2021**, *23*, 4363–4367.
- (561) Guo, H.; Tian, L.; Liu, Y.; Wan, J.-P. DMSO as a C<sub>1</sub> Source for [2 + 2 + 1] Pyrazole Ring Construction via Metal-Free Annulation with Enaminones and Hydrazines. *Org. Lett.* **2022**, *24*, 228–233.
- (562) Bernini, R.; Fabrizi, G.; Sferrazza, A.; Cacchi, S. Copper-Catalyzed C–C Bond Formation through C–H Functionalization: Synthesis of Multisubstituted Indoles from *N*-Aryl Enaminones. *Angew. Chem., Int. Ed.* **2009**, *48*, 8078–8081.
- (563) Liu, J.; Wei, W.; Zhao, T.; Liu, X.; Yu, W.; Chang, J. Iodine/Copper Iodide-Mediated C–H Functionalization: Synthesis of Imidazo[1,2-*a*]pyridines and Indoles from *N*-Aryl Enamines. *J. Org. Chem.* **2016**, *81*, 9326–9336.
- (564) Wan, J.-P.; Zhong, S.; Xie, L.; Cao, X.; Liu, Y.; Wei, L. KIO<sub>3</sub>-Catalyzed Aerobic Cross-Coupling Reactions of Enaminones and Thiophenols: Synthesis of Polyfunctionalized Alkenes by Metal-Free C–H Sulfonylation. *Org. Lett.* **2016**, *18*, 584–587.
- (565) Jiang, Y.; Liang, G.; Zhang, C.; Loh, T.-P. Palladium-Catalyzed C–S Bond Formation of Stable Enamines with Arene/Alkanethiols: Highly Regioselective Synthesis of  $\beta$ -Amino Sulfides. *Eur. J. Org. Chem.* **2016**, *2016*, 3326–3330.
- (566) Xiang, H.; Yang, C. A Facile and General Approach to 3-((Trifluoromethyl)thio)-4*H*-chromen-4-one. *Org. Lett.* **2014**, *16*, 5686–5689.
- (567) Lu, L.; Zhao, X.; Dessie, W.; Xia, X.; Duan, X.; He, J.; Wang, R.; Liu, Y.; Wu, C. Visible-Light-Promoted Trifluoromethylselenolation of *ortho*-Hydroxyarylenaminones. *Org. Biomol. Chem.* **2022**, *20*, 1754–1758.
- (568) Gao, Y.; Liu, Y.; Wan, J.-P. Visible Light-Induced Thiocyanation of Enaminone C–H Bond to Access Polyfunctionalized Alkenes and Thiocyanochromones. *J. Org. Chem.* **2019**, *84*, 2243–2251.
- (569) Duan, X.; Liu, X.; Guan, X.; Wang, L.; Liu, K.; Zhou, H.; Chen, X.; Li, H.; Wang, J. Solvent-Controlled Synthesis of Thiocyanated Enaminones and 2-Aminothiazoles from Enaminones, KSCN, and NBS. *J. Org. Chem.* **2019**, *84*, 12366–12376.
- (570) Yang, Z.; Hu, L.; Cao, T.; An, L.; Li, L.; Yang, T.; Zhou, C. PIDA-Mediated  $\alpha$ -C–H Functionalization of Enaminones: the Synthesis of Thiocyanoenaminones and Chromones in Water. *New J. Chem.* **2019**, *43*, 16441–16444.
- (571) Zhang, X.-Z.; Ge, D.-L.; Chen, S.-Y.; Yu, X.-Q. A Catalyst-Free Approach To 3-Thiocyanato-4*H*-chromen-4-ones. *RSC Adv.* **2016**, *6*, 66320–66323.
- (572) Wu, M.; Jiang, Y.; An, Z.; Qi, Z.; Yan, R. Iron-Catalyzed Synthesis of Substituted Thiazoles from Enamines and Elemental Sulfur through C–S Bond Formation. *Adv. Synth. Catal.* **2018**, *360*, 4236–4240.
- (573) Zhang, B.; Liu, D.; Sun, Y.; Zhang, Y.; Feng, J.; Fu, F. Preparation of Thiazole-2-thiones through TBPB-Promoted Oxidative Cascade Cyclization of Enaminones with Elemental Sulfur. *Org. Lett.* **2021**, *23*, 3076–3082.
- (574) Sorabad, G. S.; Maddani, M. R. Metal-Free, Green and Efficient Oxidative a Halogenation of Enaminones by Halo Acid and DMSO. *New J. Chem.* **2019**, *43*, 6563–6568.
- (575) Lin, Y.; Wan, J.-P.; Liu, Y. Synthesis of 3-Halochromones with Simple KX Halogen Sources Enabled by *in Situ* Halide Oxidation. *New J. Chem.* **2020**, *44*, 8120–8124.
- (576) Lin, Y.; Jin, J.; Wang, C.; Wan, J.-P.; Liu, Y. Electrochemical C–H Halogenations of Enaminones and Electron-Rich Arenes with Sodium Halide (NaX) as Halogen Source for the Synthesis of 3-Halochromones and Haloarenes. *J. Org. Chem.* **2021**, *86*, 12378–12385.
- (577) Wang, F.; Sun, W.; Wang, Y.; Jiang, Y.; Loh, T.-P. Highly Site-Selective Metal-Free C–H Acyloxylation of Stable Enamines. *Org. Lett.* **2018**, *20*, 1256–1260.

- (578) Guo, Y.; Xiang, Y.; Wei, L.; Wan, J.-P. Thermoinduced Free-Radical C-H Acyloxylation of Tertiary Enaminones: Catalyst-Free Synthesis of Acyloxyl Chromones and Enaminones. *Org. Lett.* **2018**, *20*, 3971–3974.
- (579) Xiang, H.; Zhao, Q.; Tang, Z.; Xiao, J.; Xia, P.; Wang, C.; Yang, C.; Chen, X.; Yang, H. Visible-Light-Driven, Radical-Triggered Tandem Cyclization of *o*-Hydroxyaryl Enaminones: Facile Access to 3-CF<sub>2</sub>/CF<sub>3</sub>-Containing Chromones. *Org. Lett.* **2017**, *19*, 146–149.
- (580) Gao, H.; Hu, B.; Dong, W.; Gao, X.; Jiang, L.; Xie, X.; Zhang, Z. Synthesis of 3-CF<sub>2</sub>-Containing Chromones via a Visible-Light-Induced Radical Cascade Reaction of *o*-Hydroxyaryl Enaminones. *ACS Omega* **2017**, *2*, 3168–3174.
- (581) Yu, Q.; Liu, Y.; Wan, J.-P. Transition Metal-Free Synthesis of 3-Trifluoromethyl Chromones via Tandem C-H Trifluoromethylation and Chromone Annulation of Enaminones. *Org. Chem. Front.* **2020**, *7*, 2770–2775.
- (582) Akram, M. O.; Bera, S.; Patil, N. T. A Facile Strategy for Accessing 3-Alkynylchromones through Gold-Catalyzed Alkynylation/Cyclization of *o*-Hydroxyarylenaminones. *Chem. Commun.* **2016**, *52*, 12306–12309.
- (583) Bagle, P. N.; Mane, M. V.; Sancheti, S. P.; Gade, A. B.; Shaikh, S. R.; Baik, M.-H.; Patil, N. T. Gold(I)-Catalyzed Hydroxy Group Assisted C(sp<sup>2</sup>)-H Alkylation of Enaminones with Diazo Compounds To Access 3-Alkyl Chromones. *Org. Lett.* **2019**, *21*, 335–339.
- (584) Mkrtchyan, S.; Iaroshenko, V. O. Visible-Light-Mediated Arylation of *ortho*-Hydroxyarylenaminones: Direct Access to Isoflavones. *Chem. Commun.* **2020**, *56*, 2606–2609.
- (585) Luo, T.; Wan, J.-P.; Liu, Y. Toward C<sub>2</sub>-Nitrogenated Chromones by Copper-Catalyzed  $\beta$ -C(sp<sup>2</sup>)-H *N*-Heteroarylation of Enaminones. *Org. Chem. Front.* **2020**, *7*, 1107–1112.
- (586) Ge, H.; Niphakis, M. J.; Georg, G. I. Palladium(II)-Catalyzed Direct Arylation of Enaminones Using Organotrifluoroborates. *J. Am. Chem. Soc.* **2008**, *130*, 3708–3709.
- (587) Kim, Y. W.; Niphakis, M. J.; Georg, G. I. Copper-Assisted Palladium(II)-Catalyzed Direct Arylation of Cyclic Enaminones with Arylboronic Acids. *J. Org. Chem.* **2012**, *77*, 9496–9503.
- (588) Kim, Y. W.; Georg, G. I. Boron-Heck Reaction of Cyclic Enaminones: Regioselective Direct Arylation via Oxidative Palladium(II) Catalysis. *Org. Lett.* **2014**, *16*, 1574–1577.
- (589) Bi, L.; Georg, G. I. Direct Hiyama Cross-Coupling of Enaminones With Triethoxy(aryl)silanes and Dimethylphenylsilanol. *Org. Lett.* **2011**, *13*, 5413–5415.
- (590) Yu, Y.-Y.; Bi, L.; Georg, G. I. Palladium-Catalyzed Direct C-H Arylation of Cyclic Enaminones with Aryl Iodides. *J. Org. Chem.* **2013**, *78*, 6163–6169.
- (591) Yu, Y.-Y.; Niphakis, M. J.; Georg, G. I. Palladium(II)-Catalyzed Dehydrogenative Alkenylation of Cyclic Enaminones via the Fujiwara-Moritani Reaction. *Org. Lett.* **2011**, *13*, 5932–5935.
- (592) Yu, Y.-Y.; Georg, G. I. Biomimetic Aerobic C-H Olefination of Cyclic Enaminones at Room Temperature: Development toward the Synthesis of 1,3,5-Trisubstituted Benzenes. *Adv. Synth. Catal.* **2014**, *356*, 1359–1369.
- (593) Stanovnik, B.; Svetec, J. Synthesis of Heterocycles from Alkyl 3-(Dimethylamino)propenoates and Related Enaminones. *Chem. Rev.* **2004**, *104*, 2433–2480.
- (594) Wan, J.-P.; Gao, Y. Domino Reactions Based on Combinatorial Bond Transformations in Electron-Deficient Tertiary Enamines. *Chem. Rec.* **2016**, *16*, 1164–1177.
- (595) Mukherjee, S.; Yang, J. W.; Hoffmann, S.; List, B. Asymmetric Enamine Catalysis. *Chem. Rev.* **2007**, *107*, 5471–5569.
- (596) Liu, X.; Cheng, R.; Zhao, F.; Zhang-Negrerie, D.; Du, Y.; Zhao, K. Direct  $\beta$ -Acyloxylation of Enamines via PhIO-Mediated Intermolecular Oxidative C-O Bond Formation and Its Application to the Synthesis of Oxazoles. *Org. Lett.* **2012**, *14*, 5480–5483.
- (597) Yuan, Y.; Hou, W.; Zhang-Negrerie, D.; Zhao, K.; Du, Y. Direct Oxidative Coupling of Enamines and Electron-Deficient Amines: TBAI/TBHP-Mediated Synthesis of Substituted Diaminoalkenes under Metal-Free Conditions. *Org. Lett.* **2014**, *16*, 5410–5413.
- (598) Zheng, C.; Wang, Y.; Fan, R. Iodine(III)-Mediated Oxidative Cross-Coupling of Enamines and Propargylamines under Metal-Free Conditions: An Alternative Way to Prepare Highly Substituted 3-Pyrrolines. *Org. Lett.* **2015**, *17*, 916–919.
- (599) Lei, Z.-Q.; Ye, J.-H.; Sun, J.; Shi, Z.-J. Direct Alkenyl C-H Functionalization of Cyclic Enamines with Carboxylic Acids via Rh Catalysis Assisted by Hydrogen Bonding. *Org. Chem. Front.* **2014**, *1*, 634–638.
- (600) Wang, Z.; Reinus, B. J.; Dong, G. Catalytic Intermolecular  $\beta$ -C-H Alkenylation of  $\alpha$ -Enamino-Ketones with Simple Alkynes. *Chem. Commun.* **2014**, *50*, 5230–5232.
- (601) Zhou, Y.; Wang, Y.; Song, Z.; Nakano, T.; Song, Q. Cu-Catalyzed C-N Bond Cleavage of 3-Aminoindazoles for the C-H Arylation of Enamines. *Org. Chem. Front.* **2020**, *7*, 25–29.
- (602) Yang, Y.-D.; Azuma, A.; Tokunaga, E.; Yamasaki, M.; Shiro, M.; Shibata, N. Trifluoromethanesulfonyl Hypervalent Iodonium Ylide for Copper-Catalyzed Trifluoromethylthiolation of Enamines, Indoles, and  $\beta$ -Keto Esters. *J. Am. Chem. Soc.* **2013**, *135*, 8782–8785.
- (603) Jiang, H.; Huang, W.; Yu, Y.; Yi, S.; Li, J.; Wu, W. Transition-Metal-Free Synthesis of  $\beta$ -Trifluoromethylated Enamines with Trifluoromethanesulfinate. *Chem. Commun.* **2017**, *53*, 7473–7476.
- (604) Sun, J.; Zhang-Negrerie, D.; Du, Y. Oxidative Coupling of Enamines and Disulfides via Tetrabutylammonium Iodide/*tert*-Butyl Hydroperoxide-Mediated Intermolecular Oxidative C(sp<sup>2</sup>)-S Bond Formation Under Transition Metal-Free Conditions. *Adv. Synth. Catal.* **2016**, *358*, 2035–2040.
- (605) Li, D.; Li, S.; Peng, C.; Lu, L.; Wang, S.; Wang, P.; Chen, Y.-H.; Cong, H.; Lei, A. Electrochemical Oxidative C-H/S-H Cross-Coupling Between Enamines and Thiophenols with H<sub>2</sub> Evolution. *Chem. Sci.* **2019**, *10*, 2791–2795.
- (606) Li, D.; Jia, J.; Zhao, X.; Zhang, Z.; Wang, H.; Li, S.; Xu, Z.; Xie, Z. Electrochemical Oxidation Cross Dehydrogenative Coupling of Enamines and Thiophenols for the Synthesis of Vinyl Sulfides. *ChemistrySelect* **2021**, *6*, 6460–6463.
- (607) Lei, T.; Liang, G.; Cheng, Y.-Y.; Chen, B.; Tung, C.-H.; Wu, L.-Z. Cobaloxime Catalysis for Enamine Phosphorylation with Hydrogen Evolution. *Org. Lett.* **2020**, *22*, 5385–5389.
- (608) Han, C.; Tian, X.; Zhang, H.; Rominger, F.; Hashmi, A. S. K. Tetrasubstituted 1,3-Enynes by Gold-Catalyzed Direct C(sp<sup>2</sup>)-H Alkynylation of Acceptor-Substituted Enamines. *Org. Lett.* **2021**, *23*, 4764–4768.
- (609) Gao, P.; Liu, J.; Wei, Y. Hypervalent Iodine(III)-Mediated Benzannulation of Enamines with Alkynes for the Synthesis of Polysubstituted Naphthalene Derivatives. *Org. Lett.* **2013**, *15*, 2872–2875.
- (610) Würtz, S.; Rakshit, S.; Neumann, J. J.; Dröge, T.; Glorius, F. Palladium-Catalyzed Oxidative Cyclization of *N*-Aryl Enamines: From Anilines to Indoles. *Angew. Chem., Int. Ed.* **2008**, *47*, 7230–7233.
- (611) Neumann, J. J.; Rakshit, S.; Dröge, T.; Würtz, S.; Glorius, F. Exploring the Oxidative Cyclization of Substituted *N*-Aryl Enamines: Pd-Catalyzed Formation of Indoles from Anilines. *Chem. - Eur. J.* **2011**, *17*, 7298–7303.
- (612) Zoller, J.; Fabry, D. C.; Ronge, M. A.; Rueping, M. Synthesis of Indoles Using Visible Light: Photoredox Catalysis for Palladium-Catalyzed C-H Activation. *Angew. Chem., Int. Ed.* **2014**, *53*, 13264–13268.
- (613) Lian, X.-L.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. Palladium-Catalyzed Oxidative Cyclization of Tertiary Enamines for Synthesis of 1,3,4-Trisubstituted Pyrroles and 1,3-Disubstituted

Indoles. *Org. Lett.* **2014**, *16*, 3360–3363.

(614) Nguyen, H. H.; Kurth, M. J. Microwave-Assisted Synthesis of 3-Nitroindoles from *N*-Aryl Enamines via Intramolecular Arene–Alkene Coupling. *Org. Lett.* **2013**, *15*, 362–365.

(615) Guan, Z.-H.; Yan, Z.-Y.; Ren, Z.-H.; Liu, X.-Y.; Liang, Y.-M. Preparation of Indoles via Iron Catalyzed Direct Oxidative Coupling. *Chem. Commun.* **2010**, *46*, 2823–2825.

(616) Drouhin, P.; Taylor, R. J. K. A Copper-Mediated Oxidative Coupling Route to 3*H*- and 1*H*-Indoles from *N*-Aryl-enamines. *Eur. J. Org. Chem.* **2015**, *2015*, 2333–2336.

(617) Hu, F.-Z.; Zhao, S.-H.; Chen, H.; Yu, S.-W.; Xu, X.-Y.; Yuan, Y.-C.; Zhang, X.-M. Facile Synthesis of 2,3-Disubstituted Indoles by NBS/CuCl Mediated Oxidative Cyclization of *N*-Aryl Enamines. *ChemistrySelect* **2017**, *2*, 1409–1412.

(618) Yu, W.; Du, Y.; Zhao, K. PIDA-Mediated Oxidative C–C Bond Formation: Novel Synthesis of Indoles from *N*-Aryl Enamines. *Org. Lett.* **2009**, *11*, 2417–2420.

(619) He, Z.; Liu, W.; Li, Z. I<sub>2</sub>-Catalyzed Indole Formation via Oxidative Cyclization of *N*-Aryl Enamines. *Chem. - Asian J.* **2011**, *6*, 1340–1343.

(620) Wu, C.-J.; Meng, Q.-Y.; Lei, T.; Zhong, J.-J.; Liu, W.-Q.; Zhao, L.-M.; Li, Z.-J.; Chen, B.; Tung, C.-H.; Wu, L.-Z. An Oxidant-Free Strategy for Indole Synthesis via Intramolecular C–C Bond Construction under Visible Light Irradiation: Cross-Coupling Hydrogen Evolution Reaction. *ACS Catal.* **2016**, *6*, 4635–4639.

(621) Tang, S.; Gao, X.; Lei, A. Electrocatalytic Intramolecular Oxidative Annulation of *N*-Aryl Enamines into Substituted Indoles Mediated by Iodides. *Chem. Commun.* **2017**, *53*, 3354–3356.

(622) Zhao, M.; Wang, F.; Li, X. Cross-Dehydrogenative Coupling between Enamino Esters and Ketones: Synthesis of Tetrasubstituted Pyrroles. *Org. Lett.* **2012**, *14*, 1412–1415.

(623) Zhang, X.-Y.; Yang, Z.-W.; Chen, Z.; Wang, J.; Yang, D.-L.; Shen, Z.; Hu, L.-L.; Xie, J.-W.; Zhang, J.; Cui, H.-L. Tandem Copper-Catalyzed Propargylation/Alkyne Azacyclization/Isomerization Reaction under Microwave Irradiation: Synthesis of Fully Substituted Pyrroles. *J. Org. Chem.* **2016**, *81*, 1778–1785.

(624) Wang, Y.; Jiang, C.-M.; Li, H.-L.; He, F.-S.; Luo, X.; Deng, W.-P. Regioselective Iodine-Catalyzed Construction of Polysubstituted Pyrroles from Allenes and Enamines. *J. Org. Chem.* **2016**, *81*, 8653–8658.

(625) Fu, L.; Wan, J.-P.; Zhou, L.; Liu, Y. Copper-Catalyzed C–H/*N*–H Annulation of Enaminones and Alkynyl Esters for Densely Substituted Pyrrole Synthesis. *Chem. Commun.* **2022**, *58*, 1808–1811.

(626) Gao, P.; Wang, J.; Bai, Z.-J.; Shen, L.; Yan, Y.-Y.; Yang, D.-S.; Fan, M.-J.; Guan, Z.-H. Synthesis of Polycarbonyl Pyrroles via K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>-Mediated Oxidative Cyclization of Enamines. *Org. Lett.* **2016**, *18*, 6074–6077.

(627) Zhao, M.-N.; Zhang, Z.-J.; Ren, Z.-H.; Yang, D.-S.; Guan, Z.-H. Copper-Catalyzed Oxidative Cyclization/1,2-Amino Migration Cascade Reaction. *Org. Lett.* **2018**, *20*, 3088–3091.

(628) Chen, Z.-W.; Zheng, L.; Liu, J. Divergent Synthesis of Multisubstituted Unsymmetric Pyrroles and Pyrrolin-4-ones from Enamino Esters via Copper-Catalyzed Aerobic Dimerization. *Eur. J. Org. Chem.* **2019**, *2019*, 3051–3060.

(629) Chen, Z.; Shi, G.; Tang, W.; Sun, J.; Wang, W. Electrochemical Oxidative Cyclization: Synthesis of Polysubstituted Pyrrole from Enamines. *Eur. J. Org. Chem.* **2021**, *2021*, 951–955.

(630) Baidya, M.; Maiti, D.; Roy, L.; Sarkar, S. D. Trifluoroethanol as a Unique Additive for the Chemoselective Electrooxidation of Enamines to Access Unsymmetrically Substituted NH-Pyrroles. *Angew. Chem., Int. Ed.* **2022**, *61*, doi.org/10.1002/anie.202111679.

(631) Li, Y.; Xu, H.; Xing, M.; Huang, F.; Jia, J.; Gao, J. Iodine-Promoted Construction of Polysubstituted 2,3-Dihydropyrroles from Chalcones and  $\beta$ -Enamine Ketones (Esters). *Org. Lett.* **2015**, *17*, 3690–3693.

(632) Neumann, J. J.; Suri, M.; Glorius, F. Efficient Synthesis of Pyrroles: Oxidative C–C/*N*–*N* Bond-Formation Cascade. *Angew. Chem., Int. Ed.* **2010**, *49*, 7790–7794.

(633) Zhang, Q.; Luo, P.; Lin, Y.; Chen, W.; Liu, X.; Zhang, Y.; Ding, Q. Palladium-Catalyzed Intermolecular Oxidative Cyclization of *N*-Aryl Enamines with Isocyanides through Double sp<sup>2</sup> C–H Bonds Cleavage: Facile Synthesis of 4-Aminoquinoline Derivatives. *Org. Biomol. Chem.* **2015**, *13*, 4657–4660.

(634) Borah, B.; Chowhan, L. R.; Recent Advances in the Transition-Metal-Free Synthesis of Quinoxalines, *RSC Adv.* **2021**, *11*, 37325–37353.

(635) Yashwantrao, G.; Saha, S. Recent Advances in the Synthesis and Reactivity of Quinoxaline. *Org. Chem. Front.* **2021**, *8*, 2820–2862.

(636) Maikhuri, V. K.; Prasad, A. K.; Jha, A.; Srivastava, S. Recent Advances in the Transition Metal Catalyzed Synthesis of Quinoxalines: A Review. *New J. Chem.* **2021**, *45*, 13214–13246.

(637) Yang, Z.-J.; Liu, C.-Z.; Hu, B.-L.; Deng, C.-L.; Zhang, X.-G. Oxidative Tandem Nitrosation/Cyclization of *N*-Aryl Enamines with Nitromethane toward 3-(Trifluoromethyl)quinoxalines. *Chem. Commun.* **2014**, *50*, 14554–14557.

(639) Jiao, Y.-X.; Wei, L.-S.; Zhao, C.-Y.; Wei, K.; Mo, D.-L.; Pan, C.-X.; Su, G.-F. Isobutyl Nitrite-Mediated Synthesis of Quinoxalines through Double C–H Bond Amination of *N*-Aryl Enamines. *Adv. Synth. Catal.* **2018**, *360*, 4446–4451.

(639) List, B. Enamine Catalysis Is a Powerful Strategy for the Catalytic Generation and Use of Carbanion Equivalents. *Acc. Chem. Res.* **2004**, *37*, 548–557.

(640) Mukherjee, S.; Yang, J. W.; Hoffmann, S.; List, B. Asymmetric Enamine Catalysis. *Chem. Rev.* **2007**, *107*, 5471–5569.

(641) Allen, A. E.; MacMillan, D. W. C. Synergistic Catalysis: A Powerful Synthetic Strategy for New Reaction Development. *Chem. Sci.* **2012**, *3*, 633–658.

(642) Du, Z.; Shao, Z. Combining Transition Metal Catalysis and Organocatalysis—An Update. *Chem. Soc. Rev.* **2013**, *42*, 1337–1378.

(643) Afewerki, S.; Córdova, A. Combinations of Aminocatalysts and Metal Catalysts: A Powerful Cooperative Approach in Selective Organic Synthesis. *Chem. Rev.* **2016**, *116*, 13512–13570.

(644) Rago, A. J.; Dong, G. Synthesis of Indoles, Indolines, and Carbazoles via Palladium-Catalyzed C–H Activation. *Green Synth. Catal.* **2021**, *2*, 216–227.

(645) Lim, H. N.; Xing, D.; Dong, G. Transition-Metal-Catalyzed Ketone  $\alpha$ -Alkylation and Alkenylation with Simple Alkenes and Alkynes through a Dual Activation Strategy. *Synlett* **2019**, *30*, 674–684.

(646) Wang, Z.; Reinus, B. J.; Dong, G. Catalytic Intermolecular C-Alkylation of 1,2-Diketones with Simple Olefins: A Recyclable Directing Group Strategy. *J. Am. Chem. Soc.* **2012**, *134*, 13954–13957.

(647) Mo, F.; Dong, G. Regioselective Ketone  $\alpha$ -Alkylation with Simple Olefins via Dual Activation. *Science* **2014**, *345*, 68–72.

(648) Ibrahim, I.; Córdova, A. Direct Catalytic Intermolecular  $\alpha$ -Allylic Alkylation of Aldehydes by Combination of Transition-Metal and Organocatalysis. *Angew. Chem., Int. Ed.* **2006**, *45*, 1952–1956.

(649) Afewerki, S.; Ibrahim, I.; Rydfjord, J.; Breistein, P.; Córdova, A. Direct Regiospecific and Highly Enantioselective Intermolecular  $\alpha$ -Allylic Alkylation of Aldehydes by a Combination of Transition-Metal and Chiral Amine Catalysts. *Chem. - Eur. J.* **2012**, *18*, 2972–2977.

(650) Tang, S.; Wu, X.; Liao, W.; Liu, K.; Liu, C.; Luo, S.; Lei, A. Synergistic Pd/Enamine Catalysis: A Strategy for the C–H/C–H Oxidative Coupling of Allylarenes with Unactivated Ketones. *Org. Lett.* **2014**, *16*, 3584–3587.

(651) Lim, H. N.; Dong, G. Catalytic Intramolecular Ketone Alkylation with Olefins by Dual Activation. *Angew. Chem., Int. Ed.* **2015**, *54*, 15294–15298.

(652) Mo, F.; Lim, H. N.; Dong, G. Bifunctional Ligand-Assisted

- Catalytic Ketone  $\alpha$ -Alkenylation with Internal Alkynes: Controlled Synthesis of Enones and Mechanistic Studies. *J. Am. Chem. Soc.* **2015**, *137*, 15518–15527.
- (653) Xu, Y.; Su, T.; Huang, Z.; Dong, G. Practical Direct  $\alpha$ -Arylation of Cyclopentanones by Palladium/Enamine Cooperative Catalysis. *Angew. Chem., Int. Ed.* **2016**, *55*, 2559–2563.
- (654) Liu, R.-H.; Li, B.-L.; Lu, J.; Shen, C.; Gao, J.-R.; Jia, Y.-X. Palladium/L-Proline-Catalyzed Enantioselective  $\alpha$ -Arylative Desymmetrization of Cyclohexanones. *J. Am. Chem. Soc.* **2016**, *138*, 5198–5201.
- (655) Hatamoto, Y.; Sakaguchi, S.; Ishii, Y. Oxidative Cross-Coupling of Acrylates with Vinyl Carboxylates Catalyzed by a Pd(OAc)<sub>2</sub>/HPMoV/O<sub>2</sub> System. *Org. Lett.* **2004**, *6*, 4623–4625.
- (656) Qiu, S.-Q.; Xu, Y.-H.; Loh, T.-P. Palladium-Catalyzed Intermolecular Oxyarylation of Vinylacetates with Retention of an Alkenyl Moiety. *Org. Lett.* **2015**, *17*, 3462–3465.
- (657) Gattinoni, S.; De Simone, C.; Dallavalle, S.; Fezza, F.; Nannei, R.; Amadio, D.; Minetti, P.; Quattrociochi, G.; Caprioli, A.; Borsini, F.; Cabri, W.; Penco, S.; Merlini, L.; Maccarrone, M. Enol Carbamates as Inhibitors of Fatty Acid Amide Hydrolase (FAAH) Endowed with High Selectivity for FAAH over the Other Targets of the Endocannabinoid System. *ChemMedChem* **2010**, *5*, 357–360.
- (658) Gong, T.-J.; Su, W.; Liu, Z.-J.; Chen, W.-M.; Xiao, B.; Fu, Y. Rh(III)-Catalyzed C–H Activation with Allenes To Synthesize Conjugated Olefins. *Org. Lett.* **2014**, *16*, 330–333.
- (659) Bouladakis-Arapinis, M.; Hopkinson, M. N.; Glorius, F. Using Rh(III)-Catalyzed C–H Activation as a Tool for the Selective Functionalization of Ketone-Containing Molecules. *Org. Lett.* **2014**, *16*, 1630–1633.
- (660) Li, T.; Zhang, J.; Yu, C.; Lu, X.; Xu, L.; Zhong, G. Ruthenium-Catalyzed Olefinic C–H Alkenylation of Enol-Carbamates: Highly Stereo-Selective Synthesis of (Z,Z) and (Z,E)-Butadienes. *Chem. Commun.* **2017**, *53*, 12926–12929.
- (661) Sharma, S.; Han, S. H.; Oh, Y.; Mishra, N. K.; Han, S.; Kwak, J. H.; Lee, S.-Y.; Jung, Y. H.; Kim, I. S. Mild and Site-Selective Allylation of Enol Carbamates with Allylic Carbonates under Rhodium Catalysis. *J. Org. Chem.* **2016**, *81*, 2243–2251.
- (662) Sharma, S.; Han, S. H.; Jo, H.; Han, S.; Mishra, N. K.; Choi, M.; Jeong, T.; Park, J.; Kim, I. S. Rhodium-Catalyzed Vinylic C–H Functionalization of Enol Carbamates with Maleimides. *Eur. J. Org. Chem.* **2016**, *2016*, 3611–3618.
- (663) Liang, Y.-R.; Si, X.-J.; Zhang, H.; Yang, D.; Niu, J.-L.; Song, M.-P. Thiocarbamate-Directed Cp\*Co(III)-Catalyzed Olefinic C–H Amidation: Facile Access to Enamines with High (Z)-Selectivity. *Eur. J. Org. Chem.* **2021**, *2021*, 694–700.
- (664) Zhou, C.-N.; Zheng, Z.-A.; Chang, G.; Xiao, Y.-C.; Shen, Y.-H.; Li, G.; Zhang, Y.-M.; Peng, W.-M.; Wang, L.; Xiao, B. Phosphorus-Containing Groups Assisted Transition Metal Catalyzed C–H Activation Reactions. *Curr. Org. Chem.* **2019**, *23*, 103–135.
- (665) Hu, X.-H.; Yang, X.-F.; Loh, T.-P. Selective Alkenylation and Hydroalkenylation of Enol Phosphates through Direct C–H Functionalization. *Angew. Chem., Int. Ed.* **2015**, *54*, 15535–15539.
- (666) Chen, D. Y.-K.; Pouwer, R. H.; Richard, J.-A. Recent Advances in the Total Synthesis of Cyclopropane-Containing Natural Products. *Chem. Soc. Rev.* **2012**, *41*, 4631–4642.
- (667) Ebner, C.; Carreira, E. M. Cyclopropanation Strategies in Recent Total Syntheses. *Chem. Rev.* **2017**, *117*, 11651–11679.
- (668) Piou, T.; Rovis, T. Rh(III)-Catalyzed Cyclopropanation Initiated by C–H Activation: Ligand Development Enables a Diastereoselective [2 + 1] Annulation of N-Enoxyphthalimides and Alkenes. *J. Am. Chem. Soc.* **2014**, *136*, 11292–11295.
- (669) Piou, T.; Romanov-Michaëlidis, F.; Ashley, M.; A.; Romanova-Michaëlidis, M.; Rovis, T. Stereodivergent Rhodium(III)-Catalyzed *cis*-Cyclopropanation Enabled by Multivariate Optimization. *J. Am. Chem. Soc.* **2018**, *140*, 9587–9593.
- (670) Phipps, E. J. T.; Rovis, T. Rh(III)-Catalyzed C–H Activation-Initiated Directed Cyclopropanation of Allylic Alcohols. *J. Am. Chem. Soc.* **2019**, *141*, 6807–6811.
- (671) Phipps, E. J. T.; Piou, T.; Rovis, T. Rhodium(III)-Catalyzed Cyclopropanation of Unactivated Olefins Initiated by C–H Activation. *Synlett* **2019**, *30*, 1787–1790.
- (672) Duchemin, C.; Cramer, N. Chiral Cyclopentadienyl Rh<sup>III</sup>-Catalyzed Enantioselective Cyclopropanation of Electron-Deficient Olefins Enable Rapid Access to UPF-648 and Oxylinin Natural Products. *Chem. Sci.* **2019**, *10*, 2773–2777.
- (673) Lahtigui, O.; Forster, D.; Duchemin, C.; Cramer, N. Enantioselective Access to 3-Azabicyclo[3.1.0]hexanes by Cp\*Rh<sup>III</sup> Catalyzed C–H Activation and Cp\*Ir<sup>III</sup> Transfer Hydrogenation. *ACS Catal.* **2022**, *12*, 6209–6215.
- (674) Piou, T.; Rovis, T. Rhodium-Catalysed *syn*-Carboamination of Alkenes via a Transient Directing Group. *Nature* **2015**, *527*, 86–90.
- (675) Duchemin, C.; Cramer, N. Enantioselective Cp\*Rh<sup>III</sup>-Catalyzed Carboaminations of Acrylates. *Angew. Chem., Int. Ed.* **2020**, *59*, 14129–14133.
- (676) Pan, L.; Bi, X.; Liu, Q. Recent Developments of Ketene Dithioacetal Chemistry. *Chem. Soc. Rev.* **2013**, *42*, 1251–1286.
- (677) Yu, H.; Jin, W.; Sun, C.; Chen, J.; Du, W.; He, S.; Yu, Z. Palladium-Catalyzed Cross-Coupling of Internal Alkenes with Terminal Alkenes to Functionalized 1,3-Butadienes Using C–H Bond Activation: Efficient Synthesis of Bicyclic Pyridones. *Angew. Chem., Int. Ed.* **2010**, *49*, 5792–5797.
- (678) Yang, X.; Liu, Z.; Sun, C.; Chen, J.; Yu, Z. Palladium-Catalyzed Oxidative Cross-Coupling of  $\alpha$ -Cyanoketene Dithioacetals with Olefins. *Chem. - Eur. J.* **2015**, *21*, 14085–14094.
- (679) Xu, Q.; Zhang, B.; Zhou, X.; Pan, L.; Liu, Q.; Li, Y. Photoinduced C(sp<sup>2</sup>)-H/C(sp<sup>2</sup>)-H Cross-Coupling of Alkenes: Direct Synthesis of 1,3-Dienes. *Org. Lett.* **2020**, *22*, 1692–1697.
- (680) Jin, W.; Yang, Q.; Wu, P.; Chen, J.; Yu, Z. Palladium-Catalyzed Oxidative Heck-Type Allylation of  $\beta,\beta$ -Disubstituted Enones with Allyl Carbonates. *Adv. Synth. Catal.* **2014**, *356*, 2097–2102.
- (681) Xu, C.; Liu, J.; Ming, W.; Liu, Y.; Liu, J.; Wang, M.; Liu, Q. In Situ Generation of PhI<sup>+</sup>CF<sub>3</sub> and Transition-Metal-Free Oxidative sp<sup>2</sup> C–H Trifluoromethylation. *Chem. - Eur. J.* **2013**, *19*, 9104–9109.
- (682) Mao, Z.; Huang, F.; Yu, H.; Chen, J.; Yu, Z.; Xu, Z. Copper-Catalyzed Trifluoromethylation of Internal Olefinic C–H Bonds: Efficient Routes to Trifluoromethylated Tetrasubstituted Olefins and N-Heterocycles. *Chem. - Eur. J.* **2014**, *20*, 3439–3445.
- (683) Gou, B.; Yang, C.; Zhang, L.; Xia, W. Visible-Light Induced Trifluoromethylation of Internal Olefinic C–H Bonds through Photoredox Catalysis. *Acta Chim. Sinica*, **2017**, *75*, 66–69.
- (684) Gu, Q.; Cheng, Z.; Zeng, X. Electrochemical Oxidative Trifluoromethylation of  $\alpha$ -Oxoketene Ketene Dithioacetals with CF<sub>3</sub>SO<sub>2</sub>Na. *Chin. J. Org. Chem.* **2022**, *42*, 1537–1544.
- (685) Tian, S.; Song, X.; Zhu, D.; Wang, M. Alternative Palladium-Catalyzed Vinylic C–H Difluoroalkylation of Ketene Dithioacetals Using Bromodifluoroacetate Derivatives. *Adv. Synth. Catal.* **2018**, *360*, 1414–1419.
- (686) Enthaler, S.; Company, A. Palladium-Catalysed Hydroxylation and Alkoxylation. *Chem. Soc. Rev.* **2011**, *40*, 4912–4924.
- (687) Liu, B.; Shi, B.-F. Transition-Metal-Catalyzed Etherification of Unactivated C–H Bonds. *Tetrahedron Lett.* **2015**, *56*, 15–22.
- (688) Moghimi, S.; Mahdavi, M.; Shafiee, A.; Foroumadi, A. Transition-Metal-Catalyzed Acyloxylation: Activation of C(sp<sup>2</sup>)-H and C(sp<sup>3</sup>)-H Bonds. *Eur. J. Org. Chem.* **2016**, *2016*, 3282–3299.
- (689) Liang, D.; Wang, M.; Dong, Y.; Guo, Y.; Liu, Q. Palladium-Catalyzed Oxidative C–O Cross-Coupling of Ketene Dithioacetals and Carboxylic Acids. *RSC Adv.* **2014**, *4*, 6564–6567.

- (690) Liu, Z.; Huang, F.; Lou, J.; Wang, Q.; Yu, Z. Copper-Promoted Direct C–H Alkoxylation of S,S-Functionalized Internal Olefins with Alcohols. *Org. Biomol. Chem.* **2017**, *15*, 5535–5540.
- (691) Shukla, G.; Srivastava, A.; Nagaraju, A.; Raghuvanshi, K.; Singh, M. S. Iodine-Mediated Copper-Catalyzed Efficient  $\alpha$ -C(sp<sup>2</sup>)-Thiomethylation of  $\alpha$ -Oxoketene Dithioacetals with Dimethyl Sulfoxide in One Pot. *Adv. Synth. Catal.* **2015**, *357*, 3969–3976.
- (692) Deng, L.; Liu, Y. Metal-Free Vinyl C–H Sulfenylation/Alkyl Thiolation of Ketene Dithioacetals for the Synthesis of Polythiolated Alkenes. *ACS Omega* **2018**, *3*, 11890–11895.
- (693) Ge, L.; Chiou, M.-F.; Li, Y.; Bao, H. Radical Azidation as a Means of Constructing C(sp<sup>3</sup>)-N<sub>3</sub> Bonds. *Green Synth. Catal.* **2020**, *1*, 86–102.
- (694) Chen, Q.; Lei, Y.; Wang, Y.; Wang, C.; Wang, Y.; Xu, Z.; Wang, H.; Wang, R. Direct Thiocyanation of Ketene Dithioacetals under Transition-Metal-Free Conditions. *Org. Chem. Front.* **2017**, *4*, 369–372.
- (695) Zhu, L.; Yu, H.; Guo, Q.; Chen, Q.; Xu, Z.; Wang, R. C–H Bonds Phosphorylation of Ketene Dithioacetals. *Org. Lett.* **2015**, *17*, 1978–1981.
- (696) Yang, Q.; Wu, P.; Chen, J.; Yu, Z. Iron-Catalyzed Alkylation of  $\alpha$ -Oxo Ketene Dithioacetals. *Chem. Commun.* **2014**, *50*, 6337–6339.
- (697) Wang, Q.; Lou, J.; Wu, P.; Wu, K.; Yu, Z. Iron-Mediated Oxidative C–H Alkylation of S,S-Functionalized Internal Olefins via C(sp<sup>2</sup>)-H/C(sp<sup>3</sup>)-H Cross-Coupling. *Adv. Synth. Catal.* **2017**, *359*, 2981–2998.
- (698) Wen, J.; Zhang, F.; Shi, W.; Lei, A. Metal-Free Direct Alkylation of Ketene Dithioacetals via Oxidative C(sp<sup>2</sup>)-H/C(sp<sup>3</sup>)-H Cross-Coupling. *Chem. - Eur. J.* **2017**, *23*, 8814–8817.
- (699) Lou, J.; He, Y.; Li, Y.; Yu, Z. Transition-Metal-Promoted Direct C–H Cyanoalkylation and Cyanoalkoxylation of Internal Alkenes via Radical C–C Bond Cleavage of Cycloketone Oxime Esters. *Adv. Synth. Catal.* **2019**, *361*, 3787–3799.
- (700) Wang, Q.; Yang, X.; Wu, P.; Yu, Z. Photoredox-Catalyzed C–H Arylation of Internal Alkenes to Tetrasubstituted Alkenes: Synthesis of Tamoxifen. *Org. Lett.* **2017**, *19*, 6248–6251.
- (701) Wang, M.; Kong, L.; Wu, Q.; Li, X. Ruthenium- and Rhodium-Catalyzed Chemodivergent Couplings of Ketene Dithioacetals and  $\alpha$ -Diazo Ketones via C–H Activation/Functionalization. *Org. Lett.* **2018**, *20*, 4597–4600.
- (702) Wang, M.; Bai, D.; Kong, L.; Liu, B.; Li, X. Ag(I)-Catalyzed Nucleophilic Addition and Friedel–Crafts Alkylation between  $\alpha$ -Oxoketene Dithioacetals and Propargyl Carbonates. *Org. Lett.* **2018**, *20*, 7775–7778.
- (703) Lou, J.; Han, W.; Liu, Z.; Xiao, J. Rhodium-Catalyzed Enone Carbonyl Directed C–H Activation for the Synthesis of Indanones Containing All-Carbon Quaternary Centers. *Org. Chem. Front.* **2021**, *8*, 1447–1453.
- (704) Bai, Y.; Zeng, J.; Cai, S.; Liu, X.-W. Palladium-Catalyzed Direct Cross-Coupling Reaction of Glycals with Activated Alkenes. *Org. Lett.* **2011**, *13*, 4394–4397.
- (705) Zhao, C.; Toste, F. D.; Bergman, R. G. Direct Michael Addition of Alkenes via a Cobalt-Dinitrosyl Mediated Vinylic C–H Functionalization Reaction. *J. Am. Chem. Soc.* **2011**, *133*, 10787–10789.
- (706) Li, S.-S.; Xia, Y.-Q.; Liu, C.-F.; Zhang, G.-T.; Su, F.; Zhang, X.-M.; Dong, L. Diverse Reactivity in a Rhodium(III)-Catalyzed Vinylic sp<sup>2</sup> C–H Bond Functionalization: Synthesis of Fused Polycyclic Heteroarenes or Conjugated Dienes. *Adv. Synth. Catal.* **2016**, *358*, 3724–3729.
- (707) Kuamr, D.; Vemula, S. R.; Cook, G. R. Merging C–H Bond Functionalization with Amide Alcoholysis: En Route to 2-Aminopyridines. *ACS Catal.* **2016**, *6*, 3531–3536.
- (708) Yang, F.; Yu, J.; Liu, Y.; Zhu, J. Rhodium(III)-Catalyzed Oxadiazole-Directed Alkenyl C–H Activation for Synthetic Access to 2-Acylamino and 2-Amino Pyridines. *J. Org. Chem.* **2017**, *82*, 9978–9987.
- (709) Yu, X.; Chen, K.; Wang, Q.; Guo, S.; Zha, S.; Zhu, J. Associative Covalent Relay: An Oxadiazolone Strategy for Rhodium(III)-Catalyzed Synthesis of Primary Pyridinylamines. *Angew. Chem., Int. Ed.* **2017**, *56*, 5222–5226.
- (710) Li, C.; Qiang, X.-Y.; Qi, Z.-C.; Cao, B.; Li, J.-Y.; Yang, S.-D. Pd-Catalyzed Heck-Type Reaction: Synthesizing Highly Diastereoselective and Multiple Aryl-Substituted P-Ligands. *Org. Lett.* **2019**, *21*, 7138–7142.
- (711) Nie, B.; Wu, W.; Ren, Q.; Wang, Z.; Zhang, J.; Zhang, Y.; Jiang, H. Access to Cycloalkeno[c]-Fused Pyridines via Pd-Catalyzed C(sp<sup>2</sup>)-H Activation and Cyclization of *N*-Acetyl Hydrazones of Acylcycloalkenes with Vinyl Azides. *Org. Lett.* **2020**, *22*, 7786–7790.
- (712) Kumar, S. V.; Banerjee, S.; Punniamurthy, T. Transition Metal-Catalyzed Coupling of Heterocyclic Alkenes via C–H Functionalization: Recent Trends and Applications. *Org. Chem. Front.* **2020**, *7*, 1527–1569.
- (713) Yu, Y.-Y.; Georg, G. I. Dehydrogenative Alkenylation of Uracils via Palladium-Catalyzed Regioselective C–H Activation. *Chem. Commun.* **2013**, *49*, 3694–3696.
- (714) Min, M.; Kim, Y.; Hong, S. Regioselective Palladium-Catalyzed Olefination of Coumarins via Aerobic Oxidative Heck Reactions. *Chem. Commun.* **2013**, *49*, 196–198.
- (715) Azpiroz, R.; Rubio-Perez, L.; Di Giuseppe, A.; Passarelli, V.; Lahoz, F. J.; Castarlenas, R.; Pérez-Torrente, J. J.; Oro, L. A. Rhodium(I)-*N*-Heterocyclic Carbene Catalyst for Selective Coupling of *N*-Vinylpyrazoles with Alkynes via C–H Activation. *ACS Catal.* **2014**, *4*, 4244–4253.
- (716) Li, J.; Ackermann, L. Cobalt(III)-Catalyzed Aryl and Alkenyl C–H Aminocarbonylation with Isocyanates and Acyl Azides. *Angew. Chem., Int. Ed.* **2015**, *54*, 8551–8554.
- (717) Boerth, J. A.; Hummel, J. R.; Ellman, J. A. Highly Stereoselective Cobalt(III)-Catalyzed Three-Component C–H Bond Addition Cascade. *Angew. Chem., Int. Ed.* **2016**, *55*, 12650–12654.
- (718) Saitou, T.; Jin, Y.; Isobe, K.; Suga, T.; Takaya, J.; Iwasawa, N. Rh-Catalyzed Direct Carboxylation of Alkenyl C–H Bonds of Alkenylpyrazoles. *Chem. - Asian J.* **2020**, *15*, 1941–1944.
- (719) Yang, L.; Steinbock, R.; Scheremetjew, A.; Kuniyil, R.; Finfer, L. H.; Messinis, A. M.; Ackermann, L. Azaruthena(II)-bicyclo[3.2.0]heptadiene: Key Intermediate for Ruthenaelectro(II/III/I)-Catalyzed Alkyne Annulations. *Angew. Chem., Int. Ed.* **2020**, *59*, 11130–11135.
- (720) Messaoudi, S.; Tréguier, B.; Hamze, A.; Provot, O.; Peyrat, J.-F.; De Losada, J. R.; Liu, J.-M.; Bignon, J.; Wdziejczak-Bakala, J.; Thoret, S.; Dubois, J.; Brion, J.-D.; Alami, M. Isocombretastatins A versus Combretastatins A: The Forgotten isoCA-4 Isomer as a Highly Promising Cytotoxic and Antitubulin Agent. *J. Med. Chem.* **2009**, *52*, 4538–4542.
- (721) Leriche, C.; He, X.; Chang, C.-W. T.; Liu, H.-W. Reversal of the Apparent Regiospecificity of NAD(P)H-Dependent Hydride Transfer: The Properties of the Difluoromethylene Group, A Carbonyl Mimic. *J. Am. Chem. Soc.* **2003**, *125*, 6348–6349.
- (722) Magueur, G.; Crousse, B.; Ourévitche, M.; Bonnet-Delpon, D.; Bégué, J.-P. Fluoro-Artemisinins: When a gem-Difluoroethylene Replaces a Carbonyl Group. *J. Fluor. Chem.* **2006**, *127*, 637–642.
- (723) Couve-Bonnaire, S.; Cahard, D.; Pannecoucke, X. Chiral Dipeptide Mimics Possessing a Fluoroolefin Moiety: A Relevant Tool for Conformational and Medicinal Studies. *Org. Biomol. Chem.* **2007**, *5*, 1151–1157.
- (724) Meanwell, N. A. Synopsis of Some Recent Tactical Application of Bioisosteres in Drug Design. *J. Med. Chem.* **2011**, *54*, 2529–2591.

- (725) Yanai, H.; Taguchi, T. Synthetic Methods for Fluorinated Olefins. *Eur. J. Org. Chem.* **2011**, *2011*, 5939–5954.
- (726) Meanwell, N. A. Fluorine and Fluorinated Motifs in the Design and Application of Bioisosteres for Drug Design. *J. Med. Chem.* **2018**, *61*, 5822–5880.
- (727) Drouin, M.; Laxio Arenas, J.; Paquin, J.-F. Incorporating a Monofluoroalkene into the Backbones of Short Peptides: Evaluating the Impact on Local Hydrophobicity. *ChemBioChem* **2019**, *20*, 1817–1826.
- (728) Landelle, G.; Bergeron, M.; Turcotte-Savard, M. O.; Paquin, J. F. Synthetic Approaches to Monofluoroalkenes. *Chem. Soc. Rev.* **2011**, *40*, 2867–2908.
- (729) Whittlesey, M. K.; Peris, E. Catalytic Hydrodefluorination with Late Transition Metal Complexes. *ACS Catal.* **2014**, *4*, 3152–3159.
- (730) Hayashi, S.; Nakai, T.; Ishikawa, N.; Burton, D. J.; Naae, D. G.; Kesling, H. S. Convenient Procedures for Conversion of Carbonyl Compounds to *gem*-Difluoroolefins and Their Selective Reductions to Monofluoroolefins. *Chem. Lett.* **1979**, *8*, 983–986.
- (731) Wu, J.; Xiao, J.; Dai, W.; Cao, S. Synthesis of Monofluoroalkenes through Selective Hydrodefluorination of *gem*-Difluoroalkenes with Red-Al<sup>®</sup>. *RSC Adv.* **2015**, *5*, 34498–34501.
- (732) Tellier, F.; Sauvêtre, R. Synthesis and Some Properties of 1-Fluoro-1-alken-3-ols. *Tetrahedron Lett.* **1995**, *36*, 4223–4226.
- (733) Tellier, F.; Sauvêtre, R. Reduction of 1,1-Difluoro-1-alken-3-ols with Lithium Tetrahydroaluminate. Application to the Synthesis of 1,1-Difluoro-2-alkenes and 2-Alkenals. *J. Fluorine Chem.* **1996**, *76*, 181–185.
- (734) Landelle, G.; Turcotte-Savard, M.-O.; Angers, L.; Paquin, J.-F. Stereoselective Synthesis of Both Stereoisomers of  $\beta$ -Fluorostyrene Derivatives from a Common Intermediate. *Org. Lett.* **2011**, *13*, 1568–1571.
- (735) Hu, J.; Han, X.; Yuan, Y.; Shi, Z. Stereoselective Synthesis of Z Fluoroalkenes through Copper-Catalyzed Hydrodefluorination of *gem*-Difluoroalkenes with Water. *Angew. Chem., Int. Ed.* **2017**, *56*, 13342–13346.
- (736) Kojima, R.; Kubota, K.; Ito, H. Stereodivergent Hydrodefluorination of *gem*-Difluoroalkenes: Selective Synthesis of (*Z*)- and (*E*)-Monofluoroalkenes. *Chem. Commun.* **2017**, *53*, 10688–10691.
- (737) Ma, Q.; Liu, C.; Tsui, G. C. Palladium-Catalyzed Stereoselective Hydrodefluorination of Tetrasubstituted *gem*-Difluoroalkenes. *Org. Lett.* **2020**, *22*, 5193–5197.
- (738) Tian, P.; Feng, C.; Loh, T.-P. Rhodium-Catalyzed C(sp<sup>2</sup>)-C(sp<sup>2</sup>) Bond Formation via C-H/C-F Activation. *Nat. Commun.* **2015**, *6*, 7472.
- (739) Liu, H.; Song, S.; Wang, C.-Q.; Feng, C.; Loh, T.-P. Redox-Neutral Rhodium-Catalyzed [4+1] Annulation through Formal Dehydrogenative Vinylidene Insertion. *ChemSusChem* **2017**, *10*, 58–61.
- (740) Ji, W.-W.; Lin, E.; Li, Q.; Wang, H. Heteroannulation Enabled by a Bimetallic Rh(III)/Ag(I) Relay Catalysis: Application in the Total Synthesis of Aristolactam BII. *Chem. Commun.* **2017**, *53*, 5665–5668.
- (741) Wu, J.-Q.; Zhang, S.-S.; Gao, H.; Qi, Z.; Zhou, C.-J.; Ji, W.-W.; Liu, Y.; Chen, Y.; Li, Q.; Li, X.; Wang, H. Experimental and Theoretical Studies on Rhodium-Catalyzed Coupling of Benzamides with 2,2-Difluorovinyl Tosylate: Diverse Synthesis of Fluorinated Heterocycles. *J. Am. Chem. Soc.* **2017**, *139*, 3537–3545.
- (742) Zhou, L.; Zhu, C.; Loh, T.-P.; Feng, C. Rh-Catalyzed C-H Bond Alkylation of Indoles with  $\alpha,\alpha$ -Difluorovinyl Tosylate via Indolyl Group Migration. *Chem. Commun.* **2018**, *54*, 5618–5621.
- (743) Song, S.; Liu, H.; Wang, L.; Zhu, C.; Loh, T.-P.; Feng, C. Rhodium-Catalyzed Defluorinative Vinylation of *gem*-Difluoroalkenes for the Synthesis of 2-Fluoro-1,3-dienes. *Chin. J. Chem.* **2019**, *37*, 1036–1040.
- (744) Tian, M.; Yang, X.; Zhang, B.; Liu, B.; Li, X. Rh(III)-Catalyzed  $\alpha$ -Fluoroalkenylation of *N*-Nitrosoanilines with 2,2-Difluorovinyl Tosylates via C-H Bond Activation. *Org. Chem. Front.* **2018**, *5*, 3406–3409.
- (745) Wang, N.; Yang, Q.; Mao, X.; Peng, Y. Rhodium-Catalyzed Merging of 2-Arylquinazolinone and 2,2-Difluorovinyl Tosylate: Diverse Synthesis of Monofluoroolefin Quinazolinone Derivatives. *ACS Omega* **2020**, *5*, 14635–14644.
- (746) Gong, T.-J.; Xu, M.-Y.; Yu, S.-H.; Yu, C.-G.; Su, W.; Lu, X.; Xiao, B.; Fu, Y. Rhodium(III)-Catalyzed Directed C-H Coupling with Methyl Trifluoroacrylate: Diverse Synthesis of Fluoroalkenes and Heterocycles. *Org. Lett.* **2018**, *20*, 570–573.
- (747) Kong, L.; Liu, B.; Zhou, X.; Wang, F.; Li, X. Rhodium(III)-Catalyzed Regio- and Stereoselective Benzylic  $\alpha$ -Fluoroalkenylation with *gem*-Difluorostyrenes. *Chem. Commun.* **2017**, *53*, 10326–10329.
- (748) Li, N.; Chang, J.; Kong, L.; Li, X. Ruthenium(II)-Catalyzed  $\alpha$ -Fluoroalkenylation of Arenes via C-H Bond Activation and C-F Bond Cleavage. *Org. Chem. Front.* **2018**, *5*, 1978–1982.
- (749) Zhang, L.; Deng, K.; Wu, G.; Yang, J.; Tang, S.; Fu, X.; Xia, C.; Ji, Y. Ruthenium(II)-Catalyzed  $\alpha$ -Fluoroalkenylation of Oxime Ethers with *gem*-Difluorostyrenes via C-H Activation and C-F Cleavage. *J. Org. Chem.* **2020**, *85*, 12670–12681.
- (750) Kong, L.; Zhou, X.; Li, X. Cobalt(III)-Catalyzed Regio- and Stereoselective  $\alpha$ -Fluoroalkenylation of Arenes with *gem*-Difluorostyrenes. *Org. Lett.* **2016**, *18*, 6320–6323.
- (751) Zell, D.; Müller, V.; Dhawa, U.; Bursch, M.; Presa, R. R.; Grimme, S.; Ackermann, L. Mild Cobalt(III)-Catalyzed Allylative C-F/C-H Functionalizations at Room Temperature. *Chem. - Eur. J.* **2017**, *23*, 12145–12148.
- (752) Murakami, N.; Yoshida, M.; Yoshino, T.; Matsunaga, S. Synthesis of Fluorine-Containing 6-Arylpurine Derivatives via Cp\*Co(III)-Catalyzed C-H Bond Activation. *Chem. Pharm. Bull.* **2018**, *66*, 51–54.
- (753) Cai, S.-H.; Ye, L.; Wang, D.-X.; Wang, Y.-Q.; Lai, L.-J.; Zhu, C.; Feng, C.; Loh, T.-P. Manganese-Catalyzed Synthesis of Monofluoroalkenes via C-H Activation and C-F Cleavage. *Chem. Commun.* **2017**, *53*, 8731–8734.
- (754) Zell, D.; Dhawa, U.; Müller, V.; Bursch, M.; Grimme, S.; Ackermann, L. C-F/C-H Functionalization by Manganese(I) Catalysis: Expedient (Per)Fluoro-Allylations and Alkenylations. *ACS Catal.* **2017**, *7*, 4209–4213.
- (755) Zhang, X.; Lin, Y.; Zhang, J.; Cao, S. Base-Mediated Direct Fluoroalkenylation of 2-Phenyl-1,3,4-oxadiazole, Benzothiazole and Benzoxazole with *gem*-Difluoroalkenes. *RSC Adv.* **2015**, *5*, 7905–7908.
- (756) Chen, K.; Chen, W.; Chen, F.; Zhang, H.; Xu, H.; Zhou, Z.; Yi, W. Synthesis of 2-Aminobenzofurans via Base-Mediated [3 + 2] Annulation of *N*-Phenoxy Amides with *gem*-Difluoroalkenes. *Org. Chem. Front.* **2021**, *8*, 4452–4458.
- (757) Lu, X.; Wang, Y.; Zhang, B.; Pi, J.-J.; Wang, X.-X.; Gong, T.-J.; Xiao, B.; Fu, Y. Nickel-Catalyzed Defluorinative Reductive Cross-Coupling of *gem*-Difluoroalkenes with Unactivated Secondary and Tertiary Alkyl Halides. *J. Am. Chem. Soc.* **2017**, *139*, 12632–12637.
- (758) Yu, L.; Tang, M.-L.; Si, C.-M.; Meng, Z.; Liang, Y.; Han, J.; Sun, X. Zinc-Mediated Decarboxylative Alkylation of *gem*-Difluoroalkenes. *Org. Lett.* **2018**, *20*, 4579–4583.
- (759) Liu, F.; Zhuang, Z.; Qian, Q.; Zhang, X.; Yang, C. Ru-Catalyzed Defluorinative Alkylation or Catalyst-Free Hydroalkylation of *gem*-Difluoroalkenes Enabled by Visible Light. *J. Org. Chem.* **2022**, *87*, 2730–2739.
- (760) Yang, L.; Ji, W.-W.; Lin, E.; Li, J.-L.; Fan, W.-X.; Li, Q.; Wang, H. Synthesis of Alkylated Monofluoroalkenes via Fe-Catalyzed Defluorinative Cross-Coupling of Donor Alkenes with *gem*-Difluoroalkenes. *Org. Lett.* **2018**, *20*, 1924–1927.

- (761) Zhou, L.; Zhu, C.; Bi, P.; Feng, C. Ni-Catalyzed Migratory Fluoro-Alkenylation of Unactivated Alkyl Bromides with *gem*-Difluoroalkenes. *Chem. Sci.* **2019**, *10*, 1144–1149.
- (762) Ma, T.; Chen, Y.; Li, Y.; Ping, Y.; Kong, W. Nickel-Catalyzed Enantioselective Reductive Aryl Fluoroalkenylation of Alkenes. *ACS Catal.* **2019**, *9*, 9127–9133.
- (763) Kondoh, A.; Koda, K.; Terada, M. Organocatalytic Nucleophilic Substitution Reaction of *gem*-Difluoroalkenes with Ketene Silyl Acetals. *Org. Lett.* **2019**, *21*, 2277–2280.
- (764) Zhang, Q.-Q.; Chen, S.-Y.; Lin, E.; Wang, H.; Li, Q. Regio- and Stereoselective Alkenylation of Allenolates with *gem*-Difluoroalkenes: Facile Access to Fluorinated 1,4-Enynes Bearing an All-Carbon Quaternary Center. *Org. Lett.* **2019**, *21*, 3123–3126.
- (765) Dai, W.; Xiao, J.; Jin, G.; Wu, J.; Cao, S. Palladium- and Nickel-Catalyzed Kumada Cross-Coupling Reactions of *gem*-Difluoroalkenes and Monofluoroalkenes with Grignard Reagents. *J. Org. Chem.* **2014**, *79*, 10537–10546.
- (766) Dai, W.; Zhang, X.; Zhang, J.; Lin, Y.; Cao, S. Synthesis of Exocyclic Trisubstituted Alkenes via Nickel-Catalyzed Kumada-Type Cross-Coupling Reaction of *gem*-Difluoroalkenes with Di-Grignard Reagents. *Adv. Synth. Catal.* **2016**, *358*, 183–187.
- (767) Yu, H.; Richey, R. N.; Carson, M. W.; Coghlan, M. J. Cyclocarbopalladation of Alkynes: A Stereoselective Method for Preparing Dibenzoxapine Containing Tetrasubstituted Exocyclic Alkenes. *Org. Lett.* **2006**, *8*, 1685–1688.
- (768) Dai, W.; Shi, H.; Zhao, X.; Cao, S. Sterically Controlled Cu-Catalyzed or Transition-Metal-Free Cross-Coupling of *gem*-Difluoroalkenes with Tertiary, Secondary, and Primary Alkyl Grignard Reagents. *Org. Lett.* **2016**, *18*, 4284–4287.
- (769) Wang, Y.; Tang, Y.; Zong, Y.; Tsui, G. C. Highly Selective C–F Bond Functionalization of Tetrasubstituted *gem*-Difluoroalkenes and Trisubstituted Monofluoroalkenes Using Grignard Reagents. *Org. Lett.* **2022**, *24*, 4087–4092.
- (770) Zhang, J.; Wang, B.; Liu, Y.; Cao, S. Stereoselective Synthesis of *Z*-Fluorostyrene Derivatives via Nickel-Catalyzed Cross-Coupling of *gem*-Difluorostyrenes with Organozinc Reagents. *Chin. J. Org. Chem.* **2019**, *39*, 249–256.
- (771) Tong, X.; Luo, S.-S.; Shen, H.; Zhang, S.; Cao, T.; Luo, Y.-P.; Huang, L.-L.; Ma, X.-T.; Liu, X.-W. Nickel-Catalyzed Defluorinative Alkylation of C(sp<sup>2</sup>)-F Bonds. *Org. Chem. Front.* **2021**, *8*, 4533–4542.
- (772) Zhu, Z.; Lin, L.; Xiao, J.; Shi, Z. Nickel-Catalyzed Stereo- and Enantioselective Cross-Coupling of *gem*-Difluoroalkenes with Carbon Electrophiles by C–F Bond Activation. *Angew. Chem., Int. Ed.* **2022**, *61*, e202113209.
- (773) Lin, H.; Jiao, W.; Chen, Z.; Han, J.; Fang, D.; Wang, M.; Liao, J. Enantioselective Cu-Catalyzed Nucleophilic Addition of Fluorinated Reagents: C–C Bond Formation for the Synthesis of Chiral Vicinal Difluorides. *Org. Lett.* **2022**, *24*, 2197–2202.
- (774) Watabe, Y.; Kanazawa, K.; Fujita, T.; Ichikawa, J. Nickel-Catalyzed Hydroalkenylation of Alkynes through C–F Bond Activation: Synthesis of 2-Fluoro-1,3-dienes. *Synthesis* **2017**, *49*, 3569–3575.
- (775) Wang, Y.; Ma, Q.; Tsui, G. C. Stereoselective Synthesis of Difluorinated 1,3-Dienes via Palladium-Catalyzed C–F Bond Activation of Tetrasubstituted *gem*-Difluoroalkenes. *Org. Lett.* **2021**, *23*, 5241–5245.
- (776) Li, M.; Wang, Y.; Tsui, G. C. Palladium-Catalyzed Stereoselective C–F Bond Vinylation and Allylation of Tetrasubstituted *gem*-Difluoroalkenes via Stille Coupling: Synthesis of Monofluorinated 1,3- and 1,4-Dienes. *Org. Lett.* **2021**, *23*, 8072–8076.
- (777) Heitz, W.; Knebelkamp, A. Synthesis of Fluorostyrenes via Palladium-Catalyzed Reactions of Aromatic Halides with Fluoroolefins. *Makromol. Chem. Rapid Commun.* **1991**, *12*, 69–75.
- (778) Fujiwara, M.; Ichikawa, J.; Okauchi, T.; Minami, T. Vinylic C–F Bond Activation with Low-Valent Zirconocene: the Generation and Cross-Coupling Reactions of 1-Fluorovinylzirconocene. *Tetrahedron Lett.* **1999**, *40*, 7261–7265.
- (779) Xiong, Y.; Huang, T.; Ji, X.; Wu, J.; Cao, S. Nickel-Catalyzed Suzuki–Miyaura Type Cross-Coupling Reactions of (2,2-Difluorovinyl)benzene Derivatives with Arylboronic Acids. *Org. Biomol. Chem.* **2015**, *13*, 7389–7392.
- (780) Thornbury, R. T.; Toste, F. D. Palladium-Catalyzed Defluorinative Coupling of 1-Aryl-2,2-Difluoroalkenes and Boronic Acids: Stereoselective Synthesis of Monofluorostilbenes. *Angew. Chem., Int. Ed.* **2016**, *55*, 11629–11632.
- (781) Wang, Y.; Qi, X.; Ma, Q.; Liu, P.; Tsui, G. C. Stereoselective Palladium-Catalyzed Base-Free Suzuki–Miyaura Cross-Coupling of Tetrasubstituted *gem*-Difluoroalkenes: An Experimental and Computational Study. *ACS Catal.* **2021**, *11*, 4799–4809.
- (782) Huang, W.; Wan, X.; Shen, Q. Enantioselective Construction of Trifluoromethoxylated Stereogenic Centers by a Nickel-Catalyzed Asymmetric Suzuki–Miyaura Coupling of Secondary Benzyl Bromides. *Angew. Chem. Int. Ed.* **2017**, *56*, 11986–11989.
- (783) Xiao, Y.; Huang, W.; Shen, Q. Stereoselective Formation of *Z*-Monofluoroalkenes by Nickel-Catalyzed Defluorinative Coupling of *gem*-Difluoroalkenes with Lithium Organoborates. *Chin. Chem. Lett.* **2022**, *33*, 4277–4280.
- (784) Ohashi, M.; Saijo, H.; Shibata, M.; Ogoshi, S. Palladium-Catalyzed Base-Free Suzuki–Miyaura Coupling Reactions of Fluorinated Alkenes and Arenes via a Palladium Fluoride Key Intermediate. *Eur. J. Org. Chem.* **2013**, *2003*, 443–447.
- (785) Kikushima, K.; Sakaguchi, H.; Saijo, H.; Ohashi, M.; Ogoshi, S. Copper-Mediated One-pot Synthesis of Trifluorostyrene Derivatives from Tetrafluoroethylene and Arylboronate. *Chem. Lett.* **2015**, *44*, 1019–1021.
- (786) Ohashi, M.; Kambara, T.; Hatanaka, T.; Saijo, H.; Doi, R.; Ogoshi, S. Palladium-Catalyzed Coupling Reactions of Tetrafluoroethylene with Arylzinc Compounds. *J. Am. Chem. Soc.* **2011**, *133*, 3256–3259.
- (787) Yamada, S.; Noma, M.; Konno, T.; Ishihara, T.; Yamanaka, H. Novel Synthesis of (*Z*)-Difluoroacrylates via a Highly Stereoselective Addition–Elimination Reaction. *Org. Lett.* **2006**, *8*, 843–845.
- (788) Jin, G.; Zhang, J.; Wu, W.; Cao, S. Stereoselective Synthesis of  $\beta$ -Fluoroenyne by the Reaction of *gem*-Difluoroalkenes with Terminal Alkynes. *J. Fluorine Chem.* **2014**, *168*, 240–246.
- (789) Ma, Q.; Wang, Y.; Tsui, G. C. Stereoselective Palladium-Catalyzed C–F Bond Alkylation of Tetrasubstituted *gem*-Difluoroalkenes. *Angew. Chem., Int. Ed.* **2020**, *59*, 11293–11297.
- (790) Rousée, K.; Bouillon, J.-P.; Couve-Bonnaire, S.; Pannecoucke, X. Stereospecific Synthesis of Tri- and Tetrasubstituted  $\alpha$ -Fluoroacrylates by Mizoroki–Heck Reaction. *Org. Lett.* **2016**, *18*, 540–543.
- (791) Yan, S.-S.; Wu, D.-S.; Ye, J.-H.; Gong, L.; Zeng, X.; Ran, C.-K.; Gui, Y.-Y.; Li, J.; Yu, D.-G. Copper-Catalyzed Carboxylation of C–F Bonds with CO<sub>2</sub>. *ACS Catal.* **2019**, *9*, 6987–6992.
- (792) Liu, J.; Nie, W.; Yu, H.; Shi, J. A Mechanistic Study on Cu(I) Catalyzed Carboxylation of the C–F Bond with CO<sub>2</sub>: a DFT Study. *Org. Biomol. Chem.* **2020**, *18*, 9065–9071.
- (793) Xie, S.-L.; Cui, X.-Y.; Gao, X.-T.; Zhou, F.; Wu, H.-L.; Zhou, J. Stereoselective Defluorinative Carboxylation of *gem*-Difluoroalkenes with Carbon Dioxide. *Org. Chem. Front.* **2019**, *6*, 3678–3682.
- (794) Xie, S.-L.; Gao, X.-T.; Wu, H.-H.; Zhou, F.; Zhou, J. Direct Electrochemical Defluorinative Carboxylation of *gem*-Difluoroalkenes with Carbon Dioxide. *Org. Lett.* **2020**, *22*, 8424–8429.
- (795) Zhang, J.; Xu, C.; Wu, W.; Cao, S. Mild and Copper-Free Stereoselective Cyanation of *gem*-Difluoroalkenes by Using Benzyl Nitrile as a Cyanating Reagent. *Chem. - Eur. J.* **2016**, *22*, 9902–9908.

- (796) Ma, Y.-C.; Zhang, Y.; Gu, C.-Z.; Du, G.-F.; He, L. Stereoselective Synthesis of  $\alpha$ -Fluoroacrylonitriles via Organocatalytic Cyanation of *gem*-Difluoroalkenes and TMSCN. *New J. Chem.* **2019**, *43*, 10985–10988.
- (797) Jiang, L.-F.; Ren, B.-T.; Li, B.; Zhang, G.-Y.; Peng, Y.; Guan, Z.-Y.; Deng, Q.-H. Nucleophilic Substitution of *gem*-Difluoroalkenes with TMSNu Promoted by Catalytic Amounts of  $\text{Cs}_2\text{CO}_3$ . *J. Org. Chem.* **2019**, *84*, 6557–6564.
- (798) Peng, J.-B.; Wu, F.-P.; Wu, X.-F. First-Row Transition-Metal-Catalyzed Carbonylative Transformations of Carbon Electrophiles. *Chem. Rev.* **2019**, *119*, 2090–2127.
- (799) Wu, F.-P.; Yuan, Y.; Liu, J.; Wu, X.-F. Pd/Cu-Catalyzed Defluorinative Carbonylative Coupling of Aryl Iodides and *gem*-Difluoroalkenes: Efficient Synthesis of  $\alpha$ -Fluoroaldehydes. *Angew. Chem., Int. Ed.* **2021**, *60*, 8818–8822.
- (800) Zhang, J.; Dai, W.; Liu, Q.; Cao, S. Cu-Catalyzed Stereoselective Borylation of *gem*-Difluoroalkenes with  $\text{B}_2\text{pin}_2$ . *Org. Lett.* **2017**, *19*, 3283–3286.
- (801) Sakaguchi, H.; Uetake, Y.; Ohashi, M.; Niwa, T.; Ogoshi, S.; Hosoya, T. Copper-Catalyzed Regioselective Monodefluoroborylation of Polyfluoroalkenes en Route to Diverse Fluoroalkenes. *J. Am. Chem. Soc.* **2017**, *139*, 12855–12862.
- (802) Ito, H.; Seo, T.; Kojima, R.; Kubota, K. Copper(I)-Catalyzed Stereoselective Defluoroborylation of Aliphatic *gem*-Difluoroalkenes. *Chem. Lett.* **2018**, *47*, 1330–1332.
- (803) Hu, J.; Zhao, Y.; Shi, Z. Highly Tunable Multi-Borylation of *gem*-Difluoroalkenes via Copper Catalysis. *Nat. Catal.* **2018**, *1*, 860–869.
- (804) Tan, D.-H.; Lin, E.; Ji, W.-W.; Zeng, Y.-F.; Fan, W.-X.; Li, Q.; Gao, H.; Wang, H. Copper-Catalyzed Stereoselective Defluorinative Borylation and Silylation of *gem*-Difluoroalkenes. *Adv. Synth. Catal.* **2018**, *360*, 1032–1037.
- (805) Sakaguchi, H.; Ohashi, M.; Ogoshi, S. Fluorinated Vinylsilanes from the Copper-Catalyzed Defluorosilylation of Fluoroalkene Feedstocks. *Angew. Chem., Int. Ed.* **2018**, *57*, 328–332.
- (806) Gao, P.; Wang, G.; Xi, L.; Wang, M.; Li, S.; Shi, Z. Transition-Metal-Free Defluorosilylation of Fluoroalkenes with Silylboronates. *Chin. J. Chem.* **2019**, *37*, 1009–1014.
- (807) Zhou, S.; Pu, Y.; Liu, Z.; Zhang, X.; Zhu, J.; Feng, Z. Iron-Catalyzed Diborylation of Unactivated Aliphatic *gem*-Dihalogenoalkenes: Synthesis of 1,2-Bis(boryl)alkanes. *Org. Lett.* **2021**, *23*, 5565–5570.
- (808) Zhang, H.; Wang, E.; Geng, S.; Liu, Z.; He, Y.; Peng, Q.; Feng, Z. Experimental and Computational Studies of the Iron-Catalyzed Selective and Controllable Defluorosilylation of Unactivated Aliphatic *gem*-Difluoroalkenes. *Angew. Chem., Int. Ed.* **2021**, *60*, 10211–10218.
- (809) Xiong, Y.; Zhang, X.; Huang, T.; Cao, S. Synthesis of *N*-( $\alpha$ -Fluorovinyl)azoles by the Reaction of Difluoroalkenes with Azoles. *J. Org. Chem.* **2014**, *79*, 6395–6402.
- (810) Wang, M.; Liang, F.; Xiong, Y.; Cao, S. Synthesis of Fluorovinyl Aryl Ethers by a Three-Component Reaction of *gem*-Difluoroalkenes with Arylboronic Acids and Oxygen. *RSC Adv.* **2015**, *5*, 11996–11999.
- (811) Huang, T.; Zhao, X.; Ji, X.; Wu, W.; Cao, S. Synthesis of Fluorovinyl Pyrazolyl(thio)ethers by the Reaction of *gem*-Difluoroalkenes with Pyrazolin-5-ones(thiones). *J. Fluorine Chem.* **2016**, *182*, 61–68.
- (812) Cong, Z.-S.; Li, Y.-G.; Chen, L.; Xing, F.; Du, G.-F.; Gu, C.-Z.; He, L. N-Heterocyclic Carbene-Catalyzed Stereoselective Construction of Olefinic Carbon–Sulfur Bonds via Cross-Coupling Reaction of *gem*-Difluoroalkenes and Thiols. *Org. Biomol. Chem.* **2017**, *15*, 3863–3868.
- (813) Li, J.; Rao, W.; Wang, S.-Y.; Ji, S.-J. Nickel-Catalyzed Defluorinative Reductive Cross-Coupling Reaction of *gem*-Difluoroalkenes with Thiosulfonate or Selenium Sulfonate. *J. Org. Chem.* **2019**, *84*, 11542–11552.
- (814) Li, Y.; Li, X.; Li, X.; Shi, D. Highly E-Selective Synthesis of  $\alpha$ -Fluoro- $\beta$ -arylalkenyl Sulfones from *gem*-Difluoroalkenes with Sodium Sulfonates. *J. Org. Chem.* **2021**, *86*, 6983–6993.
- (815) Li, Z.; Qiu, X.; Jiang, L.; Wang, Q. Progress in Visible-Light Catalyzed C–F Bond Functionalization of *gem*-Difluoroalkenes. *Chin. J. Org. Chem.* **2021**, *41*, 4192–4207.
- (816) Xie, J.; Yu, J.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. Monofluoroalkenylation of Dimethylamino Compounds through Radical–Radical Cross-Coupling. *Angew. Chem., Int. Ed.* **2016**, *55*, 9416–9421.
- (817) Hu, Y.; Liu, X.; Ren, Z.; Hu, B.; Li, J.  $\text{Csp}^3\text{-H}$  Monofluoroalkenylation via Stereoselective C–F Bond Cleavage. *Chem. Commun.* **2022**, *58*, 2734–2737.
- (818) Li, L.; Xiao, T.; Chen, H.; Zhou, L. Visible Light-mediated Twofold Unsymmetrical  $\text{sp}^3$  C–H Functionalization and Double C–F Substitution. *Chem. - Eur. J.* **2017**, *23*, 2249–2254.
- (819) Chen, H.; Xiao, T.; Li, L.; Anand, D.; He, Y.; Zhou, L. Synthesis of Fluorinated Benzo[a]quinolizidines via Visible Light-induced Tandem Substitution of Two Fluorine Atoms in a  $\text{CF}_3$  Group. *Adv. Synth. Catal.* **2017**, *359*, 3642–3647.
- (820) Chen, H.; He, Y.; Zhou, L. A Photocatalytic Decarboxylative/Defluorinative [4 + 3] Annulation of *o*-Hydroxyphenylacetic Acids and Trifluoromethyl Alkenes: Synthesis of Fluorinated Dihydrobenzoxepines. *Org. Chem. Front.* **2018**, *5*, 3240–3244.
- (821) Fuchibe, K.; Takahashi, M.; Ichikawa, J. Substitution of Two Fluorine Atoms in a Trifluoromethyl Group: Regioselective Synthesis of 3-Fluoropyrazoles. *Angew. Chem., Int. Ed.* **2012**, *51*, 12059–12062.
- (822) Ichitsuka, T.; Fujita, T.; Arita, T.; Ichikawa, J. Double C–F Bond Activation through  $\beta$ -Fluorine Elimination: Nickel-Mediated [3+2] Cycloaddition of 2-Trifluoromethyl-1-alkenes with Alkynes. *Angew. Chem., Int. Ed.* **2014**, *53*, 7564–7568.
- (823) Fujita, T.; Takazawa, M.; Sugiyama, K.; Suzuki, N.; Ichikawa, J. Domino C–F Bond Activation of the  $\text{CF}_3$  Group: Synthesis of Fluorinated Dibenzo[a,c][7]annulenes from 2-(Trifluoromethyl)-1-alkenes and 2,2'-Diceriobiaryls. *Org. Lett.* **2017**, *19*, 588–591.
- (824) Yang, J.; Mao, A.; Yue, Z.; Zhu, W.; Luo, X.; Zhu, C.; Xiao, Y.; Zhang, J. A Simple Base-Mediated Synthesis of Diverse Functionalized Ring-Fluorinated 4*H*-Pyran via Double Direct C–F Substitutions. *Chem. Commun.* **2015**, *51*, 8326–8329.
- (825) Yang, J.; Zhou, X.; Zeng, Y.; Huang, C.; Xiao, Y.; Zhang, J. Synthesis of 2-Fluoro-2-pyrrolines via Tandem Reaction of  $\alpha$ -Trifluoromethyl- $\alpha,\beta$ -unsaturated Carbonyl Compounds with *N*-Tosylated 2-Aminomalonates. *Chem. Commun.* **2016**, *52*, 4922–4925.
- (826) Wu, L.-H.; Cheng, J.-K.; Shen, L.; Shen, Z.-L.; Loh, T.-P. Visible Light-Mediated Trifluoromethylation of Fluorinated Alkenes via C–F Bond Cleavage. *Adv. Synth. Catal.* **2018**, *360*, 3894–3899.
- (827) Wang, Q.; Qu, Y.; Tian, H.; Liu, Y.; Song, H.; Wang, Q. Trifluoromethylation and Monofluoroalkenylation of Alkenes Through Radical–Radical Cross-Coupling. *Chem. - Eur. J.* **2019**, *25*, 8686–8690.
- (828) Li, J.; Lefebvre, Q.; Yang, H.; Zhao, Y.; Fu, H. Visible Light Photocatalytic Decarboxylative Monofluoroalkenylation of  $\alpha$ -Amino Acids with *gem*-Difluoroalkenes. *Chem. Commun.* **2017**, *53*, 10299–10302.
- (829) Yang, H.; Tian, C.; Qiu, D.; Tian, H.; An, G.; Li, G. Organic photoredox catalytic decarboxylative cross-coupling of *gem*-difluoroalkenes with unactivated carboxylic acids. *Org. Chem. Front.* **2019**, *6*, 2365–2370.
- (830) Zhu, C.; Zhang, Y.-F.; Liu, Z.-Y.; Zhou, L.; Liu, H.; Feng, C. Selective C–F Bond Carboxylation of *gem*-Difluoroalkenes with  $\text{CO}_2$  by Photoredox/Palladium Dual Catalysis. *Chem. Sci.* **2019**, *10*, 6721–6726.

- (831) Tian, H.; Xia, Q.; Wang, Q.; Dong, J.; Liu, Y.; Wang, Q. Direct  $\alpha$ -Monofluoroalkenylation of Heteroatomic Alkanes via a Combination of Photoredox Catalysis and Hydrogen-Atom-Transfer Catalysis. *Org. Lett.* **2019**, *21*, 4585–4589.
- (832) Cao, C.-L.; Zhang, G.-X.; Xue, F.; Deng, H.-P. Photoinduced C–H Monofluoroalkenylation with *gem*-Difluoroalkenes through Hydrogen Atom Transfer under Batch and Flow Conditions. *Org. Chem. Front.* **2022**, *9*, 959–965.
- (833) Du, H.-W.; Sun, J.; Gao, Q.-S.; Wang, J.-Y.; Wang, H.; Xu, Z.; Zhou, M.-D. Synthesis of Monofluoroalkenes through Visible-Light-Promoted Defluorinative Alkylation of *gem*-Difluoroalkenes with 4-Alkyl-1,4-dihydropyridines. *Org. Lett.* **2020**, *22*, 1542–1546.
- (834) Tian, H.; Yang, S.; Wang, X.; Xu, W.; Liu, Y.; Li, Y.; Wang, Q. Dehalogenative Cross-Coupling of *gem*-Difluoroalkenes with Alkyl Halides via a Silyl Radical-Mediated Process. *J. Org. Chem.* **2021**, *86*, 12772–12782.
- (835) Nambo, M.; Ghosh, K.; Yim, J. C.-H.; Tahara, Y.; Inai, N.; Yanai, T.; Crudden, C. M. Desulfonylative Coupling of Alkylsulfones with *gem*-Difluoroalkenes by Visible-Light Photoredox Catalysis. *ACS Catal.* **2022**, *12*, 9526–9532.
- (836) Wang, J.; Huang, B.; Yang, C.; Xia, W. Visible-Light-Mediated Defluorinative Cross-Coupling of *gem*-Difluoroalkenes with Thiols. *Chem. Commun.* **2019**, *55*, 1103–1106.
- (837) Li, Y.; Li, X.; Li, X.; Shi, D. Visible-Light-Promoted *E*-Selective Synthesis of  $\alpha$ -Fluoro- $\beta$ -arylalkenyl Sulfides via the Deoxygenation/Isomerization Process. *Chem. Commun.* **2021**, *57*, 2152–2155.
- (838) Wu, W.; Jiang, H.; Leng, J.; Ong, H.-W.; Wu, J. Visible-Light-Induced Selective Defluoroborylation of Polyfluoroarenes, *gem*-Difluoroalkenes, and Trifluoromethylalkenes. *Angew. Chem., Int. Ed.* **2020**, *59*, 4009–4016.
- (839) Ichikawa, J.; Wada, Y.; Okauchi, T.; Minami, T. 5-*Endo*-Trigonal Cyclization of *o*-Substituted *gem*-Difluorostyrenes: Syntheses of 2-Fluorinated Indoles, Benzo[*b*]furans and Benzo[*b*]thiophenes. *Chem. Commun.* **1997**, 1537–1538.
- (840) Ichikawa, J.; Fujiwara, M.; Wada, Y.; Okauchi, T.; Minami, T. The Nucleophilic 5-*Endo*-*Trig* Cyclization of *gem*-Difluoroolefins with Homoallylic Functional Groups: Syntheses of Ring-Fluorinated Dihydroheteroaromatics. *Chem. Commun.* **2000**, 1887–1888.
- (841) Wada, Y.; Ichikawa, J.; Katsume, T.; Nohiro, T.; Okauchi, T.; Minami, T. Intramolecular Cyclizations of *o*-Substituted  $\beta,\beta$ -Difluorostyrenes: Synthesis of 3-Fluorinated Isochromenes and Isothiochromenes. *Bull. Chem. Soc. Jpn.* **2001**, *74*, 971–977.
- (842) Ichikawa, J.; Wada, Y.; Fujiwara, M.; Sakoka, K. The Nucleophilic 5-*Endo*-*Trig* Cyclization of 1,1-Difluoro-1-alkenes: Ring-Fluorinated Hetero- and Carbocycle Synthesis and Remarkable Effect of the Vinyl Fluorines on the Disfavored Process. *Synthesis* **2002**, *13*, 1917–1936.
- (843) Sakoda, K.; Mihara, J.; Ichikawa, Heck-Type 5-*Endo*-*Trig* Cyclization Promoted by Vinylic Fluorines: Synthesis of 5-Fluoro-3*H*-pyrroles. *Chem. Commun.* **2005**, 4684–4686.
- (844) Yokota, M.; Fujita, D.; Ichikawa, J. Activation of 1,1-Difluoro-1-alkenes with a Transition-Metal Complex: Palladium(II)-Catalyzed Friedel-Crafts-Type Cyclization of 4,4-(Difluorohomoallyl)arenes. *Org. Lett.* **2007**, *9*, 4639–4642.
- (845) Zhang, X.; Dai, W.; Wu, W.; Cao, S. Copper-Catalyzed Coupling Cyclization of *gem*-Difluoroalkenes with Activated Methylene Carbonyl Compounds: Facile Domino Access to Polysubstituted Furans. *Org. Lett.* **2015**, *17*, 2708–2711.
- (846) Zhang, X.; Wu, M.; Zhang, J.; Cao, S. Synthesis of *N,N*-Disubstituted 2-Aminothiophenes by the Cyclization of *gem*-Difluoroalkenes with  $\beta$ -Keto Thioamides. *Org. Biomol. Chem.* **2017**, *15*, 2436–2442.
- (847) Zhang, X.; He, J.; Cao, S.; Facile Synthesis of Unsymmetrical 2,5-Disubstituted 1,3,4-Oxadiazoles by the Cyclization of *gem*-Difluoroalkenes with Acyl Hydrazides. *Asian J. Org. Chem.* **2019**, *8*, 279–282.
- (848) Fujita, T.; Watabe, Y.; Yamashita, S.; Tanabe, H.; Nojima, T.; Ichikawa, J. Silver-Catalyzed Vinylic C–F Bond Activation: Synthesis of 2-Fluoroindoles from  $\beta,\beta$ -Difluoro-*o*-sulfonamidostyrenes. *Chem. Lett.* **2016**, *45*, 964–966.
- (849) Zhang, B.; Zhang, X.; Hao, J.; Yang, C. Direct Approach to *N*-Substituted-2-Fluoroindoles by Sequential Construction of C–N Bonds from *gem*-Difluorostyrenes. *Org. Lett.* **2017**, *19*, 1780–1783.
- (850) Clavier, G.; Audebert, P. *s*-Tetrazines as Building Blocks for New Functional Molecules and Molecular Materials. *Chem. Rev.* **2010**, *110*, 3299–3314.
- (851) Fang, Z.; Hu, W.-L.; Liu, D.-Y.; Yu, C.-Y.; Hu, X.-G. Synthesis of Tetrazines from *gem*-Difluoroalkenes under Aerobic Conditions at Room Temperature. *Green Chem.* **2017**, *19*, 1299–1302.
- (852) Ausekle, E.; Ehlers, P.; Villinger, A.; Langer, P. One-Pot Synthesis of Dibenzo[*b,d*]oxepines via Olefinic C–F Bond Functionalization and Intramolecular Pd-Catalyzed C–H Arylation. *J. Org. Chem.* **2018**, *83*, 14195–14202.
- (853) Lu, C.-J.; Yu, X.; Chen, Y.-T.; Song, Q.-B.; Wang, H. Indolizine Synthesis via Copper-Catalyzed Cyclization of *gem*-Difluoroalkenes and 2-(Pyridin-2-yl)acetate Derivatives. *Org. Chem. Front.* **2020**, *7*, 2313–2318.
- (854) Fang, X.; Wu, Y.-M.; Deng, J.; Wang, S.-W. Synthesis of Monofluorinated Indolizines and Their Derivatives by the 1,3-Dipolar Reaction of *N*-Ylides with Fluorinated Vinyl Tosylates. *Tetrahedron* **2004**, *60*, 5487–5493.
- (855) Zhang, J. -Q.; Hu, D.; Song, J.; Ren, H. [3 + 2]-Annulation of *gem*-Difluoroalkenes and Pyridinium Ylides: Access to Functionalized 2-Fluoroindolizines. *J. Org. Chem.* **2021**, *86*, 4646–4660.
- (856) Yang, L.-M.; Zhang, Y.-Y.; Deng, J.-T.; Ma, A.-J.; Zhang, X.-Z.; Zhang, S.-Y.; Peng, J.-B. Oxidative [3+2] Annulation of Pyridinium Salts with *gem*-Difluoroalkenes: Synthesis of 2-Fluoroindolizines. *Asian J. Org. Chem.* **2021**, *10*, 1679–1682.
- (857) Li, J.; Liu, S.; Zhong, R.; Yang, Y.; Xu, J.; Yang, J.; Ding, H.; Wang, Z. Cascade Cyclization of Azadienes with Difluoroenoxy silanes: A One-Pot Formal [4 + 2] Approach to Fluorinated Polyfused Heterocycles. *Org. Lett.* **2021**, *23*, 9526–9532.
- (858) Swartz, C. R.; Parkin, S. R.; Bullock, J. E.; Anthony, J. E.; Mayer, A. C.; Malliaras, G. G. Synthesis and Characterization of Electron-Deficient Pentacenes. *Org. Lett.* **2005**, *7*, 3163–3166.
- (859) Fuchibe, K.; Morikawa, T.; Ueda, R.; Okauchi, T.; Ichikawa, J. Pinpoint-Fluorinated Phenanthrene Synthesis Based on C–F Bond Activation of Difluoroalkenes. *J. Fluorine Chem.* **2015**, *179*, 106–115.
- (860) Fuchibe, K.; Morikawa, T.; Shigeno, K.; Fujita, T.; Ichikawa, J. Pinpoint-Fluorinated Phenacenes: New Synthesis and Solubility Enhancement Strategies. *Org. Lett.* **2015**, *17*, 1126–1129.
- (861) Fuchibe, K.; Shigeno, K.; Zhao, N.; Aihara, H.; Akisaka, R.; Morikawa, T.; Fujita, T.; Yamakawa, K.; Shimada, T.; Ichikawa, J. Pinpoint-fluorinated polycyclic aromatic hydrocarbons (F-PAHs): Syntheses of difluorinated subfamily and their properties. *J. Fluorine Chem.* **2017**, *203*, 173–184.
- (862) R. P. Hughes, Conversion of Carbon–Fluorine Bonds  $\alpha$  to Transition Metal Centers to Carbon–Hydrogen, Carbon–Carbon, and Carbon–Heteroatom Bonds. *Eur. J. Inorg. Chem.* **2009**, *2009*, 4591–4606.
- (863) Karimzadeh-Younjali, M.; Wendt, O. F.  $\alpha$ - and  $\beta$ -Eliminations in Transition Metal Complexes: Strategies to Cleave Unstrained C–C and C–F Bonds. *Helv. Chim. Acta* **2021**, *104*, e210014.
- (864) Takachi, M.; Kita, Y.; Tobisu, M.; Fukumoto, Y.; Chatani, N. Nickel-Catalyzed Cyclization of Difluoro-Substituted 1,6-Enynes with Organozinc Reagents through the Stereoselective Activation of C–F Bonds: Synthesis of Bicyclo[3.2.0]heptene

Derivatives. *Angew. Chem., Int. Ed.* **2010**, *49*, 8717–8720.

(865) Fujita, T.; Watabe, Y.; Ichitsuka, T.; Ichikawa, J. Ni-Catalyzed Synthesis of Fluoroarenes via [2 + 2 + 2] Cycloaddition Involving  $\alpha$ -Fluorine Elimination. *Chem. - Eur. J.* **2015**, *21*, 13225–13228.

(866) Ohashi, M.; Ueda, Y.; Ogoshi, S. Nickel(o)-Mediated Transformation of Tetrafluoroethylene and Vinylarenes into Fluorinated Cyclobutyl Compounds. *Angew. Chem., Int. Ed.* **2017**, *56*, 2435–2439.

## Table of Contents

