

**Optical Measures of Cerebral Arterial Stiffness are Associated with White Matter Signal  
Abnormalities and Cognitive Performance in Normal Aging**

Chin Hong Tan<sup>a,b,c\*</sup>, Kathy A. Low<sup>a</sup>, Antonio M. Chiarelli<sup>a,d</sup>, Mark A. Fletcher<sup>a</sup>, Riccardo Navarra<sup>d</sup>,  
Agnieszka Z. Burzynska<sup>e</sup>, Tania S. Kong<sup>a,g</sup>, Benjamin Zimmerman<sup>a</sup>, Edward L. Maclin<sup>a</sup>, Bradley P.  
Sutton<sup>a,f</sup>, Gabriele Gratton<sup>a,g</sup> & Monica Fabiani<sup>a,g\*</sup>

<sup>a</sup>Beckman Institute, University of Illinois at Urbana-Champaign

<sup>b</sup>Division of Psychology, Nanyang Technological University

<sup>c</sup>Department of Pharmacology, National University of Singapore

<sup>d</sup>Department of Neuroscience, Imaging and Clinical Sciences, University G. D'Annunzio of Chieti-  
Pescara

<sup>e</sup>Department of Human Development and Family Studies, Molecular, Cellular and Integrative  
Neurosciences, Colorado State University

<sup>f</sup>Department of Bioengineering, University of Illinois at Urbana-Champaign

<sup>g</sup>Department of Psychology, University of Illinois at Urbana-Champaign

\*Corresponding Authors:

Prof. Monica Fabiani  
University of Illinois at Urbana-Champaign  
Beckman Institute  
405 N. Mathews Ave.  
Urbana, IL 61801-2325  
Email: [mfabiani@illinois.edu](mailto:mfabiani@illinois.edu)

Dr. Chin Hong Tan  
Division of Psychology  
Nanyang Technological University, Singapore  
48 Nanyang Avenue  
Singapore 639818  
[chinhong.tan@ntu.edu.sg](mailto:chinhong.tan@ntu.edu.sg)

### Abstract

Decline in fluid abilities in normal aging is associated with increased white matter lesions, measured on T1-weighted images as white matter signal abnormalities (WMSAs). WMSAs are particularly evident in hypertensive older adults, suggesting vascular involvement. However, since hypertension is assessed systemically, the specific role of *cerebral* arterial stiffening in WMSAs has yet to be demonstrated. In 93 cognitively normal adults (aged 18–87), we employed a novel method to measure cerebral arterial elasticity (pulse relaxation function, PReFx) with diffuse optical tomography (pulse-DOT) and investigated its association with WMSA, age, and cognition. PReFx was associated with WMSA, with older adults with low PReFx showing the greatest WMSA burden. PReFx in brain regions perfused by the middle cerebral artery showed the largest associations with WMSAs, and partially mediated the relationship between age and WMSA. Lastly, WMSA partially mediated the relationship between PReFx and fluid, but not crystallized abilities scores. Taken together, these findings suggest that loss of cerebral arterial elasticity is associated with cerebral white matter lesions and age-related cognitive decline.

**Keywords:** Aging; Fluid abilities; Cerebrovascular health; Arterial elasticity; Diffuse optical tomography (DOT); Pulse-DOT.

## 1. Introduction

Normal aging is often accompanied by decreases in cognitive functions (Fabiani, 2012; Park & Reuter-Lorenz, 2009; Salthouse, 2010). Associations have been found between age-related cognitive declines and white matter lesion burden (e.g., DeCarli et al., 1995; Leaper et al., 2001; Rabbitt et al., 2007; Raz et al., 2007). This age-related white matter degradation manifests as white matter signal abnormalities (WMSAs), which appear as *hypointensities* on T1-weighted magnetic resonance images (MRI) and *hyperintensities* on T2-weighted images (Brickman et al., 2011; Debette & Markus, 2010; Wardlaw, 2015). Clinical and pathophysiological data suggest that greater WMSA volume is associated with chronic hypoperfusion and/or lacunar lesions (Behrouz, Malek & Torbey, 2012; Pantoni & Simoni, 2003; VanDijk et al., 2008), which may be the consequence of age-related arteriosclerosis (i.e., the hardening of arteries, characterized by a loss of their elastic properties).

Studies investigating the relationship between arterial stiffness and WMSAs typically assess pulse pressure (i.e., the difference between systolic and diastolic blood pressure, an index of arteriosclerosis) systemically rather than within the brain (e.g., Kennedy & Raz, 2009; Leritz et al., 2014; Salat et al., 2012). As such, a specific link between white matter degeneration and cerebral arterial elasticity is not yet well established. Although Transcranial Doppler (TCD) ultrasound methods have been used to quantify intracranial cerebrovascular function using pulsatility and resistivity indices (see Keage et al., 2012 for a review), they are typically restricted to the insonation of one, or at most a few, large arteries, which limit its field of view. A broad field of view is essential to map elasticity across the cortical mantle and investigate its regional effects. Here we used a newly developed measure of the cerebral arterial pulse relaxation function (PReFx, also referred to as arterial compliance in previous papers) obtained throughout the cortex, based on diffuse optical tomography (pulse-DOT; Fabiani et al., 2014b; Tan et al., 2016; 2017; Chiarelli et al., 2017; 2019) to index cerebral arterial elasticity in cognitively normal adults.

In previous research, PReFx averaged across the whole head was found to be negatively associated with age and positively associated with estimated cardiorespiratory fitness (eCRF), brain volume (cortical gray and white matter and subcortical gray matter), and cognitive flexibility in healthy older adults

(Fabiani et al., 2014b; Tan et al., 2017). Recent methodological improvements now allow for the derivation of tomographic regional PReFx measurements in at least 50 (out of 70) cerebral cortical regions obtained from Freesurfer (Chiarelli et al., 2017).

As evidenced by studies investigating age-related changes in glial (e.g., Duncombe et al., 2017; Hutchison et al., 2013) and vascular function (e.g., Restom et al., 2007; Chen et al., 2014; Ances et al., 2009; Mohtasib et al., 2012), the tight link between neural activity and vascular properties (i.e., neurovascular coupling, see Fabiani et al., 2014) may play important roles in the hemodynamic changes observed in neurocognitive aging studies. Other studies have also established the role of white matter health on cognition in aging (e.g., Salami et al., 2012; Coxon et al., 2012; Madden et al., 2012; Charlton et al., 2009). As such, the primary goal of the current study was to investigate the relationships between cerebral arterial elasticity (measured with PReFx), macrostructural measures of white matter lesions (WMSAs) and cognitive function in healthy, cognitively normal younger and older adults.

Specifically, we hypothesized that cerebral arterial elasticity (measured with PReFx) should be associated with WMSAs, and that these effects should be particularly evident in older adults with low PReFx. We further investigated whether *regional* PReFx was associated with WMSAs. Lastly, we tested whether PReFx mediated the relationship between age and WMSAs, and whether WMSAs mediated the relationship between PReFx and fluid intelligence.

## **2. Methods**

The majority of the methods and procedures reported here overlap with Fabiani et al., (2014b) and Tan et al. (2017), so we are including only an abbreviated methods section, as per journal guidelines. Please note that all analyses reported here are new and not previously published.

### **2.1 Participants**

Ninety-three unique adult participants (age range = 18-87 years, mean age = 58.8, 46 females) were included in the current study. These are the same participants included in Fabiani et al. (2014b) and Tan et al. (2017), but none of the analyses presented here nor the combination of both samples were previously published. Supplementary Figure S1 shows the age distribution of this combined sample. Participants

were recruited through a mix of advertisements in local newspapers, gyms, retirement homes and community centers, in addition to campus-wide emails. Sample demographics are summarized in Table 1. Twenty-one participants reported taking blood pressure medications and 17 reported taking cholesterol medications. Six individuals showed signs of hypertension (i.e.  $>140/90$  mmHg) based on sphygmomanometer blood pressure measurements. Note, however, that PReFx measures are normalized (see Methods) by cerebral peak-systole to peak-diastole amplitude (a measure of cerebral pulse pressure, see Tan et al., 2016, 2017). Thus, pulse pressure and PReFx are substantially independent.

All participants reported themselves in good health and were free of depression and dementia, as assessed by the Beck's depression inventory (Beck, Steer, & Brown 1996) and the modified Mini-Mental Status Exam (mMMSE; Mayeux, Stern, Rosen, & Leventhal, 1981). All procedures were approved by the University of Illinois Institutional Review Board.

**INSERT TABLE 1 ABOUT HERE**

## **2.2 Assessment of cognitive function**

Participants underwent a battery of neuropsychological tests (see Fabiani et al., 2014; Tan et al., 2017). To reduce the number of comparisons, using an approach similar to Chan et al. (2014), we sorted the neuropsychological scales into two different domains, which we labeled “fluid” and “crystallized” abilities, respectively. The fluid abilities domain included the following tests: Raven’s matrices (Raven, Raven, & Court, 2003), forward and backward digit span (Mayeux, Stern, Rosen, & Leventhal, 1981), Operation-Span (O-Span; Altamura et al., 2007), trail A, trail B, and trail B-A (Corrigan & Hinkeldey, 1987). The crystallized abilities domain included Shipley’s vocabulary (Shipley, 1940) and a test of verbal fluency (letters CFL, Benton & Hamsher, 1989). For each test, each participant’s scores were standardized, and, when needed, their sign was changed so that positive values always indicated a higher-than-average performance and negative values indicated a lower-than-average performance. Then, for each participant, the standardized values for the tests in each domain were averaged together (effectively giving equal weight to each of them). These yielded two scores for each individual, a “fluid abilities” score and a “crystallized abilities” score. These scores were used for all further analyses.

## **2.3 Electrocardiogram (EKG) acquisition and processing**

Brain-Vision™ recorder and a Brain-Vision professional BrainAmp™ integrated amplifier system (Brain Products GmbH, Germany) were used to record Lead I of the EKG (left wrist referenced to right wrist; impedance < 20 kOhm; sampling rate=1000 Hz). EKG data were used for time-locking the optical pulse data to the R-wave, to ensure that the same pulse cycle was measured at all scalp locations.

## **2.4 Optical recording and analysis**

Optical data were acquired using a multi-channel frequency-domain oximeter (ISS Imagent™, Champaign, Illinois) equipped with 128 laser diodes (64 emitting light at 690 nm and 64 at 830 nm) and 24 photo-multiplier tubes (PMTs) at a sampling rate of 39.0625 Hz. Time-division multiplexing was used such that each detector picked up light from 16 different sources at different times within a multiplexing cycle. The light was transported to the scalp with optic fibers (0.4 mm core) and from the scalp back to the detectors using optic fiber bundles (3 mm diameter). Fibers were securely attached using custom-built

helmets of varying sizes. A resting-state paradigm (e.g., Eggebrecht et al., 2014) was used during recording. Optical data were recorded for a total of 8 blocks (6 minutes each) using 4 different optical montages (2 blocks, or 12 min recording time per montage), with the sequence of 4 montages counterbalanced across subjects. 384 channels (192 at 830 nm and 192 at 690 nm) were acquired for each montage (total=1536 channels, encompassing most of the scalp surface; source-detector distances = 15-80 mm). These procedures are similar to what was described in Tan et al., (2017) and Fabiani et al., (2014), with the exception that data from Fabiani et al. (2014) were based on 384 channels and 72 seconds of recording time. Comparisons between the PReFx values obtained in the two studies revealed no significant difference in mean and standard deviation, when participants with overlapping age ranges were considered, lending strong support to the merging of these two data sets.

DC optical intensity data were normalized, movement corrected (Chiarelli et al. 2015), and band-pass filtered between 0.5–5 Hz. As cerebral arterial blood is highly (>95%) oxygen saturated, only the 830nm light intensity data were used for pulse estimation, as this wavelength is more sensitive to oxygenated hemoglobin (Fabiani et al., 2014b). Optical pulse waveforms for each channel were derived by averaging the DC light intensity time-locked to the peak of the EKG R wave (signaling the depolarization of the ventricular myocardium). Sensitivity to superficial artifacts was reduced by only using channels with inter-optode distance >25 mm. Channels with inter-optode distance >70 mm were also excluded to increase signal-to-noise ratio (as light intensity is too low at those distances to be reliable).

3D reconstruction of the pulse waveform was conducted with a model of light propagation within the head (forward model) and an inverse procedure. A Finite Element Method (FEM) applied to the diffusion equation (Ishimaru, 1989; Paulsen & Jiang, 1995) was used to estimate the forward model. The FEM software NIRFAST (Dehghani et al., 2009; Eggebrecht et al., 2014) was used to model light propagation through heterogeneous head models and to compute Jacobian (sensitivity) matrices of DC light intensity reflecting absorption changes. “Fine” meshes (maximum tetrahedral volume = 2 mm<sup>3</sup>) were generated for FEM using the MATLAB software iso2mesh (Fang & Boas, 2009). The heterogeneous head models were based on segmented T1-weighted 3D anatomical images. Segmentation of the skull and scalp,

cerebrospinal fluid (CSF), white matter, and gray matter was performed using Statistical Parametric Mapping (SPM, Friston et al., 1994) functions applied to T1-weighted images (Penny et al., 2011). Baseline optical properties (absorption  $\mu_a$ , reduced scattering coefficient  $\mu_s'$  and refraction index  $\eta$ ) of tissues at the relevant wavelength were taken from Tian and Liu (2014) and Chiarelli et al. (2016). The optical values at the wavelength of interest (830 nm) were set to: scalp and skull:  $\mu_a=0.014 \text{ mm}^{-1}$ ,  $\mu_s'=0.84 \text{ mm}^{-1}$ ; CSF:  $\mu_a=0.004 \text{ mm}^{-1}$ ,  $\mu_s'=0.3 \text{ mm}^{-1}$ ; grey matter:  $\mu_a=0.019 \text{ mm}^{-1}$ ,  $\mu_s'=0.673 \text{ mm}^{-1}$ ; white matter:  $\mu_a=0.021 \text{ mm}^{-1}$ ,  $\mu_s'=1.01 \text{ mm}^{-1}$ . The refraction index was set equal for all the simulations and mediums considered at  $\eta=1.4$ . An inverse procedure was used to convert intensity changes on individual channels to absorption changes in voxel space (Chiarelli et al., 2016), which allows for an unbiased localization of absorption fluctuations up to a depth of 30 mm from the scalp.

PReFx (Fabiani et al., 2014b; Tan et al., 2017; Chiarelli et al., 2017) is a measure of the shape of the pulse during the interval between a peak systole and peak diastole. It represents the way in which arteries return to their original size after dilating to accommodate the blood bolus generated by a heart pulsation. If this curve is decelerated, the artery can be considered to be elastic, whereas acceleration of this function is an indication of arterial stiffness. As such, arterial elasticity is computed as the area under the pulse waveform during this interval, after subtracting a triangle corresponding to a linear relaxation. In order to ensure that this measure of elasticity is independent of peak amplitude (which is related to pulse pressure) and heart rate, the values are normalized (setting the peak systole = 1 and peak diastole = 0; and dividing them by the duration of the systole-diastole interval). PReFx was estimated for each voxel in which sensitivity (measured by the average Jacobian) was greater than 1/1000 (60 dB) of the maximum value. This allowed for the exclusion of voxels too deep to provide useful data (approximately > 3 cm from the scalp), as well as voxels beyond the coverage of the optical montages. In addition, only voxels within the cortex (as identified by FreeSurfer) were considered. *Regional* arterial elasticity was computed for 50 of the 70 FreeSurfer regions (25 per hemisphere), because at least 10% of their volume was sampled by the optical montage. The remaining 20 regions (10 per hemisphere) were either too deep or insufficiently covered by the optical montages, preventing reliable measurements. The 50 usable regions (25 per

hemisphere) were averaged to compute a measure of *global* PReFx for each subject. In addition, we computed 3 composite PReFx scores by averaging regions that were expected to be perfused by the anterior, middle, or posterior cerebral artery.

#### 2.4.1 Optodes co-registration with structural MRI

For each participant, optical sources and detectors' locations on the scalp were digitized using a Polhemus "3Space" FASTRAK 3D digitizer (Polhemus, Colchester, VT), with the nasion and preauricular points as references. The Polhemus digitization points were then co-registered with the MR based on procedures describe by Chiarelli et al. (2015; see also Whalen et al., 2008).

#### 2.5 MRI Acquisition and WMSA volume estimation

Structural images were collected for each participant using a 3T Siemens Trio full body scanner. No other MRI sequences (e.g. T2/FLAIR), which are commonly used for examining WMSAs, were collected. A high resolution, 3D MPRAGE protocol was used flip angle = 9°, TE = 2.32 ms, TR = 1900 ms, inversion time = 900 ms, 192 sagittal slices, voxel size 0.9 x 0.9 x 0.9 mm with a field of view of 172.8 x 230 x 230 mm. Cortical reconstruction and volumetric segmentation were performed with the FreeSurfer 6.0 image analysis suite (<http://surfer.nmr.mgh.harvard.edu/>; Fischl & Dale, 2000). Estimated total intracranial volume (eTIV) was also extracted from the structural images derived from the same automated procedures in FreeSurfer (Buckner et al., 2004). WMSAs (appearing as hypointense on T1-weighted images) were labeled automatically based on FreeSurfer 6.0's probabilistic procedure (Fischl et al., 2002) and the automatic segmentation was extensively examined for errors, which were corrected following <http://surfer.nmr.mgh.harvard.edu/fswiki/FsTutorial/TroubleshootingData>. Intraclass correlations between T1-weighted WMSAs and manually segmented white matter hyperintensities have also been reported to be as high as  $r=0.91$  (Smith et al., 2011). Here we focused on the overall WMSA volume as an index of white matter lesions, to avoid over-interpreting the T1-weighted hypointensities. WMSA volumes were log-transformed to reduce the skewing of the distribution (e.g., Atwood et al., 2004; Jefferson et al., 2007) and adjusted for sex and eTIV to obtain the final WMSA scores used in the current study.

## 2.7 Mediation Analysis

In two simple mediation analyses, we investigated whether PReFx mediated the relationship between age and WMSA, and whether WMSA mediated the relationships between PReFx and scores in the fluid and crystallized domains of cognition. Mediations were performed by employing PROCESS model 4 macro in SPSS (IBM, 2013) based on a series of ordinary least square (OLS) regression models (Hayes, 2013). This analysis was chosen for its directional path approach, which is the best option here, given the limitations of a cross-sectional study. We tested the indirect effect by using a bias-corrected bootstrapping procedure based on 5000 samples, chosen because it is considered the most trustworthy and provides more power given our relatively small sample size (Hayes & Scharkow, 2013).

## 3. Results

### 3.1 Relationships between global PReFx, WMSA, and age

Higher global PReFx was associated with lower WMSA burden ( $\beta = -0.16$ ,  $SE = 0.03$ ,  $p = 7.90 \times 10^{-7}$ , Figure 1). Post-hoc power analysis using R package “pwr” showed that the analysis is powered at 99.9%. The Kolmogorov-Smirnov test indicated that the model residuals were normally distributed ( $p > .05$ ). Testing for age-associated difference in this relationship, we found a significant age by PReFx interaction ( $\beta = -0.05$ ,  $SE = 0.02$ ,  $p = 0.04$ ), with older adults showing a significant association ( $\beta = -0.11$ ,  $SE = 0.04$ ,  $p = 0.01$ ), but not younger adults ( $\beta = -0.01$ ,  $SE = 0.04$ ,  $p > .05$ , Figure 2).

**INSERT FIGURES 1-2 ABOUT HERE**

### 3.2 Relationships between regional PReFx, WMSAs, and age.

Higher WMSA burden was associated with lower regional PReFx in 24 out of the 25 brain regions (average of left and right) identified on the basis of the FreeSurfer atlas, corrected for multiple comparisons at a FDR < .05. The only region that did not reach statistical significance was the frontal pole. The largest associations were found in the banks of the superior temporal sulcus, surpramarginal gyrus, and post and precentral gyrus (Figure 3, Supplemental Table S1), all areas that are likely perfused by the middle cerebral artery. None of the interactions between age and regional PReFx on WMSA burden reached statistical significance at FDR < .05.

**INSERT FIGURE 3 ABOUT HERE****3.3 Mediating effects of PReFx and WMSA**

We found that global PReFx did not mediate the relationship between age and WMSA (95% CI: -.006 - .196). However, the composite PReFx from brain regions supplied by the middle cerebral artery partially mediated the relationship between age and WMSA (95% CI: .01 - .22, Figure 4A), meaning that the direct effect (path c'), although significantly reduced in value, remains significant after accounting for the effects of PReFx. Post-hoc power analysis using R package “WebPower” showed that this analysis is powered at 91.5%. PReFx computed from brain regions supplied by the anterior and posterior cerebral artery, however, did not significantly mediate the age-WMSA relationship. For the cognitive function mediation, we found that WMSA partially mediated the relationship between global PReFx and fluid abilities (95% CI: .045 - .258, Figure 4B) such that the direct effect (path c') is reduced but remains significant after accounting for the effects of WMSA. Post-hoc power analysis showed that this analysis is powered at 90.0%. WMSA did not significantly mediate the relationship between PReFx and verbal scores (95% CI: -.2213 - .017).

**INSERT FIGURE 4 ABOUT HERE****4. Discussion**

The results presented here support the presence of an association between reduced cerebral arterial elasticity measured with PReFx pulse-DOT in the cortex and the extent of white matter abnormalities (WMSAs) in normally aging adults. This relationship was particularly evident for older participants, and was not significant in younger adults. Although correlations between systemic indices of arterial status (such as pulse pressure) and WMSA volume have been reported in the literature (Kennedy & Raz, 2009; Leritz et al., 2014; Salat et al., 2012), our results indicates that WMSA volume is also associated with a more direct assessment of the status of the *cerebral* vasculature. The regional associations between PReFx and WMSA also suggest that PReFx in regions in the frontal, temporal and parietal cortices, which are perfused by the middle cerebral artery, are most indicative of the degree of overall WMSA burden.

That said, it should be noted that the PReFx measures were obtained in cortical regions, rather than in deep periventricular white matter, a limitation due to the poor penetration of diffuse optical measures. Although this initial correlational finding would need to be replicated on a much larger sample, to determine the actual sensitivity and specificity of the relationship, the results suggest a strong association between arterial elasticity in the cortex and WMSAs (a possible indication of cerebral small vessel disease, CSVD; Sachdev, Wen, Chen & Brodaty, 2007; Shi & Wardlaw; 2016). This also suggests that there may be potential value in using pulse-DOT measurements as a screening tool for CSVD. The validity and effectiveness of such screening procedures should be the subject of further larger scale investigations.

The data reported here also suggest that arterial elasticity in regions perfused by the middle cerebral artery likely plays a mediator role in age-related increases in WMSA burden in normal aging. These findings are consistent with the general consensus that the emergence of WMSA lesions stems from cerebral vascular pathology, which is exacerbated in aging (Salat, 2014), and are also consistent with our regional PreFx results showing that the largest correlations are in temporal and parietal cortices.

We additionally found that WMSA burden partially mediated the relationship between global PReFx and fluid abilities but not crystallized abilities scores. While verbal ability, which largely contributes to crystallized intelligence is relatively preserved in aging, our results suggest that reductions in PReFx may lead to declines in fluid abilities through increases in WMSA (Baltes & Lindenberger, 1997). Taken together, the mediation analyses suggest that PReFx may reflect a mechanistic chain of events linking arterial dysfunction with white matter damage and cognitive decline. Future research will be needed to further investigate the independent and interacting influences of neural activity and vascular properties on age-related decline in central processing.

The partial mediations found also suggest a role for unexamined variables such as cardiorespiratory fitness, which may influence the effects of both arterial elasticity and WMSA volume (see Cotman, Berchtold & Christie, 2007; Barnes, 2015). These variables could be investigated in larger studies in the future. Future studies may also investigate whether regional PReFx and WMSA could act as mediators of

age-related decline in cognitive function and determine the regional specificity of the mediation results found in the current study.

#### 4.1 Limitations

The findings presented here must be interpreted within the inherently limited depth of penetration of diffuse optical imaging. A consequence of this is that the PreFx measures are not taken directly from the periventricular areas where the majority of white matter abnormalities are found. That said, it should be noted that PreFx reflects the speed of propagation of the recoil wave from the periphery (muscular arteries and arterioles providing peripheral resistance) back to the point of measurement (reflected-wave, see Izzo and Shykoff, 2001). In other words, PReFx is a measure of arterial function *downstream* relative to the measurement point, functionally extending the field of view of pulse-DOT. Further, the measurements used in the current study do use recent advancements in 3D reconstruction (Chiarelli et al., 2016), which have increased the effective optical depth penetration to 30-35 mm, and therefore provide the best estimation of cerebral arterial elasticity possible given the current state of the technology.

The regional measures have limited spatial resolution, which makes it difficult to distinguish the specific arterial collaterals that feed into any specific WMSA cluster. Magnetic resonance angiography (MRA) may be used to aid the spatial mapping of regional PReFx with the cerebrovasculature. Although newer dynamic ASL methods for quantifying vascular elasticity based on changes in cerebral blood volume and arterial pressure can be extracted from deep structures (Yan et al., 2016), they also suffer from lack of adequate spatial resolution. Further, diffuse optical measures for quantifying arterial elasticity and their association with a host of factors such as age, fitness, brain volume have been replicated and found to be highly robust (Fabiani et al., 2014b, Tan et al; 2017). Whether dynamic ASL methods can serve as a mediator of age-related declines in cognition remains to be tested. In the future, it would be very fruitful to combine ASL, TCD, and/or MRA with pulse-DOT methods to address some of these spatial specificity issues. There has also not been any systematic study comparing T1-weighted WMSAs with T2/FLAIR WMHs, demonstrating that they share the same pathophysiology. Future research using pulse-DOT with both types of measures should explore this issue.

## 4.2 Summary and conclusions

Despite these limitations, the current study opens up the novel possibility of providing an extensive assessment of global and regional cerebral arterial elasticity, and of using arterial elasticity as a target for intervention. Although vascular risk factors accumulate over the life span, they are amenable to changes by increasing physical activity (e.g., Hawkins et al., 2014) and/or by maintaining healthy dietary practices (e.g., Santos-Parker, LaRocca & Seals, 2014). A recent review of studies investigating the association between white matter lesions and physical activity concluded that greater physical activity was associated with a lower amount of white matter lesions (Torres et al., 2015; see also Burzynska et al., 2014; Fletcher et al., 2016; Gordon et al., 2008). This suggests that early interventions may help slow or prevent progressive CSVD from developing and manifesting as WMSAs. This preventive approach is especially important given the general consensus that lesions in the white matter are non-reversible, although revascularization surgery may help (Komatsu et al., 2016). Therefore, in addition to using WMSAs as an indirect marker of CSVD severity (Sachdev, Wen, Chen & Brodaty, 2007), measures of arterial elasticity may also be useful as an early risk-screening tool in cognitively normal individuals, for selective intervention before the onset of widespread WMSAs.

### **Acknowledgements**

This work was supported by NIA grant R01AG059878 to M. Fabiani and G. Gratton; NIH grant 1R56MH097973-01 to G. Gratton and M. Fabiani and by NCRR grant S10-RR029294 to G. Gratton. This work was conducted in part at the Biomedical Imaging Center of the Beckman Institute for Advanced Science and Technology at the University of Illinois at Urbana-Champaign (UIUC-BI-BIC).

### **Disclosure Statement**

The authors declare no actual or potential conflicts of interest. The views expressed in this paper are solely those of the authors, and all authors have approved this manuscript.

## References

- Adams, R. D., Victor, M., & Ropper, A. H. (1997) Cerebrovascular Disease. In Adams, R. D., Victor, M., & Ropper, A. H. (Eds.) *Principles of Neurology*, 777-873. New York: McGraw-Hill.
- Alosco, M. L., Gunstad, J., Jerskey, B. A., Xu, X., Clark, U. S., Hassenstab, J., ... Sweet, L. H. (2013). The adverse effects of reduced cerebral perfusion on cognition and brain structure in older adults with cardiovascular disease. *Brain and Behavior*, 3(6), 626–636. <http://doi.org/10.1002/brb3.171>
- [Altamura, M., Elvevåg, B., Blasi, G., Bertolino, A., Callicott, J. H., Weinberger, D. R., ... Goldberg, T. E. \(2007\). Dissociating the effects of Sternberg working memory demands in prefrontal cortex. \*Psychiatry Research\*, 154\(2\), 103–114. <http://doi.org/10.1016/j.psychres.2006.08.002>](#)
- Ances, B. M., Liang, C. L., Leontiev, O., Perthen, J. E., Fleisher, A. S., Lansing, A. E. & Buxton, R. B. (2009). Effects of aging on cerebral blood flow, oxygen metabolism, and blood oxygenation level dependent responses to visual stimulation. *Human Brain Mapping*, 30:1120–1132
- Atwood, L. D., Wolf, P. A., Heard-Costa, N. L., Massaro, J. M., Beiser, A., D'Agostino, R. B., & DeCarli, C. (2004). Genetic variation in white matter hyperintensity volume in the Framingham Study. *Stroke; a Journal of Cerebral Circulation*, 35(7), 1609–1613. <http://doi.org/10.1161/01.STR.0000129643.77045.10>
- Baltes, P. B., & Lindenberger, U. (1997). Emergence of a powerful connection between sensory and cognitive functions across the adult life span: a new window to the study of cognitive aging? *Psychology and Aging*, 12, 12–21.
- Beason-Held, L. L., Moghekar, A., Zondeman, A. B., Kraut, M. A., & Resnik, S. M. (2007) Longitudinal changes in cerebral blood flow in the older hypertensive brain. *Stroke*, 38(6), 1766-1773.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory* (2nd ed.). San Antonio, TX: The Psychological Corporation.
- Behrouz, R., Malek, A. R., & Torbey, M. T. (2012). Small vessel cerebrovascular disease: The past, present, and future. *Stroke Research and Treatment*, 839151. doi: 10.1155/2012/839151
- Bennett, I. J., & Madden, D. J. (2014). Disconnected aging: cerebral white matter integrity and age-related differences in cognition. *Neuroscience*, 276, 187–205. <http://doi.org/10.1016/j.neuroscience.2013.11.026>
- Benton, A. L., & Hamsher, K. (1989). *Multilingual aphasia examination manual*. Iowa City, IA: University of Iowa.
- Brickman, A. M., Siedlecki, K. L., Muraskin, J., Manly, J. J., Luchsinger, J. A., Yeung, L.-K., ... Stern, Y. (2011). White matter hyperintensities and cognition: testing the reserve hypothesis. *Neurobiology of Aging*, 32(9), 1588–1598. <http://doi.org/10.1016/j.neurobiolaging.2009.10.013>

- Brickman, A. M., Zahra, A., Muraskin, J., Steffener, J., Holland, C. M., Habeck, C., ... Stern, Y. (2009). Reduction in cerebral blood flow in areas appearing as white matter hyperintensities on magnetic resonance imaging. *Psychiatry Research*, *172*(2), 117–120. <http://doi.org/10.1016/j.psychresns.2008.11.006>
- Buckner, R. L., Head, D., Parker, J., Fortenos, A. F., Marcus, D., Morris, J. C., & Snyder A. Z. (2004) A unified approach for morphometric and functional data analysis in young, old, and demented adults using automated atlas-based head size normalization: reliability and validation against manual measurement of total intracranial volume. *Neuroimage*, *23*(2), 723 - 738
- Burzynska, A.Z., Chaddock-Heyman, L., Voss, M. W., Wong, C. N., Gothe, N. P., ... Kramer, A. F. (2014) Physical activity and cardiorespiratory fitness are beneficial for white matter in low-fit older adults *PLoS One*, *9*, Article e107413
- Chan, M. Y., Park, D. C., Savalia, N. K., Petersen, S. E., & Wig, G. S. (2014). Decreased segregation of brain systems across the healthy adult lifespan. *Proceedings of the National Academy of Sciences*, *111*(46), E4997-E5006.
- Charlton, R. A., Barrick, T. R., Markus, H. S. & Morris, R. G. (2009). Theory of mind associations with other cognitive functions and brain imaging in normal aging. *Psychol Aging*, *24*(2):338-48. doi: 10.1037/a0015225.
- Chiarelli, A. M., Maclin, E. L., Low, K. A., Fabiani, M., & Gratton, G. (2015). Comparison of procedures for co-registering scalp-recording locations to anatomical magnetic resonance images. *Journal of Biomedical Optics*, *20*(1), 16009. <http://doi.org/10.1117/1.JBO.20.1.016009>
- Chiarelli, A. M., Maclin, E. L., Low, K. A., Mathewson, K. E., Fabiani, M., & Gratton, G. (2016). Combining energy and Laplacian regularization to accurately retrieve the depth of brain activity of diffuse optical tomographic data. *Journal of Biomedical Optics*, *21*(3), 36008. <http://doi.org/10.1117/1.JBO.21.3.036008>
- Chiarelli, A.M., Fletcher, M.A., Tan, C.H., Low, K.A., Maclin, E.L., Zimmerman, B., Kong T., Gorsuch, A., Gratton, G. & Fabiani, M. (2017). Individual Differences in Regional Cortical Volumes are Associated with Regional Optical Measures of Arterial Elasticity in Healthy Older Adults. *Neuroimage*, *162*, PMID: 28866349.
- Chen. S. T., Siddarth, P., Ercoli, L. M., Merrill, D. A., Torres-Gil, F. & Small, G. W. (2014). Modifiable risk factors for Alzheimer disease and subjective memory impairment across age groups. *PLoS One*, *9*:e98630.
- Corrigan, J. D., & Hinkeldey, N. S. (1987). Relationships between parts A and B of the Trail Making Test. *Journal of Clinical Psychology*, *43*(4), 402–409.

- Coxon, J. P., Van Impe, A., Wenderoth, N. & Swinnen, S. P. (2012). Aging and inhibitory control of action: cortico-subthalamic connection strength predicts stopping performance. *J Neurosci* 32: 8401–8412.
- DeBette, S., & Markus, H. S. (2010). The clinical importance of white matter hyperintensities on brain magnetic resonance imaging: systematic review and meta-analysis. *BMJ (Clinical Research Ed.)*, 341, c3666.
- DeCarli, C., Murphy, D. G., Tranh, M., Grady, C. L., Haxby, J. V., Gillette, J. A., ... Rapoport, S. I. (1995). The effect of white matter hyperintensity volume on brain structure, cognitive performance, and cerebral metabolism of glucose in 51 healthy adults. *Neurology*, 45(11), 2077–2084.
- Dehghani, H., Eames, M. E., Yalavarthy, P. K., Davis, S. C., Srinivasan, S., Carpenter, C. M., Pogue, B. W., Paulsen, K. D. (2009) Near infrared optical tomography using NIRFAST: Algorithm for numerical model and image reconstruction. *Communications in Numerical Methods in Engineering*, 25:711-732.
- Duncombe, J., Lennen, R. J., Jansen, M. A., Marshall, I., Wardlaw, J. M. & Horsburgh, K. (2017). Ageing causes prominent neurovascular dysfunction associated with loss of astrocytic contacts and gliosis. *Neuropathol Appl Neurobiol*. 43(6):477-491. doi: 10.1111/nan.12375
- Eggebrecht, A. T., Ferradal, S. L., Robichaux-Viehoever, A., Hassanpour, M. S., Dehghani, H., Snyder, A. Z., Hershey, T., Culver, J. P. (2014) Mapping distributed brain function and networks with diffuse optical tomography. *Nature photonics*, 8:448-454.
- Elias, P. K., Elias, M. F., Robbins, M. A., & Budge, M. M. (2004). Blood pressure-related cognitive decline: does age make a difference? *Hypertension*, 44(5), 631-636.
- Fabiani, M. (2012). SPR 2008 Presidential Address. It was the best of times, it was the worst of times: A psychophysiological's view of cognitive aging. *Psychophysiology*, 49(3), 283-304. <http://dx.doi.org/10.1111/j.1469-8986.2011.01331.x>
- Fabiani, M., Gordon, B. A., Maclin, E. L., Pearson, M. A., Brumback-Peltz, C. R., Low, K. A., ... Gratton G (2014a). Neurovascular coupling in normal aging: a combined optical, ERP and fMRI study. *Neuroimage*, 85, 592-607, doi: 10.1016/j.neuroimage.2013.04.113
- Fabiani, M., Low, K. A., Tan, C.-H., Zimmerman, B., Fletcher, M. A., Schneider-Garces, N., ... Gratton, G. (2014b). Taking the pulse of aging: mapping pulse pressure and elasticity in cerebral arteries with optical methods. *Psychophysiology*, 51(11), 1072–1088. <http://doi.org/10.1111/psyp.12288>
- Fang, Q. & Boas, D. A. (2009) Tetrahedral mesh generation from volumetric binary and grayscale images. *Proceedings of IEEE International Symposium on Biomedical Imaging*. 1142-1145.
- Fischl, B., Dale, A.M. (2000) Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A*, 97:11050-5.

- Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., ... Dale, A. M. (2002). Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*, 33(3), 341–355.
- Fletcher, M., Low, K. A., Boyd, R., Zimmerman, B., Gordon, B., Tan, C. H., Schneider-Garces, N., ... Fabiani, M. (2016). Comparing aging and fitness effects on brain anatomy. *Frontiers in Human Neuroscience*, 10(286) doi:10.3389/fnhum.2016.00286
- Friston, K. J., Holmes, A. P., Worsley, K. J., Poline, J. P., Frith, C.D., Frackowiak, R. S. J. (1994) Statistical parametric maps in functional imaging: A general linear approach. *Human Brain Mapping*, 2:189-210.
- Gordon, B. A., Rykhlevskaia, E. I., Brumback, C. R., Lee, Y., Elavsky, S., Konopack, J. F., McAuley, E., Kramer, A. F., Colcombe, S., Gratton, G., Fabiani, M. (2008) Neuroanatomical correlates of aging, cardiopulmonary fitness level, and education. *Psychophysiology*, 45:825-38.
- Hatazawa, J., Shimosegawa, E., Satoh, T., Toyoshima, H., Okudera, T. (1997). Subcortical hypoperfusion associated with asymptomatic white matter lesions on magnetic resonance imaging. *Stroke; a Journal of Cerebral Circulation*, 28(10), 1944–1947.
- Hawkins, M., Gabriel, K. P., Cooper, J., Storti, K. L., Sutton-Tyrrell, K., & Kriska, A. (2014). The impact of change in physical activity on change in arterial stiffness in overweight or obese sedentary young adults. *Vascular Medicine*, 19(4), 257–263. <http://doi.org/10.1177/1358863X14536630>
- Hayes, A. F. (2013). *An introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. New York, NY: Guilford Press.
- Hayes, A. F., & Scharkow, M. (2013). The relative trustworthiness of inferential tests of the indirect effect in statistical mediation analysis: does method really matter? *Psychological Science*, 24(10), 1918–1927. <http://doi.org/10.1177/0956797613480187>
- Hutchison, E. R., Kawamoto, E. M., Taub, D. D., Lal, A., Abdelmohsen, K., Zhang, Y., ... Mattson MP. (2013). Evidence for miR-181 involvement in neuroinflammatory responses of astrocytes. *Glia*, 61(7):1018-28. doi: 10.1002/glia.22483
- Ishimaru, A. (1989) Diffusion of light in turbid material. *Applied Optics*, 28(12), 2210-2215.
- Jefferson, A. L., Tate, D. F., Poppas, A., Brickman, A. M., Paul, R. H., Gunstad, J., & Cohen, R. A. (2007). Lower cardiac output is associated with greater white matter hyperintensities in older adults with cardiovascular disease. *Journal of the American Geriatrics Society*, 55(7), 1044–1048. <http://doi.org/10.1111/j.1532-5415.2007.01226.x>
- Jubran, A. (1999) Pulse oximetry. *Critical Care*, 3(2), 11-17. Doi: 10.1186/cc341.
- Keage, H. A. D., Kurylowicz, L., Lavrencuc, L. M., Churches, O. F., Flitton, A., Hofmann, J., ... Badcock, N. A. (2015). Cerebrovascular function associated with fluid, not crystallized, abilities in

- older adults: A transcranial Doppler study. *Psychology and Aging*, 30(3), 613-23. doi: 10.1037/pag0000026
- Keage, H. A. D., Churches, O. F., Kohler, M., Pomeroy, D., Luppino, R., Bartolo, M. L. & Elliott, S. (2012). Cerebrovascular function in aging and dementia: A systematic review of transcranial Doppler studies. *Dementia and Geriatric Cognitive Disorders*, 2, 258 – 270. doi: 10.1159/000339234
- Kennedy, K. M., & Raz, N. (2009). Aging White Matter and Cognition: Differential Effects of Regional Variations in Diffusion Properties on Memory, Executive Functions, and Speed. *Neuropsychologia*, 47(3), 916–927. <http://doi.org/10.1016/j.neuropsychologia.2009.01.001>
- Komatsu, K., Mikami, T., Noshiro, S., Miyata, K., Wanibuchi, M., & Mikuni, N. (2016). Reversibility of White Matter Hyperintensity by Revascularization Surgery in Moyamoya Disease. *Journal of Stroke and Cerebrovascular Diseases*, 25(6), 1495–1502. <http://doi.org/10.1016/j.jstrokecerebrovasdis.2016.02.035>
- Leaper, S. A., Murray, A. D., Lemmon, H. A., Staff, R. T., Deary, I. J., Crawford, J. R., & Whalley, L. J. (2001). Neuropsychologic correlates of brain white matter lesions depicted on MR images: 1921 Aberdeen Birth Cohort. *Radiology*, 221(1), 51–55. <https://doi.org/10.1148/radiol.2211010086>
- Leritz, E. C., Shepel, J., Williams, V. J., Lipsitz, L. A., McGlinchey, R. E., Milberg, W. P., & Salat, D. H. (2014). Associations between T1 white matter lesion volume and regional white matter microstructure in aging. *Human Brain Mapping*, 35(3), 1085–1100. <http://doi.org/10.1002/hbm.22236>
- Madden, D. J., Bennett, I. J., Burzynska, A., Potter, G. G., Chen, N. K. & Song, A. W. (2012). Diffusion tensor imaging of cerebral white matter integrity in cognitive aging. *Biochim Biophys Acta*, 1822, 386–400.
- Mayeux, R., Stern, Y., Rosen, J., & Leventhal, J. (1981). Depression, intellectual impairment, and Parkinson disease. *Neurology*, 31(6), 645–650
- Marstrand, J. R., Garde, E., Rostrup, E., Ring, P., Rosenbaum, S., Mortensen, E. L., & Larsson, H. B. W. (2002). Cerebral perfusion and cerebrovascular reactivity are reduced in white matter hyperintensities. *Stroke; a Journal of Cerebral Circulation*, 33(4), 972–976.
- Mohtasib, R. S., Lumley, G., Goodwin, J. A., Emsley, H, C, A., Sluming, V. & Parkes, L. M. Calibrated fMRI during a cognitive Stroop task reveals reduced metabolic response with increasing age. *Neuroimage*, 59, 1143–1151.
- Muller, M., van der Graaf, Y., Visseren, F. L., Mai, W. P., Geerlings, M. I., & SMART Study Group. (2012) *Annals of Neurology*, 71(6), 825-833. doi: 10.1002/ana.23554

- Mutsaerts, H. J. M. M., van Dalen, J. W., Heijtel, D. F. R., Groot, P. F. C., Majoie, C. B. L. M., Petersen, E. T., ... Nederveen, A. J. (2015). Cerebral Perfusion Measurements in Elderly with Hypertension Using Arterial Spin Labeling. *PloS One*, *10*(8), e0133717.
- Panigrahy, A., Barnes, P. D., Robertson, R. L., Back, S. A., Sleeper, L. A., Sayre, J. W., ... Volpe, J. J. (2001). Volumetric brain differences in children with periventricular T2-signal hyperintensities: a grouping by gestational age at birth. *AJR. American Journal of Roentgenology*, *177*(3), 695–702. <http://doi.org/10.2214/ajr.177.3.1770695>
- Pantoni, L., & Simoni, M. (2003). Pathophysiology of cerebral small vessels in vascular cognitive impairment. *International Psychogeriatrics / IPA*, *15* Suppl 1, 59–65. <http://doi.org/10.1017/S1041610203008974>
- Park, D. C., & Reuter-Lorenz, P. (2009). The adaptive brain: aging and neurocognitive scaffolding. *Annual Review of Psychology*, *60*, 173–196. <http://doi.org/10.1146/annurev.psych.59.103006.093656>
- Paulsen, K.D., & Jiang, H. (1995) Spatially varying optical property reconstruction using a finite element diffusion equation approximation. *Medical Physics*, *22*, 691-701.
- Peltz, C. B., Gratton, G., & Fabiani, M. (2011). Age-related changes in electrophysiological and neuropsychological indices of working memory, attention control, and cognitive flexibility. *Frontiers in Psychology*, *18*;2:190. <http://dx.doi.org/10.3389/fpsyg.2011.00190>
- Penny, W. D., Friston, K. J., Ashburner, J. T., Kiebel, S. J. & Nichols, T. E. (2011) Statistical parametric mapping: the analysis of functional brain images: the analysis of functional brain images. Academic press.
- Rabbitt, P., Scott, M., Lunn, M., Thacker, N., Lowe, C., Pendleton, N., ... Jackson, A. (2007). White matter lesions account for all age-related declines in speed but not in intelligence. *Neuropsychology*, *21*(3), 363–370. <https://doi.org/10.1037/0894-4105.21.3.363>
- Raven, J., Raven, J.C., & Court, J.H. (2003) *Manual for Raven's Progressive Matrices and Vocabulary Scales*. San Antonio, TX: Harcourt Assessment.
- Raz, N., Lindenberger, U., Ghisletta, P., Rodrigue, K. M., Kennedy, K. M., & Acker, J. D. (2008). Neuroanatomical correlates of fluid intelligence in healthy adults and persons with vascular risk factors. *Cerebral Cortex*, *18*, 718–726.
- Raz, N., & Rodrigue, K. M. (2006) Differential aging of the brain: patterns, cognitive correlates and modifiers. *Neuroscience and biobehavioral reviews*, *30*(6), 730-48. <https://doi.org/10.1016/j.neubiorev.2006.07.001>
- Raz, N., Rodrigue, K. M., Kennedy, K. M., & Acker, J. D. (2007). Vascular health and longitudinal changes in brain and cognition in middle-aged and older adults. *Neuropsychology*, *21*(2), 149–157. <http://doi.org/10.1037/0894-4105.21.2.149>

- Restom, K., Bangen, K. J., Bondi, M. W., Perthen, J. E., & Liu, T. T. (2007). Cerebral blood flow and BOLD responses to a memory encoding task: a comparison between healthy young and elderly adults. *Neuroimage* 37, 430–439. doi: 10.1016/j.neuroimage.2007.05.024
- Resnick, S. M., Pham, D. L., Kraut, M. A., Zonderman, A. B., & Davatzikos, C. (2003). Longitudinal magnetic resonance imaging studies of older adults: a shrinking brain. *The Journal of Neuroscience*, 23(8), 3295–3301.
- Sachdev, P., Wen, W., Chen, X., & Brodaty, H. (2007). Progression of white matter hyperintensities in elderly individuals over 3 years. *Neurology*, 68(3), 214–222. <http://doi.org/10.1212/01.wnl.0000251302.55202.73>
- Salami, A., Eriksson, J., Nilsson, L. G. & Nyberg, L. Age-related white matter microstructural differences partly mediate age-related decline in processing speed but not cognition. *Biochim. Biophys. Acta*, 1822, 408–415 .
- Salat, D. H. (2014). Imaging small vessel-associated white matter changes in aging. *Neuroscience*, 276, 174–186. <http://doi.org/10.1016/j.neuroscience.2013.11.041>
- Salat, D. H., Tuch, D. S., van der Kouwe, A. J. W., Greve, D. N., Pappu, V., Lee, S. Y., ... Rosas, H. D. (2010). White matter pathology isolates the hippocampal formation in Alzheimer's disease. *Neurobiology of Aging*, 31(2), 244–256. <http://doi.org/10.1016/j.neurobiolaging.2008.03.013>
- Salat, D. H., Williams, V. J., Leritz, E. C., Schnyer, D. M., Rudolph, J. L., Lipsitz, L. A., ... Milberg, W. P. (2012). Inter-individual variation in blood pressure is associated with regional white matter integrity in generally healthy older adults. *NeuroImage*, 59(1), 181–192. <http://doi.org/10.1016/j.neuroimage.2011.07.033>
- Salthouse, T. A. (2010). Selective review of cognitive aging. *Journal of the International Neuropsychological Society : JINS*, 16(5), 754–760. <http://doi.org/10.1017/S1355617710000706>
- Salthouse, T. A., & Pink, J. E. (2008). Why is working memory related to fluid intelligence? *Psychonomic Bulletin & Review*, 15, 364–371. doi:10.3758/PBR.15.2.364
- Sanchez-Cubillo, I., Perianez, J. A., Adrover-Roig, D., Rodriguez-Sanchez, J. M., Rios-Lago, M., Tirapu, J., & Barcelo, F. (2009) Construct validity of the Trail Making Test: Role of task-switching, working memory, inhibition/interference control, and visuomotor abilities. *Journal of the International Neuropsychological Society*, 15:438–450.
- Santos-Parker, J. R., LaRocca, T. J., & Seals, D. R. (2014). Aerobic exercise and other healthy lifestyle factors that influence vascular aging. *Advances in Physiology Education*, 38(4), 296–307. <https://doi.org/10.1152/advan.00088.2014>
- Schiffmann, R., & van der Knaap, M. S. (2009). An MRI-based approach to the diagnosis of white matter disorders. *Neurology*, 72(8), 750–759. <http://doi.org/10.1212/01.wnl.0000343049.00540.c8>

- [Shi, Y., & Wardlaw, J. M. \(2016\). Update on cerebral small vessel disease: a dynamic whole-brain disease. \*BMJ\*, 1\(3\), 83–92. <https://doi.org/10.1136/svn-2016-000035>](#)
- Shipley, W. C. (1940). A self-administering scale for measuring intellectual impairment and deterioration. *Journal of Psychology*, 9, 371–377.
- Smith, E. E., Salat, D. H., Jeng, J., McCreary, C. R., Fischl, B., Schmahmann, J. D., ... Greenberg, S. M. (2011). Correlations between MRI white matter lesion location and executive function and episodic memory. *Neurology*, 76(17), 1492–1499. <http://doi.org/10.1212/WNL.0b013e318217e7c8>
- Tan, C. H., Low, K. A., Kong, T. S., Fletcher, M., Zimmerman, B., Maclin, E.L., Gratton, G. & Fabiani, M. (2017) Mapping cerebral pulse pressure and arterial compliance over the adult lifespan with optical imaging. *PloSONE* 12(2): e0171305. doi:10.1371/journal.pone.0171305
- Tan, C. H., Low, K. A., Schneider-Garces, N., Fletcher, M., Zimmerman, B., Maclin, E.L., Chiarelli, A. M., Gratton, G. & Fabiani, M. (2016). Optical measures of changes in cerebral vascular tone during voluntary breath holding and a Sternberg memory task. *Biological Psychology*, 118. doi:10.1016/j.biopsycho.2016.05.008
- Tian, F., & Liu, H. (2014) Depth-compensated diffuse optical tomography enhanced by general linear model analysis and an anatomical atlas of human head. *Neuroimage*, 85, Part 1:166-180.
- Torres, E. R., Strack, E. F., Fernandez, C. E., Tumey, T. A., Hitchcock, M. E. (2015) Physical activity and white matter hyperintensities: A systematic review of quantitative studies. *Prev Med Reports*, 2: 319-325.
- Unsworth, N., Fukuda, K., Awh, E., and Vogel, E. K. (2014). Working memory and fluid intelligence: capacity, attention control, and secondary memory retrieval. *Cogn. Psychol.* 71, 1–26. doi: 10.1016/j.cogpsych.2014.01.003
- Unsworth, N., Heitz, R. P., Schrock, J. C., & Engle, R. W. (2005). An automated version of the operation span task. *Behavior Research Methods*, 37(3), 498–505.
- Van Dijk, E. J., Prins, N. D., Vrooman, H. A., Hofman, A., Koudstaal, P. J. & Breteler, M. M. B (2008). Progression of cerebral small vessel disease in relation to risk factors and cognitive consequences: Rotterdam scan study. *Stroke*, 39, 2712-2719. doi: 10.1161/STROKEAHA.107.513176
- Waldstein, S. R., Lefkowitz, D. M., Siegal, E. L., Rosenberger, W. F., Spencer, R. J., Tankard, C. F., ... Katzel, L. I. (2010) *Journal of Hypertension*, 28(5), 993-998
- Wardlaw, J. M., Valdés Hernández, M. C., & Muñoz-Maniega, S. (2015). What are White Matter Hyperintensities Made of? *Journal of the American Heart Association: Cardiovascular and Cerebrovascular Disease*, 4(6). <http://doi.org/10.1161/JAHA.114.001140>

- Whalen, C., Maclin, E. L., Fabiani, M., & Gratton, G. (2008). Validation of a method for coregistering scalp recording locations with 3D structural MR images. *Human Brain Mapping, 29*(11), 1288–1301. <http://doi.org/10.1002/hbm.20465>
- Yan, L., Liu, C. Y., Smith, R. X., Jog, M., Langham, M., Krasileva, K., ... Wang, D. J. J. (2016). Assessing intracranial vascular compliance using dynamic arterial spin labeling. *NeuroImage, 124*(Pt A), 433–441. <http://doi.org/10.1016/j.neuroimage.2015.09.008>
- Zimmerman, B., Sutton, B.P., Low, K.A., Fletcher, M.A., Tan, C.H., Schneider-Garces, N., Li, Y., Ouyang, C., Maclin, E.L., Gratton, G., & Fabiani, M. (2014). Cardiorespiratory fitness mediates the effects of aging on cerebral blood flow. *Frontiers in Aging Neuroscience, 6*:59. <http://dx.doi.org/10.3389/fnagi.2014.00059>

**Table 1. Demographic characteristics of the participants**

Variable	N = 93 Mean (SD)
Age (years)	58.8 (17.9)
Education (years)	16.8 (2.5)
mMMSE	55.5 (1.3)
Beck's Depression Index	2.8 (3.4)
PReFx	0.08 (0.02)
WMSA (log)	3.1 (0.3)
Systolic blood pressure	133.1 (13.2)
Diastolic blood pressure	78.6 (7.9)

### Figure Captions

Figure 1. Scatter plot showing the relationship between arterial elasticity (PReFx, averaged across all cortical regions) and WMSA burden. Color scale reflects age of the participants.

Figure 2. Interaction between global PReFx and age on WMSA. Older individuals with low PReFx had the highest amount of WMSA burden.

Figure 3. Associations of regional PReFx with WMSA burden. All associations except frontal pole were statistically significant after correction for multiple comparisons at  $FDR < .05$ .

Figure 4. A) PReFx from regions perfused by the middle cerebral artery partially mediated the relationship between age and WMSA. B) WMSA partially mediated the relationship between global PReFx and fluid abilities scores.

This manuscript has been approved by all authors.

None of the authors have financial or other conflicts of interest regarding the contents of this paper.

This work was supported by NIA grant R01AG059878 to M. Fabiani and G. Gratton; NIH grant 1R56MH097973-01 to G. Gratton and M. Fabiani and by NCCR grant S10-RR029294 to G. Gratton.

All research reported in this paper was approved by the IRB of the University of Illinois.

Also please note that this paper is not under consideration at any other journal.

Figure 1




Figure 2




Figure 3




Figure 4

